

Brucella in Himalayan Marmots (*Marmota himalayana*)

Authors: Yan, Bin, Zhu, Qingfeng, Xu, Jun, Zhao, Shanshan, Piao, Dongri, et al.

Source: Journal of Wildlife Diseases, 56(3) : 730-732

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/2019-09-237>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Brucella* in Himalayan Marmots (*Marmota himalayana*)**

Bin Yan,^{1,2,10} Qingfeng Zhu,^{3,4,10} Jun Xu,^{5,10} Shanshan Zhao,¹ Dongri Piao,⁶ Chuangfu Chen,⁷ Quan Liu,⁸ Sándor Hornok,⁹ Yuanzhi Wang,^{1,11} Baoju Wang,^{3,10} and Hai Jiang^{6,10} ¹School of Medicine, Shihezi University, Shihezi, Xinjiang Uygur Autonomous Region 832002, People's Republic of China; ²Clinical Laboratory Center, Zaozhuang Municipal Hospital, Zaozhuang, Shandong 277100, People's Republic of China; ³Department of Infectious Diseases, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430022, People's Republic of China; ⁴Department of Infectious Diseases, First Affiliated Hospital of the School of Medicine, Shihezi University, Shihezi, Xinjiang Uygur Autonomous Region 832002, People's Republic of China; ⁵Urumqi Customs, Urumqi, Xinjiang Uygur Autonomous Region 833400, People's Republic of China; ⁶State Key Laboratory for Infectious Disease Prevention and Control, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102200, People's Republic of China; ⁷College of Animal Science & Technology, Shihezi University, Shihezi, Xinjiang Uygur Autonomous Region 832000, People's Republic of China; ⁸School of Life Sciences and Engineering, Foshan University, Foshan, Guangdong 528225, People's Republic of China; ⁹Department of Parasitology and Zoology, University of Veterinary Medicine, Budapest 1078, Istvan u. 2, Hungary; ¹⁰These authors contributed equally to this work; ¹¹Corresponding author (email: wangyuanzhi621@126.com)

ABSTRACT: *Brucella abortus* biovar 1 and atypical rough *Brucella* were isolated from Himalayan marmot (*Marmota himalayana*). Multiple-locus variable-number tandem-repeat analysis-16 typing indicated that the isolates both for smooth and atypical rough phenotypes had a common novel multiple-locus variable-number tandem-repeat analysis-16 type (Panel 1: 3-2-7-5-3-4-12-3; Panel 2A: 6-40-8; Panel 2B: 5-2-3-3-3).

Brucellosis is a highly contagious bacterial zoonosis. It is an important public health problem in many parts of the world (Godfroid et al. 2005). Currently, the genus *Brucella* includes at least 10 species, such as *Brucella melitensis*, *Brucella abortus*, *Brucella suis*, and *Brucella canis*, according to serology, morphology, staining, metabolic phenotype, culture characteristics, and phage typing (Corbel 1997; Scholz and Vergnaud 2013). The natural maintenance of *Brucella* in wildlife is a great obstacle to the control and eradication of brucellosis. For instance, *B. abortus* has been isolated in bison (*Bison bison*), coyotes (*Canis latrans*), bighorn sheep (*Ovis canadensis*), Chinese water deer (*Cervus unicolor deejani*), and gorals (*Naemorhedus goral*). In addition, some atypical *Brucella* strains have been found in amphibians, murine rodents, blue-spotted stingrays (*Dasyatis akajei*), sea otters (*Enhydra lutris*), and red foxes (*Vulpes vulpes*) (Godfroid 2002).

The Xinjiang Uygur Autonomous Region (XUAR), located in northwestern China, is considered an epidemic area of brucellosis

caused by *B. abortus*. In this region, a large rodent species, the Himalayan marmot (*Marmota himalayana*), inhabits alpine grasslands at an altitude of 2,500–5,200 m, and is considered to be the principal host of plague (Wang et al. 1983). In the southern XUAR, the Himalayan marmot often comes into close contact with local yaks (*Bos grunniens*) and yak hybrids (i.e., dzo), which prefer to graze in marmot habitats (Poudel et al. 2016). From August to September 2016–18, a total of 226 wild Himalayan marmots were captured (as approved by hunting license) by cages in Qira county (3,753 m above sea level; 37°52'N latitude, 81°57'E longitude, southern region of the XUAR). Zoletil 50 (Virbac, Paris, France) was used in anesthesia by intramuscular injection, blood was collected via the dorsalis pedis vein, serum was separated immediately by Test Kit for Plague F1 Antibody (Colloidal Gold, Beijing Zhuangdi-haohe Biomedical Technology Company Limited, Beijing, China) and by smooth/rough rose bengal plate test for antibodies to *Brucella* lipopolysaccharide (LPS) in the field. No plague F1 antibody-positive samples were detected in 226 marmots, and 33 *Brucella*-seropositive animals were detected cumulatively (1 case in 2016, 6 cases in 2017, and 26 cases in 2018). The latter result was further confirmed by a standard tube agglutination test to *Brucella* smooth/rough LPS. A total of 27 smooth and 6 rough to *Brucella* LPS were




FIGURE 1. Dendrogram based on the multiple-locus variable-number tandem-repeat analysis-16 genotyping assay showing relationships of *Brucella* isolates from the Himalayan marmot (*Marmota himalayana*) and other strains in this study. In the columns, the following data for isolates are indicated: ID=strain name; year=year of isolation; place=geographic region; host=source of *Brucella* isolation.

screened out, and their antibody titers ranged from 1:100 to 1:400.

To detect and isolate *Brucella*, 33 *Brucella*-seropositive marmots were all enrolled, and their DNA samples of heart, liver, spleen, kidney, lung, large intestine, small intestine, and muscle were respectively extracted and individually amplified by two genetic markers (*omp22* and *IS711*; Wang et al. 2018). The PCR results are shown in Supplementary Material Table 1. Relevant sequences were deposited in the GenBank database (*omp22*: MH316168 and MH316169; *IS711*: MH316166 and MH316167). According to PCR and sequencing results, the procedures of classical *Brucella* isolation and phenotypic identification were conducted (Yang et al. 2018). Briefly, the livers and spleens from 10 *Brucella* DNA-positive marmots were respectively homogenized, then their suspensions were individually inoculated onto *Brucella*-selective agar plates (Becton, Dickinson and

Company, Sparke, Maryland, USA). All cultures were incubated at 37 C with 5% carbon dioxide for 5 d. The suspected *Brucella* colonies were identified by the Brucellosis Laboratory, National Institute for Communicable Disease Control and Prevention (Beijing, China), and the Chinese Center for Disease Control and Prevention (Beijing, China). Consequently, six isolates of *B. abortus* biovar 1 and two isolates of an atypical rough *Brucella* strain were obtained (Supplementary Table 2). To further investigate genetic characteristics of *Brucella* isolates, a multiple-locus variable-number tandem-repeat analysis (MLVA) typing assay was carried out (Maquart et al. 2009). The MLVA-16 typing indicated that the isolates both for smooth and atypical rough phenotypes had a common novel MLVA-16 type (Panel 1: 3-2-7-5-3-4-12-3; Panel 2A: 6-40-8; Panel 2B: 5-2-3-3-3). The MLVA-16 phylogenetic analysis showed that the isolates from marmots shared

a common branch with *B. abortus* Ma51 and Ma08 strains from yaks and humans in Qinghai Province (Fig. 1).

In this study, *B. abortus* biovar 1 and atypical rough *Brucella* were isolated from the Himalayan marmot. Previously, atypical *Brucella* strains had been reported from sheep (*Ovis aries*), blue sheep (bharal, *Pseudois nayaur*), Mongolian gazelle (*Procapra przewalskii*), cattle (*Bos taurus*), yak, reindeer (*Rangifer tarandus*), dogs (*Canis lupus familiaris*), and humans in China (Cui et al. 2010; Godfroid et al. 2013). According to our retrospective investigation, *M. himalayana* and the yak and the dzo share a common habitat, raising the possibility of *Brucella* transmission between these hosts. In previous work, 5.9% (33/557) of yaks in Tibet Autonomous Region and 10.5% (50/474) of dzos in Qinghai Province were serologically positive to smooth *Brucella* antigen (Zhang et al. 2011; Wang et al. 2013). *Brucella abortus* can be naturally transmitted between wild ruminants and cattle (Truong et al. 2011). Therefore, it is an important task for the future to isolate and characterize brucellae from yaks and dzos. Serologic investigations should be carried out in shepherd dogs and wild wolves (*Canis lupus*), which eat the Himalayan marmot. In addition, further studies should determine if *Brucella* isolates in the Himalayan marmot originate from local yaks or dzos or naturally circulate in the Himalayan marmot, and the genomic characteristics of *Brucella* isolates in the Himalayan marmot.

SUPPLEMENTARY MATERIAL

Supplementary material for this article is online at <http://dx.doi.org/10.7589/2019-09-237>.

LITERATURE CITED

- Corbel MJ. 1997. Brucellosis: An overview. *Emerg Infect Dis* 3:213–221.

- Cui LJ, Wang XL, Wang YC. 2010. Brucellosis in wild animals. *Chin J Zoonoses* 26:283–288. [In Chinese.]
- Godfroid J. 2002. Brucellosis in wildlife. *Rev Sci Tech* 21: 277–286.
- Godfroid J, Cloeckaert A, Liautard JP, Kohler S, Fretin D, Walravens K, Garin-Bastuji B, Letesson JJ. 2005. From the discovery of the Malta fever's agent to the discovery of a marine mammal reservoir, brucellosis has continuously been a re-emerging zoonosis. *Vet Res* 36:313–326.
- Godfroid J, Garin-Bastuji B, Saegerman C, Blasco JM. 2013. Brucellosis in terrestrial wildlife. *Rev Sci Tech* 32:27–42.
- Maquart M, Le Flèche P, Foster G, Tryland M, Ramisse F, Dønne B, Al Dahouk S, Jacques I, Neubauer H, Walravens K, et al. 2009. MLVA-16 typing of 295 marine mammal *Brucella* isolates from different animal and geographic origins identifies 7 major groups within *Brucella ceti* and *Brucella pinnipedialis*. *BMC Microbiol* 9:145.
- Poudel BS, Spooner PG, Matthews A. 2016. Behavioural changes in marmots in relation to livestock grazing disturbance: An experimental test. *Eur J Wildl Res* 62:491–495.
- Scholz HC, Vergnaud G. 2013. Molecular characterisation of *Brucella* species. *Rev Sci Tech* 32:149–162.
- Truong LQ, Kim JT, Yoon BI, Her M, Jung SC, Hahn TW. 2011. Epidemiological survey for *Brucella* in wildlife and stray dogs, a cat and rodents captured on farms. *J Vet Med Sci* 73:1597–1601.
- Wang CL, Gao LY, Li GC, Wang QJ. 2013. Serological investigation of dzo brucellosis. In Delingha Zone, Qinghai Province. *Chin Qinghai J Anim Vet Sci* 43: 28. [In Chinese.]
- Wang Q, Zhao SS, Wureli H, Xie SS, Chen CF, Wei QQ, Cui BY, Tu CC, Wang YZ. 2018. *Brucella melitensis* and *B. abortus* in eggs, larvae and engorged females of *Dermacentor marginatus*. *Ticks Tick Borne Dis* 9: 1045–1048.
- Wang SB, Yang GY. 1983. *Rodent records of Xinjiang*. Xinjiang People's Publishing House, Urumqi, China, 223 pp. Chinese.
- Yang HX, Feng JJ, Zhang QX, Hao RE, Yao SX, Zhao R, Piao DR, Cui BY, Jiang H. 2018. A case report of spontaneous abortion caused by *Brucella melitensis* biovar 3. *Infect Dis Poverty* 7:31.
- Zhang KR, Gong G, Suolang SZ, Danba CR, Li JK, Guo AZ. 2011. The primary reports on serological investigation of yak brucellosis in several areas of Tibet. *China Dly Cattle* 24:48–49. [In Chinese.]

Submitted for publication 26 September 2019.

Accepted 11 December 2019.