

ISOLATION AND ANTIMICROBIAL SUSCEPTIBILITIES OF NONTUBERCULOUS MYCOBACTERIA FROM WILDLIFE IN JAPAN

Authors: Odoi, Justice Opare, Ohya, Kenji, Moribe, Junji, Takashima, Yasuhiro, Sawai, Kotaro, et al.

Source: Journal of Wildlife Diseases, 56(4) : 851-862

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/2019-10-261>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISOLATION AND ANTIMICROBIAL SUSCEPTIBILITIES OF NONTUBERCULOUS MYCOBACTERIA FROM WILDLIFE IN JAPAN

Justice Opare Odoi,^{1,7} Kenji Ohya,^{1,2,3,7} Junji Moribe,^{2,4} Yasuhiro Takashima,^{1,2,3} Kotaro Sawai,² Kyoichiro Taguchi,² Hideto Fukushi,^{1,2} Takayuki Wada,⁵ Shiomi Yoshida,⁶ and Tetsuo Asai^{1,3,8}

¹ Department of Applied Veterinary Sciences, United Graduate School of Veterinary Sciences, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

² Faculty of Applied Biological Sciences, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

³ Education and Research Center for Food Animal Health, Gifu University (GeFAH), 1-1 Yanagido, Gifu 501-1193, Japan

⁴ Research Center for Wildlife Management, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

⁵ Department of International Health, Institute of Tropical Medicine, Nagasaki University, Nagasaki 852-8523, Japan

⁶ Clinical Research Centre, National Hospital Organization Kinki-chuo Chest Medical Centre, Sakai, Osaka 591-8555, Japan

⁷ These authors contributed equally to this work.

⁸ Corresponding author (email: tasai@gifu-u.ac.jp)

ABSTRACT: Nontuberculous mycobacteria (NTM) are opportunistic pathogens of humans and animals and are transmitted among the environment, wildlife, livestock, and humans. The aim of this study was to investigate the rate of isolation and antimicrobial susceptibility of NTM in wildlife. In total, 178 samples of feces ($n=131$) and tissues ($n=47$) were collected from 11 wildlife species in Gifu Prefecture and Mie Prefecture, Japan, between June 2016 and October 2018. We isolated NTM from 15.3% (20/131) of fecal samples using Ogawa medium, and isolates were identified by sequencing the *rpoB* and *hsp65* genes. The *rpoB* sequences were compared with those from other strains of human and environmental origin. The NTM isolates were obtained from sika deer (*Cervus nippon*), wild boar (*Sus scrofa*), Japanese monkey (*Macaca fuscata*), raccoon dog (*Nyctereutes procyonoides*), masked palm civet (*Paguma larvata*), and Japanese weasel (*Mustela itatsi*) and were classified as rapidly growing mycobacteria (RGM) and slowly growing mycobacteria (SGM). The 12 RGM identified were *Mycolicibacterium peregrinum* ($n=5$), *Mycolicibacterium fortuitum* ($n=3$), *Mycolicibacterium septicum* ($n=3$), and *Mycolicibacterium thermoresistibile* ($n=1$), and the eight SGM were *Mycobacterium paraense* ($n=4$), *Mycolicibacter arupensis* ($n=2$), *Mycolicibacter virginianensis* ($n=1$), and *Mycobacterium nebraskense* ($n=1$). The NTM from wildlife showed $\geq 99\%$ similarity with strains from different sources including humans. The RGM were susceptible to the antimicrobial agents tested except for *M. fortuitum*, which was resistant to azithromycin and clarithromycin. The SGM showed multiple drug resistance qualities but were susceptible to amikacin, clarithromycin, and rifabutin. These results indicate that wildlife may be reservoir hosts of NTM in Japan. The presence of antimicrobial-resistant NTM in wildlife suggests that the trends of NTM antimicrobial susceptibility in wildlife should be monitored.

Key words: Antimicrobial resistance, fecal sample, nontuberculous mycobacteria, wildlife.

INTRODUCTION

Nontuberculous mycobacteria (NTM) are organisms residing in natural environments to which humans and animals are exposed (Falkinham 1996). Although they do not cause tuberculosis or leprosy, they are known to be opportunistic pathogens of humans and animals (Biet et al. 2005). Among NTM, *Mycobacterium avium*–*Mycobacterium intracellulare* complex (MAC) is the most important etiologic agent to humans and animals (Primm et al. 2004). In fact, NTM have been isolated from humans and livestock, but more attention has been given to MAC, since it is

frequently associated with infections in humans (Nishiuchi et al. 2009). The importance of NTM, other than MAC, as an opportunistic infection in humans and animals has also been increasing (Donohue 2018). For example, there have been recent reports of the isolation of *Mycolicibacterium thermoresistibile* from feline panniculitis and isolation of *Mycolicibacterium peregrinum* and *Mycobacterium genavense* from migratory birds (Biet and Boschiroli 2014; Patiño et al. 2018).

Based on a recent comparative and phylogenomic study of the genus *Mycobacterium*, the division of the genus into five distinct monophyletic clades has been proposed. The

suggested modifications include an emended genus *Mycobacterium* (Tuberculosis-Simiae clade) and four other new genera, *Mycolicibacterium* gen. nov. (Fortuitum-Vaccae clade), *Mycolicibacter* gen. nov. (Terraе clade), *Mycolicibacillus* gen. nov. (Triviale clade), and *Mycobacteroides* gen. nov. (Abscessus-Chelonae clade; Gupta et al. 2018). This comparative analysis has improved our understanding of NTM and clearly places all the rapidly growing mycobacteria (RGM) and the slowly growing mycobacteria (SGM) in distinct clades.

To provide more information on the ecology and etiology of NTM, we previously presented the genome sequences of *Mycolicibacter virginensis*, *Mycolicibacter senuensis*, and *Mycobacterium colombiense* isolated from the environment surrounding pigs in Japan (Ito et al. 2018a, b). In addition, we analyzed the genome sequences of *Mycolicibacterium peregrinum* isolated from the lymph nodes of a pig and from the surrounding soil by in silico DNA-DNA hybridization, and the results showed that the two isolates were 100% identical (Komatsu et al. 2019), implying circulation of NTM between the animal and the surrounding environment. Although NTM cases have been recorded in both humans and animals, the focus has been on humans more than on animals, especially wildlife.

Antimicrobial resistance in NTM from wildlife is critical, since there is the possibility of spreading to humans and livestock. However, the antimicrobial resistance pattern varies according to the species of NTM and often hinders the efficacy of therapeutic regimens (van Ingen et al. 2012). In Japan, antimicrobial resistance has been observed in clinical NTM isolates (Hatakeyama et al. 2017), but antimicrobial susceptibility testing of NTM from wildlife has not been reported.

Wildlife can acquire environmental mycobacteria through contaminated water and food (Kazda et al. 2009). The bacteria then inhabit favorable niches within the animals and can be transmitted through their feces to the environment (Pavlik et al. 2009). We investigated the rate of isolation and antimicrobial susceptibility of NTM from wildlife in two prefectures (Gifu and Mie) in Japan and




FIGURE 1. The samples were collected from wildlife in Gifu Prefecture and Mie Prefecture, located in central Japan. The two prefectures provide habitats for various wildlife species. The animals were hunted with traps by authorized hunters between June 2016 and October 2018. The study sites were selected according to the population density of wildlife, with a focus on mammals as potential reservoir hosts in proximity to the human environment. The dark shaded portion of the map represents the location of wildlife species from which samples were used.

examined the genetic relationship between NTM isolated from wildlife and other sources.

MATERIALS AND METHODS

Study area and design

The samples were collected from wildlife (Table 1) in Gifu Prefecture and Mie Prefecture, located in central Japan (Fig. 1). The two prefectures provide habitats for various wildlife species. The study sites were selected according to the population density of wildlife, with a focus on mammals as potential reservoir hosts in proximity to the human environment. Some wildlife species were annually hunted as part of culling practices in the fields, forests, and rivers in the sampling areas (Table 2). Other animals (e.g., fox and some weasels) were found as roadkill or had died of natural causes. The animals were hunted with traps by authorized hunters between June 2016 and October 2018 in Gifu Prefecture and Mie Prefecture, Japan, with adherence to the Protection and Control of Wild

TABLE 1. Rate of isolation of nontuberculous mycobacteria (NTM) from wildlife species included in this study in Japan. The animals were hunted with traps by authorized hunters between June 2016 and October 2018 in Gifu Prefecture and Mie Prefecture, Japan. Fecal and tissue samples were collected at a game-handling establishment during gralloching (removal of the gut) and carcass inspection and subjected to NTM isolation using 2% Ogawa slant culture tube.

Order	Family	English name	Scientific name	No. samples		No. positive for NTM (%)	
				Feces	Tissue	Feces	Tissue
Primates	Cercopithecidae	Japanese monkey	<i>Macaca fuscata</i>	31	10	1 (3)	0
Rodentia	Myocastoridae	Nutria	<i>Myocastor coypus</i>	1	0	0	0
Carnivora	Canidae	Red fox	<i>Vulpes vulpes</i>	1	0	0	0
		Raccoon dog	<i>Nyctereutes viverrinus</i>	22	13	1 (5)	0
	Procyonidae	Raccoon	<i>Procyon lotor</i>	6	10	0	0
	Mustelidae	Japanese weasel	<i>Mustela itatsi</i>	2	0	1 (50)	0
		Siberian weasel	<i>Mustela sibirica</i>	4	0	0	0
	Viverridae	Masked palm civet	<i>Paguma larvata</i>	9	6	1 (11)	0
Artiodactyla	Cervidae	Sika deer	<i>Cervus nippon</i>	41	8	13 (32)	0
	Suidae	Wild boar	<i>Sus scrofa</i>	12	0	3 (25)	0
Anseriformes	Anatidae	Wild duck (mallard)	<i>Anas platyrhynchos</i>	2	0	0	0
Total				131	47	20 (15.3)	0

Birds and Mammals and Hunting Management Law (law no. 88, 2002). Samples used in this study were collected at a game-handling establishment during gralloching (removal of the gut) and carcass inspection. Because animals were not killed specifically for this study, no ethical approval was required from the Gifu University ethical committee. However, the Protection and Control of Wild Birds and Mammals Hunting Management Law (law no. 88, 2002) was followed during sample collection. The sample size was dependent on availability and the hunting period.

Sample collection

In total, 178 samples were investigated, which included 131 fecal samples from the large intestine and 47 tissue samples of the lymph node ($n=11$), lung ($n=9$), tonsil ($n=9$), liver ($n=9$), spleen ($n=6$), and kidney ($n=3$). The samples were collected from Japanese monkey (*Macaca fuscata*), nutria (*Myocastor coypus*), red fox (*Vulpes vulpes*), raccoon dog (*Nyctereutes viverrinus*), raccoon (*Procyon lotor*), Japanese weasel (*Mustela itatsi*), Siberian weasel (*Mustela sibirica*), masked palm civet (*Paguma larvata*), sika deer (*Cervus nippon*), wild boar (*Sus scrofa*), and mallard (*Anas platyrhynchos*) as shown in Table 1. Classification and scientific names of the wildlife species were attributed according to previous literature (Ohdachi et al. 2015).

Isolation of bacteria

Sample collection was conducted by veterinary wildlife specialists to avoid contamination of samples during transportation and at the laboratory (Pate et al. 2016). The fecal and tissue samples were immediately processed using 2% sodium hydroxide (Hibiya et al. 2010; Ito et al. 2018b). The inocula obtained from both tissues and feces were inoculated onto a 2% Ogawa slant culture tube (Kyokuto Pharmacy, Tokyo, Japan) and into BBL™ (Difco Laboratories, Sparks, Maryland, USA) mycobacterium growth indicator tube liquid media supplemented with polymyxin B, amphotericin B, nalidixic acid, trimethoprim, and azlocillin (Becton Dickinson, Franklin Lakes, New Jersey, USA) and incubated at 37 C for 4 wk. Each single colony formed on the slant tube was subcultured on Middlebrook 7H11 agar (Difco) to obtain distinct colonies (Ito et al. 2018b).

DNA extraction, sequencing, and phylogenetic analysis

We extracted DNA using the boiling method (Iwamoto et al. 2012). The first round of PCR was performed to determine the samples that were positive for mycobacteria (Chen et al. 1996). To identify the isolated mycobacteria, sequencing of the *rpoB* and *hsp65* genes (Telenti et al. 1993; Adekambi et al. 2003) was performed. Because of its high discrimination ability, the partial sequence of the *rpoB* gene was used to compare

TABLE 2. Species of nontuberculous mycobacteria (NTM) identified from wildlife in Japan. Bacteria identification was achieved by partial sequencing of *rpoB* and *hsp65* gene (Telenti et al. 1993; Adekambi et al. 2003). Except when otherwise stated, all of the isolates were made from samples from animals from Gifu Prefecture, Japan.

Wildlife species	Identified NTM	Isolate ID	City	GenBank accession no.	
				<i>rpoB</i>	<i>hsp65</i>
Sika deer	<i>Mycolicibacterium peregrinum</i>	wg41	Ibigawa	MK341497	MK341516
	<i>Mycolicibacterium septicum</i>	wg56	Ibigawa	MK341503	MK341522
		wg60	Ogaki	MK341504	MK341523
	<i>Mycolicibacterium fortuitum</i>	wg7	Ibigawa	MK341491	MK341511
		wg21	Ibigawa	MK341494	MK341514
		wg86	Ibigawa	MK341508	MK341527
	<i>Mycolicibacterium thermoresistibile</i>	wg48(i)	Motosu	MK341499	MK341518
	<i>Mycolicibacter arupensis</i>	wg48(ii)	Motosu	MK341500	MK341519
		wg68	Ibigawa	MK341506	MK341525
	<i>Mycobacterium nebraskense</i>	wg8	Ibigawa	MK341492	MK341512
	<i>Mycobacterium paraense</i>	wg4	Ibigawa	MK341490	MK341510
		wg42	Ibigawa	MK341498	MK341517
		wg54	Ibigawa	MK341502	MK341521
		wg18	Ibigawa	MK341493	MK341513
Wild boar	<i>Mycobacterium paraense</i>	wg29	Motosu	MK341495	MK341515
	<i>Mycolicibacter virginiensis</i>	wg30	Motosu	MK341496	MK587449
Japanese monkey	<i>Mycolicibacterium peregrinum</i>	wg51	Ibigawa	MK341501	MK341520
Raccoon dog	<i>Mycolicibacterium peregrinum</i>	wg90	Ibigawa	MK341509	MK341528
Masked palm civet	<i>Mycolicibacterium septicum</i>	wg72 ^a	Miegun	MK341507	MK341526
Japanese weasel	<i>Mycolicibacterium peregrinum</i>	wg62	Ibigawa	MK341505	MK341524

^a Isolate was obtained from an animal from Mie Prefecture.

the isolates obtained from wildlife (Table 2) with previously characterized NTM sequences downloaded from GenBank; isolates with $\geq 99\%$ identities were considered closely related (de Zwaan et al. 2014). The obtained sequences were analyzed using nucleotide BLAST (National Center for Biotechnology Information 2018), and the sequences with a similarity percentage above 98% were downloaded for phylogenetic analysis. The phylogenetic analysis was conducted according to the neighbour-joining method (Saitou and Nei 1987) under the total gap removal and Kimura's two-parameter substitution model (Kimura 1980), and it was evaluated by bootstrap analysis based on 1,000 replicates using MEGA software version 7 (Kumar et al. 2016). The tree was rooted using *Rhodococcus equi* (GenBank no. AY492285) as the outgroup.

Antimicrobial susceptibility testing of RGM

We used broth microdilution plates containing antimicrobial agents and cation-supplemented Müeller-Hinton broth (Eiken Chemical Co., Ltd., Tokyo, Japan). The final concentrations of

the antimicrobial agents evaluated for RGM isolates were: clarithromycin (CAM) 0.0015 to 64 $\mu\text{g/mL}$, azithromycin (AZM) 0.0015 to 64 $\mu\text{g/mL}$, sitafloxacin 0.0015 to 64 $\mu\text{g/mL}$, moxifloxacin 0.0015 to 64 $\mu\text{g/mL}$, ciprofloxacin 0.0015 to 64 $\mu\text{g/mL}$, imipenem 1 to 64 $\mu\text{g/mL}$, meropenem 1 to 64 $\mu\text{g/mL}$, amikacin (AMK) 2 to 64 $\mu\text{g/mL}$, and linezolid 2 to 64 $\mu\text{g/mL}$. An inoculum with a final concentration ranging from 1×10^5 to 5×10^5 colony-forming units/mL was prepared according to the Clinical and Laboratory Standards Institute (CLSI) guidelines (2011). The plates were inoculated and incubated at 37 C. The minimum inhibitory concentration (MIC) was determined on day 3, 4, or 5 (Hatakeyama et al. 2017). However, to detect inducible resistance in the case of CAM, the incubation period was extended, and the final MIC was determined on day 14 (Brown-Elliott et al. 2012). The MIC breakpoints were interpreted according to CLSI criteria (CLSI 2011) with the exception of sitafloxacin, which was interpreted using the breakpoint proposed by O'Grady et al. (2001).

Antimicrobial susceptibility testing of SGM

The BrothMIC NTM system (Kyokuto Pharmaceutical Industrial Co. Ltd., Tokyo, Japan) was used to evaluate the susceptibility of SGM to streptomycin 0.06 to 128 µg/mL, ethambutol (EB) 0.06 to 128 µg/mL, kanamycin 0.06 to 128 µg/mL, rifampicin 0.06 to 32 µg/mL, rifabutin (RBT) 0.06 to 16 µg/mL, levofloxacin 0.06 to 32 µg/mL, CAM 0.06 to 32 µg/mL, ethionamide 0.06 to 16 µg/mL, and AMK 0.5 to 16 µg/mL, based on the manufacturer's guidelines, which were in accordance with the standard protocol of CLSI. The suspension for inoculation was prepared as described elsewhere (Pang et al. 2017). The plates were incubated at 37 C for 7 d, and the growth end point was determined visually. The MIC breakpoints were interpreted following the guidelines of CLSI (2011) and of the World Health Organization (2008).

RESULTS

Rate of isolation of NTM in wildlife species

Twenty NTM were isolated from fecal samples of six wildlife species, sika deer (32%, 13/41), wild boar (25%, 3/12), Japanese monkey (3%, 1/31), raccoon dog (5%, 1/22), masked palm civet (11%, 1/9), and Japanese weasel (50%, 1/2), resulting in an estimated isolation rate of 15.3% (Table 1). The isolated NTM included four RGM, *Mycolicibacterium peregrinum* (n=5), *Mycolicibacterium fortuitum* (n=3), *Mycolicibacterium septicum* (n=3), and *Mycolicibacterium thermoresistibile* (n=1), and four SGM, *Mycobacterium paraense* (n=4), *Mycolicibacter arupensis* (n=2), *Mycolicibacter virginiensis* (n=1), and *Mycobacterium nebraskense* (n=1), as shown in Figures 2 and 3.

The phylogenetic tree of the NTM identified was based on the 430-base pair *hsp65* gene (Fig. 3). The *hsp65* gene could not clearly differentiate between some members of the *Mycolicibacter terrae* clade. The sequences of both *rpoB* and *hsp65* were deposited in GenBank (accession numbers shown in Table 2).

Comparison of NTM isolated from wildlife with NTM from other sources—*Fortuitum-Vaccae* clade

The NTM from wildlife were compared with those from other sources to investigate




FIGURE 2. Neighbor-joining phylogenetic tree of nontuberculous mycobacteria (NTM) isolates based on the alignment of partial nucleotide sequence (720-base pair) of *rpoB* gene sequenced using primers described by Adekambi et al. (2003). The representative isolates of NTM identified in this study are preceded with the letters wg, and the source is indicated with the obtained accession numbers in parentheses. The reference NTM strains downloaded from the National Center for Biotechnology Information are in italics, with their respective accession numbers in parentheses. Branching with greater than 55% support from 1,000 bootstrap replicates is shown at the nodes. Horizontal distances indicate genetic distance, and the scale bar represents nucleotide substitution per site.

their genetic relatedness. We found similarity among all the NTM from different sources (Fig. 4). The isolates of *Mycolicibacterium peregrinum* were 100% identical to each other and were also 100% identical to the reference strain *Mycolicibacterium peregrinum* strain Y7 (KU861853) isolated from water in Iran. Similarly, the isolates from wildlife identified in this study as *Mycolicibacterium fortuitum* were 100% identical to each other and to the *Mycolicibacterium fortuitum* strain garg (KC242461) isolated from water in India. The *Mycolicibacterium septicum* isolated from wildlife and the reference strain *Mycolicibacterium septicum* strain 14MF (KY400670) isolated from soil in China were 99% similar. The *Mycolicibacterium thermoresistibile* isolated from wildlife was 100% identical to isolates from humans (France) and soil (UK).




FIGURE 3. Neighbor-joining phylogenetic tree of nontuberculous mycobacteria (NTM) isolates based on the alignment of partial nucleotide sequence (430-base pair) of *hsp65* gene sequenced using primers described by Telenti et al. (1993). The isolates of NTM identified in this study are preceded with the letters wg, and the source is indicated with the obtained accession numbers in parentheses. The reference NTM strains downloaded from the National Center for Biotechnology Information are in italics, with their respective accession numbers in parentheses. Horizontal distances indicate genetic distance, and the scale bar represents nucleotide substitution per site.

Comparison of NTM isolated from wildlife and NTM from other sources—*Terrae* and *Tuberculosis-Simiae* clades

Mycobacterium paraense and *Mycobacterium nebraskense* are members of the Tuberculosis-Simiae clade, while the *Terrae* clade includes the *Mycolicibacter arupensis* and *Mycolicibacter virginiensis* isolates identified in this study. Five isolates were members of the Tuberculosis-Simiae, and three were members of the *Terrae* clade. The two clades were analyzed together because of the relatively small number of each. *Mycobacterium paraense* isolated in this study was 100% identical to *Mycobacterium paraense* strain IEC23 isolated from humans in Brazil. *Mycobacterium nebraskense* was 100% identical to *Mycobacterium nebraskense* isolated from humans in the US. *Mycolicibacter arupensis*




FIGURE 4. Comparison of the nontuberculous mycobacteria (NTM) from the Fortuitum-Vacciae clade isolated from wildlife (this study) and other sources, including human, soil, and water based on nucleotide sequence (720-base pair) of *rpoB* gene (Adekambi et al. 2003). The isolates of NTM identified in this study are preceded with the letters wg, and the source is indicated with the obtained accession numbers in parentheses. The reference NTM strains downloaded from the National Center for Biotechnology Information are in italics, with their respective source and accession numbers in parentheses.

and *Mycolicibacter virginiensis* were also 100% identical to the *Mycolicibacter arupensis* strain FI-06239 isolated from a human in Italy and the *Mycolicibacter virginiensis* strain 12MF isolated from soil in China (Fig. 5).

Antimicrobial susceptibility testing of RGM and SGM

Most of the RGM isolated from wildlife were susceptible to all nine antimicrobial agents used in this study (Table 3); the exception was *Mycolicibacterium fortuitum*, which showed resistance to CAM and AZM. Table 4 shows the MIC and the resistance pattern of the SGM to the antimicrobial agents tested. All the SGM investigated were resistant to more than two antimicrobial agents but susceptible to CAM, RBT, and AMK.

DISCUSSION

In this study, we successfully isolated 20 NTM from the large intestine of six wildlife




FIGURE 5. Comparison of the nontuberculous mycobacteria (NTM) belonging to Terrae and Tuberculosis-Simiae clades isolated from wildlife (this study) and other sources, including human, soil, and water, based on the nucleotide sequence (720-base pair) of the *rpoB* gene (Adekambi et al. 2003). The isolates of NTM identified in this study are preceded with the letters wg, and the source is indicated with the obtained accession numbers in parentheses. The reference NTM strains downloaded from the National Center for Biotechnology Information are in italics, with their respective source and accession numbers in parentheses.

species. The isolated NTM included four RGM (*Mycolicibacterium peregriinum*, *Mycolicibacterium fortuitum*, *Mycolicibacterium septicum*, and *Mycolicibacterium thermoresistibile*) and four SGM (*Mycobacterium paraense*, *Mycobacterium nebraskense*, *Mycolicibacter virginianus*, and *Mycolicibacter arupensis*). The isolation rate of NTM in feces in this study was 15.3%, which is higher than the rates of 11.8% and 8.4% observed in Slovenia (Pate et al. 2016) and Hungary (Ronai et al. 2016), respectively. The occurrence of NTM in wildlife may be due to their extensive interaction with the environment, indicating that these animals may be reservoir hosts for NTM.

Mycolicibacterium peregriinum was the most frequently isolated RGM from the different wildlife species—sika deer, wild boar, Japanese monkey, Japanese weasel, masked palm civet, and raccoon dog—which strongly suggested that wildlife may acquire *Mycolicibacterium peregriinum* from their

environment. Studies conducted in Japan revealed that the draft genome sequence of *Mycolicibacterium peregriinum* isolated from a pig affected with lymphadenitis shares close identity with that of *Mycolicibacterium peregriinum* isolated from the soil of the pig's environment (Komatsu et al. 2019). Moreover, the fact that *Mycolicibacterium peregriinum* was isolated from other animals (e.g., lion; *Panthera leo*), and from drinking water (Gcebe and Hlokwe 2017; Mohajeri et al. 2017) indicates that *Mycolicibacterium peregriinum* can inhabit several environments. In our study, the *rpoB* gene of the *Mycolicibacterium peregriinum* isolated from wildlife showed 100% similarity to that of *Mycolicibacterium peregriinum* isolated from drinking water (Mohajeri et al. 2017).

All the *Mycolicibacterium fortuitum* isolates in this study were from deer fecal samples, although *Mycolicibacterium fortuitum* was isolated from other animals, including monkey, in previous studies (Gcebe and Hlokwe 2017). The isolates of *Mycolicibacterium fortuitum* from wildlife in this study were identical to *Mycolicibacterium fortuitum* isolated from water (Fig. 4) and formed a cluster with other *Mycolicibacterium fortuitum* strains isolated from cats (Higgins et al. 2011) and hospital water (Khosravi et al. 2016), which suggests that, regardless of the location, NTM, in particular *Mycolicibacterium fortuitum*, were widespread in the environment, including among wildlife (Honda et al. 2018). *Mycolicibacterium septicum* and *Mycolicibacterium thermoresistibile* isolated from wildlife were 100% identical to other isolates from different sources, implying high relatedness among the isolates. In Japan, *Mycolicibacterium thermoresistibile* isolated from dust was not associated with infection (Tsukamura et al. 1985). However, both *Mycolicibacterium septicum* and *Mycolicibacterium thermoresistibile* have been reported in human and veterinary infection in other countries (Schinsky et al. 2000; Vishkautsan et al. 2016).

Among the wildlife fecal samples observed, diverse NTM were isolated from sika deer. Because NTM are known to inhabit the roots

TABLE 3. Results of the antimicrobial susceptibility test for rapidly growing mycobacteria (RGM). Minimum inhibitory concentration (MIC) range values were recorded along with the resistance percentage of the various RGM tested. Antimicrobial susceptibility testing of RGM was performed by broth microdilution in accordance with Clinical and Laboratory Standards Institute guidelines (CLSI 2011). Except where indicated, MIC values were interpreted based on breakpoints by CLSI (2011).

Antibiotic ^a	Reference MIC (μg/L)	MIC (μg/mL) range RGM isolates (no.)							
		<i>Mycobacterium peregrinum</i> (5)		<i>Mycobacterium fortuitum</i> (3)		<i>Mycobacterium septicum</i> (3)		<i>Mycobacterium thermoresistibile</i> (1)	
		MIC	% Resistant	MIC	% Resistant	MIC	% Resistant	MIC	% Resistant
CAM	≥8 ^b	0.03–2	0	8	100	0.06–0.09	0	0.03	0
AZM	≥8 ^b	0.06–4	0	8	100	0.16–0.38	0	0.02	0
STFX	≥30 ^c	0.03–0.25	0	0.02–0.03	0	0.02–0.04	0	0.02	0
MFLX	≥4 ^b	0.02–0.5	0	0.03–0.12	0	0.06–0.25	0	0.03	0
CPFX	≥4 ^b	0.03–2	0	0.06–1	0	0.06–0.25	0	0.03	0
IPM	≥32 ^b	1–4	0	≤1	0	≤1	0	≤1	0
MEPM	≥32 ^b	1–2	0	≤2	0	≤2	0	≤2	0
AMK	≥64 ^b	≤2	0	≤2	0	≤2	0	≤2	0
LZD	≥32 ^b	2–4	0	2–4	0	≤2	0	≤2	0

^a CAM = clarithromycin; AZM = azithromycin; STFX = sitafloxacin; MFLX = moxifloxacin; CPFX = ciprofloxacin; IPM = imipenem; MEPM = meropenem; AMK = amikacin; LZD = linezolid.

^b Resistance breakpoint (μg/mL) based on CLSI guidelines (2011).

^c Resistance breakpoint based on study by O’Grady et al. (2001).

and surface of leaves (Pavlik et al. 2009), it is possible that the high diversity of NTM detected in deer is related to the herbivorous feeding habits of the species. In addition, isolation of NTM, including *Mycobacterium fortuitum* and *Mycobacter arupensis*, from deer has been reported (de Lisle and Havill 1985; Mombeni et al. 2016). Three different NTM (*Mycobacterium peregrinum*, *Mycobacter virginiensis*, and *Mycobacterium paraense*) were isolated from wild boar samples because wild boar, as omnivores, can acquire NTM from infected carcasses they feed on as well as from the environment including plants (Pavlik et al. 2009). Moreover, *Mycobacterium paraense* was the most frequently isolated SGM and was isolated from sika deer, wild boar, and Japanese monkey. This finding is enlightening because most of the reported *Mycobacterium paraense* cases concern humans (Chin’ombe et al. 2016; Poonawala et al. 2017). *Mycobacterium paraense* was isolated from a human patient in Brazil and was described in 2015 as a novel species that is closely related to *Mycobacteri-*

um interjectum (da Costa et al. 2015). Given the different sources of isolation in this study, it is possible that *Mycobacterium paraense* has a broad host range.

Mycobacter arupensis and *Mycobacter virginiensis* identified in this study were clustered together (Fig. 5), since they belong to the Terrae clade as described previously (Gupta et al. 2018). *Mycobacter arupensis* was isolated from sika deer fecal samples, while *Mycobacter virginiensis* was isolated from wild boar. The *Mycobacter virginiensis* from wildlife in our study was identical to the *Mycobacter virginiensis* of human and soil origin (Fig. 5). However, it was not identical to the *Mycobacter virginiensis* strain GF75 isolated in Japan (Ito et al. 2018a). The partial *rpoB* gene—with a region known to have high discrimination power (Higgins et al. 2011; de Zwaan et al. 2014)—was able to identify and highlight the diversity within NTM. The lack of MAC isolation could be due to its absence in the samples or to growth inhibition due to decontamination (Sattar et al. 2018). In this study, we could

TABLE 4. Results of antimicrobial susceptibility test for slowly growing mycobacteria (SGM). Minimum inhibitory concentration (MIC) range values were recorded along with the resistance percentage of the various SGM tested. Antimicrobial susceptibility testing of SGM was performed by broth microdilution in accordance with Clinical and Laboratory Standards Institute guidelines (CLSI 2011). Except where indicated, MIC values were interpreted based on breakpoints by CLSI (2011).

Antibiotic ^a	Reference MIC (μg/L)	MIC (μg/mL) range SGM isolates (no.)							
		<i>Mycobacterium paraense</i> (4)		<i>Mycolicibacter arupensis</i> (2)		<i>Mycolicibacter virginiensis</i> (1)		<i>Mycobacterium nebraskense</i> (1)	
		MIC	% Resistant	MIC	% Resistant	MIC	% Resistant	MIC	% Resistant
SM	≥5 ^b	16–64	100	64–128	100	128	100	16	100
EB	≥4 ^b	32–128	100	0.5–1	0	2	0	16	100
KM	≥4 ^c	2–8	25	128	100	128	100	32	100
RFP	≥1 ^b	0.25–2	25	16–32	100	8	100	1	100
RBT	≥2 ^b	0.25–1	0	0.25–1	0	0.5	0	1	0
LVFX	≥8 ^b	1–4	0	32	100	32	100	2	0
CAM	≥32 ^b	0.25–1	0	1	0	1	0	0.25	0
TH	≥5 ^c	8–16	100	16	100	8	100	16	100
AMK	≥64 ^b	2–8	0	16	0	16	0	16	0

^a SM = streptomycin; EB = ethambutol; KM = kanamycin; RFP = rifampicin; RBT = rifabutin; LVFX = levofloxacin; CAM = clarithromycin; TH = ethionamide; AMK = amikacin.

^b Resistance breakpoint (μg/mL) based on CLSI guidelines (2011).

^c Resistance breakpoint (μg/mL) based on World Health Organization guidelines (2008).

not isolate NTM from wildlife tissue samples (liver, lungs, lymph node, tonsils, kidney, and spleen), unlike in other studies (Katale et al. 2014; Ghielmetti et al. 2018). This may have been due to the absence of lesions when the tissue samples were macroscopically observed. Considering that identical NTM were isolated from different wildlife species, there is a possibility of wildlife transmitting NTM to the environment of domestic animals. Since wildlife, humans, and other domestic animals can interact, the use of whole-genome analysis will provide additional evidence and deepen our understanding of the genetic relatedness and possible transfer of NTM among humans, domestic animals, and wildlife.

All the 12 RGM tested were susceptible to the fluoroquinolones, aminoglycosides, and carbapenems used in this study, which is in line with other reports indicating that aminoglycosides, such as amikacin, were effective against RGM (Li et al. 2013; Pang et al. 2015). The observed resistance of *Mycolicibacterium fortuitum* to CAM and AZM may be due to the presence of erythromycin ribosomal

methylese (*erm*; Brown-Elliott et al. 2012). The *Mycolicibacterium peregrinum*, *Mycolicibacterium septicum*, and *Mycolicibacterium thermoresistibile* isolates in this study were susceptible to all the antimicrobial agents tested in vitro (Table 3); however, in vitro antimicrobial susceptibility is not entirely conclusive.

All the *Mycobacterium paraense* isolates identified in this study were resistant to streptomycin, EB, and ethionamide (Table 4), similar to the resistance pattern observed in *Mycobacterium paraense* isolated from humans (da Costa et al. 2015). *Mycolicibacter arupensis* and *Mycolicibacter virginiensis* showed a similar resistance pattern and were susceptible to EB, CAM, RBT, and AMK. This finding is in agreement with the results of other studies in which EB, CAM, and RBT were found to be effective against the species of the *Terrae* clade (Vasiredy et al. 2017).

The study has some limitations. More fecal samples than tissue samples were investigated. However, this does not alter the conclusions of this study. In addition, we did not examine

the mechanism underlying the resistance of NTM to the individual antimicrobial agents tested. Furthermore, the partial sequencing of *rpoB* and *hsp65* genes for identification and comparison with other NTM species was not sufficient to establish a strong relationship among NTM from different sources; whole-genome sequences are required to establish this relationship. Thus, this study can be considered a preliminary study.

In conclusion, the isolation of diverse NTM from wildlife fecal samples may not be directly related to high exposure. However, these findings provide insight into the occurrence of NTM in wildlife and indicate that wildlife may be reservoir hosts. Notably, the presence of antimicrobial-resistant NTM is important with respect to human and veterinary medicine, since interaction with wildlife is plausible. Furthermore, the isolation of antimicrobial-resistant NTM in wildlife indicates that the trends of NTM antimicrobial susceptibility in wildlife should be carefully considered and monitored for emergence of resistant strains.

ACKNOWLEDGMENTS

This study was supported by the Japan Racing Association (JRA) through JRA livestock industry promotion project grants attributed to K.O., Y.T., and J.M. (H28-29_239) and to K.O. and Y.T. (H29-30_7). The Ito Foundation is kindly acknowledged for funding K.O. through research grants for Meat and Meat Products (H30-60). Japan Society for the Promotion of Science (JSPS) KAKENHI (JP18H04073) funded J.M. and T.A. We thank the Japanese Ministry of Education, Culture, Sports, Science, and Technology (MEXT) for the Joint Research Program of the Research Center for Zoonosis Control, Hokkaido University. J.O.O. received a scholarship from the Japanese government (MEXT).

LITERATURE CITED

- Adekambi T, Colson P, Drancourt M. 2003. *rpoB*-based identification of nonpigmented and late-pigmenting rapidly growing mycobacteria. *J Clin Microbiol* 41: 5699–5708.
- Biet F, Boschiroli ML. 2014. Non-tuberculous mycobacterial infections of veterinary relevance. *Res Vet Sci* 97:S69–S77.
- Biet F, Boschiroli ML, Thorel MF, Guilloteau LA. 2005. Zoonotic aspects of *Mycobacterium bovis* and *Mycobacterium avium-intracellulare* complex (MAC). *Vet Res* 36:411–436.
- Brown-Elliott BA, Nash KA, Wallace RJ Jr. 2012. Antimicrobial susceptibility testing, drug resistance mechanisms, and therapy of infections with nontuberculous mycobacteria. *Clin Microbiol Rev* 25:545–582.
- Chen ZH, Butler WR, Baumstark BR, Ahearn DG. 1996. Identification and differentiation of *Mycobacterium avium* and *M. intracellulare* by PCR. *J Clin Microbiol* 34:1267–1269.
- Chin'ombe N, Muzivizi B, Munemo E, Nziramasanga P. 2016. Molecular identification of nontuberculous mycobacteria in humans in Zimbabwe using 16S ribosequencing. *Open Microbiol J* 10:113–123.
- CLSI (Clinical and Laboratory Standards Institute). 2011. *Susceptibility testing of Mycobacteria, Nocardiae, and other aerobic actinomycetes; Approved Standard*; 2nd Ed., M24-A2. Clinical and Laboratory Standards Institute, Wayne, Pennsylvania, pp. 19–43.
- da Costa AR, Fedrizzi T, Lopes ML, Pecorari M, Da Costa WL, Giacobazzi E, da Costa Bahia JR, De Sanctis V, Lima KV, Bertorelli R, et al. 2015. Characterization of 17 strains belonging to the *Mycobacterium simiae* complex and description of *Mycobacterium paraense* sp. nov. *Int J Syst Evol Microbiol* 65:656–662.
- de Lisle GW, Havill PF. 1985. Mycobacteria isolated from deer in New Zealand from 1970–1983. *New Zeal Vet J* 33:138–140.
- de Zwaan R, van Ingen J, van Soolingen D. 2014. Utility of *rpoB* gene sequencing for the identification of nontuberculous mycobacteria in the Netherlands. *J Clin Microbiol* 52:2544–2551.
- Donohue MJ. 2018. Increasing nontuberculous mycobacteria reporting rates and species diversity identified in clinical laboratory reports. *BMC Infect Dis* 18:163.
- Falkinham JO III. 1996. Epidemiology of infection by nontuberculous mycobacteria. *Clin Microbiol Rev* 9: 177–215.
- Gcebe N, Hlokwé TM. 2017. Non-tuberculous mycobacteria in South African wildlife: Neglected pathogens and potential impediments for bovine tuberculosis diagnosis. *Front Cell Infect Microbiol* 7:15.
- Ghielmetti G, Friedel U, Scherrer S, Sarno E, Landolt P, Dietz O, Hilbe M, Zweifel C, Stephan R. 2018. Nontuberculous mycobacteria isolated from lymph nodes and faecal samples of healthy slaughtered cattle and the abattoir environment. *Transbound Emerg Dis* 65: 711–718.
- Gupta RS, Lo B, Son J. 2018. Phylogenomics and comparative genomic studies robustly support division of the genus *Mycobacterium* into an emended genus *Mycobacterium* and four novel genera. *Front Microbiol* 9:67.
- Hatakeyama S, Ohama Y, Okazaki M, Nukui Y, Moriya K. 2017. Antimicrobial susceptibility testing of rapidly growing mycobacteria isolated in Japan. *BMC Infect Dis* 17:197.

- Hibiya K, Kazumi Y, Nishiuchi Y, Sugawara I, Miyagi K, Oda Y, Oda E, Fujita J. 2010. Descriptive analysis of the prevalence and the molecular epidemiology of *Mycobacterium avium* complex-infected pigs that were slaughtered on the main island of Okinawa. *Comp Immunol Microbiol Infect Dis* 33:401–421.
- Higgins J, Camp P, Farrell D, Bravo D, Pate M, Robbe-Austerman S. 2011. Identification of *Mycobacterium* spp. of veterinary importance using *rpoB* gene sequencing. *BMC Vet Res* 7:77.
- Honda JR, Viridi R, Chan ED. 2018. Global environmental nontuberculous mycobacteria and their contemporaneous man-made and natural niches. *Front Microbiol* 9:2029.
- Ito T, Maruyama F, Sawai K, Nozaki K, Otsu K, Ohya K. 2018a. Draft genome sequence of *Mycobacterium virginiense* strain GF75, isolated from the mud of a swine farm in Japan. *Genome Announc* 6:e00362-18.
- Ito T, Sawai K, Kawai M, Nozaki K, Otsu K, Fukushima H, Ohya K, Maruyama F. 2018b. Draft genome sequences of *Mycobacterium senuensis* isolate GF74 and *Mycobacterium colombiense* isolates GF28 and GF76 from a swine farm in Japan. *Microbiol Resour Announc* 7:e00936-18.
- Iwamoto T, Nakajima C, Nishiuchi Y, Kato T, Yoshida S, Nakanishi N, Tamaru A, Tamura Y, Suzuki Y, Nasu M. 2012. Genetic diversity of *Mycobacterium avium* subsp. *hominissuis* strains isolated from humans, pigs, and human living environment. *Infect Genet Evol* 12:846–852.
- Katale BZ, Mbugi EV, Botha L, Keyyu JD, Kendall S, Dockrell HM, Michel AL, Kazwala RR, Rweyemamu MM, Van Helden P, et al. 2014. Species diversity of non-tuberculous mycobacteria isolated from humans, livestock and wildlife in the Serengeti ecosystem, Tanzania. *BMC Infect Dis* 14:616.
- Kazda J, Pavlik I, Falkinham JO III, Hruska K. 2009. The occurrence of pathogenic and potential pathogenic mycobacteria in animals and the role of environment in the spread of infection. In: *The ecology of mycobacteria: Impact on animal's and human's health*, Kazda J, Pavlik I, Falkinham JO III, Hruska K, editors. Springer Science and Business Media, Berlin, Germany, pp. 247–259.
- Khosravi AD, Hashemi Shahraki A, Hashemzadeh M, Sheini Mehrabzadeh R, Teimoori A. 2016. Prevalence of non-tuberculous mycobacteria in hospital waters of major cities of Khuzestan Province, Iran. *Front Cell Infect Microbiol* 6:42.
- Kimura M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16:111–120.
- Komatsu T, Ohya K, Sawai K, Odoi JO, Otsu K, Ota A, Ito T, Kawai M, Maruyama F. 2019. Draft genome sequences of *Mycobacterium peregrinum* isolated from a pig with lymphadenitis and from soil on the same Japanese pig farm. *BMC Res Notes* 12:341.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870–1874.
- Li G, Lian LL, Wan L, Zhang J, Zhao X, Jiang Y, Zhao LL, Liu H, Wan K. 2013. Antimicrobial susceptibility of standard strains of nontuberculous mycobacteria by microplate Alamar Blue assay. *PLoS One* 8:e84065.
- Mohajeri P, Yazdani L, Shahraki AH, Alvandi A, Atashi S, Farahani A, Almasi A, Rezaei M. 2017. Verification of frequency in species of nontuberculous mycobacteria in Kermanshah drinking water supplies using the PCR-sequencing method. *Microb Drug Resist* 23:359–364.
- Mombeni EG, Mosavari N, Gravand MM, Rezai AA, Keshavarz R, Tadayon K, Bakhshi R, Behmanesh R. 2016. First report of *Mycobacterium bovis* isolation from a European fallow deer (*Dama dama dama*) in Iran. *Iran J Public Health* 45:814–816.
- National Center for Biotechnology Information. 2018. Basic local alignment search tool (BLAST). <http://blast.ncbi.nlm.nih.gov/Blast.cgi>. Accessed October 2018.
- Nishiuchi Y, Tamura A, Kitada S, Taguri T, Matsumoto S, Tateishi Y, Yoshimura M, Ozeki Y, Matsumura N, Ogura H, et al. 2009. *Mycobacterium avium* complex organisms predominantly colonize in the bathtub inlets of patients' bathrooms. *Jpn J Infect Dis* 62:182–186.
- O'Grady J, Briggs A, Atarashi S, Kobayashi H, Smith RL, Ward J, Ward C, Milatovic D. 2001. Pharmacokinetics and absolute bioavailability of sitafloxacin, a new fluoroquinolone antibiotic, in healthy male and female Caucasian subjects. *Xenobiotica* 31:811–822.
- Ohdachi S, Ishibashi Y, Iwasa MA, Fukui D, Saitoh T. 2015. Chapters Primates; Rodentia; Carnivora; Artiodactyla. In: *The wild mammals of Japan*, 2nd Ed., Ohdachi S, Ishibashi Y, Iwasa MA, Saitoh T, editors. Shoukadoh Book Sellers, Kyoto, Japan, pp. 128–296.
- Pang H, Jiang Y, Wan K. 2017. Drug susceptibility testing of 33 reference strains of slowly growing mycobacteria to 19 antimicrobial agents. *Biomed Res Int* 2017:1584658.
- Pang H, Li G, Zhao X, Liu H, Wan K, Yu P. 2015. Drug susceptibility testing of 31 antimicrobial agents on rapidly growing mycobacteria isolates from China. *Biomed Res Int* 2015:419392.
- Pate M, Zajc U, Kušar D, Žele D, Vengušt G, Pirš T, Ocepek M. 2016. *Mycobacterium* spp. in wild game in Slovenia. *Vet J* 208:93–95.
- Patino W LC, Monge O, Suzán G, Gutiérrez-Espeleta G, Chaves A. 2018. Molecular detection of *Mycobacterium avium* and *Mycobacterium genavense* in feces of free-living scarlet macaws (*Ara macao*) in Costa Rica. *J Wildl Dis* 54:357–361.
- Pavlik I, Falkinham J, Kazda J. 2009. Transmission of mycobacteria from the environment to susceptible hosts. In: *The ecology of mycobacteria: Impact on animal's and human's health*, Kazda J, Pavlik I, Falkinham JO III, Hruska K, editors. Springer Science and Business Media, Berlin, Germany, pp. 283–312.

- Poonawala H, Piscitelli VE, Ladutko L, Campbell S. 2017. Misidentification of *Mycobacterium paraense* as *Mycobacterium avium* complex by Accuprobe. *J Clin Microbiol* 55:2283–2284.
- Primm TP, Lucero CA, Falkinham JO III. 2004. Health impacts of environmental mycobacteria. *Clin Microbiol Rev* 17:98–106.
- Ronai Z, Eszterbauer E, Csivincsik A, Guti CF, Dencsó L, Janosi S, Dán Á. 2016. Detection of wide genetic diversity and several novel strains among non-avian nontuberculous mycobacteria isolated from farmed and wild animals in Hungary. *J Appl Microbiol* 12:41–54.
- Saitou N, Nei M. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425.
- Sattar A, Zakaria Z, Abu J, Aziz SA, Gabriel RP. 2018. Evaluation of six decontamination procedures for isolation of *Mycobacterium avium* complex from avian feces. *PLoS One* 13:e0202034.
- Schinsky MF, McNeil MM, Whitney AM, Steigerwalt AG, Lasker BA, Floyd MM, Hogg GG, Brenner DJ, Brown JM. 2000. *Mycobacterium septicum* sp. nov., a new rapidly growing species associated with catheter-related bacteraemia. *Int J Syst Evol Microbiol* 50: 575–581.
- Telenti A, Marchesi F, Balz M, Bally F, Böttger EC, Bodmer T. 1993. Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *J Clin Microbiol* 31: 175–178.
- Tsukamura M, Mizuno S, Toyama H. 1985. *Mycobacterium thermoresistibile* from dust of Japanese houses. *Microbiol Immunol* 29:467–473.
- van Ingen J, Boeree MJ, Van Soolingen D, Mouton JW. 2012. Resistance mechanisms and drug susceptibility testing of nontuberculous mycobacteria. *Drug Resist Updat* 15:149–161.
- Vasireddy R, Vasireddy S, Brown-Elliott BA, Wengenack NL, Eke UA, Benwill JL, Turenne C, Wallace RJ Jr. 2017. *Mycobacterium arupense*, *Mycobacterium heraklionense*, and a newly proposed species, “*Mycobacterium virginienne*” sp. nov., but not *Mycobacterium nonchromogenicum*, as species of the *Mycobacterium terrae* complex causing tenosynovitis and osteomyelitis. *J Clin Microbiol* 54:1340–1351.
- Vishkautsan P, Reagan KL, Keel MK, Sykes JE. 2016. Mycobacterial panniculitis caused by *Mycobacterium thermoresistibile* in a cat. *JFMS Open Rep* 2: 2055116916672786.
- World Health Organization. 2008. *Policy guidance on drug-susceptibility testing (DST) of second-line anti-tuberculosis drugs*. World Health Organization, Geneva, Switzerland, 20 pp.

Submitted for publication 23 October 2019.

Accepted 25 February 2020.