

## **Authors' Reply to “Host Taxonomy is Critical in Zoonotic Disease Surveillance and Reporting”**

Source: Journal of Wildlife Diseases, 60(2) : 555-557

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/JWD-D-23-00178b>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Authors' Reply to "Host Taxonomy is Critical in Zoonotic Disease Surveillance and Reporting"

Dear Editor,

We appreciate the feedback from Olson and Juman (2024) on our manuscript (Hareza et al. 2023). After conducting a thorough review of the original surveillance data, we acknowledge that both rabid marmots in question were indeed identified in states along the East Coast, and we have supplied a corrected table to address this.

While we concur with the significance of precise taxonomic nomenclature, it is imperative to note certain nuances regarding the functioning of the United States National Rabies Surveillance System (NRSS), which may not have been fully appreciated in the Letter by Olson and Juman. The NRSS compiles data from over 130 rabies laboratories across the United States, representing 54 reporting jurisdictions (Ma et al. 2022). These data are received in unstandardized formats and are collated by the Centers for Disease Control and Prevention (CDC). CDC is often unable to impute missing data elements (like species) or verify the taxonomic naming that reporting jurisdictions have provided.

It is pertinent to highlight that the Council for State and Territorial Epidemiologists (CSTE) requests states to submit species-level information for any animal tested for rabies. However,

the availability of such information, particularly for bats, is not consistently ensured. This issue is described in various citations included in our original article, namely, "Rabies Surveillance in the United States" and the CSTE Position Statement (CSTE 2011; Ma et al. 2022).

Addressing the concerns raised regarding De Benedictis et al (De Benedictis et al. 2022), we find it necessary to emphasize the challenges associated with implementing costly technologies within large surveillance systems or publicly funded initiatives. Factoring in all associated expenses of DNA barcoding assays, inclusion of this as a routine test in a rabies diagnostic lab would likely cost \$75 USD per sample. The NRSS laboratory network tests nearly 100,000 animals annually (CDC 2011) which would add \$7.5 million in public funding if the suggested cytochrome B testing were to be adopted. It is crucial to contextualize this against the backdrop of the current cost for rabies diagnostic services across the 130 US rabies laboratories participating in the NRSS, which is estimated at \$6 million USD (ADHS 2017). Selective DNA barcoding technology may have a role, but we must remember that the NRSS operates to prevent rabies infections

TABLE 1. Rabies in rodents and lagomorphs in the USA and Puerto Rico, 2011–2020.

Rodent or Lagomorph	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Total positive	Total tested	Percent positive
Groundhog ( <i>Marmota monax</i> )	45	42	37	43	25	44	33	23	38	38	368	9,084	4.1%
Beaver ( <i>Castor canadensis</i> )	3	4	0	2	2	5	0	4	1	0	21	283	7.4%
Rabbit (Family Leporidae)	0	0	1	0	7	0	0	0	0	0	8	1,364	0.6%
<i>Marmota</i> spp. <sup>a</sup>	0	0	2	0	0	0	0	0	0	0	2	109	1.8%
Squirrel (Family Sciuridae)	0	0	0	0	1	0	0	0	1	0	2	6,195	0.0%
Rat ( <i>Rattus</i> spp.)	0	0	0	0	0	0	0	0	0	0	0	1,052	0.0%
House mouse ( <i>Mus musculus</i> )	0	0	0	0	0	0	0	0	0	0	0	840	0.0%
Muskrat ( <i>Onychia zibethicus</i> )	0	0	0	0	0	0	0	0	0	0	0	835	0.0%
Chipmunk ( <i>Tamias striatus</i> )	0	0	0	0	0	0	0	0	0	0	0	658	0.0%
All others <sup>b</sup>	0	0	0	0	0	0	0	0	0	0	0	1,505	0.0%
<b>Grand total</b>	<b>48</b>	<b>46</b>	<b>40</b>	<b>45</b>	<b>35</b>	<b>49</b>	<b>33</b>	<b>27</b>	<b>40</b>	<b>38</b>	<b>401</b>	<b>21,925</b>	<b>1.8%</b>
All tested	2,516	2,504	2,243	2,309	2,273	2,168	2,090	1,904	1,828	1,825	21,925		
Percent positive	1.9%	1.8%	1.8%	1.9%	1.5%	2.3%	1.6%	1.3%	2.1%	2.1%	1.8%		

<sup>a</sup> Includes animals recorded only as 'Marmota' or 'Marmota' sp., which may have included *Marmota monax* individuals.  
<sup>b</sup> All others (small mammals not rodents or lagomorphs): Moles (e.g., Subfamily Scalopinidae) and shrews (Family Soricidae); note that surveillance records are usually applied with common names only.

TABLE 2. Rabies percent positivity in rodents, lagomorphs and raccoons in the USA and Puerto Rico from 1985 to 2020.

Timespan	Raccoon ( <i>Procyon lotor</i> )			Groundhog ( <i>Marmota monax</i> )			Beaver ( <i>Castor canadensis</i> )			Rabbit (Family Leporidae)			Other rodents and lagomorphs <sup>b</sup>		
	Tested	Positive	% Pos	Tested	Positive	% Pos	Tested	Positive	% Pos	Tested	Positive	% Pos	Tested	Positive	% Pos
1985–1994	122,394	27,284	22.29%				206	12	5.82%	3,380	17	0.50%	52,266	19	0.04%
1995–2010	202,426	46,637	23.04%	14,051	663	4.72%	551	31	5.63%	5,502	25	0.45%	53,571	16	0.03%
2011–2020	123,704	16,172	13.07%	9,084	368	4.05%	283	21	7.42%	1,364	8	0.58%	11,194	4	0.04%
Total	448,524	90,093	20.09%	29,120	1,358	4.66%	1,040	64	6.15%	10,246	50	0.49%	117,031	39	0.03%

<sup>a</sup> Other rodents and lagomorphs: Marmot (*Marmota* spp.), squirrel (*Family Sciuridae*), rat (*Rattus* sp.), house mouse (*Mus musculus*), muskrat (*Onychia zibethicus*), chipmunk (*Tamias striatus*), all other.  
<sup>b</sup> All rodents and lagomorphs: Groundhog, beaver, rabbit, marmot, squirrel, rat, mouse, muskrat, chipmunk, all other.

in people and animals, epidemiologic studies such as this are conducted opportunistically, taking advantage of the data generated by this large and routine public health system. Until truly low cost and accurate means of taxonomic identification are available, large-scale, publicly funded surveillance systems will struggle to implement the suggestions provided by Olson and Juman.

In light of the aforementioned challenges and constraints, we propose a pragmatic solution by correcting the table to reflect “*Marmota* sp.” to prevent any future misinterpretations. Additionally, we recommend replacing the specific taxonomies for rabbit and squirrel with more general taxonomic names, as indicated in the corrected table. Furthermore, we would like to clarify that discrepancies raised by Olson and Juman are due to differences in data included in our analysis (we did not include guinea pigs or chinchilla in our study, for example, so we did not include these animals from previous publications in our analyses).

While we acknowledge the significance of addressing errors and ensuring accuracy, we assert that the issues raised by Olson and Juman primarily pertain to a straightforward erratum in Table 1 (please see Table 2 for corrections). The challenges associated with working with a vast surveillance database and the limited accessibility of accurate taxonomic identification tools in public health laboratories contributed to the issue raised by Olson and Juman.

**Dariusz A. Hareza, Ricky Langley, Xiaoyue Ma, Ryan Wallace, and Charles E. Rupprecht**

*Disclaimer: The findings and conclusions in this letter are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.*

#### LITERATURE CITED

- Arizona Department of Health Services (ADHS). 2017. *Manual for Rabies Control and Bite Management*. <https://www.pinal.gov/DocumentCenter/View/2102/Compendium-of-Animal-Rabies-Prevention-and-Control-PDF?bidId=>. Accessed January 2024.
- Centers for Disease Control and Prevention (CDC). 2011. *Diagnosis in animals and humans*. <https://www.cdc.gov/rabies/diagnosis/animals-humans.html>. Accessed January 2024.
- Council of State and Territorial Epidemiologists (CSTE). 2009. *Public health reporting and national notification for animal rabies*. <https://cdn.ymaws.com/www.cste.org/resource/resmgr/PS/09-ID-12.pdf>. Accessed January 2024.
- De Benedictis P, Leopardi S, Markotter W, Velasco-Villa A. 2022. The importance of accurate host species identification in the framework of rabies surveillance, control and elimination. *Viruses* 14:492.
- Hareza DA, Langley R, Ma X, Wallace R, Rupprecht CE. 2023. Rabies in rodents and lagomorphs in the USA, 2011–20. *J Wildl Dis* 59:734–742.
- Ma X, Bonaparte S, Toro M, Orciari LA, Gigante CM, Kirby JD, Chipman RB, Fehlner-Gardiner C, Gutiérrez Cedillo V, et al. 2022. Rabies surveillance in the United States during 2020. *J Am Vet Med Assoc* 260:1157–1165.
- Olson LE, Juman MM. 2024. Host taxonomy is critical to zoonotic disease remorting. *J Wildl Dis* 60:554–555.