

Long-Term Trends in Vegetation on Bureau of Land Management Rangelands in the Western United States★

Authors: Kleinhesselink, Andrew R., Kachergis, Emily J., McCord, Sarah E., Shirley, Justin, Hupp, Nicole R., et al.

Source: Rangeland Ecology and Management, 87(1) : 1-12

Published By: Society for Range Management

URL: <https://doi.org/10.1016/j.rama.2022.11.004>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Original Research

Long-Term Trends in Vegetation on Bureau of Land Management Rangelands in the Western United States[☆]

Andrew R. Kleinhesselink^{1,*}, Emily J. Kachergis², Sarah E. McCord³, Justin Shirley⁴, Nicole R. Hupp⁵, Jennifer Walker⁵, John C. Carlson^{5,8}, Scott L. Morford¹, Matthew O. Jones^{1,7}, Joseph T. Smith¹, Brady W. Allred⁶, David E. Naugle⁶

¹ Numerical Terradynamic Simulation Group, University of Montana, Missoula, MT 59812, USA

² National Assessment, Inventory, and Monitoring Coordinator, Bureau of Land Management, Lakewood, CO 80215, USA

³ Jornada Experimental Range, US Department of Agriculture–Agricultural Research Service, Las Cruces, NM 88003, USA

⁴ Rangeland Management Specialist, Bureau of Land Management HQ 220, Boise, ID 83705, USA

⁵ Montana-Dakotas State Office, Bureau of Land Management, Billings, MT 59101, USA

⁶ University of Montana, W. A. Franke College of Forestry and Conservation, Missoula, MT 59812, USA

⁷ Regrow Agriculture, Durham, NH 03842, USA

⁸ US Fish and Wildlife Service Mountain/Prairie Region (R6), Billings, MT 59105, USA



ARTICLE INFO

Article history:

Received 11 July 2022

Revised 8 November 2022

Accepted 15 November 2022

Keywords:

Exotic annual grasses

Monitoring

Public lands

Remote sensing

Woodland expansion

ABSTRACT

The US Bureau of Land Management (BLM) manages nearly 1 million km² of public lands that support recreation, livestock production, and wildlife habitat. Monitoring the condition of vegetation on these lands is crucial for sound management but has historically been difficult to do at scale. Here we used newly developed remote-sensing tools to conduct an unprecedented assessment of trends in vegetation cover and production for all BLM rangelands from 1991 to 2020. We found widespread increases in cover and production of annual grasses and forbs, declines in herbaceous perennial cover, and expansion of trees. Cover and production of annual plants now exceed that of perennials on > 21 million ha of BLM rangeland, marking a fundamental shift in the ecology of these lands. This trend was most dramatic in the Western Cold Desert of Nevada and parts of surrounding states where aboveground production of annuals has more than tripled. Trends in annuals were negatively correlated with trends in bare ground but not with trends in perennials, suggesting that annuals are filling in bare ground rather than displacing perennials. Tree cover increased in half of ecoregions affecting some 44 million ha and underscoring the threat of woodland expansion for western rangelands. A multiscale variance partitioning analysis found that trends often varied the most at the finest spatial scale. This result reinforces the need to combine plot-level field data with moderate-resolution remote sensing to accurately quantify vegetation changes in heterogeneous rangelands. The long-term changes in vegetation on public rangelands argue for a more hands-on approach to management, emphasizing preventative treatment and restoration to preserve rangeland habitat and functioning. Our work shows the power of new remote-sensing tools for monitoring public rangelands and developing effective strategies for adaptive management and conservation.

© 2022 The Authors. Published by Elsevier Inc. on behalf of The Society for Range Management. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Introduction

Rangelands cover half of earth's terrestrial surface and provide wildlife habitat, areas for recreation, food production, carbon storage, flood control, and water purification (Havstad et al. 2007). In the western United States, the Bureau of Land Management (BLM) oversees > 945 000 km² of land (Dombeck 1996), of which most

[☆] Funding for this project was provided by the Montana-Dakotas State Office of the Bureau of Land Management. Working Lands for Wildlife, part of the Natural Resources Conservation Service, provided financial support to David E. Naugle as their National Science Advisor during this research. Findings and conclusions in this article are those of the authors and do not necessarily represent the views of the US Fish and Wildlife Service, US Department of Agriculture (USDA)—Agricultural Research Service, or the Bureau of Land Management. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the US government.

* Correspondence: Andrew R. Kleinhesselink, NTSG, 32 Campus Dr, University of Montana, Missoula, MT 59812, USA
<https://doi.org/10.1016/j.rama.2022.11.004>
1550-7424/© 2022 The Authors. Published by Elsevier Inc. on behalf of The Society for Range Management. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

E-mail address: andy.kleinhesselink@umont.edu (A.R. Kleinhesselink).

Table 1
Summary of allotment data used in the analysis. Number of Bureau of Land Management districts, field offices and grazing allotments, total allotment area, average allotment area, and percent land ownership within allotments is given for each ecoregion. Nonrangeland area masked out before analysis is given in % Masked.

Ecoregion	Districts	Field offices	Allotments	Total area (10 ⁶ ha)	Avg. area (ha)	% BLM	% Private	% Other	% Masked
AZ/NM Highlands	7	12	562	2.50	4,446	40	32	27.5	1.3
E Cold Deserts	20	37	4,574	23.20	5,072	69	19	10.4	0.8
Forested Mts	32	67	5,086	8.26	1,623	55	33	10.0	1.3
Marine West Coast Forest	3	4	10	0.01	521	89	5	0.1	6.2
Mediterranean California	3	7	232	0.43	1,842	54	40	0.4	5.8
N Great Plains	5	14	5,066	11.82	2,334	27	60	7.2	6.5
S Great Plains	6	9	839	2.51	2,990	29	55	15.6	1.1
W Cold Deserts	19	38	3,471	34.74	10,008	83	12	3.5	1.3
Warm Deserts	12	21	1,171	10.91	9,315	63	15	19.2	2.4
Total	43*	109*	21,011	94.37	4,491	65	24	9.0	2.0

*Total number is less than the sum of values in the table as the same field office or district may have allotments in more than one ecoregion.




Figure 1. Map of Bureau of Land Management (BLM) rangelands included in the analysis. Filled regions show BLM allotments colored by ecoregion. Lines display state and field office boundaries.

is rangeland. The management of BLM lands is divided among 10 state-level offices, 120 regional field offices excluding Alaska and the eastern United States, and > 21 000 local grazing allotments (Table 1; Fig. 1). Rangelands across western North America are being altered by tree expansion (Ratajczak et al. 2012; Nackley et al. 2017), annual plant invasions (Coates et al. 2016), increased wild-

fires (Li et al. 2021), and climate change (McIntosh et al. 2019; Brookshire et al. 2020); understanding these changes calls for a multiscale assessment of vegetation trends across all BLM rangelands.

Vegetation monitoring provides data essential to understanding the causes and consequences of environmental change (McCord




Figure 2. Three spatial scales of vegetation trend analysis. **A**, Each ecoregion (e.g., Western Cold Deserts) contains many Bureau of Land Management (BLM) field offices, boundaries shown with thin gray lines. **B**, A single field office (blue outline in A) contains dozens to hundreds of grazing allotments, boundaries shown with white lines. **C**, A single grazing allotment (blue outline in B) with 3 yr of Rangeland Analysis Platform (RAP) cover data. Each BLM grazing allotment contains many thousands of individual 30-m RAP cover and production pixels.

and Pilliod 2022) and has long been central to sustainable grazing management and restoration (West 2003; Jones et al. 2020; Kachergis et al. 2022). Monitoring requires the repeated collection of vegetation data at the same locations so that trends, directional changes in vegetation over time, can be assessed (West 2003; Toevs et al. 2011b). Quantifying vegetation trends on BLM rangelands is often a prerequisite for developing and implementing resource management plans that meet the agency's multiple use and sustained yield mandates under the Federal Land Policy and Management Act (Federal Land Policy and Management Act 1976). Collecting sufficient data to quantify vegetation trends across BLM rangelands is inherently a challenge due to their heterogeneity and enormous area. Indeed, during the latter half of the 20th century, the adequacy of BLM vegetation data became a point of contention among grazing permittees, environmental advocacy groups, and government staff (Fernandez-Gimenez et al. 2005) and federal audits repeatedly recommended that the BLM improve and expand monitoring (GAO 1992). This issue persisted into recent years, when a review found that monitoring data for many BLM allotments were missing or incomplete (Veblen et al. 2014).

Recognizing the need for more consistent vegetation data, in 2011 the BLM began an ambitious data collection program for its lands: the Assessment, Inventory and Monitoring (AIM) strategy (Toevs et al. 2011a; Kachergis et al. 2022). The AIM strategy defined a standard set of monitoring protocols and a rigorous approach for rangeland monitoring so that the data being collected could inform decision making across local, regional, and national scales (Toevs et al. 2011b; Taylor et al. 2014). Since implementation, nearly 50 000 plots have been monitored using standardized AIM field methods. These data have expanded the capacity for scientists and rangeland managers to evaluate the effectiveness of land management plans, identify critical habitat, and monitor the effects of global change (Kachergis et al. 2022). Combined with the Natural Resources Conservation Service's National Resource Inventory (NRI), which applies the same standardized methods to private lands, data on the status of rangeland vegetation are now available at > 80 000 locations across the western United States (Spaeth et al. 2003).

One of AIM's principal goals is to collect field data that can inform the development of new remote-sensing (RS) technology (Boyte et al. 2019; Jones et al. 2020; Allred et al. 2021; Rigge et al. 2021). Field data from AIM and NRI have been instrumental in training and validating RS algorithms that translate moderate resolution satellite reflectance data into estimates of rangeland cover and production (Allred et al. 2021; Rigge et al. 2021). New RS products complement field-based monitoring with advantages that overcome some of the limitations inherent to field-based data collection (West and Wu 2003). In particular, RS data provide new opportunities to measure vegetation trends at the most appropriate spatial scales for management (Kachergis et al. 2022): at the ecoregion scale (Fig. 2A), assessing vegetation trends across BLM lands plays an important role in planning biome-scale conservation; at the field office scale (see Fig. 2B), comparing trends between local units of land management such as grazing allotments can help identify areas for restoration; and at the local scale (see Fig. 2C), BLM field staff may examine trends in cover within a single pasture, grazing allotment, or project boundary to understand patterns of grazing pressure, annual grass invasion, woody plant encroachment, or postfire recovery (Toevs et al. 2011a). Moreover, RS data going back to the 1980s may be the only source of long-term vegetation data for BLM allotments with limited historical field sampling.

Here we use recently developed 30-m RS products to measure trends in vegetation cover and aboveground production on BLM rangelands between 1991 and 2020. Our analysis of vegetation trends is organized around four spatial scales relevant to the ecology and management of BLM rangelands: ecoregions, field offices, individual grazing allotments, and 30-m remote sensing pixels (see Fig. 2). Our objectives were to 1) measure trends in vegetation cover and rangeland production at the ecoregion scale; 2) quantify variance in trends within ecoregions at the field office, allotment, and pixel scales; 3) investigate correlations between trends at each scale; and 4) discuss the implications of our results for the management of BLM rangelands.

Methods

Spatial data collection and processing

We downloaded publicly available boundary data for all BLM grazing allotments, BLM field offices, and EPA ecoregions from government websites (Appendix A, Table A1). We used geoprocessing tools in R and QGIS to remove duplicate allotment boundaries and removed allotments < 1 ha in area (R Core Team 2022). We assigned each allotment to one of nine ecoregions derived from the EPA ecoregion levels I–III: AZ/NM Highlands, Marine West Coast Forests, Eastern Cold Deserts, Forested Mountains, Northern Great Plains, Southern Great Plains, Mediterranean California, Warm Deserts, and Western Cold Deserts (see Fig. 1; Table A2). We excluded allotments from the Marine West Coast Forest ecoregion from further analyses due to small sample size and area (see Table 1).

We use BLM grazing allotments as a practical unit for summarizing local-scale rangeland trends in our analysis. While allotment boundaries are delineated by the BLM grazing program, the public lands within them are open to and managed for multiple uses. We do not evaluate the effects of livestock grazing or any other specific land use. Allotments often include private rangelands surrounding BLM parcels, and we used data from the US Geological Survey Protected Areas Database (PAD-US) to find the area of BLM and private lands within each allotment. Allotments averaged 4 491 ha in area and comprised 65% BLM, 24% private, and 9% other public lands (see Table 1). We retained private lands within allotment boundaries in our analysis so that trends are representative of these local administrative units. After processing, the allotment data included 21 011 grazing allotments totaling 94.37 10⁶ ha of rangeland (see Table 1).

We used estimates of vegetation cover and aboveground herbage production from the Rangeland Analysis Platform (RAP). The RAP uses machine learning to estimate cover of annual grasses and forbs (annuals), herbaceous perennial grasses and forbs (perennials), shrubs, trees, and bare ground, as well as aboveground production of annuals and perennials in kg ha⁻¹ on US rangelands from 30-m RS data (Allred et al. 2021; Jones et al. 2021). As described in Jones et al. (2018), RAP cover estimates correspond to first-hit cover as measured by the AIM and NRI protocols and species are grouped according to characteristics in the US Department of Agriculture (USDA) plants database (USDA, NRCS 2018). In the remainder of the paper, we refer to RAP cover and production categories (i.e., annuals, perennials, shrubs, trees, and bare ground) as vegetation functional groups. We masked out nonrangelands and areas of hay, alfalfa, and idle cropland within allotments using the National Land Cover Database (Homer et al. 2020) and the USDA National Agricultural Statistics Service Cropland Data Layer (USDA NASS 2016). After masking, we calculated allotment-level average cover and production using the RAP v.3.0 dataset for each yr from 1986 to 2020. Masking and allotment-level vegetation averages were calculated in Google Earth Engine (Gorelick et al. 2017).

Modeling long-term vegetation trends

We analyzed long-term trends in vegetation cover and production using separate linear mixed-effects models for each functional group. We fitted fixed effects for ecoregion and time (calendar year) and their interaction (year by ecoregion). Mixed-effects models were fitted with the lme4 package (Bates et al. 2015) using the following model formula in lme4 notation:

$$y \sim E + t + E:t + (t|E:O) + (t|A), \quad (1)$$

where y is yearly allotment-level cover or production, E is ecoregion, t is year, O is BLM field office, and A is allotment. We included random intercepts and slopes at the level of allotment and BLM field office nested within the ecoregion. For each model, we included 30 yr of data from 1991 to 2020. We log-transformed cover and production values before analysis. To avoid log-transforming values of zero, we excluded allotments that for any year had average cover of < 0.25%; in the herbaceous production models, we excluded allotments that for any year had production of < 0.25 kg ha⁻¹. This resulted in dropping several hundred allotments from most analyses and several thousand from the annual cover and production and tree cover models (see Table A3–A9). Adding a small increment to cover and production values before log-transformation and including all allotments in the analyses resulted in similar trend estimates. To avoid identifiability problems, we excluded a handful of allotments from field offices that had only a single allotment. After fitting models, we extracted trend estimates and 95% confidence intervals for each ecoregion using the “emmeans” package in R (Lenth 2022). We interpret trends as significant when 95% confidence intervals for an ecoregional trend do not overlap zero.

Spatial heterogeneity in vegetation trends across scales

To quantify spatial variation in vegetation trends across scales, we fit separate linear-mixed effects models for cover and production of each functional group as described in the *Modeling long-term vegetation trends* section but with an additional term to capture variance in trends at the pixel scale:

$$y \sim E + t + E:t + (t|E:O) + (t|A) + (t|P), \quad (2)$$

where P identifies individual pixels as a grouping factor. Fitting this model to all pixels within all BLM allotments was prohibitively slow, so we used a subset of data. We randomly selected 2069 allotments (~10% of total) and extracted yearly cover and production values from eight randomly selected 30-m pixels within each ($n = 16\,541$). As in the allotment-level analysis, cover and production values were log-transformed before analysis, and we excluded pixels that had fewer than 20 yr of nonzero cover and production data. After fitting the models, we extracted group-level variance estimates for vegetation trends (slopes) at the field office, allotment, and pixel-scales.

Correlations between vegetation trends across scales

At each of the four spatial scales, we examined correlations between cover trends for annuals, herbaceous perennials, bare ground, and shrubs. Trend estimates for each ecoregion, field office, and allotment were drawn from models fitted in the *Modeling long-term vegetation trends* section. Trend estimates for pixels were drawn from the models fitted in the *Spatial heterogeneity in vegetation trends across scales* section. We used random effects slopes (Best Linear Unbiased Predictions) as trend estimates for individual field offices, allotments, and pixels. Since random effects were nested, (pixels in allotments, allotments in field offices and field offices in ecoregions), vegetation trend estimates at each scale represent the residual trend not captured by trends at larger spatial scales.

Supplemental analyses of annual invasion and woodland expansion

We performed three post-hoc analyses to further investigate trends in annual plants and trees. First, we quantified the number and proportion of allotments with average cover and production of annuals greater than that of perennials for each ecoregion

for each decade. Second, we conducted a follow-up nonparametric test of tree cover trends among all allotments (including those below the 0.25% cover threshold used in the *Modeling long-term vegetation trends* section). We report the number and proportion of allotments in each ecoregion with positive and negative trends in tree cover as determined by the Mann-Kendall test for significant trends ($P < 0.05$). Thirdly, we used field data from the AIM program to identify which annual and tree taxa were most common in each ecoregion. This does not directly reveal which species are responsible for trends but can provide context for interpreting RS trends in overall annual and tree cover.

Results

Status and trends of vegetation cover and production by ecoregion

Herbaceous perennial vegetation dominated allotments in the AZ/NM Highlands, Northern Great Plains, Southern Great Plains, Forested Mountains, and Western Cold Deserts (Fig. 3). Bare ground cover was dominant or codominant in Eastern Cold Deserts and Warm Deserts, while annuals dominated Mediterranean California. Average shrub cover of allotments was $> 10\%$ in all ecoregions except for the Northern Great Plains. Average tree cover was below 10% in all ecoregions except for the Forested Mountains ($\sim 15\%$). Average herbaceous perennial cover and aboveground production were generally several times that of annuals in all ecoregions ex-

cept Mediterranean California, where annuals were dominant (see Figs. 3 and 4).

We found positive trends in annual cover and aboveground production in all ecoregions (see Figs. 3 and 4). In contrast, there were declining trends in perennial cover in all ecoregions except the Northern Great Plains—with the most rapid declines in Mediterranean California and Warm Deserts (see Fig. 3). Perennial production trends were negative in the Eastern Cold Deserts, Forested Mountains, Mediterranean California, Western Cold Deserts, and Warm Deserts (see Fig. 4). Bare ground cover was declining in four ecoregions, most rapidly in the Northern Great Plains and Western Cold Deserts, but has increased slightly in Mediterranean California (see Fig. 3). Woody shrub cover declined in the Western Cold Deserts but increased in the Northern Great Plains and Mediterranean California (see Fig. 3). Since we modeled log-transformed cover and aboveground production, the trend coefficients (slopes) shown in Figure 5 can be interpreted as annual proportional rates of change. For example, a 0.028 rate of increase for annual grass cover in the Western Cold Deserts (see Fig. 5) indicates an increase of 2.8% per yr, equivalent to a doubling of cover in 25 yr.

The number of allotments with annual cover and production greater than herbaceous perennials has increased over time in the AZ/NM Highlands, Eastern Cold Deserts, Forested Mountains, Mediterranean California, Western Cold Deserts, and Warm Deserts (see Tables A10, A11). And across all ecoregions, the proportion of allotments with annual cover surpassing that of perennials increased from 4.9% ($n = 1\,025$) in the 1990s to 11% ($n = 2\,320$) dur-




Figure 3. Mean vegetation cover of Bureau of Land Management allotments from 1991 to 2020. Each column shows a separate ecoregion and each row a separate functional group. Colored ribbons show interquartile range of allotment-level average cover. Trend lines from the mixed effects models are shown where significant. Y-axis is on a log scale.




Figure 4. Mean aboveground herbaceous production of Bureau of Land Management allotments from 1991 to 2020. Each panel shows a separate ecoregion. Colored ribbons show interquartile range of allotment-level average aboveground production for annuals and herbaceous perennials separately. Trend lines from the mixed effects models are shown where significant. Y-axis is on a log scale.

ing the past decade. This shift to annual dominance was most dramatic in the Western Cold Deserts of Nevada and surrounding states, where more than a third of allotments ($n = 1\,277$), representing some 18.2 million ha, now have more annual than perennial cover. Likewise, nearly one quarter of allotments in the Western Cold Deserts now have more production from annuals than perennials (see Table A11). The AIM data from 31 908 plots located within 6 744 allotments showed that exotic bromes such as cheatgrass (*Bromus tectorum*) and Japanese brome (*Bromus japonicus/arvensis*) were common in many ecoregions (see Table A12). For example, in the Western Cold Deserts, Northern Great Plains, and Eastern Cold Deserts, exotic annual bromes were found in 94%, 85%, and 65% of allotments, respectively. Native annuals were dominant in three ecoregions: the AZ/NM Highlands, Southern Great Plains, and Warm Deserts.

The average tree cover of allotments increased in the Eastern Cold Deserts, Forested Mountains, Northern Great Plains, and Western Cold Deserts (see Fig. 3). In the Northern Great Plains, tree cover increased at a rate of 2.2% per year, nearly doubling from 1.8% cover in the early 1990s to 3.5% cover in the past few years (see Fig. 5). Nonparametric tests showed that 52% of all allotments had increases in tree cover and only 5% had decreases (see Table A13). In the Northern Great Plains and Forested Mountains, tree cover increased in 65% and 60% of allotments, respectively. Notably, Mediterranean California was the only region where there were more declines than increases in tree cover. Field data from AIM showed that native conifers were the most common trees in many ecoregions (see Table A14). Junipers (e.g., *Juniperus monosperma*, *J. osteosperma*, *J. occidentalis*, *J. scopulorum*) were found on 30–50% of allotments in the AZ/NM Highlands, Eastern Cold Deserts, Forested Mountains, Southern Great Plains, and Western Cold Deserts and on nearly one fifth of allotments in the Northern Great Plains. Mesquite species (*Prosopis glandulosa*, *P. veletuna*) were also prevalent in the AZ/NM Highlands, Warm Deserts, and Southern Great Plains.

Spatial heterogeneity in vegetation trends at field office, allotment, and pixel-scales

Average vegetation trends reported for each ecoregion bely considerable variation in trends at field office, allotment, and pixel scales (Fig. 6). Except for annuals, trends varied more between pixels within the same allotment than between allotments in the same field office, or between field offices in the same ecoregion. Trend variance was especially pronounced for tree cover trends, and 59% of group-level variance occurred at the pixel scale. Trends for annual cover and production were unique in having proportion-

ally more variance at the allotment scale and less at the pixel scale. Compared with cover trends, production trends showed a higher proportion of variance at the field office scale.

Correlations between vegetation trends across scales

We further examined correlations between trends for different functional groups (Fig. 7). Annual cover trends were negatively correlated with bare ground trends at the ecoregion, field office, allotment, and pixel scales, with the strongest correlations at the ecoregion and field office scales (see Fig. 7A–D). Annual and herbaceous perennial trends were positively correlated at the field office, allotment, and pixel scales (see Fig. 7E–H). Annual trends were weakly negatively correlated with shrub trends at the allotment and pixel scales (see Fig. A1A). Bare ground trends were negatively correlated with perennial trends and positively correlated with shrub trends at subecoregion scales (Fig. A1B and C). The correlation between shrub and perennial trends was negative at the field office scale but switched to positive at the pixel scale (Fig. A1D).

Discussion

New RS tools enabled us to conduct an unprecedented multiscale assessment of trends in vegetation cover and production on all rangeland grazing allotments administered by the BLM—encompassing > 10% of the contiguous United States. Trends revealed striking increases in cover and production of annual grasses and forbs and widespread expansion of trees. Emblematic of this rapid invasion was a doubling of allotments in which annual cover and production now exceed that of herbaceous perennials (Tables A10 and A11). Correlations between trends provide inference that increases in annual cover have driven a loss of bare ground in several ecoregions, with annuals invading the interstitial areas between larger perennials and shrubs (see Fig. 7A). Tree cover also rapidly expanded in many ecoregions, particularly in the Northern Great Plains, where it will increase from 3.5% to 6.8% by 2050 if current trends continue (see Figs. 3 and 5). The preponderance of variance in trends at the suballotment scale reinforces the idea that rangelands are defined by their heterogeneity in space and time (see Fig. 6). Remotely sensed vegetation data provide vital information for understanding the threats facing BLM rangelands and designing effective management strategies to confront them.

Trends in vegetation cover and production by ecoregion

Our BLM-specific analyses reinforce the vexing challenge of managing annual plant invasions on western rangelands (*Maestas*




Figure 5. Vegetation cover and aboveground production trends for Bureau of Land Management allotments from 1991 to 2020 estimated by mixed-effects models. Trend magnitudes are shown on the x-axis surrounded by 95% confidence intervals. Trends represent annual rate of change in log cover (left) or log aboveground production (right).

et al. 2022; Tarbox et al., 2022). The continued increase in annual cover and loss of bare ground is evidence that rangelands are accumulating fine fuels that can further increase wildfire (Davies et al. 2012; Smith et al. 2022a). Exacerbating frequency and severity of wildfires can devastate ecosystems by removing keystone plants (e.g., big sagebrush *Artemisia tridentata*) and imperiling endemic wildlife such as greater-sage grouse (*Centrocercus urophasianus*; Coates et al. 2016). Even though exotic annuals began invading western rangelands over a century ago (Mack 1981), their continued spread shows little evidence of slowing (Smith et al. 2022b).

While annuals are not the dominant vegetation group in most ecoregions, a growing number of allotments now have annual cover and production exceeding that of perennials. Today, annuals are the dominant herbaceous component on 1 691 allotments en-

compassing 21 million ha of rangeland (Tables A10 and A11). The most dramatic rise in annuals was in the Western Cold Deserts of Nevada and parts of adjoining states where annual aboveground production more than tripled from roughly 90 kg ha^{-1} in the early 1990s to $> 300 \text{ kg ha}^{-1}$ over the past few years (see Fig. 4). This corresponds to an increasing trend of nearly 5% per year and highlights the growing amount of fine fuel available for destructive wildfires (see Fig. 5). In ecoregions where annual cover was increasing most rapidly, exotic bromes such as cheatgrass were the most common annual species in AIM field data (Table A12)—strong evidence that these species are driving increases in annual cover. Climate change, specifically regional warming, may be facilitating the invasion of exotic annuals into higher elevations and cooler climates (Compagnoni and Adler 2014; Blumenthal et al. 2016; Smith




Figure 6. Estimated variance in vegetation trends at three spatial scales. Text within bars shows percent of group-level variance found at each scale. Trend variances are estimated from mixed effects models fit to log-transformed cover or aboveground production.

et al. 2022b). However, hotter and drier conditions may be detrimental at lower elevations (Larson et al. 2017).

Another more subtle yet concerning set of results were widespread loss of herbaceous perennial cover and production (see Fig. 5). Perennial grasses and forbs play a central role in forage production, carbon storage, biodiversity, and as indicators of overall rangeland health (West 2003; Pellant et al. 2005; Koteen et al. 2011). The epicenters of perennial loss were in Mediterranean California and the Warm Deserts (see Fig. 5). In Mediterranean California, perennial production has declined at a rate of 5.9% per year and is roughly one seventh what it was 30 yr ago (see Figs. 4 and 5). Our findings are consistent with a loss of cover and mortality of perennial herbs and bunchgrasses after prolonged drought in

the Southwest and elsewhere (Munson et al. 2013; Brookshire and Weaver 2015; McIntosh et al. 2019; Winkler et al. 2019; Williams et al. 2022). The exception to perennial declines was in the Northern Great Plains, where a lengthened growing season and increased CO₂ concentrations could be offsetting the detrimental effects of heat or drought (Reeves et al. 2014; Hufkens et al. 2016; Brookshire et al. 2020).

Despite a strong shift from perennial to annual dominance in many allotments (Tables A10–11), we did not find strong evidence that increases in annuals were directly responsible for declines in perennials. In general, areas with more rapid increases in annuals tended to show slower declines in perennials (positive correlations in Fig. 7E–H). This result is counter to the pattern we would expect if these groups strongly limited one another's abundances (Humphrey and Schupp 2004; Parkinson et al. 2013; Reisner et al. 2013; Chambers et al. 2016). To explain this, we hypothesize that perennials and annuals often respond similarly to variation in climate. For instance, if both perennials and annuals are water limited, they may show correlated trends across regions that have become drier or wetter over the past three decades (Munson et al. 2013; Larson et al. 2017). Positive correlations between trends were strongest at the field office scale, (see Fig. 7), a pattern that supports a role for climate variation rather than local interactions.

The proliferation of trees on BLM rangelands (see Fig. 3) is consistent with other studies showing the widespread expansion of trees into rangelands, both in the United States and worldwide (Nackley et al. 2017; Filippelli et al. 2020; Morford et al. 2022). In arid regions of North America, the expansion of native conifers into rangelands has been attributed to wildfire suppression and natural and anthropogenic climate change (Miller et al. 2008; Romme et al. 2009). Tree cover expansion can reduce plant species richness (Ratajczak et al. 2012) and habitat for wildlife that depend




Figure 7. Correlations between cover trends (x-axis) for annuals and bare ground (A–D) or between annuals and herbaceous perennials (E–H) at four spatial scales. Trends have been rescaled between 0 and 1 to facilitate visual comparison. Trends for ecoregions, field offices, and allotments are estimated from mixed-effects models fit to allotment-level average cover. Pixel-level trends are from the mixed-effect models fit cover at the individual pixel-scale. Blue lines and Pearson's r are shown for pairs of trends with a significant correlation at $P < 0.05$ level.




Figure 8. Trends in annual cover (A–C) and tree cover (D–F) across three management scales. **A**, Field office-scale trends in annual cover across the western Cold Deserts. **B**, Allotment-scale trends in annual cover for allotments in the Jarbidge Field Office in Idaho (blue outline in A). **C**, Pixel-scale trends in annual cover within a single allotment (blue outline in B). **D**, Field office-scale trends in tree cover for field offices within the Western Cold Deserts. **E**, Allotment-scale trends in tree cover for allotments in the Jarbidge Field Office in Idaho (no trends were calculated for allotments in gray, where tree cover is $\approx 0\%$). **F**, Pixel-scale trends in tree cover within a single allotment.

on treeless rangelands (Archer and Predick 2014). For example, even low abundances of tree cover may result in the extirpation of grouse (Lautenbach et al. 2017; Nackley et al. 2017) and other grassland birds (Brennan and Kuvlesky Jr. 2005). Given broad concern over the 50% reduction of grassland bird populations in the United States (Rosenberg et al. 2019), tracking further expansion of trees into rangelands will be key for future biodiversity conservation.

The big picture afforded by RS revealed the vulnerability of rangelands to continued tree expansion in the Northern Great Plains (see Fig. 5). Despite having $< 5\%$ tree cover today, expansion has occurred on two thirds of allotments in the Northern Great Plains during our 30-yr analysis window (Table A13). Increases in tree cover represent both infilling of existing stands and recruitment of new trees into formerly treeless rangelands (Filippelli et al. 2020). The latter process may be the most consequential for conservation of grasslands as trees rapidly mature and become seed sources for continued invasion. Using RS-derived maps to focus management on recently invaded rangelands can be an efficient strategy for preventing tree invasion into the most vulnerable habitats (Twidwell et al. 2021). While we cannot determine which species are responsible for tree cover increases from the RS data alone, the AIM data showed that junipers were the most common trees in all four ecoregions with significant increases in tree cover (Table A14). Where junipers have recently invaded rangelands, targeted removal can have benefits for local shrub and grassland associated species (Olsen et al. 2021; Roberts et al. 2022), increase soil moisture availability (Roundy et al. 2014), and help maintain herbaceous productivity (Morford et al. 2022).

Shrub cover was increasing in the Northern Great Plains and Mediterranean California and declining in the Western Cold Deserts (see Fig. 3). Shrub losses in the Western Cold Deserts likely reflect losses of sagebrush to increased wildfire and highlight the threat facing sagebrush-dependent wildlife (Coates et al. 2016). This region also saw both rapid increases in annuals and declines in bare ground (see Fig. 5), trends that are correlated with one another (see Fig. 7) and loss of shrubs (Fig. A1). Taken together, our results support the idea that exotic annual grasses are reducing bare ground, fueling more frequent and severe wildfires, and ultimately driving losses of cover of sagebrush and other shrubs (Chambers et al. 2014; Coates et al. 2016). Notably, we did not observe significant increases in shrubs in the AZ/NM Highlands and Warm Deserts, where previous studies raised concerns of shrub expansion (Eldridge et al. 2011). Satellite-based results do not disprove earlier cases of shrub expansion in the Southwest; however, they may indicate that the process of shrub invasion has stopped in recent decades or that it is a local phenomenon rather than ecoregion wide.

Quantifying spatial heterogeneity in vegetation trends

Heterogeneity and scale present grand challenges in monitoring rangelands across space and time (Fuhlendorf et al. 2017; Sayre 2017). Our findings show that trends varied greatly at all subecoregion scales: between field offices, allotments, and especially pixels within the same allotment (see Fig. 6). For rangeland managers, a critical implication of this result is that trends measured at a single field plot are unlikely to represent trends across an entire

allotment—let alone a field office or ecoregion (West 2003; West and Wu 2003). Fine-scale spatial heterogeneity may be especially problematic for quantifying trends in tree, shrub, and perennial cover, all of which had the majority of trend variance at the pixel scale (see Fig. 6). On the other end of the spectrum, annual cover and production trends showed more variation at the field office and allotment scales than at the pixel scale (see Fig. 6). This may indicate the importance of larger regional-scale factors such as climate in determining the direction and rate of annual trends. In general, the large proportion of total trend variance at the 30-m pixel scale cautions against extrapolating plot-level trends to larger spatial scales. Failure to consider the bigger picture provided by RS may result in poorly targeted management actions that fail to add up to beneficial outcomes at larger scales. The wall-to-wall coverage provided by RS data allows today's rangeland scientists to pinpoint hotspots of vegetation change and develop a spatially targeted strategy for defending intact habitat from annual invasion and woodland expansion (Doherty et al. 2022; Maestas et al. 2022; Tarbox et al. 2022).

In this analysis, we summarize vegetation trends at the ecoregion scale; however, BLM managers will frequently want to examine trends at much finer spatial scales. In Figure 8, we illustrate how vegetation trends derived from RS could be visualized at the appropriate scale for management. For instance, trends aggregated at the scale of BLM field offices highlight subregions where invasions by annuals and expansion of trees are most rapid (see Fig. 8A, 8D). Allotment-scale trends within a single field office (Fig. 8B, 8E) help locate priority areas for reviewing land use authorizations and identify core areas of intact rangelands for conservation (Maestas et al. 2022). At an even finer scale, trends measured at the pixel scale (Fig. 8C, 8F) make it clear which areas within an allotment have experienced increases in annuals and trees and which areas have experienced no change or even declines. Despite their utility, the large size of RS datasets can make them difficult to work with in desktop GIS software (Tarbox et al. 2022). To facilitate adoption of RS data in rangeland management, stakeholders may wish to prioritize the development of online dashboards for displaying trends at field office, allotment, and pixel scales (see Fig. 8). Examples of web apps that display spatial data at a variety of spatial and temporal scales are the U.S. Forest Service's Rangeland Production Monitoring Service (US Forest Service, RMRS 2018) and the US Drought Monitor (University of Nebraska, Lincoln 2022).

Management Implications

Over the past 30 yr, annuals have dramatically increased on BLM rangelands, while herbaceous perennials have declined, and tree cover has increased. These vegetation changes threaten the ecological function and habitat value of millions of acres of public lands. Unfortunately, the processes, such as climate change, that are likely driving these trends are largely outside of the BLM's control. Maintaining the ecological function of rangeland ecosystems in the future will be an enormous challenge and may require the BLM to adopt a more hands-on approach to vegetation management with a greater focus on restoration (Boyd et al. 2014; Maestas et al. 2022).

A newly released Sagebrush Conservation Design (SCD) provides one potential model for conserving western rangelands in the 21st century (Doherty et al. 2022). Instead of investing limited resources into areas that have already been invaded by trees or exotic annuals, the SCD prioritizes protecting the most intact areas of rangeland and expanding their extent by restoring surrounding areas. This strategy relies on RS data to map sagebrush-dominated ecosystems across both public and private rangelands. Mirroring the findings of our analysis, the SCD showed that 73% of the threats that intact sagebrush rangelands face are problems of

ecosystem function, such as invasion by exotic annuals. In contrast, only 27% of threats are the direct consequences of human activity, such as agricultural development (Doherty et al. 2022).

Although RS data are a powerful new tool for measuring rangeland status and trends, the continued collection of field data through the AIM program will remain essential for understanding BLM rangelands (Kachergis et al. 2022). Field monitoring captures important details about vegetation, such as species identity, that RS may never be able to resolve. Moreover, new field data will remain a precious commodity essential for the development and improvement of RS algorithms (Allred et al. 2021). Field data also represent the ground truth for land managers who may be skeptical of the accuracy of RS data (Tarbox et al. 2022). To gain the trust of stakeholders, RS data should be regularly compared against field data collected from a range of vegetation types so that discrepancies between the two sources of data can be better understood (Applestein and Germino 2022). Rather than thinking of satellite-based data as an alternative to field data, the two are complementary and both will play an important role in managing BLM rangelands in the future.

Data Availability

The remote sensing data used in this analysis are freely available from the Rangeland Analysis Platform (<https://rangelands.app/>). Computer code used to perform the statistical analysis reported in this study is available on Zenodo at <https://doi.org/10.5281/zenodo.7388703>.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We thank the Intermountain West Joint Venture, part of the United States Fish and Wildlife Service, for their contracting expertise. We thank Eric Jensen and two anonymous reviewers for providing helpful feedback on drafts of this manuscript.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1016/j.rama.2022.11.004](https://doi.org/10.1016/j.rama.2022.11.004).

References

- Allred, B.W., Bestelmeyer, B.T., Boyd, C.S., Brown, C., Davies, K.W., Duniway, M.C., Ellsworth, L.M., Erickson, T.A., Fuhlendorf, S.D., Griffiths, T.V., Jansen, V., Jones, M.O., Karl, J., Knight, A., Maestas, J.D., Maynard, J.J., McCord, S.E., Nangle, D.E., Starns, H.D., Twidwell, D., Uden, D.R., 2021. Improving Landsat predictions of rangeland fractional cover with multitask learning and uncertainty. *Methods in Ecology and Evolution* 12, 841–849.
- Applestein, C., Germino, M.J., 2022. How do accuracy and model agreement vary with versioning, scale, and landscape heterogeneity for satellite-derived vegetation maps in sagebrush steppe? *Ecological Indicators* 139, 108935.
- Archer, S.R., Predick, K.L., 2014. An ecosystem services perspective on brush management: research priorities for competing land-use objectives. *Journal of Ecology* 102, 1394–1407.
- Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67, 1–48.
- Blumenthal, D.M., Kray, J.A., Ortman, W., Ziska, L.H., Pendall, E., 2016. Cheatgrass is favored by warming but not CO₂ enrichment in a semi-arid grassland. *Global Changes in Biology* 22, 3026–3038.
- Boyd, C.S., Johnson, D.D., Kerby, J.D., Svejcar, T.J., Davies, K.W., 2014. Of grouse and golden eggs: can ecosystems be managed within a species-based regulatory framework? *Rangeland Ecology & Management* 67, 358–368.
- Boyle, S.P., Wylie, B.K., Major, D.J., 2019. Validating a time series of annual grass percent cover in the sagebrush ecosystem. *Rangeland Ecology & Management* 72, 347–359.

- Brennan, L.A., Kuvlesky Jr, W.P., 2005. North American grassland birds: an unfolding conservation crisis? *Journal of Wildlife Management* 69, 1–13.
- Brookshire, E.N.J., Stoy, P.C., Currey, B., Finney, B., 2020. The greening of the Northern Great Plains and its biogeochemical precursors. *Global Changes in Biology* 26, 5404–5413.
- Brookshire, E.N.J., Weaver, T., 2015. Long-term decline in grassland productivity driven by increasing dryness. *Nature Communications* 6, 7148.
- Chambers, J.C., Bradley, B.A., Brown, C.S., D'Antonio, C., Germino, M.J., Grace, J.B., Hardegree, S.P., Miller, R.F., Pyke, D.A., 2014. Resilience to stress and disturbance, and resistance to *Bromus tectorum* L. invasion in cold desert shrublands of western North America. *Ecosystems* 17, 360–375.
- Chambers, J.C., Germino, M.J., Belnap, J., Brown, C.S., Schupp, E.W., Clair, S.B., 2016. Plant community resistance to invasion by bromus species: the roles of community attributes, *Bromus* interactions with plant communities, and *Bromus* traits. In: Germino, M.J., Chambers, J.C., Brown, C.S. (Eds.), *Exotic brome-grasses in arid and semiarid ecosystems of the western US: causes, consequences, and management implications*. Springer Series on Environmental Management. Cham, Switzerland: Springer International Publishing, pp. 275–304.
- Coates, P.S., Ricca, M.A., Prochazka, B.G., Brooks, M.L., Doherty, K.E., Kroger, T., Blomberg, E.J., Hagen, C.A., Casazza, M.L., 2016. Wildfire, climate, and invasive grass interactions negatively impact an indicator species by reshaping sagebrush ecosystems. *Proceedings of the National Academy of Sciences* 113, 12745–12750.
- Compagnoni, A., Adler, P.B., 2014. Warming, soil moisture, and loss of snow increase *Bromus tectorum* 's population growth rate. *Elementa: Science of the Anthropocene* 2, 000020.
- Davies, K.W., Nafus, A.M., Davies, K.W., Nafus, A.M., 2012. Exotic annual grass invasion alters fuel amounts, continuity and moisture content. *International Journal of Wildland Fire* 22, 353–358.
- Doherty, K., Theobald, D. M., Bradford, J. B., Wiechman, L. A., Bedrosian, G., Boyd, C. S., Cahill, M., Coates, P. S., Creutzburg, M. K., Crist, M. R., Finn, S. P., Kumar, A. V., Littlefield, C. E., Maestas, J. D., Prentice, K. L., Prochazka, B. G., Remington, T. E., Sparklin, W. D., Tull, J. C., Wurtzbach, Z., and Zeller, K. A., 2022. A sagebrush conservation design to proactively restore America's sagebrush biome (no. 2022–1081), Open-File Report. US Geological Survey. Available at: <https://doi.org/10.3133/ofr20221081>. Accessed September 23, 2022.
- Dombeck, M.P., 1996. Thinking like a mountain: BLM's approach to ecosystem management. *Ecological Applications* 6, 699–702.
- Eldridge, D.J., Bowker, M.A., Maestre, F.T., Roger, E., Reynolds, J.F., Whitford, W.G., 2011. Impacts of shrub encroachment on ecosystem structure and functioning: towards a global synthesis. *Ecology Letters* 14, 709–722.
- Fernandez-Gimenez, M.E., McClaran, S.J., Ruyle, G., 2005. Arizona permittee and land management agency employee attitudes toward rangeland monitoring by permittees. *Rangeland Ecology & Management* 58, 344–351.
- Filippelli, S.K., Falkowski, M.J., Hudak, A.T., Fekety, P.A., Vogeler, J.C., Khalyani, A.H., Rau, B.M., Strand, E.K., 2020. Monitoring pinyon-juniper cover and aboveground biomass across the Great Basin. *Environmental Research Letters* 15, 025004.
- Federal Lands Policy and Management Act 1976. USC §§ 1701–1736, 1737–1782.
- Fuhlendorf, S.D., Fynn, R.W., McGranahan, D.A., Twidwell, D., 2017. Heterogeneity as the basis for rangeland management. In: Briske, D.D. (Ed.), *Rangeland Systems: Processes, Management and Challenges*. Springer Nature, Cham, Switzerland, pp. 169–196.
- GAO, 1992. Interior's monitoring has fallen short of agency requirements (no. GAO/RCED-92-51), GAO/RCED. US General Accounting Office, Washington, DC, USA. Available at: <https://www.gao.gov/assets/rced-92-51.pdf>. Accessed June 4, 2022.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., Moore, R., 2017. Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sensing and the Environment* 202, 18–27.
- Havstad, K.M., Peters, D.P.C., Skaggs, R., Brown, J., Bestelmeyer, B., Fredrickson, E., Herrick, J., Wright, J., 2007. Ecological services to and from rangelands of the United States. *Ecology and the Economy, Special Section—Ecosystem Services and Agriculture* 64, 261–268.
- Homer, C., Dewitz, J., Jin, S., Xian, G., Costello, C., Danielson, P., Gass, L., Funk, M., Wickham, J., Stehman, S., Auch, R., Ritters, K., 2020. Conterminous United States land cover change patterns 2001–2016 from the 2016 National Land Cover Database. *ISPRS Journal of Photogrammetry and Remote Sensing* 162, 184–199.
- Hufkens, K., Keenan, T.F., Flanagan, L.B., Scott, R.L., Bernacchi, C.J., Joo, E., Brunsell, N.A., Verfaillie, J., Richardson, A.D., 2016. Productivity of North American grasslands is increased under future climate scenarios despite rising aridity. *Nature and Climate Change* 6, 710–714.
- Humphrey, L.D., Schupp, E.W., 2004. Competition as a barrier to establishment of a native perennial grass (*Elymus elymoides*) in alien annual grass (*Bromus tectorum*) communities. *Journal of Arid Environment* 58, 405–422.
- Jones, M.O., Allred, B.W., Naugle, D.E., Maestas, J.D., Donnelly, P., Metz, L.J., Karl, J., Smith, R., Bestelmeyer, B., Boyd, C., Kerby, J.D., McIver, J.D., 2018. Innovation in rangeland monitoring: annual, 30 m, plant functional type percent cover maps for U.S. rangelands, 1984–2017. *Ecosphere* 9, e02430.
- Jones, M.O., Naugle, D.E., Twidwell, D., Uden, D.R., Maestas, J.D., Allred, B.W., 2020. Beyond inventories: emergence of a new era in rangeland monitoring. *Rangeland Ecology & Management* 73, 577–583.
- Jones, M.O., Robinson, N.P., Naugle, D.E., Maestas, J.D., Reeves, M.C., Lankston, R.W., Allred, B.W., 2021. Annual and 16-day rangeland production estimates for the western United States. *Rangeland Ecology & Management* 77, 112–117.
- Kachergis, E., Miller, S.W., McCord, S.E., Dickard, M., Savage, S., Reynolds, L.V., Lepak, N., Dietrich, C., Green, A., Nafus, A., Prentice, K., Davidson, Z., 2022. Adaptive monitoring for multiscale land management: lessons learned from the assessment, inventory, and monitoring (AIM) principles. *Rangelands* 44, 50–63. doi:10.1016/j.rala.2021.08.006.
- Koteen, L.E., Baldocchi, D.D., Harte, J., 2011. Invasion of non-native grasses causes a drop in soil carbon storage in California grasslands. *Environmental Research Letters* 6, 044001.
- Larson, C.D., Lehnhoff, E.A., Rew, L.J., 2017. A warmer and drier climate in the northern sagebrush biome does not promote cheatgrass invasion or change its response to fire. *Oecologia* 185, 763–774.
- Lautenbach, J.M., Plumb, R.T., Robinson, S.G., Hagen, C.A., Haukos, D.A., Pitman, J.C., 2017. Lesser prairie-chicken avoidance of trees in a grassland landscape. *Rangeland Ecology & Management* 70, 78–86.
- Lenth, R. V. 2022. Emmeans: estimated marginal means, aka least-squares means. Available at: <https://CRAN.R-project.org/package=emmeans>. Accessed June 4, 2022.
- Li, Z., Angerer, J.P., Wu, X.B., 2021. Temporal patterns of large wildfires and their burn severity in rangelands of western United States. *Geophysical Research Letters* 48, e2020GL091636.
- Mack, R.N., 1981. Invasion of *Bromus tectorum* L. into western North America: an ecological chronicle. *Agro-Ecosystems* 7, 145–165.
- Maestas, J.D., Porter, M., Cahill, M., Twidwell, D., 2022. Defend the core: Maintaining intact rangelands by reducing vulnerability to invasive annual grasses. *Rangelands* 44, 181–186. doi:10.1016/j.rala.2021.12.008.
- McCord, S.E., Pilliod, D.S., 2022. Adaptive monitoring in support of adaptive management in rangelands. *Rangelands* 44, 1–7.
- McIntosh, M.M., Holechek, J.L., Spiegel, S.A., Cibils, A.F., Estell, R.E., 2019. Long-term declining trends in Chihuahuan Desert forage production in relation to precipitation and ambient temperature. *Rangeland Ecology & Management* 72, 976–987.
- Miller, R.F., Tausch, R.J., McArthur, E.D., Johnson, D.D., Sanderson, S.C., 2008. Age structure and expansion of pinon-juniper woodlands: a regional perspective in the Intermountain West. Res Pap RMRS-RP-69 Fort Collins CO US Dep. Agric. For. Serv. Rocky Mt. Res. Stn. 15, 69.
- Morford, S.L., Allred, B.W., Twidwell, D., Jones, M.O., Maestas, J.D., Roberts, C.P., Naugle, D.E., 2022. Herbaceous production lost to tree encroachment in United States rangelands. *Journal of Applied Ecology* 59, 2971–2982.
- Munson, S.M., Muldavin, E.H., Belnap, J., Peters, D.P.C., Anderson, J.P., Reiser, M.H., Gallo, K., Melgoza-Castillo, A., Herrick, J.E., Christiansen, T.A., 2013. Regional signatures of plant response to drought and elevated temperature across a desert ecosystem. *Ecology* 94, 2030–2041.
- Nackley, L.L., West, A.G., Skowno, A.L., Bond, W.J., 2017. The nebulous ecology of native invasions. *Trends in Ecology and Evolution* 32, 814–824.
- Olsen, A.C., Severson, J.P., Maestas, J.D., Naugle, D.E., Smith, J.T., Tack, J.D., Yates, K.H., Hagen, C.A., 2021. Reversing tree expansion in sagebrush steppe yields population-level benefit for imperiled grouse. *Ecosphere* 12, e03551.
- Parkinson, H., Zabinski, C., Shaw, N., 2013. Impact of native grasses and cheatgrass (*Bromus tectorum*) on Great Basin forb seedling growth. *Rangeland Ecology & Management* 66, 174–180.
- Pellant, M.L., Shaver, P., Pyke, D.A., Herrick, J.E., 2005. Interpreting indicators of rangeland health, version 4. Technical Reference 1734-6. U.S. Department of the Interior, Bureau of Land Management, National Science and Technology Center, Denver, CO.
- R Core Team, 2022. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <https://www.R-project.org/>. Accessed June 4, 2022.
- Ratajczak, Z., Nippert, J.B., Collins, S.L., 2012. Woody encroachment decreases diversity across North American grasslands and savannas. *Ecology* 93, 697–703.
- Reeves, M.C., Moreno, A.L., Bagne, K.E., Running, S.W., 2014. Estimating climate change effects on net primary production of rangelands in the United States. *Climate Change* 126, 429–442.
- Reisner, M.D., Grace, J.B., Pyke, D.A., Doescher, P.S., 2013. Conditions favouring *Bromus tectorum* dominance of endangered sagebrush steppe ecosystems. *Journal of Applied Ecology* 50, 1039–1049.
- Rigge, M., Homer, C., Shi, H., Meyer, D., Bunde, B., Granneman, B., Postma, K., Danielson, P., Case, A., Xian, G., 2021. Rangeland fractional components across the western United States from 1985 to 2018. *Remote Sensing* 13, 813.
- Roberts, C.P., Scholtz, R., Fogarty, D.T., Twidwell, D., Walker Jr, T.L., 2022. Large-scale fire management restores grassland bird richness for a private lands ecoregion. *Ecological Solutions and Evidence* 3, e12119.
- Romme, W.H., Allen, C.D., Bailey, J.D., Baker, W.L., Bestelmeyer, B.T., Brown, P.M., Eisenhart, K.S., Floyd, M.L., Huffman, D.W., Jacobs, B.F., Miller, R.F., Muldavin, E.H., Swetnam, T.W., Tausch, R.J., Weisberg, P.J., 2009. Historical and modern disturbance regimes, stand structures, and landscape dynamics in piñon-juniper vegetation of the western United States. *Rangeland Ecology & Management* 62, 203–222.
- Rosenberg, K.V., Dokter, A.M., Blancher, P.J., Sauer, J.R., Smith, A.C., Smith, P.A., Stanton, J.C., Panjabi, A., Helft, L., Parr, M., Marra, P.P., 2019. Decline of the North American avifauna. *Science* 366, 120–124.
- Roundy, B.A., Young, K., Cline, N., Hulet, A., Miller, R.F., Tausch, R.J., Chambers, J.C., Rau, B., 2014. Piñon-juniper reduction increases soil water availability of the resource growth pool. *Rangeland Ecology & Management* 67, 495–505.
- Sayre, N.F., 2017. The politics of scale. University of Chicago Press, Chicago, IL, USA.
- Smith, J.T., Allred, B.W., Boyd, C.S., Davies, K.W., Jones, M.O., Kleinhesselink, A.R., Maestas, J.D., Naugle, D.E., 2022. Where there's smoke, there's fuel: dynamic

- vegetation data improve predictions of wildfire hazard in the Great Basin. *Rangeland Ecology & Management* doi:10.1016/j.rama.2022.07.005.
- Smith, J.T., Allred, B.W., Boyd, C.S., Davies, K.W., Jones, M.O., Kleinhesselink, A.R., Maestas, J.D., Morford, S.L., Naugle, D.E., 2022b. The elevational ascent and spread of exotic annual grass dominance in the Great Basin, USA. *Divers Distrib* 28, 83–96.
- Spaeth, K.E., Pierson, F.B., Herrick, J.E., Shaver, P.L., Pyke, D.A., Pellant, M., Thompson, D., Dayton, B., 2003. New proposed national resources inventory protocols on nonfederal rangelands. *Journal of Soil Water Conservation* 58, 18A–21A.
- Tarbox, B.C., Van Schmidt, N.D., Shyvers, J.E., Saher, D.J., Heinrichs, J.A., Aldridge, C.L., 2022. Bridging the gap between spatial modeling and management of invasive annual grasses in the imperiled sagebrush biome. *Rangeland Ecology & Management* 82, 104–115.
- Taylor, J.T., Toevs, G.R., Karl, J.W., Bobo, M., Karl, M.S., Miller, S., Spurrier, S., 2014. AIM-Monitoring: a component of the BLM assessment, inventory, and monitoring strategy. Government Publication/Report. Technical Note 445. U.S. Department of Interior, Bureau of Land Management, National Operations Center, Denver, CO (Technical Note No. 445).
- Toevs, G.R., Karl, J.W., Taylor, J.J., Spurrier, C.S., Karl, M., Bobo, M.R., Herrick, J.E., 2011a. Consistent Indicators and Methods and a Scalable Sample Design to Meet Assessment, Inventory, and Monitoring Information Needs Across Scales. *Rangelands* 33, 14–20.
- Toevs, G.R., Taylor, J.J., Spurrier, C.S., MacKinnon, W.C., Bobo, M.R., 2011b. Bureau of Land Management assessment, inventory, and monitoring strategy: for integrated renewable resources management. US Department of the Interior Bureau of Land Management National Operations Center, Denver, CO, USA.
- Twidwell, D., Fogarty, D.T., Weir, J.R., 2021. Reducing woody encroachment in grasslands: a guide for understanding risk and vulnerability. Oklahoma Cooperative Extension Service. (2021), Publication E-1054. Available at <https://handle.nal.usda.gov/10113/7548409>. Accessed June 4, 2022.
- University of Nebraska, Lincoln. 2022. US drought monitor. Available at: <https://droughtmonitor.unl.edu/>. Accessed June 4, 2022.
- US Forest Service, RMRS. 2018. Development of the Rangeland Production Monitoring Service could improve rangeland management. Rangeland Production Monitoring Service. Available at: <https://www.fs.usda.gov/rmrs/projects/development-rangeland-production-monitoring-service-could-improve-rangeland-management>. Accessed June 4, 2022.
- USDA NASS, 2016. National Agricultural Statistics Service Cropland Data Layer. USDA-NASS, Washington, DC. Available at <https://nassgeodata.gmu.edu/CropScape/>. Accessed June 4, 2022.
- USDA NRCS, 2022. The PLANTS Database National Plant Data Team, Greensboro, NC USA. Available at <http://plants.usda.gov>. Accessed June 4, 2022.
- Veblen, K.E., Pyke, D.A., Aldridge, C.L., Casazza, M.L., Assal, T.J., Farinha, M.A., 2014. Monitoring of livestock grazing effects on Bureau of Land Management Land. *Rangeland Ecology & Management* 67, 68–77.
- West, N., Wu, B., 2003. New alternatives for monitoring rangelands. *Rangelands* 25 (1), 24–26.
- West, N.E., 2003. History of rangeland monitoring in the U.S.A. *Arid Land Research and Management* 17, 495–545.
- Williams, A.P., Cook, B.I., Smerdon, J.E., 2022. Rapid intensification of the emerging southwestern North American megadrought in 2020–2021. *Nature Climate Change* 12, 232–234.
- Winkler, D.E., Belnap, J., Hoover, D., Reed, S.C., Duniway, M.C., 2019. Shrub persistence and increased grass mortality in response to drought in dryland systems. *Global Change Biology* 25, 3121–3135.