

Sicklepod [Senna obtusifolia (L.) H. S. Irwin & Barneby] "Getting sleepy?"

Authors: Sosnoskie, Lynn M., Steckel, Sandy, and Steckel, Lawrence E.

Source: Weed Technology, 35(6) : 1052-1058

Published By: Weed Science Society of America

URL: https://doi.org/10.1017/wet.2021.70

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

www.cambridge.org/wet

Intriguing World of Weeds

Cite this article: Sosnoskie LM, Steckel S, Steckel LE (2021) Sicklepod [*Senna obtusifolia* (L.) H. S. Irwin & Barneby] "Getting sleepy?". Weed Technol. **35**: 1052–1058. doi: 10.1017/ wet.2021.70

Received: 28 April 2021 Revised: 31 July 2021 Accepted: 18 August 2021 First published online: 6 September 2021

Associate Editor:

Daniel Stephenson, Louisana State University Agricultural Center

Author for correspondence:

Larry Steckel, Professor, University of Tennessee, 605 Airways Blvd., Jackson, TN 38301. Email: lsteckel@utk.edu

© The Author(s), 2021. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons.Attribution licence (https:// creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Sicklepod [*Senna obtusifolia* (L.) H. S. Irwin & Barneby] "Getting sleepy?"

Lynn M. Sosnoskie¹, Sandy Steckel² and Lawrence E. Steckel³

¹Assistant Professor, Cornell University, Geneva, NY, USA; ²Extension Assistant, University of Tennessee, Jackson, TN, USA and ³Professor, University of Tennessee, Jackson, TN, USA

There is a Reaper, whose name is Death, And, with his sickle keen, He reaps the bearded grain at a breath, And the flowers that grow between. – Henry Wadsworth Longfellow, "The Reaper and the Flowers"

I love sleep. My life has a tendency to fall apart when I'm awake, you know. - Ernest Hemingway

Introduction

Sicklepod [Senna obtusifolia (L.) H. S. Irwin & Barneby], also known as Cassia obtusifolia (L.), is an annual, herbaceous, dicotyledonous plant in the Fabaceae (Leguminosae) family, which is commonly known as the bean, legume, and pea family. The Fabaceae consist of herbs, shrubs, vines, or trees; the family has a cosmopolitan distribution with members numbering approximately 751 genera and 19,500 species (Christenhusz and Byng 2016). Characteristics of the Fabaceae include alternate, stipulate, and compound leaves. Leaflets often have pulvini (i.e., cushion-like swellings at the base of leaves that are subject to changes in turgor pressure), which are responsible for growth-independent or "sleep" movements. Another interesting anatomical feature exhibited by many species in the family is the formation of parenchymatous root nodules that are generated in association with nitrogen-fixing bacteria (Zomlefer 1994). The ovary of the Fabaceae usually develops into a dehiscent legume (e.g., pod). Although some Fabaceae may be weedy pests, others are important food crops [e.g., soybean, Glycine max (L.) Merr.] and fodder and forage plants (e.g., alfalfa, Medicago sativa L.). Some members of the Fabaceae produce valuable gums [e.g., gum arabic, Acacia senegal (L.) Britton] and dyes (e.g., indigo, Indigofera tinctoria L.), whereas others are prized as desirable ornamentals (e.g., eastern redbud, Cercis canadensis L.). Many species in the Fabaceae produce alkaloids or cyanogenic glycosides in different plant structures. Rotenone, an isoflavone insecticide, is derived from Derris eclipta (Wall.) Benth.

Etymology

Carl von Linné (Linnaeus) in 1753 first described sicklepod as *Cassia obtusifolia*. *Obtusifolia* is derived from the Latin words obtusus meaning dull or blunt, and folium meaning leaf, both of which refer to leaflet shape. Irwin and Barneby (1982) undertook a revision of *Cassia* and suggested that the species be moved to the genus *Senna*, emphasizing the importance of floral morphology for classification as opposed to fruit structure (e.g., Bentham, 1871; Marazzi et al. 2006). Further studies, including molecular examinations, have confirmed the separation of *Senna* from *Cassia* (Marazzi et al. 2006; Randell 1988). The Weed Science Society of America (WSSA) currently uses *Senna obtusifolia* (L.) H. S. Irwin and Barneby as the preferred scientific designation for sicklepod (WSSA 2021a). In preparing this manuscript, the authors found that the genus name *Cassia* was used in many cited references until roughly the mid-1990s, after which sicklepod was referred to as *Senna*.

Linnaeus also described what he considered a separate species, sickle senna, as *Cassia tora* L. [now *Senna tora* (L.) Roxb.]. More than one hundred years later, Bentham (1871) argued that *C. obtusifolia* and *C. tora* were the same species and suggested they both be referred to as *C. tora*. From this time on there has been confusion in the literature and debate among taxonomists about whether *S. obtusifolia* and *S. tora* are the same or independent species (Crawford et al. 1990). Some publications state that *S. obtusifolia* is synonymous with *S. tora* (e.g., Holm et al. 1979, 1997), while others distinguish them as separate species (e.g., Brenan 1958; Irwin and Barneby 1982; Randell 1995).

While it is commonly acknowledged that *S. obtusifolia* and *S. tora* are related and are morphologically similar, they do vary in appearance and distribution. Documented taxonomical differences between species occur in anther shape and structure, seed structure, shape and weight, number of glands on leaf rachis, and length of pedicels (Brenan 1958; Randell 1988). Irwin and Turner (1960) also found high levels of infertility resulting from experimental crosses



Sicklepod flowering.

between the two species, which suggested they are, indeed, distinctly different. Upadhaya and Singh (1986) also reported that *S. obtusifolia* and *S. tora* failed to hybridize.

With respect to range, sicklepod occurs within the core distribution area of the series (South America), whereas sickle senna is found primarily in China, India, the Philippines, and other parts of Asia (Randell 1995). However, *S. obtusifolia* does co-occur in some areas with *S. tora* (e.g., India and southern Asia; Randell 1995). Randell (1995) noted it was significant that those who thought the taxa were synonymous worked in America or Europe, where only herbarium material was available for *S. tora*. Researchers in Asia who had access to living specimens uniformly declared the taxa to be separate. Randell (1995) hypothesized that *S. obtusifolia* made its way to Africa from the Caribbean and spread from there to Asia and gave rise to *S. tora*. Irwin and Turner (1960) also state that *S. tora* likely arose from *S. obtusifolia* and that the wide distribution of sicklepod occurred at least in part by human intervention.

The common name for *S. obtusifolia* is sicklepod (WSSA 2021a), which alludes to its long, strongly curved (sickle-shaped), leguminous fruit. Other common names include sickle-pod, sickle senna, java-bean, coffeebean, coffee bean, coffee-pod, coffee-weed, and indigo (Bryson and DeFelice 2009; Burrows and Tyrl 2006; Cock and Evans 1984; McCormack and Neisler 1980; Nowick 2015; Page et al. 1977 Robertson and Lee 1976; USDA 2020). Adding to the confusion over nomenclature, *S. tora* is also known by the common name sicklepod. *Senna occidentalis* (L.) Link is referred to as coffee senna, which is similar to several common names of *S. obtusifolia* (ITIS 2021; WSSA 2021b). Both *S. tora* and *S. occidentalis* are called coffee weed (Nowick 2015).

Description

The following descriptive information is summarized from works published by Bryson and DeFelice (2009), Creel et al. (1968), Gleason and Cronquist (1963), Hilty (2018), Irwin and Barneby (1982), and Zomlefer (1994). Sicklepod is an erect (but sometimes sprawling), annual (sometimes biennial) herb. Cotyledons are rounded, with three to five distinct veins. Stems are erect, 0.3 to 2.5 m tall, branched, green, round, and glabrous. Leaves are alternate, pinnately compound with two or three (most commonly three) pairs of leaflets. Leaflets are elliptic to obovate with blunt or rounded tips. Leaflets are 2.0 to 7.0 cm long with the terminal

1053

Leaflets of young plants may be wrinkled. A gland (sometimes two) is present on the rachis between the lower leaflets. Petioles are 10 to 15 mm long. Leaves and stems have a distinct, rank odor when crushed or cut. Mature plants have a fibrous root system originating from a taproot. However, unlike many members of the Fabaceae family, sicklepod is non-nodulating and therefore does not form symbiotic associations with nitrogen-fixing bacteria.

Sicklepod plants bloom sparingly. One or two inflorescences are borne on short (3 to 4 mm) peduncles in leaf axils. Flowers consist of five rounded, asymmetrical petals that are yellow, showy, and 0.8 to 2.0 cm long. Sepals are unequal, 4.0 to 10.0 cm long and 2.0 to 5.0 mm wide. There are 10 stamens (three are infertile) and ovaries are finely hairy. Sicklepod is self-pollinated, and fertilization typically occurs before flowers open (Irwin and Barneby 1982; Retzinger 1984). The fruit is a legume, 8.0 to 20.0 cm long, 3.0 to 6.0 mm wide, slender, strongly curved, glabrous, rounded in cross-section, and terete or rod-like. Fruits are green turning to brown at maturity and dehiscing elastically along both sutures. Fruit set at the bottom portion of plants matures first; maturation progresses up the plant even while the younger, uppermost nodes are still flowering (Turner and Karlander 1975). Seeds are thick, rhomboidal or angular, 4.0 to 5.0 mm long, brownish in color, and shiny. Seeds have a waxy covering about 0.1 mm thick and a closed, elliptical pleurogram. Sicklepod has a hard seed coat, which is a barrier to germination. Unlike most other species in the genera, the embryo of sicklepod has folded cotyledons (Robertson and Lee 1976).

There are two forms of sicklepod in the Western hemisphere (Irwin and Barneby 1982). The form that occurs in the Caribbean and the southeastern United States has one extrafloral nectary on the upper surface of the rachis and a chromosome count of 2n = 28; the other form, which originated in northern South America, has two extrafloral nectaries and a chromosome count of 2n = 26 (Irwin and Turner 1960; Retzinger 1984). Retzinger (1984) reported that 10 sicklepod selections from nine southeastern U.S. states all produced plants that had one petiolar gland located on the upper side of the rachis between the two lower leaflets, which agrees with that reported by Irwin and Barneby (1982). The variant in the United States also has a broader pod compared with the South American form and is the subject of this paper.

In addition to *S. tora*, sicklepod may be easily confused with coffee senna; both species often grow together in the southern United States (McCormack and Neisler 1980). However, leaves of coffee senna produce four to six pairs of leaflets compared with sicklepod, which has two to three pairs (Bryson and DeFelice 2009). Leaflets of coffee senna are ovate or oblong-lanceolate with a pointed tip, unlike the blunted or round-tipped obovate leaflets of sicklepod. Cotyledons of coffee senna are glabrous above, but have white trichomes below, whereas trichomes are not found on sicklepod. The legume of coffee senna is more linear, straight or slightly curved, as compared to the strongly curved, sickle-shaped fruit of sicklepod. Coffee senna seeds are also elliptical and dull brown, whereas sicklepod seeds are shiny brown and angular.

An interesting characteristic of many Fabaceae plants is the pulvinus, which is a vegetative thickening at each petiole and petiolule base that is associated with leaf movement. Changes in turgor resulting from water movement into or out of extracellular spaces causes leaflets to "open" during the day and "close" at night, and is sometimes referred to as "sleep movements" (Andersen and Koukkari 1979; Zomlefer 1994). This phenomenon was observed and recorded for some plants as early as the time of Alexander the Great (Cumming and Wagner 1968). Linnaeus even included a list of plants that "sleep" in his essay titled "Somnus Plantarum" published in 1755. Kraatz and Andersen (1980) studied the difference in percent projected leaf area (PPLA) of sicklepod, defined as "the projected leaf area expressed as a percentage of total actual leaf areas". Sicklepod cotyledons and leaflets open to an almost horizontal position during the day and fold tightly together to an almost vertical position at night resulting in a 7-fold difference between the highest (daytime) and lowest (nighttime) PPLA values (Andersen and Koukkari 1979; Kraatz and Andersen 1980). Sleep movements are a circadian phenomenon that is endogenously controlled and occurs even if the sicklepod plants are kept in continuous light or darkness (Andersen and Koukkari 1979; Kraatz and Andersen 1980). This phenomenon can have an impact of weed control success; percent control of sicklepod with linuron was highly correlated with PPLA in an herbicide application timeof-day study (Kraatz and Andersen 1980).

Sicklepod development is influenced by environmental factors such as moisture availability, temperature, pH, and light. Sicklepod has a relatively high stomatal resistance and a low threshold leaf water potential and is characterized as a "water saver" (Patterson and Flint 1983), which may explain the species' drought tolerance (Hilty 2018). Maximum dry weight, leaf area, plant height, node number, and leaf number for sicklepod occur when day/night temperature regimes of 29/26 C and 34/26 C occur. Leaf production ceases when aerial temperatures are below 13 C. Daytime temperature has more effect on main stem node number, leaf number, leaf weight, and total leaf area than night temperature (Patterson 1993). Night temperature has more influence on plant height, root weight, and total plant weight (Patterson 1993). Optimal temperatures for sicklepod root growth are reported to be in the range of 32 to 39 C or higher (Teem et al. 1974). These findings may explain why sicklepod thrives in the higher temperatures of July and August in the southeastern United States (Creel et al. 1968; Jones and Walker 1993 Patterson 1993; Patterson and Flint 1983; Shibles et al. 1975; Teem et al. 1974, 1980; Wright et al. 1999).

Although sicklepod root growth can occur over a wide range of soil pH values, in one study, primary root elongation was greater when the soil pH was 6.5 compared with pH 5.5 or 5.1 (Teem et al. 1974). Creel et al. (1968) reported that sicklepod plants grew well in soils with a pH range of 3.2 to 7.9 with an optimal pH range of 5.5 to 6.0. Sicklepod responded similarly to cotton [*Gossypium hirsu-tum* L.] in response to nitrogen, phosphorus, and potassium levels in soil (Creel et al. 1968). Sicklepod was found to be more tolerant to low potassium levels in soil than to low phosphorous levels (Hoveland et al. 1976). Sicklepod plants are also tolerant to a range of soil types (Dirar 1984; Hilty 2018).

With respect to root architecture, sicklepod is reported to have two times more roots than soybean [*Glycine max* (L.) Merr.] among plants with similar root fresh weights (Wright et al. 1999). Sicklepod roots were 2.9 times longer, although smaller in diameter, than soybean roots, providing the species with significantly more root surface area. Sicklepod seedling roots occupy a much larger volume of soil compared to that of soybean (Wright et al. 1999). Combined, these factors would, presumably, give sicklepod an advantage for acquiring water and nutrients when in competition with soybean.

Sicklepod can emerge continually beneath a soybean canopy, and it grows well under trees, which suggests shade tolerance (Becker 1983; Norsworthy and Oliveira 2007). The presence of a soybean canopy stimulated aboveground biomass elongation and growth of sicklepod when roots competed (Jones and Walker 1993). Sicklepod is a short-day plant with a critical day length of 13 to 14 h for flower initiation (Patterson 1993; Senseman and Oliver 1993; Shibles et al. 1975; Turner and Karlander 1975; Wright et al. 1999). Sicklepod in Arkansas initiated flowering 70 to 77 d after emergence even with a 2-wk difference in planting date, affirming a photoperiodic relationship to flowering (Senseman and Oliver 1993). Flowering ceased when the photoperiod reached 12 h (Senseman and Oliver 1993). No flowers developed in 15- or 16-h photoperiods (Patterson 1993).

Sicklepod plants grown without competition produced on average 1,500 to 16,000 seeds per plant (Retzinger 1984; Senseman and Oliver 1993). Average numbers of seeds per pod ranged from 24.8 to 30 (Bridges and Walker 1985; Retzinger 1984). Bridges and Walker (1985) reported that sicklepod plants produced 2.5×10^7 and 4.5×10^7 seeds ha⁻¹ at a subcompetitive density of 0.45 and 0.90 plants m⁻², respectively. Sicklepod produced 145 and 995 pods per plant in Arkansas, which resulted in 2,000 to 11,000 seeds per plant, depending on growing conditions (Senseman and Oliver 1993). Sicklepod plants produced 14×10^7 seeds ha⁻¹ under intraspecific interference and 7.0 × 10^7 seeds ha⁻¹ under interspecific interference (Senseman and Oliver 1993). Soybean competition reduced sicklepod seeds per plant by 49% to 74% at a weed density of 1 plant per 30-cm row (Sims and Oliver 1990).

Seed dormancy, which is enforced by the hard and waxy coat, resulted in only 15% germination over a 12-mo study (Creel et al. 1968). Similarly, Retzinger (1984) reported that the germination of hand-harvested, nonscarified sicklepod seed grown from 10 different seed selections ranged from 2% to 23%. Germination in the laboratory may be increased by mechanical scarification or abrasion, puncturing the seed coat with a needle, or treatment with sulfuric acid (Creel et al. 1968; Teem et al. 1974, 1980). For example, Egley and Chandler (1978) found that piercing sicklepod seed coats resulted in dramatically increased germination (pierced $96\% \pm 3\%$) relative to the untreated check $(4\% \pm 2\%)$ by allowing imbibition of water. Bararpour and Oliver (1998) and Vencill and Banks (1994) reported that sicklepod fares better in a no-till system than in a conventionally tilled system because the seedbank will not be actively depleted. Repeated disking favors sicklepod germination and emergence (perhaps due to the mechanical scarification), and the resulting plants produced more pods per plant in a conventional system (Bridges and Walker 1985).

Creel et al. (1968) and Teem et al. (1980) found that sicklepod germination occurs between 15 and 36 C with an optimum germination range of 24 to 36 C; rapid seedling growth happens between 30 and 36 C. More recently, Norsworthy and Oliveira (2006) reported sicklepod germination occurred over a range of constant temperatures from 15 to 50 C with an optimum germination temperature range of 15 and 30 C for seed sourced from South Carolina. No germination was observed at 12.5 C or below (Norsworthy and Oliveira 2006).

Sicklepod seeds were found to be viable after dry, low-temperature storage up to 30 mo (Egley and Chandler 1978). Even though sicklepod has a hard seed coat, seed longevity was less than what was reported for similarly hard-seeded species such as velvetleaf (*Abutilon theophrasti* Medik.), spurred anoda [*Anoda cristata* (L.) Schiltdl.], or purple moonflower [*Ipomoea muricata* (L.) Jacq.]. Egley and Chandler (1983) reported sicklepod seeds in the soil lost their hardness relatively quickly; after 5.5 yr of burial, only 6% of seeds were still hard, although all remained viable. Although sicklepod emergence is greater at shallow burial depths, seedlings could



Sicklepod competing with soybean.

still emerge from seeds buried to a depth of 12.7 cm (Norsworthy and Oliveira 2006; Teem et al. 1980). Seedling emergence was observed to occur from April through November in South Carolina (Norsworthy and Oliveira 2006).

Sicklepod seeds were more tolerant than soybean under simulated drought conditions with some seed germinating at an osmotic pressure as high as 10 bars (10-bar water solutions of polyethylene glycol; Hoveland and Buchanan 1973). Norsworthy and Oliveira (2006) reported a relationship between soil solution osmotic potential and soil temperature effects on sicklepod germination. For example, soil solution osmotic potential had no effect on sicklepod germination at a soil temperature of 15 C; however, at 30 C, when osmotic potential was low (no stress), sicklepod germination was less than when osmotic potential was high (more stress; Norsworthy and Oliveira 2006). This suggests that sicklepod is more likely to germinate when temperatures are high and the seed is under drought stress, which occurs more frequently in areas such as the southeastern United States, where it is a common weed.

Natural light, and red- to far-red light ratio did not affect germination, although Norsworthy and Oliveira (2006) reported having scarified the seed and did not study whether this influenced their results. Norsworthy and Oliveira (2006) also reported that germination was less at 2% oxygen than at 20% oxygen. They went on to note that sicklepod germinated from solution pH 3 to pH 9, with a predicted maximum germination at a solution pH of 6.1. Their research suggests that sicklepod germinates most readily near neutral pH but it can germinate under both acidic and basic conditions.

Various attempts have been made to reduce the weed seed soil seedbank, including use of the integrated Harrington

Seed Destructor mill to destroy seed at harvest. Sicklepod seeds had <1% survival from this process (Schwartz-Lazaro and Norsworthy 2018).

History and Distribution

Senna is a large, widespread, and diverse genus with approximately 350 species; of these, 80% occur in tropical and warm, temperate climates of the Americas (Irwin and Barneby 1982; Mabberley 1997; Marazzi et al. 2006). S. obtusifolia is native to tropical America and the southeastern United States and is now widespread throughout most tropical and warm-temperate regions. Its range in the United States has expanded and is now found in southern states from Texas to Florida, north to New Jersey and New York, and west to Nebraska, Wisconsin, and California. Herbarium records reveal that it has also been introduced to Hawaii (Staples et al. 2003). Sicklepod can be found growing in waste places, annual cropping systems (especially no-till), pastures, fallow land, orchards, riparian zones, roadsides and railways, and moist, open woodlands (Bryson and DeFelice 2009; Hilty 2018; Vencill and Banks 1994). Sicklepod seeds do not have specialized dispersal mechanisms and most fall close to the parent plant. Seed movement may occur via machinery, or through contamination of grain, hay, fodder, or forage. Sicklepod most commonly grows at elevations of 0 to 500 m, but may be found up to 1,680 m (Irwin and Barneby 1982).

Characteristics

Weediness and Control

In a 1998 survey of weeds in Georgia fields of corn (*Zea mays* L.), cotton, forages and pastures, peanut (*Arachis hypogaea* L.), small grains, soybean, tobacco (*Nicotiana tabacum* L.), and vegetables, sicklepod was identified as the most troublesome across the state (Webster and MacDonald 2001). It was found throughout all nine climate gradients of the state and was present in all the crops examined.

Densities of 7.7 sicklepod plants per square meter reduced soybean yield by as much as 35% (Thurlow and Buchanan 1972). A sicklepod density of 1.6 plants 1-m row caused a reduction in crop height and a 5% yield loss in soybean (Shurtleff and Coble 1985). One sicklepod plant per 2-m row (0.6 m⁻²) caused a 20% yield reduction in soybean in Florida (Currey et al. 1981).

Height and weight of sicklepod was reduced when soybeans were grown on narrow rows (25 cm) compared with wider rows (102 cm; Monks and Oliver 1988). Weed weight at harvest was inversely correlated with soybean yield, and sicklepod was more competitive when grown 15 to 30 cm from the drill versus in the drill row. Soybean competing with sicklepod plants that were spaced 10 and 30 cm apart yielded 25% to 35% less than when grown alone or with sicklepod spaced 90 cm apart, indicating that 90 cm is outside the area of influence of sicklepod (Bozsa et al. 1989). This suggests that sicklepod is only competitive with soybean at high weed density (Monks and Oliver 1988).

Soybean yield was reduced if sicklepod competed with the crop for as little as 4 wk (McWhorter and Sciumbato 1988), although when in competition with soybean, sicklepod's greatest growth occurs later in the season (Monks and Oliver 1988). At 6 wk after emergence (WAE), the major portion of sicklepod leaf area index (LAI) was in the lower 30 cm of the sicklepod canopy (Sims and Oliver 1990. By 10 WAE, LAI was uniform throughout the plant canopy, and as sicklepod height increased through 14 WAE, a greater percentage of LAI was in the upper portion of the plant (Sims and Oliver 1990). Reduction in soybean dry matter to sicklepod competition was first detected 6 WAE (Sims and Oliver 1990). Sicklepod reduced soybean dry matter weights 20% by 14 WAE, indicating that it became more competitive during the reproductive stage of soybean (Sims and Oliver 1990). By 10 WAE, sicklepod was taller than soybean and had formed a partial canopy over the crop during its reproductive stages. S. obtusifolia reduced soybean yield more than johnsongrass [Sorghum halapense (L.) Pers.] when competing individually and was similar when both weeds were present together (Sims and Oliver 1990). Individual sicklepod plants exhibited the greatest increase in weight and growth of axillary branches from 72 to 96 d after emergence. The increase in weight was attributed to an almost 8-fold increase in leaf number during this period. Axillary branches continued to grow rapidly from 96 to 120 d after emergence and about 30% to 60% of sicklepod seed production occurred at these sites. Sicklepod height increased most rapidly from 96 to 120 d after emergence. Soybean grew little in height from 35 to 84 d after emergence, though the competing sicklepod continued to grow until they were 30 to 45 cm taller than the crop (Bozsa et al. 1989).

Sicklepod is very competitive with cotton. Seed cotton yield in a study in Alabama was reduced 10% to 40% from a sicklepod density of 1.09 plants row m^{-1} (Buchanan and Burns 1971). Increasing the weed density to 6.6 plants row m^{-1} resulted in a 45% to 80% yield loss (Buchanan and Burns 1971). Additional studies have reported that for every sicklepod plant per 15 m^{-1} of row, cotton yield is reduced by 24 to 49 kg ha⁻¹ (Buchanan et al. 1980; Street et al. 1981). Additionally, the percentage of trash in mechanically harvested seed cotton typically increased with higher sicklepod densities (Buchanan et al. 1980).

Sicklepod did not reduce yields of two peanut cultivars when the crop was kept weed-free for 4 wk after crop emergence (Hauser et al. 1975). Other reports suggest that the weed-free requirement is even longer (5 to 6 wk) to adequately preserve peanut yields (Buchanan and Hauser 1980; Hauser and Buchanan 1981). Sicklepod plants in competition with peanut grew taller than the crop if weeds emerged within 4 wk of crop planting (Hauser and Buchanan 1981). Although sicklepod seedlings were often able to emerge after the fourth week, few were able to break through the peanut canopy (Hauser et al. 1975). Hauser et al. (1975) reported that sicklepod had to compete for a duration of 10 wk or more before peanut yield was reduced. With respect to density, Hauser et al. (1982) found a 6.1 to 22.3 kg ha⁻¹ reduction in peanut yield for each sicklepod plant 10 m⁻². Planting peanuts in narrow rows (20.3 cm < 40.6 cm < 81.2 cm) reduced sicklepod vegetative growth (Buchanan and Hauser 1980).

Sicklepod densities >10.8 weeds m^{-2} frequently reduced yield of lima bean [*Phaseolus lunatus* (L.); Glaze and Mullinix 1984]. Average yield of lima bean decreased if sicklepod competed for more than 6 wk (Glaze and Mullinix 1984). However, losses due to decreased harvest efficiency or contamination in green vegetable crops by weeds could be greater than those from competition.

Toxicity

All parts of *S. obtusifolia* are toxic, especially the seeds; this is attributed to various derivatives of anthraquinones (Burrows and Tyrl 2006; Crawford et al. 1990). Because sicklepod seeds have a higher density than corn they tend to settle at the bottom of freight cars during transport; Page et al. (1977) reported grain samples

containing as much as 40% sicklepod. Large amounts of sicklepod seeds should be removed prior to feeding grain to livestock (Crawford et al. 1990). In addition to anthraquinones, vegetative sicklepod has a high calcium-to-phosphorus ratio, which could result in metabolic disorders if these weeds were used as a sole feed source (Bosworth et al. 1980). McCormack and Neisler (1980) reported sicklepod to be the leading cause of livestock poisoning in the southern United States. While cattle do not usually forage on sicklepod plants if other, more desirable forage is available, problems may occur if animals are fed hay or green chop containing an abundance of sicklepod plant material or fed grain that is heavily contaminated with seeds (Cock and Evans 1984; McCormack and Neisler 1980). Often the first signs of Senna intoxication in cattle are diarrhea and anorexia, which occur within hours of ingestion (Burrows and Tyrl 2006; McCormack and Neisler 1980). The toxicity also affects the skeletal muscles (especially those in the hind legs), kidneys, and liver of cattle (McCormack and Neisler 1980). Weakness, uncoordinated movement, and muscular trembling are often noticeable before the animal becomes recumbent (Burrows and Tyrl 2006; McCormack and Neisler 1980). Cattle often die from congestive heart failure within 2 to 3 d after recumbency (McCormack and Neisler 1980). Feeding Holstein calves rations including S. obtusifolia seed resulted in intermittent feed refusal, decreased feed consumption, intermittent diarrhea, and decreased weight gain (Putnam et al. 1988). The average daily weight gains were significantly greater for the calves in the control group than for calves in either group that were fed sicklepod seed in their ration.

Ingestion of feed contaminated by sicklepod seeds also poses a serious threat to pigs and chickens (Burrows and Tyrl 2006). White leghorn chicken hens consumed less feed and laid fewer eggs when sicklepod seed was ground and mixed into feed at levels of 2%, 5%, and 10% (Page et al. 1977). Egg size was also reduced, and egg yolks were discolored. Feed consumption and egg production did not return to control levels for weeks after the contaminated feed was stopped, with the effects more pronounced in younger hens. Cock and Evans (1984) reported that sicklepod seeds were harmful to chickens due to the presence of a trypsin inhibitor.

Uses

Conversely, sicklepod has been reported to be an important food plant for human nutrition in northern Senegal when other food sources are scarce (Becker 1983). The leaves are harvested and cooked like greens and supply several nutrients including Vitamin B₂. In Sudan, green leaves harvested at the flowering and fruiting stages of sicklepod growth are ground, fermented, dried, and consumed in a stew as a meat substitute called kawal (Dirar 1984). Though it has a strong, repugnant odor, kawal consumption has spread in that area. The foliage of *S. obtusifolia* has also been reported as a purgative drug, and poultices are made for treating sores, ulcers, and insect bites (Crawford et al. 1990; Irwin and Barneby 1982). Though seeds are not used for human consumption in Sudan, Dirar (1984) reported claims that they could be a cure for jaundice (hepatitis), and a tea made from seeds could be imbibed as a treatment for headache, stomachache, or fatigue.

An extract of sicklepod tops inhibited germination of cotton and oats (*Avena sativa* L.); the germination of cotton seed was also reduced when sicklepod residues were incorporated into soil (Creel et al. 1968). Additionally, sicklepod appeared to release a phytotoxic residue from its roots that reduced the growth of cotton (Creel et al. 1968). Ground whole sicklepod seed meal incorporated into soil at 5% had no adverse effects on tomato plants but populations of *Meloidogyne chitwoodi* (Columbia root knot nematode) were reduced (Abbott et al. 1998). An extract from sicklepod has recently been investigated as a possible deer repellant in soybean and for insecticidal activity against soybean looper [*Chrysodeixis includens* Walker; Yue et al. 2020].

Sicklepod is tolerant to anthracnose (*Colletotrichum dematium* f. *truncate*) but is susceptible to seedling blight caused by *Alternaria cassia* (A. M. M. Juriar & A. Khan), which was investigated as a mycoherbicide (Teem et al. 1980; Walker 1982, 1983; Walker and Riley 1982). *Alternaria alternata* (Fr.) Keissl. and *Pseudocercospora nigricans* (Cooke) Deighton were also identified as suitable pathogens for further study for biocontrol of sicklepod (Hofmeister 1987; Mello et al. 2001). A beetle in the Bruchidae family, *Sennius instabilis*, feeds on *S. obtusifolia* seeds in tropical countries in the Western Hemisphere and could be considered for biological control of weedy *Senna* species elsewhere (Cock and Evans 1984).

Sicklepod may serve as an alternate host to some crop pests. It was identified as an excellent host for reniform nematode [Rotylenchulus reniformis (Linford and Oliveira)] in Georgia, Alabama, and the Mississippi Delta region with populations greater than or equal to those developing on susceptible cotton plants (Davis and Webster 2005; Lawrence et al. 2008; Molin and Stetina 2016). The ability of sicklepod to sustain root-knot nematode (Meloidogyne spp.) populations was also reported (Singh et al. 2010). Cowpea curculio (Chalcodermus aeneus Boheman), is an important pest of cowpea [Vigna unguiculata (L.) Walp.] in the southeastern United States, and adult insects were observed feeding on sicklepod pods and stems during the cowpea growing season, although no eggs or larvae were found in pods (Sudbrink et al. 1998). The whitefly, Bemisia tabaci (Gennadius), damaged tomato in Florida. Smith et al. (2014) found oviposition on S. obtusifolia supporting high B. tabaci populations. Sicklepod extrafloral nectaries attract ants primarily, but also some species of wasps, flies, and small bees (Hilty 2018; Marazzi et al. 2006). The pollen attracts honeybees, bumblebees, miner bees, and leaf-cutting bees (Hilty 2018; Retzinger 1984). The caterpillars of several species of sulphur butterflies have been observed to feed on foliage of Senna species, although it is unclear whether sicklepod is one of them (Hilty 2018). Sicklepod has no wildlife value due to its toxicity (Miller and Miller 1999).

Sicklepod has been the subject of extensive studies in the United States, despite not having a history of evolving herbicide resistance. This in part speaks to it distribution, toxicity, and competitiveness with important agronomic crops. Additionally, sicklepod possesses intrinsic characteristics that make it interesting to study, among these are its sleep movements. Good night!

References

- Abbott TP, Vaughn SF, Dowd PF, Mojtahedi H, Wilson RF (1998) Potential uses of sicklepod (*Cassia obtusifolia*). Industrial Crops Prod 8:77–82
- Andersen RN, Koukkari WL (1979) Rhythmic leaf movements of some common weeds. Weed Sci 27:401–415
- Bararpour MT, Oliver LR (1998) Effect of tillage and interference on common cocklebur (*Xanthium strumarium*) and sicklepod (*Senna obtusifolia*) population, seed production and seedbank. Weed Sci 16:424–431
- Becker B (1983) The contribution of wild plants to human nutrition in the Ferlo (Northern Senegal). Agroforest Syst 1:257–267
- Bentham G (1871) Revision of the genus *Cassia*. Trans Linnean Soc London 27:503–591
- Bosworth SC, Hoveland CS, Buchanan GA, Anthony WB (1980) Forage quality of selected warm-season weed species. Agron J 72:1050–1054

- Bozsa RC, Oliver LR, Driver TL (1989) Intraspecific and interspecific sicklepod (*Cassia obtusifolia*) interference. Weed Sci 37:670–673
- Brenan JPM (1958) New and noteworthy cassias from tropical Africa. Kew Bull 13:231–252
- Bridges DA, Walker RH (1985) Influence of weed management and cropping systems on sicklepod (*Cassia obtusifolia*) seed in the soil. Weed Sci 33: 800–804
- Bryson CT, DeFelice MS (2009) Weeds of the South. Athens, GA: University of Georgia. 468 p
- Buchanan GA, Burns ER (1971) Weed competition in cotton. I. Sicklepod and tall morningglory. Weed Sci 19:576–579
- Buchanan GA, Crowley RH, Street JE, McGuire JA (1980) Competition of sicklepod (*Cassia obtusifolia*) and redroot pigweed (*Amaranthus retroflexus*) with cotton (*Gossypium hirsutum*). Weed Sci 28:258–262
- Buchanan GA, Hauser EW (1980) Influence of row spacing on competitiveness and yield of peanuts (Arachis hypogaea). Weed Sci 28:401–409
- Burrows GE, Tyrl RJ (2006) Handbook of Toxic Plants of North America. Ames, IA: Blackwell Publishing. 307 p
- Christenhusz MJM, Byng JW (2016) The number of known plant species in the world and its annual increase. Phytotaxa 261:201–217
- Cock MJW, Evans HC (1984) Possibilities for biological control of *Cassia tora* and C. *obtusifolia*. Trop Pest Mgt 30:339–350
- Crawford L, McDonald GM, Friedman M (1990) Composition of sicklepod (Cassia obtusifolia) toxic weed seeds. J Agric Food Chem 38:2169–2175
- Creel JM Jr, Hoveland CS, Buchanan GA (1968) Germination, growth, and ecology of sicklepod. Weed Sci 16:396–400
- Cumming BG, Wagner E (1968) Rhythmic processes in plants. Ann Rev Plant Physiol 19:381–416
- Currey WL, Teem DH, Jordan JH (1981) Sicklepod competition and control programs in Florida soybeans. Page 66 *in* Proceedings of the Southern Weed Science Society Annual Meeting. Dallas, Texas, January 20–21, 1981
- Davis RF, Webster TM (2005) Relative host status of selected weeds and crops for Meloidogyne incognita and Rotylenchulus reniformis. J Cotton Sci 9:41–46
- Dirar HA (1984) Kawal, meat substitute from fermented *Cassia obtusifolia* leaves. Econ Bot 38:342–349
- Egley GH, Chandler JM (1978) Germination and viability of weed seeds after 2.5 years in a 50-year buried seed study. Weed Sci 26:230–239
- Egley GH, Chandler JM (1983) Germination and viability of weed seeds after 5.5 years in the Stoneville 50-year buried seed study. Weed Sci 31:264–270
- Glaze NC, Mullinix BG Jr (1984) Competitive effects of sicklepod on lima beans. Weed Sci 32:1–3
- Gleason HA, Cronquist A (1963) Manual of Vascular Plants of Northeastern United States and Adjacent Canada. Boston: Willard Grant Press. 810 p
- Hauser EW, Buchanan GA, Ethredge WJ (1975) Competition of Florida beggarweed and sicklepod with peanuts: I. Effects of periods of weed-free maintenance or weed competition. Weed Sci 23:368–372
- Hauser EW, Buchanan GA (1981) Influence of row spacing, seeding rates and herbicide systems on the competitiveness and yield of peanut. Peanut Sci 8:74–81
- Hauser EW, Buchanan GA, Nichols RL, Patterson RM (1982) Effects of Florida beggarweed (*Desmodium tortuosum*) and sicklepod (*Cassia obtusifolia*) on peanut (*Arachis hypogaea*) yield. Weed Sci 30:602–604
- Hilty J (2018) Illinois wildflowers. Last updated October 24, 2018. https://www. illinoiswildflowers.info/prairie/plantx/sicklepodex.htm. Accessed: January 22, 2021
- Hofmeister FM (1987) *Pseudocercospora nigricans*, a pathogen of sicklepod (*Cassia obtusifolia*) with biocontrol potential. Plant Dis 71:44
- Holm LG, Pancho JV, Herberger JP, Plunknett DL (1979) A Geographical Atlas of World Weeds. Malabar, FL: Krieger Publishing. 391 p
- Holm L, Doll J, Holm E, Pancho J, Herberger J (1997) World Weeds; Natural Histories and Distribution. New York: John Wiley & Sons 1,129 p
- Hoveland CS, Buchanan GA (1973) Weed seed germination under simulated drought. Weed Sci 21:322–324
- Hoveland CS, Buchanan GA, Harris MC (1976) Response of weeds to soil phosphorous and potassium. Weed Sci 24:194–201
- Irwin HS, Barneby RC (1982) The American Cassiinae. New York: New York Botanical Garden 35:1–918

- Irwin HS, Turner BL (1960) Chromosomal relationships and taxonomic considerations in the genus *Cassia*. Am J Bot 47:309–318
- ITIS (2021) Integrated Taxonomic Information System. *S. occidentalis*. https:// www.itis.gov/servlet/SingleRpt/SingleRpt#null. Accessed: February 26, 2021
- Jones RE, Jr, Walker RH (1993) Effect of interspecific interference, light intensity, and soil moisture on soybean (*Glycine max*), common cocklebur (*Xanthium strumarium*), and sicklepod (*Cassia obtusifolia*) water uptake. Weed Sci 41:534–540
- Kraatz GW, Andersen RN (1980) Leaf movements in sicklepod (*Cassia obtu-sifolia*) in relation to herbicide response. Weed Sci 28:551–556
- Lawrence KS, Price AJ, Lawrence GW, Jones JR, Akridge JR (2008) Weed hosts for *Rotylenchulus reniformis* in cotton fields rotated with corn in the southeast of the United States. Nematropica 40:275–288
- Mabberley DJ (1997) The Plant-Book. Second ed., Cambridge, UK: Cambridge University Press. 858 p
- Marazzi B, Endress PK, Paganucci De Queiroz L, Conti E (2006) Phylogenetic relationships within *Senna* (Leguminosae, Cassiinae) based on three chloroplast DNA regions: patterns in evolution of floral symmetry and extrafloral nectaries. Am J Bot 93:288–303
- McCormack JE, Neisler WE (1980) *Cassia obtusifolia* (sicklepod) toxicity in a dairy herd. Vet Med Small Anim Clin 75:1849–1851
- McWhorter CG, Sciumbato GL (1988) Effects of row spacing, benomyl, and duration of sicklepod (*Cassia obtusifolia*) interference on soybean (*Glycine max*) yields. Weed Sci 36:254–259
- Mello SCM, Ribeiro ZMA, Sousa GR, Tigano M, Nachtigal GDF, Fontes EMG (2001) Isoenzyme patterns and morphology of isolates of *Alternaria* species pathogenic to *Senna obtusifolia*. Fitopatologia Brasileira 26:667–669
- Miller JH, Miller KV (1999) Forest Plants of the Southeast and Their Wildlife Uses. Athens: University of Georgia Press. 464 p
- Molin WT, Stetina SR (2016) Weed hosts and relative weed and cover crop susceptibility to *Rotylenchulus reniformis* in the Mississippi Delta. Nematropica 46:121–131
- Monks DW, Oliver LR (1988) Interactions between soybean (*Glycine max*) cultivars and selected weeds. Weed Sci 36:770–774
- Norsworthy JK, Oliveira MJ (2006) Sicklepod (*Senna obtusifolia*) germination and emergence as affected by environmental factors and seeding depth. Weed Sci 54:903–909
- Norsworthy JK, Oliveira MJ (2007) Effect of tillage and soyabean on *Ipomoea* lacumosa and Senna obtusifolia emergence. Weed Res 47:499–508
- Nowick E (2015) Historical Common Names of Great Plains Plants with Scientific Names Index. Lincoln: University of Nebraska Lincoln Libraries. 452 p
- Page RK, Vezey S, Charles OW, Hollifield T (1977) Effects on feed consumption and egg production of coffee bean seed (*Cassia obtusifolia*) fed to white leghorn chickens. Avian Dis 21:90–96
- Patterson DT (1993) Effects of temperature and photoperiod on growth and development of sicklepod (*Cassia obtusifolia*). Weed Sci 41:574–582
- Patterson DT, Flint EP (1983) Comparative water relations, photosynthesis, and growth of soybean (*Glycine max*) and seven associated weeds. Weed Sci 31:318–323
- Putnam MR, Boosinger T, Spano J, Wright J, Wiggins A (1988) Evaluation of *Cassia obtusifolia* (sicklepod) seed consumption in Holstein calves. Vet Hum Toxicol 30:316–318
- Randell BR (1988) Revision of the Cassiinae in Australia. I. Senna sect. Chamaefistula. J Adelaide Bot Gard 11:19–49
- Randell BR (1995) Taxonomy and evolution of Senna obtusifolia and S. tora. J Adelaide Bot Gard 16:55–58
- Retzinger EJ Jr (1984) Growth and development of sicklepod (*Cassia obtusifo-lia*) selections. Weed Sci 32:608–611
- Robertson KR, Lee YT (1976) The genera of Caesalpinioideae (Leguminosae) in the southeastern United States. J Arnold Arbor 57:1–53
- Schwartz-Lazaro LM, Norsworthy JK (2018) Research Series Arkansas Agricultural Experiment Station, Issue 648, pp 80–82 https://agcomm. uark.edu/agnews/publications/648_Arkansas_Soybean_Research_Studies_ 2016.pdf. Accessed: January 30, 2021

- Senseman SA, Oliver LA (1993) Flowering patterns, seed production, and somatic polymorphism of 3 weed species. Weed Sci 41:418–425
- Shibles RM, Anderson IC, Gibson AH (1975) Soybean. Pages 151–189 in LY Evans, ed. Crop Physiology: Some Case Histories. London: Cambridge University Press
- Shurtleff JL, Coble HD (1985) Interference of certain broadleaf weed species in soybeans (*Glycine max*). Weed Sci 33:654–657
- Sims BD, Oliver LR (1990) Mutual influences of seedling johnsongrass (Sorghum halapense), sicklepod (Cassia obtusifolia), and soybean (Glycine max). Weed Sci 38:139–147
- Singh SK, Khurma UR, Lockhart PJ (2010) Weed hosts of root-knot nematodes and their distribution in Fiji. Weed Technol 24:607–612
- Smith HA, Nagle CA, Evans GA (2014) Densities of eggs and nymphs and percent parasitism of *Bemisia tabaci* (Hemiptera:Aleyrodidae) on common weeds in West Central Florida. Insects 5:860–876
- Staples GW, Imada CT, Herbs DR (2003) New Hawaiian plant records for 2001. Bishop Mus Occasional Papers 74:7–21
- Street JE, Buchanan GA, Crowley RH, McGuire JA (1981) Influence of cotton (Gossypium hirsutum) densities on competitiveness of pigweed (Amaranthus spp.) and sicklepod (Cassia obtusifolia). Weed Sci 29:253–256
- Sudbrink DL, Mack TP, Zehnder GW (1998) Alternate host plants of cowpea curculio, (Coleoptera: Curculionidae) in Alabama. Florida Entomologist 81:373–383
- Teem DH, Hoveland CS, Buchanan GA (1974) Primary root elongation of three weed species. Weed Sci 22:47–50
- Teem DH, Hoveland CS, Buchanan GA (1980) Sicklepod (*Cassia obtusifolia*) and coffee senna (*Cassia occidentalis*): Geographic distribution, germination, and emergence. Weed Sci 28:68–71
- Thurlow DL, Buchanan GA (1972) Competition of sicklepod with soybeans. Weed Sci 20:379–384
- Turner BC, Karlander EP (1975) Photoperiodic control of floral initiation in sicklepod (*Cassia obtusifolia* L.). Bot Gaz 136:1-4
- Upadhyaya SK, Singh V (1986) Phytochemical evaluation of *Cassia obtusifolia* L. and *Cassia tora* L. Proc Indian Acad Sci Plant Sci 96:321–326
- USDA (2020) The Plants database. *Senna obtusifolia*. Last updated October 29, 2020. https://plants.sc.egov.usda.gov/core/profile?symbol=SEOB4. Accessed: February 20, 2021
- Vencill WK, Banks PA (1994) Effects of tillage systems and weed management on weed populations in grain sorghum. Weed Sci 42:541–547
- Walker HL (1983) Biocontrol of sicklepod with Alternaria cassiae. Page 139 in Proceedings of the Southern Weed Science Society Annual Meeting. Biloxi, Mississippi: January 18–20, 1983
- Walker HL (1982) Seedling blight of sicklepod caused by *Alternaria cassia*. Plant Dis 66:426–428
- Walker HL, Riley JA (1982) Evaluation of *Alternaria cassiae* for the biocontrol of sicklepod (*Cassia obtusifolia*). Weed Sci 30:651–654
- Webster TM, MacDonald GE (2001) A survey of weeds in various crops in Georgia. Weed Technol 15:771-790
- Wright SR, Jennette MW, Coble HD, Rufty TW Jr (1999) Root morphology of young *Glycine max, Senna obtusifolia*, and *Amaranthus palmeri*. Weed Sci 47:706–711
- [WSSA] Weed Science Society of America (2021a) Composite list of weeds, *S. obtusifolia*. https://wssa.net/wssa/weed/composite-list-ofweeds/?commonname=&latinname=senna+obtusifolia&bayercode= &uscode=&search=Search. Accessed: January 19, 2021
- [WSSA] Weed Science Society of America (2021b) Composite list of weeds, S. occidentalis. https://wssa.net/wssa/weed/composite-list-ofweeds/?commonname&latinname=senna%20occidentalis&bayercode& uscode&search=Search&last=126. Accessed: January 19, 2021
- Yue Z, Tseng T, Krishnan N (2020) Antifeedant effects of sicklepod (Senna obtusifolia) extract on soybean looper Chrysodeixis includens (Lepidoptera: Noctuidae). Page 198 in Proceedings of the Southern Weed Science Society Annual Meeting. Biloxi, Mississippi, January 27–30, 2020
- Zomlefer WB (1994) Guide to Flowering Plant Families. Chapel Hill, NC: University of North Carolina Press. 430 p