

## **Self Reported Hearing Impairments and Associated Risk Factors Among Metal and Woodwork Workers in Gondar Town, North West Ethiopia**

Authors: Worede, Eshetu Abera, Yalew, Walelegn Worku, and Wami, Sintayehu Daba

Source: Environmental Health Insights, 16(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/11786302221084868>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Self Reported Hearing Impairments and Associated Risk Factors Among Metal and Woodwork Workers in Gondar Town, North West Ethiopia

Environmental Health Insights  
Volume 16: 1–7  
© The Author(s) 2022  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/11786302221084868



Eshetu Abera Worede<sup>ib</sup>, Walelegn Worku Yalew and Sintayehu Daba Wami

Department of Environmental and Occupational Health and Safety, Institute of Public Health, University of Gondar, Gondar, Ethiopia.

## ABSTRACT

**BACKGROUND:** The global prevalence of occupational noise-induced hearing loss ranges between 16% and 24%. The wood and metal-work industries have recently expanded in Ethiopia. This study aims to determine the level of noise exposure and the prevalence of self-reported hearing impairments and associated risk factors among metal and woodworkers in Gondar town Ethiopia.

**MATERIAL AND METHODS:** An institutional-based cross-sectional study was conducted on 580 metal and woodwork workers from February 10 to March 25/2020. The data were collected through an interviewer-led questioner and the noise level measurement. Multivariate Poisson regression models were used. *P*-values less than .05 and adjusted prevalence ratios with 95%CI were used to declare the presence and strength of an association respectively.

**RESULT:** The mean (SD) average noise exposure level in the wood and metalworking industries was  $96.9 \pm 3.5$  dBA and  $96.2 \pm 4$  dBA, respectively. The overall prevalence of self-reported hearing impairment was 20.7% [95%CI: (17.4-24)]. In an adjusted Poisson regression, listening to music with earphones for more than 2 hours per day (PR = 2.95, 95%CI: 1.32, 6.21) and listening to music at maximum volume (PR = 2.24, 95%CI: 1.05, 4.79) were associated with hearing impairments.

**CONCLUSION:** The majority of workers are exposed to noise levels that exceed OSHA's permissible exposure limit value. A hearing conservation program should be implemented to reduce noise exposure levels in the wood and metal work industries. Workers should be aware of the duration and volume of recreational noise exposure.

**KEYWORDS:** Hearing impairment, noise exposure, risk factor, listening to music

**RECEIVED:** November 17, 2021. **ACCEPTED:** February 2, 2022.

**TYPE:** Learning from Failure in Environmental and Public Health Research – Original Research

**FUNDING:** The author(s) received no financial support for the research, authorship, and/or publication of this article.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**CORRESPONDING AUTHOR:** Eshetu Abera Worede, Department of Environmental and Occupational Health and Safety, Institute of Public Health, University of Gondar, Gondar 196, Ethiopia. Email: aberaeshetu44@gmail.com

## Introduction

Hearing loss affects 1.3 billion people worldwide<sup>1</sup> and is the fourth leading cause of disability, with an estimated annual cost of more than 750 billion dollars.<sup>2</sup> In the United States, 22 to 30 million workers are potentially exposed to noise levels from both occupational and non-occupational sources.<sup>2</sup> According to a global hearing report, 1.1 billion young people are at risk of permanent hearing loss from listening to music at high volumes for extended periods of time.<sup>3</sup>


The metal and woodworking factories have a high level of noise exposure.<sup>4-9</sup> Occupational noise exposure has been documented since the 18th century when copper miners developed hearing loss as a result of noise from hammering on metal.<sup>1,10</sup> Hearing loss caused by work-related noise exposure is known as occupational noise-induced hearing loss (NIHL),<sup>6,8,11</sup> and its global prevalence is estimated to be 16% to 24%.<sup>11,12</sup> Occupational noise-induced hearing loss is the second-most common self-reported occupational injury or illness, accounting for 7% and 21% of all reported injuries or illnesses in developed and developing countries, respectively.<sup>2,13</sup> Over 4 million disability-adjusted

life years (DALYs) are attributed to ONIHL.<sup>14</sup> Noise-induced hearing loss (NIHL) is the most common form of occupational disease in Malaysia, accounting for 78.1% of all diseases reported in Malaysian industries in 2013 and 2015.<sup>15</sup>

Approximately 600 million workers globally are exposed to occupational noise.<sup>16</sup> In the United States, the prevalence of workplace noise exposure was highest in mining (76%), followed by woodworking factories (55%).<sup>11</sup> All mean noise levels in all studied metalwork factories and 50% of studied woodwork industries in Saudi Arabia were higher than the standard level of 85 dBA.<sup>6</sup> The average noise level in Greek wood industries was found to be above the acceptable limit values.<sup>17</sup> In Nepal, 30.4% of metalworkers developed noise-induced hearing loss (NIHL),<sup>18</sup> and in the Southeast Asian furniture industry, 34.7% respondents experienced permanent threshold shifts.<sup>8</sup> Noise-induced hearing loss (NIHL) is 1 of the top 5 occupational illnesses in Zimbabwe, and the mining industry has a high prevalence of NIHL due to excessive noise levels.<sup>19</sup> 44% of study participants in Ghana's quarry industry had a hearing threshold greater than 25 dBA.<sup>20</sup>



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without



**Figure 1.** Schematic presentation of sampling procedure both metal and woodwork.

According to different research, factors which influence the occurrence of hearing impairments include loud sounds<sup>2,21</sup> with duration of exposure,<sup>8,22</sup> gender,<sup>23</sup> age,<sup>18,24</sup> use of hearing protective devices,<sup>20</sup> smoking, and alcohol.<sup>18,24,25</sup> Prior Noise exposure from garage (a repair shop for automotive vehicles), construction, armed services,<sup>26,27</sup> patient-related factors like family history of hearing loss, ear infection, and injury,<sup>24,28</sup> ototoxic medicines,<sup>29</sup> and vibration.<sup>14,29,30</sup>

Industrialization poses a public health risk throughout Sub-Saharan Africa, including Ethiopia.<sup>18</sup> Despite the fact that the number of metal and woodworking factories in Ethiopia is growing, the level of noise exposure remains unknown, particularly in the study area. This study is designed to determine the noise exposure level, the prevalence of self-reported hearing impairments, and associated risk factors among metal and woodworkers in Gondar town.

## Materials and Methods

### Study design, settings, and period

An institutional-based cross-sectional study was conducted on 580 metal and woodwork workers from February 10 to March 25/2020 in Gondar town. Gondar is the capital city of the central Gondar zone in the Amhara regional state and one of Ethiopia's historical towns.

### Source and study populations

All metal and woodworkers in Gondar town were the study's source population, and workers who had worked for at least 6 months were included.<sup>31</sup>

### Sample size determination

The sample size was done for both first and second objectives. Assumptions for the first objective was 95%CI, 30.4% prevalence of hearing impairment among metalwork worker,<sup>18</sup> and 4% margin of error and for the second objective 95%CI, and power 80% and factors that have strong significant relation

with hearing impairment.<sup>16,23</sup> After adding a 10% non-response rate, sample size for the first and second objectives was 599 and 644 respectively.

### Sampling procedure

As the total sample size of this study is close to the entire target population, a survey sampling procedure Figure 1 was used to select study participants.

### Data collection tool and procedure

An interviewer-led structured questionnaire was used to collect data on socio-demographic characteristics and risk factors such as current and previous occupational noise exposure, work experience, behavioral factors such as listening to music, drinking alcohol, using hearing protective devices, and patient-related factors such as a family history of hearing loss, ear infection, ear injury, and signs and symptoms of noise-induced hearing impairments. The level of workplace noise exposure was measured using a sound level meter (IEC 651, type II, Taiwan), which is recommended for field measurements due to its precision or provides a scale for noise level as perceived by the human ear.<sup>32</sup> Sound Pressure Level (SPL) measurements were taken at workers' head level over 15 minutes at one-minute intervals, and this noise level represented workers' exposure.<sup>6</sup> The average sound pressure level was calculated using the logarithmic formula shown below.

$$\text{Average LP} = \frac{10 \log_{10} \left[ \frac{10^{LP1/10} + 10^{LP2/10} + 10^{LP3/10} + \dots + 10^{LP15/10}}{15} \right]}{15}$$

**Operational definitions.** Hearing impairment was defined by using signs and symptoms of noise-induced hearing loss such as difficulty hearing when people speak, difficulty understanding conversation, tinnitus, and workers who showed at least one of these signs and symptoms were considered to have hearing impairments.<sup>10,14,22,33</sup>

**Ear infection:** This was ascertained by the history of ear infection under the age of 18 years and in this regard certain viral infections in the inner ear destroy the cochlea, producing total deafness.<sup>31</sup>

#### *Data quality control*

To assure the quality of data, careful design, translation, and retranslation of the questionnaire were done. Pre-test was conducted on 5% of the sample population from other sites of similar industries. Two-day training was also given for data collectors. Proper categorization and coding of the data were done. Reliability of the questioner was assessed and its Cronbach's alpha value was .71.

#### *Data management and analysis*

The data was entered into the EPI Info version 7 software and exported to SPSS Version 20 software for further analysis. The mean, standard deviation (SD), and average mean noise exposure levels were computed as descriptive statistics. Multivariate Poisson regression models with robust variance were employed to estimate the prevalence ratios (PR) and corresponding 95% confidence intervals (CI).<sup>34</sup> To control for potential confounder variables, independent variables with *P*-values less than .25 in bivariable analysis were included in multivariate Poisson regression model. To declare the presence and strength of an association, *P*-values less than .05 and adjusted prevalence ratios with 95%CI were used.

## Results

### *Socio-demographic characteristics of study participants*

From a total of 626 eligible workers, 580 respondents with a response rate of 92.7% were fully participated in the study. The mean age of respondents was 26.32 ( $\pm 7.32$ ) years old. 53.8% of study participants worked in the woodwork industries, and nearly three-fourths (72%) were exposed to noise levels greater than 95dBA. Half of the participants (50.9%) had less than 3 years of work experience (Table 1).

### *Behavioral characteristics of study participants*

More than three-quarters (80.9%) of study participants had music listening habits. Only 6.9% of respondents used hearing protection devices. The majority of respondents reported a lack of provision and comfort issues as the main reasons for non-utilization of hearing protective devices respectively.

### *Comparative noise exposure level between wood and metalwork industries*

The average noise exposure level in the wood and metalworking industries was  $96.9 \pm 3.5$  dBA and  $96.2 \pm 4$  dBA

respectively. In the woodworking industry, the maximum noise level was 114dBA (circular saw), and in the metalworking industry, the maximum noise level was 108dBA (cutters and welding machines). There was a significant difference in sound level between the metal and woodworking industries ( $\chi^2 = 15.1$ , *df* = 4, *P* = .005).

### *The prevalence of hearing impairment among the wood and metalwork industry*

In this study, the overall prevalence of hearing impairment among wood and metalworkers was 20.7% [95%CI: 17.4%-24%]. Communication difficulties and tinnitus were reported by 32.9% and 26.4% of total study participants, respectively. Furthermore, 16.4% and 8.4% of workers reported difficulty hearing in the left and right ear without the use of a hearing aid, respectively.

*Prevalence of hearing impairments in terms of industry.* According to this study, the prevalence of hearing impairment was comparable among woodworker (20.8%) and metalworkers (20.5%). There is no statistically significant difference in hearing impairments between the wood and metalworking industries ( $\chi^2 = 0.008$ , *df* = 1, *P*-value = .5) (Table 2).

### *Factors associated with hearing impairments*

In adjusted Poisson regression analysis, listening to music at maximum volume and listening to music using earphones for more than 2 hours per day were significantly associated with hearing impairments. Wood and metalworkers who listen to music at maximum volume were 2.24 times more likely to have hearing impairments compared to workers who listen to music at lower volume (PR = 2.24; 95%CI: 1.05, 4.79), and workers who listen to music using earphones for more than 2 hours per day were 2.95 times more likely to have hearing impairments compared to their counterparts (PR = 2.95, 95%CI: 1.32, 6.21) (Table 3).

## Discussion

According to this study, the prevalence of hearing impairment was comparable among metal and woodwork workers. The overall prevalence of self-reported hearing impairment was 20.7% [95%CI: 17.4%-24%]. This finding is consistent with studies conducted in the United States woodwork industry (20.86%),<sup>23</sup> Ethiopia's metalwork industry (22%),<sup>35</sup> and the South Thailand sawmill industry (22.8%).<sup>36</sup> This similarity could be attributed to similar methods used (cross-sectional and retrospective cross-sectional in the case of the US) and worker characteristics. However, the findings of this study are lower than those of previous studies in Rwanda's wood and metalwork industries (36%),<sup>5</sup> and Nepal's woodwork industry (31%).<sup>8</sup> This difference could be attributed to differences in methods used (audiogram tests in Rwanda may have increased the prevalence), operational definition of hearing impairments, use of hearing protective devices (0.5% in Rwanda and 6.6% in

**Table 1.** Socio-demographic characteristics of wood and metalwork workers in Gondar Town (n=580).

VARIABLES	CATEGORY	FREQUENCY	PERCENT
Sex	Male	543	93.6
	Female	37	6.4
Age	15-24	265	46.2
	25-34	234	40.3
	35-44	59	10.2
	>44	22	3.8
Religion	Orthodox Christian	476	82.1
	Muslim	96	16.2
	Protestant	6	1
	Others	2	0.3
Marital status	Single	432	74.5
	Married	148	25.5
Educational level	Primary education	115	19.8
	Secondary education	304	52.4
	Diploma and certificate	138	23.8
	Degree and above	23	4
Work experience in current occupation (y)	1-3	295	50.9
	4-8	190	32.8
	9-12	63	10.3
	>12	32	5.5
Types of occupation	Woodwork	312	53.8
	Metalwork	268	46.2
Noise exposure level	<95 dBA	162	27.9
	95.0-99.0 dBA	276	47.6
	99.00-100 dBA	52	9
	>100dBA	90	15.5

Others; 7th day Adventist, Hawariyaw.

**Table 2.** Noise-induced hearing impairment and work categories.

HEARING IMPAIRMENT	WORK CATEGORIES			P-VALUE
	WOODWORK	METALWORK	TOTAL	
No	247 (79.2%)	213 (79.5%)	460 (79.3%)	>.05
Yes	65 (20.8%)	55 (20.5%)	120 (20.7%)	

our study), and duration of exposure or work experience, as well as worker characteristics. This finding, on the other hand, is higher than that of a study conducted in Brazil Metalworking

Company (15.9%),<sup>37</sup> and this difference could be attributed to high noise exposure levels and a lack of enforcement of occupational health and safety regulations in Ethiopia.

**Table 3.** Prevalence ratios of factors associated with self reported hearing impairments among wood and metalwork workers in Gondar town, northwest Ethiopia (n=580).

VARIABLES	PREVALENCE RATIO (PR)	95%CI		P-VALUE
		LOWER	UPPER	
Intercept	0.07	0.015	0.32	.001
Educational level				
Primary education	1.25	0.25	6.21	.78
Secondary education	1.57	0.78	3.17	.2
Diploma and certificate	0.83	0.46	1.5	.53
Degree and above	1			
Monthly salary				
<1500	0.49	0.23	1.65	.07
1500-2000	0.82	0.44	1.53	.54
2001-3200	0.67	0.34	1.34	.26
>3200	1			
Noise exposure level				
<95 dBA	1			
95.0-99 dBA	1.0.4	0.48	2.63	.7
99.0-100 dBA	1.13	0.48	23	.9
>100 dBA	0.85	0.36	2.03	.7
Use of ear protective device				
No	2.07	0.52	8.31	.302
Yes	1			
Volume of music listening				
Quite	1			
Moderate	1.76	0.94	3.31	.08
Maximum volume	2.24	1.05	4.79	.037
Duration of using earphone to listen music				
<-2 h	1			
>2 h	2.95	1.32	6.21	.008
Involve in dance concert				
Yes	1.49	0.82	2.74	.19
No	1			
Ear infection under age of 18				
Yes	1.93	0.97	3.83	.06
No	1			

1 = Reference group, Model fittest ( $P = .82$ ).

In this study, 16% and 8.4% of study participants reported difficulty hearing in the left and right ear without the use of a hearing aid, respectively. According to this finding, the left ear is more affected than the right ear, and this result is supported by various studies, such as Iran,<sup>38</sup> the Swedish wood processing industry,<sup>39</sup> New York,<sup>40</sup> Iran,<sup>41</sup> the United States of Louisiana,<sup>42</sup> and air force pilots.<sup>43</sup> This similarity could be due to noise shielding in one ear, unequal recovery after severe noise exposure, and unequal sensitivity of the ears and direction of noise exposure.<sup>44</sup>

In this study, listening to music with earphones was significantly associated with hearing impairments, and workers who listened to music with earphones more than 2 hours per day were 2.95 times more likely to have hearing impairments compared to their counterparts. This finding is consistent with a study in Singapore, where 1 in every 6 young people is at risk of developing leisure NIHL from music delivered via earphones,<sup>45</sup> and in Taiwan, listening music through headphones for 3 hours at maximum level showed transient shifts of 10 and 30 dB and returned to normal within 24 hours and can cause of noise-induced hearing impairments.<sup>46</sup> Workers who listened to music at maximum volume were 2.24 times more likely to have hearing impairments than workers who listened to music at low volume. This finding is consistent with a study that found that exposure to loud leisure noise is associated with hearing loss and tinnitus, with the risk increasing as noise exposure increases,<sup>47</sup> and another study done among young people found that listening too loudly for an extended period of time on personal listening devices (PLDs) such as CDs, iPods, and other MP3 players is a potential contributor to NIHL.<sup>48</sup>

In this study, one of the variables of interest was noise exposure level in the wood and metalwork industries, but found insignificant factors for noise-induced hearing impairments. This finding is consistent with the findings of a study conducted in Ethiopia's metalworking industry.<sup>35</sup> However, in other studies, the noise level was identified as a significant risk factor for NIHL.<sup>16,49,50</sup> This disparity could be attributed to a young labor force with limited work experience. According to the findings of this study, nearly three-fourths (72%) of respondents were exposed to average noise levels greater than 95 dBA, which is above the OSHA permissible exposure limit value for 8 hours of working time.<sup>51</sup> Similar findings were found in Rwanda's wood (99.4 dBA) and metalwork (105.4 dBA) industries, where 99.5% of all participants were not protected during work time.<sup>5</sup> Based on OSHA, workers exposed to noise exposures equal or exceed an 8-hour TWA of 85 dBA must be in a hearing conservation program comprised of exposure monitoring, audiometric testing, hearing protection, employee training, and record keeping.<sup>52</sup> This research found that workers are exposed to high average noise levels for an extended period of time (8 hours), despite the country having exposure limits to continuous noise at 90, 92, 95, 95, 97, 100, 102, 105, 110, 115 dBA to a period of 8, 6, 4, 3, 2, 1 and 12, 1, 12, 1/4, hours, respectively.<sup>53</sup>

## Limitation of the Study

The inability to use a noise dosimeter to measure personal noise exposure levels.

There is no audiogram test to assess the level of hearing loss.

There is no control group for comparisons.

## Conclusion

This study found that the prevalence of hearing impairment is comparable in the metal and woodwork industries, but the woodwork industry has a higher noise exposure level, and the majorities (72%) of employees in both industries are exposed to noise levels above the OSHA permissible exposure limit value. Listening to music with earphones for more than 2 hours per day, as well as listening to music at maximum volume, were found to be significant risk factors for hearing impairment. As a result, a hearing conservation program must be implemented in the woodworking and metalworking industries, and workers must be aware of the duration and volume of recreational noise exposure.

## Future research

Future research should include a noise dosimeter, an audiogram test, and a control group.

## Acknowledgements

First, I would like to thank my research advisors, Dr. Walelegn Worku (Ph.D., Assoc. pro) and Mr. Sintayehu Daba (MPH, Assis. pro) for their friendly support and involvement throughout the proposal development to research thesis work. Second, I would like to thank the University of Gondar and the Environmental and occupational safety and health department for every effort they exerted. I would like also to thank Gondar town industrial development office, owners, and study participants for their positive involvement during the data collection period. Last but not least, I would like to thank data collectors, supervisors, and my families and friends who have contributed directly or indirectly to the successful completion of the research.

## Author Contributions

Eshetu Abera conceived of the study and contributed to its design, data collection, data analysis, results interpretation, and manuscript writing. Dr. Walelegn worked on, commented on, and edited the statistical output interpretation. Mr. Sintayehu Daba contributed to data analysis, commented on and edited statistical output interpretation, wrote up the manuscript, and all authors approved the submitted version of the manuscript.

## Availability of Data and Materials

All data generated for this study are included in this article. The data are also available from the corresponding author upon reasonable request.

## Ethics Approval and Consent to Participate

The Ethical Review Committee of the Institute of Public Health, College of Medicine and Health Sciences, University of

Gondar, evaluated the ethical issue of this research and approved it as ethically sound research by the Rf No IPH/837/06/2020 and date 13/06/2020, and participants were informed about the purpose of the study, the importance of their participation, and their right to withdraw at any time, and written consent was obtained from each participant during data collection

## ORCID iD

Eshetu Abera Worede  <https://orcid.org/0000-0002-2939-8329>

## REFERENCES

- Stucken EZ, Hong RS. Noise-induced hearing loss: an occupational medicine perspective. *Curr Opin Otolaryngol Head Neck Surg*. 2014;22:388-393.
- World Health Organization. *Addressing the Rising Prevalence of Hearing Loss*. WHO; February 2018.
- World Health Organization. *World Report on Hearing*. World Health Organization; 2021.
- Brühl P, Grenner J. Impact noise exposure from sheet-metal presses: manual versus automated production. *Acta Acust United Acust*. 2000;86:378-380.
- Bukuru J, Byiringiro F, Mukara BK, et al. Prevalence of occupational noise induced hearing loss among wood and metal workers of Gakiriro, Kigali city. *Rw. Public Health Bul*. 2019;1:20-25.
- Noweir MH, Bafail AO, Jomoah IM. Noise pollution in metalwork and woodwork industries in the Kingdom of Saudi Arabia. *Int J Occup Saf Ergon*. 2014;20:661-670.
- Nyarubeli IP, Brätveit M, Tungu AM, Mamuya SH, Moen BE. Temporary threshold shifts among iron and steel factory workers in Tanzania: A pre-interventional study. *Ann Global Health*. 2021;87:35.
- Robinson T, Whittaker J, Acharya A, Singh D, Smith M. Prevalence of noise-induced hearing loss among woodworkers in Nepal: a pilot study. *Int J Occup Environ Health*. 2015;21:1-15.
- Zheng Y-P, Juang Y-J, Yiin L-M. Modeling of Woodworkers' exposure to occupational noises by integrating frequency spectra generated by power tools: a pilot study. *Appl Sci*. 2020;10:6453.
- Hong O, Kerr MJ, Poling GL, Dhar S. Understanding and preventing noise-induced hearing loss. *Dis Month*. 2013;59:110-118.
- Tak S, Davis RR, Calvert GM. Exposure to hazardous workplace noise and use of hearing protection devices among US workers—NHANES, 1999-2004. *Am J Ind Med*. 2009;52:358-371.
- Nelson DI, Nelson RY, Concha-Barrientos M, Fingerhut M. The global burden of occupational noise-induced hearing loss. *Am J Ind Med*. 2005;48:446-458.
- Reddy R, Welch D, Ameratunga S, Thorne P. Development of the hearing protection assessment (HPA-2) questionnaire. *Occup Med*. 2014;64:198-205.
- Garshick E, Segal MR, Worobec TG, Salekin CMS, Miller MJ. Alcohol consumption and chronic obstructive pulmonary disease. *Am Rev Respir Dis*. 1989;140:373-378.
- Wali NY. Influence of varying degree of wood dust exposure on pulmonary function and respiratory symptoms among wood workers in Kano, North Western Nigeria. *Niger J Physiol Sci*. 2020;35:161-165.
- Ferrite S, Santana V. Joint effects of smoking, noise exposure and age on hearing loss. *Occup Med*. 2005;55:48-53.
- Bougoudis I, Dimou V, Liadis L. Computational Intelligence modeling and assessment of industrial noise: the case of wood manufacturing. *Int J Eng Intell Syst Electr Eng Commun*. 2014.
- Whittaker JD, Robinson T, Acharya A, Singh D, Smith M. Noise-induced hearing loss in small-scale metal industry in Nepal. *J Laryngol Otol*. 2014;128:871-880.
- Chadambuka A, Mususa F, Muteti S. Prevalence of noise induced hearing loss among employees at a mining industry in Zimbabwe. *Afr Health Sci*. 2014;13:899-906.
- Gyamfi CK, Amankwaa I, Owusu Sekyere F, Boateng D. Noise exposure and hearing capabilities of quarry workers in Ghana: a cross-sectional study. *J Environ Public Health*. 2016;2016:7054276.
- Armitage CJ, Loughran MT, Munro KJ. Epidemiology of the extent of recreational noise exposure and hearing protection use: cross-sectional survey in a nationally representative UK adult population sample. *BMC Public Health*. 2020;20:1529.
- Penafel E. *Developing a Questionnaire to Assess Noise Exposure in Children and Teens*. The Ohio State University; 2007.
- Masterson EA, Tak S, Themann CL, et al. Prevalence of hearing loss in the United States by industry. *Am J Ind Med*. 2013;56:670-681.
- Daniel E. Noise and hearing loss: a review. *J Sch Health*. 2007;77:225-231.
- Le Prell CG, Henderson D, Fay RR, Popper AN. *Noise-Induced Hearing Loss*. Springer; 2011.
- Pawlaczyk-Iuszczynska M, Dudarewicz A, Czaja N, Bortkiewicz A. Do hearing threshold levels in workers of the furniture industry reflect their exposure to noise? *Med Pr*. 2016;67:337-351.
- World Health Organization. *Hearing Loss Due to Recreational Exposure to Loud Sounds: A Review*. WHO; 2015.
- Masilamani R, Rasib A, Darus A, Ting AS. Noise-induced hearing loss and associated factors among vector control workers in a Malaysian state. *Asia Pac J Public Health*. 2014;26:642-650.
- Turcot A, Girard SA, Courteau M, Baril J, Larocque R. Noise-induced hearing loss and combined noise and vibration exposure. *Occup Med*. 2015;65:238-244.
- Issever H, Aksoy C, Sabuncu H, Karan A. Vibration and its effects on the body. *Med Princ Pract*. 2003;12:34-38.
- Goelzer B, Hansen CH, Sehrndt G. *Occupational Exposure to Noise: Evaluation, Prevention and Control*. World Health Organisation; 2001.
- Campopiano A, Basili F, Angelosanto F, et al. Field comparison of two inhalable samplers used in Italy to measure the wood dust exposure. *Int J Occup Environ Health*. 2016;22:159-166.
- Muca A, Standafer E, Apawu AK, et al. Tinnitus and temporary hearing loss result in differential noise-induced spatial reorganization of brain activity. *Brain Struct Funct*. 2018;223:2343-2360.
- Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol*. 2003;3:21.
- Hailu A. *Assessment of Noise Induced Hearing Loss and Associated Factors Among Workers in Akaki Basic Metal Industry*. Addis Ababa, Ethiopia. Addis Ababa University; 2015.
- Thepaksorn P, Koizumi A, Harada K, Siri Wong W, Neitzel RL. Occupational noise exposure and hearing defects among sawmill workers in the south of Thailand. *Int J Occup Saf Ergon*. 2019;25:458-466.
- Guerra MR, Lourenço PM, Bustamante-Teixeira MT, Alves MJ. [Prevalence of noise-induced hearing loss in metallurgical company]. *Rev Saude Publica*. 2005;39:238-244.
- Jabbari K, Nassiri P, Monazzam Esmacelpour MR, Azam K, Faridan M, Heidari L. The relationship between occupational noise exposure and noise induced hearing loss (NIHL) in small-scale industries: a case study in the city of Damavand, Iran. *Biotechnol Health Sci*. 2016;3:49-56.
- Johansson M, Arlinger S. The development of noise-induced hearing loss in the Swedish County of Östergötland in the 1980s and 1990s. *Noise Health*. 2001;3:15-28.
- Marvel ME, Pratt DS, Marvel LH, Regan M, May JJ. Occupational hearing loss in New York dairy farmers. *Am J Ind Med*. 1991;20:517-531.
- Janghorbani M, Sheykhi A, Pourabdian S. The prevalence and correlates of hearing loss in drivers in Isfahan, Iran. *Arch Iran Med*. 2009;12:128-134.
- Carruth A, Robert AE, Hurley A, Currie PS. The impact of hearing impairment, perceptions and attitudes about hearing loss, and noise exposure risk patterns on hearing handicap among farm family members. *AAOHN J*. 2007;55:227-234.
- Muhr P, Johnson A-C, Selander J, Svensson E, Rosenhall U. Noise exposure and hearing impairment in air force pilots. *Aerosp Med Hum Perform*. 2019;90:757-763.
- Alberti PW, Symons F, Hyde ML. Occupational hearing loss the significance of asymmetrical hearing thresholds. *Acta Otolaryngol*. 1979;87:255-263.
- Lee J, Lim M, Kuan Y, Teo H, Tan H, Low W. The music listening preferences and habits of youths in Singapore and its relation to leisure noise-induced hearing loss. *Singapore Med J*. 2014;55:72-77.
- Tung C-Y, Chao K-P. Effect of recreational noise exposure on hearing impairment among teenage students. *Res Dev Disabil*. 2013;34:126-132.
- Holgers KM, Pettersson B. Noise exposure and subjective hearing symptoms among school children in Sweden. *Noise Health*. 2005;7:27-37.
- Levey S, Fligor BJ, Ginocchi C, Kagimbi L. The effects of noise-induced hearing loss on children and young adults. *Contemp Issues Commun Sci Disord*. 2012;39:76-83.
- Belachew A, Berhane Y. Noise-induced hearing loss among textile workers. *Ethiop J Health Dev*. 1999;13:69-75.
- Ologe FE, Akande TM, Olajide TG. Occupational noise exposure and sensorineural hearing loss among workers of a steel rolling mill. *Eur Arch Otorhinolaryngol*. 2006;263:618-621.
- Shaikh GH. Occupational noise exposure limits for developing countries. *Appl Acoust*. 1999;57:89-92.
- Plog BA, Quinlan PJ. *Fundamentals of Industrial Hygiene*. 5th ed. NSC Press; 2001.
- Molsa E. *Occupational Safety and Health Directive*. International Labour Office; 2008.