

## **Comments to Recent Studies Showing Systemic Mechanisms Enabling Drosophila Larvae to Recover From Stress-Induced Damages**

Author: Hayakawa, Yoichi

Source: International Journal of Insect Science, 10(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1179543318795894>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.


---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Comments to Recent Studies Showing Systemic Mechanisms Enabling *Drosophila* Larvae to Recover From Stress-Induced Damages

Yoichi Hayakawa

Department of Applied Biological Sciences, Saga University, Saga, Japan.

International Journal of Insect Science  
Volume 10: 1–3  
© The Author(s) 2018  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1179543318795894  


**ABSTRACT:** Compensational recovery from the damage created by stressors is important for all animals. However, how organisms recover from stress-induced negative impacts has been poorly understood. An 1-hour exposure to heat stress at 35°C led to reduced feeding activity of *Drosophila melanogaster* larvae, which caused reduction in body weight 2 hours after the stress, but not at other times. Such weight losses seem to be rescued by following enhanced feeding activities. We investigated the mechanisms underlying the accelerated feeding activity after the stress-induced reduction in feeding behavior. Our data showed increased expression of sweet taste gustatory receptor genes (*Grs*) and concomitant decreased expression of bitter taste *Grs* in the mouth parts 2 to 4 hours after the heat treatment for 1 hour. However, nontypical taste *Gr* expression was not changed. Furthermore, integration of both messenger RNA and protein expression analysis revealed that expression levels of tropomyosin and ATP (adenosine triphosphate) synthase  $\beta$  subunit were significantly increased in their mouths 3 to 5 hours after the heat stress. The increased expression of these genes would contribute to accelerated muscular movement of the mouth hooks. This study indicated that *Drosophila* larvae possess an efficient systemic mechanism that enables them to recover from growth delay caused by stress conditions.

**KEYWORDS:** Stress, recovery, gustatory receptors, tropomyosin, ATP synthase  $\beta$  subunit

**RECEIVED:** July 16, 2018. **ACCEPTED:** July 18, 2018.

**TYPE:** Commentary

**FUNDING:** The author(s) received no financial support for the research, authorship, and/or publication of this article.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**CORRESPONDING AUTHOR:** Yoichi Hayakawa, Department of Applied Biological Sciences, Saga University, Saga 840-8502, Japan. Email: hayakayo@cc.saga-u.ac.jp

In natural environments, wild animals seldom experience fulfillment of all their biotic and abiotic requirements and are routinely exposed to a variety of environmental stressors. To combat these stressors, they have evolved highly successful strategies.<sup>1,2</sup> Much of the field of stress biology has focused on multiple physiological changes of test organisms to reveal such strategies that enable them to survive under stressful environments. It is becoming apparent that various cellular and molecular mechanisms essential for stress resistance and tolerance occur in organisms. A representative example of the research is the finding that heat shock proteins (Hsps) are conserved in a wide variety of organisms not only in the ways they are induced by stress but also in their primary structures.<sup>3,4</sup> A prolonged production of unfolded or misfolded proteins triggers the expression of Hsps, which represents an essential cellular strategy in all organisms to deal with stress conditions such as high temperature.<sup>5,6</sup> Despite substantial attention to stress resistance mechanisms, the processes of the recovery phase after stress have not been actively explored yet. It is therefore poorly understood whether animals possess a specific mechanism to recover from the damage done by environmental stresses.

We recently investigated the effects of acute heat stress on feeding activities and body weights of *Drosophila melanogaster* larvae of the y w strain during the poststress period to test our hypothesis that animals possess systemic mechanisms to cope with and survive the damage caused by environmental stressors.<sup>7</sup> Feeding rates of *Drosophila* larvae were shown to decrease soon after the stress at 35°C for 1 hour, which caused reduction in body weight 2 hours after the stress. Thereafter, feeding


activities increased to higher levels compared with control larvae at 3 hours after the heat stress. This enhanced feeding action led to weight restoration by around 4 hours after heat stress. Those observations enabled us to surmise that *Drosophila* larvae possess a compensational recovery mechanism during poststress periods.

We examined the mechanism by which animals increase their feeding activity after the stress treatment (at 35°C for 1 hour). It is generally known that taste neurons expressing gustatory receptor genes (*Grs*) in insect gustatory organs detect tastants by direct interactions between *Grs* and ligands.<sup>8,9</sup> We expected that expression levels of *Grs* in mouths would be changed before and after the heat stress, which affects larval feeding activities. The results showed that expression of all sweet taste *Grs* such as *Gr5a*, *Gr64a*, *Gr64f*, and *Gr43a* was elevated at around 4 hours after the heat stress. By contrast, expression of bitter taste *Grs* such as *Gr66a* and *Gr33a* was depressed after the heat stress (Figure 1). No change was observed in expression of a nontypical taste *Gr*, *Gr68a*, sensing acoustic signals after the heat treatment<sup>10</sup> (Figure 1). Similar changes in expression levels of sweet taste *Gr5a* and bitter taste *Gr66a* genes were recorded in the mouths of test larvae after cold-stress treatment 4°C for 12 hours. We interpreted these results as an indication that enhanced expression of sweet taste *Grs* and depressed expression of bitter taste *Grs* would contribute to the increased feeding activities of *Drosophila* larvae.

Although we have not examined how expression of those *Grs* is controlled during the poststress period, we might take some hints from prior reports on starvation.<sup>11</sup> It is undoubted



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).  
Downloaded From: <https://complete.bioone.org/journals/International-Journal-of-Insect-Science> on 13 Jul 2025  
Terms of Use: <https://complete.bioone.org/terms-of-use>




**Figure 1.** Tendency curves of sweet taste, bitter taste, and nontypical taste *Gr* expression changes after heat stress at 35°C for 1 hour.

that starved animals have increased appetite; at that time, they must become less selective in their food choices. Increasing their sweet taste sensitivity enhances their appetite.<sup>12,13</sup> In *Drosophila*, it has been reported that hunger elevates gustatory sensitivity to sweet, in part, via enhanced dopamine release in *Gr5a*-expressing neurons, which rises calcium responses to *Gr* activation.<sup>14,15</sup> Hunger also reduces sensitivity to inedible and toxic bitter compounds. It is possible that the upregulated sweet taste *Gr* expression and downregulated bitter taste *Gr* expression concomitantly occur in the gustatory organs of starved animals. We speculate that sugar-sensing neurons with increased sweet taste *Gr* expression and bitter-sensing neurons with decreased bitter taste *Gr* expression are, respectively, activated and inactivated via dopaminergic regulation in *Drosophila* larvae during the poststress phase just as in starved larvae.

Quantitative analysis of protein and messenger RNA species in the mouths of *Drosophila* larvae showed elevated expression of tropomyosin and ATP (adenosine triphosphate) synthase  $\beta$  subunit 3 to 5 hours after heat stress. In insects, feeding responses are physically controlled by synchronous muscle movement in which one nerve excitation leads to one muscle contraction. The muscle contraction unit structure is extremely conserved within the animal kingdom, together with a similar conservation of the major myofibrillar proteins chiefly consisted of actin, myosin, troponin, and tropomyosin.<sup>16</sup> Muscle tension is caused by interaction of myosin crossbridges on thick filaments with actin in thin filaments. Tropomyosin directly interacts with both actin and myosin and plays a pivotal role in regulating the actin-myosin interaction.<sup>17</sup> Therefore, elevated expression of tropomyosin in the mouth parts must enable *Drosophila* larvae to move the mouth hooks for acceleration of diet intake. Furthermore, enhanced expression of the

## Gene expression 3-4 h after heat stress in mouth



**Figure 2.** Graphic summary showing upregulated and downregulated expression of genes in the mouth part of *Drosophila* larvae after heat stress at 35°C for 1 hour.

ATP synthase  $\beta$  subunit would contribute to supplying ATP more efficiently because it has been reported that the ATP synthase  $\beta$  subunit serves as a key component in ATP synthesis in rat pancreatic  $\beta$  cells.<sup>18,19</sup> Therefore, it is reasonable to assume that both enhanced expressions of tropomyosin and ATP synthase  $\beta$  subunit are essential for *Drosophila* larvae to accelerate the diet intake by speedily moving the mouth hooks (Figure 2).

In summary, our recent study supported our hypothesis that insects have evolved a systemic mechanism that promotes their recovery from the damage done by stressors during poststress periods. The 2 new findings mentioned above, which are interesting in terms of physiological aspects, allow us to propose that this recovery system is composed of multiple functional components. First, we found the elevated taste-sensing capacities in test larvae after heat stress. Upregulation of sweet taste *Gr* expression and concomitant downregulation of bitter taste *Gr* expression during the accelerated feeding periods led us to propose that insect feeding activities would be controlled by expression patterns of sweet and bitter taste *Gr* genes. Although starvation has been found to facilitate odor representation in odor receptor-expressing neurons that process olfactory input,<sup>20</sup> it has not been reported any *Gr* expression changes in starved *Drosophila* larvae. Therefore, this must be the first observation implying that population of both sweet and bitter taste *Grs* in the mouths directly or indirectly controls feeding activities of insects. Second, we found the enhanced mouth movement ability of *Drosophila* larvae after heat stress. This is due to elevated expression of the mouth myofibrillar protein, tropomyosin, and the key enzyme in ATP synthesis, ATP synthase  $\beta$  subunit. Our data suggest that enhancement of feeding activity during

poststress periods would be supported by supplied mouth muscle fiber component together with the fuel ATP.

The fact that insects have evolved such a complicated and highly elaborate system is regarded as a testament to the importance of the mechanisms that overcome the damages caused by stressors. If the stress-induced damages significantly retard larval growth rates, their reproductive maturation and the synchronized emergence of matured male and female adults would be affected. Given that this is applied to animals with short life spans such as insects, it is reasonable to assume that the recovery mechanism characterized in this study crucially contributes to preventing their population from decreasing.

### Author Contributions

This article commentary is written and prepared by YH.

### REFERENCES

1. Aragon T, van Anken E, Pincus D, et al. Messenger RNA targeting to endoplasmic reticulum stress signalling sites. *Nature*. 2009;457:736–740.
2. Quiros PM, Mottis A, Auwerx J. Mitonuclear communication in homeostasis and stress. *Nat Rev Mol Cell Biol*. 2016;17:213–226.
3. Feder ME, Hofmann GE. Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu Rev Physiol*. 1999;61:243–282.
4. Huang LH, Wang HS, Kang L. Different evolutionary lineages of large and small heat shock proteins in eukaryotes. *Cell Res*. 2008;18:1074–1076.
5. Mayer MP, Bukau B. Hsp70 chaperones: cellular functions and molecular mechanism. *Cell Mol Life Sci*. 2005;62:670–684.
6. Thanaphum S, Haymer DS. A member of the hsp70 gene family from the Mediterranean fruit fly, *Ceratitidis capitata*. *Insect Mol Biol*. 1998;7:63–72.
7. Ryuda M, Tabuchi M, Matsumoto H, Matsumura T, Ochiai M, Hayakawa Y. A gene-driven recovery mechanism: *Drosophila* larvae increase feeding activity for post-stress weight recovery [published online ahead of print December 8, 2018]. *Arch Insect Biochem Physiol*. doi:10.1002/arch.21440.
8. Amrein H, Thorne N. Gustatory perception and behavior in *Drosophila melanogaster*. *Curr Biol*. 2005;15:R673–R684.
9. Joseph RM, Carlson JR. *Drosophila* chemoreceptors: a molecular interface between the chemical world and the brain. *Trends Genet*. 2015;31:683–695.
10. Ejima A, Griffith LC. Courtship initiation is stimulated by acoustic signals in *Drosophila melanogaster*. *PLoS ONE*. 2008;3:e3246.
11. Barton-Browne L. Regulatory mechanisms in insect feeding. *Adv Insect Physiol*. 1975;11:1–116.
12. Itskov PM, Ribeiro C. The dilemmas of the gourmet fly: the molecular and neuronal mechanisms of feeding and nutrient decision making in *Drosophila*. *Front Neurosci*. 2013;7:12.
13. Pool AH, Scott K. Feeding regulation in *Drosophila*. *Curr Opin Neurobiol*. 2014;29:57–63.
14. Inagaki HK, de-Leon SB-T, Wong AM, et al. Visualizing neuromodulation in vivo: TANGO-mapping of dopamine signaling reveals appetite control of sugar sensing. *Cell*. 2012;148:583–595.
15. Marella S, Mann K, Scott K. Dopaminergic modulation of sucrose acceptance behavior in *Drosophila*. *Neuron*. 2012;73:941–950.
16. Pringle JW. The Croonian Lecture, 1977. Stretch activation of muscle: function and mechanism. *Proc R Soc Lond B Biol Sci*. 1978;201:107–130.
17. Bullard B, Pastore A. Regulating the contraction of insect flight muscle. *J Muscle Res Cell Motil*. 2011;32:303–313.
18. Wang C, Chen Z, Li S, et al. Hepatic overexpression of ATP synthase  $\beta$  subunit activates PI3K/Akt pathway to ameliorate hyperglycemia of diabetic mice. *Diabetes*. 2014;63:947–959.
19. Yang J, Wong RK, Wang X, et al. Leucine culture reveals that ATP synthase functions as a fuel sensor in pancreatic beta-cells. *J Biol Chem*. 2004;279:53915–53923.
20. Root CM, Ko KI, Jafari A, Wang JW. Presynaptic facilitation by neuropeptide signaling mediates odor-driven food search. *Cell*. 2011;145:133–144.