

## **Assortments of Digestive Enzymes Induced in First Instar Larvae of *Busseola fusca* Feeding on Different Plants**

Authors: Juma, Gérald, Le Ru, Bruno, and Calatayud, Paul-André

Source: International Journal of Insect Science, 11(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1179543319843521>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.


Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Assortments of Digestive Enzymes Induced in First Instar Larvae of *Busseola fusca* Feeding on Different Plants

Gérald Juma<sup>1</sup>, Bruno Le Ru<sup>2,3</sup> and Paul-André Calatayud<sup>2,3</sup> 

<sup>1</sup>Department of Biochemistry, University of Nairobi, Nairobi, Kenya. <sup>2</sup>UMR EGCE (Evolution, Génome, Comportement, Ecologie), CNRS-IRD-Univ. Paris-Sud, IDEEV, Université Paris-Saclay, Gif-sur-Yvette Cedex, France. <sup>3</sup>International Centre of Insect Physiology and Ecology (*icipe*), Nairobi, Kenya.

International Journal of Insect Science  
Volume 11: 1–6  
© The Author(s) 2019  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1179543319843521



**ABSTRACT:** The stem borer *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae) is an important pest of maize and sorghum in sub-Saharan Africa. This insect has oligophagous feeding habits, feeding mostly on maize and sorghum with a narrow range of wild Poaceous plant species. We hypothesised that first instar *B. fusca* larvae, the critical stage for successful establishment on a host plant, can establish and then grow on a particular plant as a result of induction of a complement of digestive enzymes that mediates host acceptance at first instars. A fast semi-quantitative analysis of potentially digestive enzymatic activities present in the first larvae previously fed for 4 days on leaves of host and non-host plants was performed using the API-ZYM kit system able to detect a multiplex of enzyme activities. Regardless of the plant species, the larvae exhibited higher activities of the carbohydrate metabolising enzymes than of aminopeptidases and proteases. In addition, highest activities of carbohydrates degrading enzymes were exhibited by larvae that consumed leaves of the most preferred plant species of *B. fusca*. Conversely, esterases were only detected in neonate larvae that consumed leaves of the less preferred and non-host plants. No alkaline phosphatase and lipase activities were detected. The significance of these results was discussed in terms of food requirements of first instar larvae when settling on a plant.

**KEYWORDS:** Lepidoptera, stem borers, maize pest, sub-Saharan Africa, digestive enzymes

**RECEIVED:** March 11, 2019. **ACCEPTED:** March 20, 2019.

**TYPE:** Original Research

**FUNDING:** The author(s) received no financial support for the research, authorship, and/or publication of this article.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**CORRESPONDING AUTHOR:** Paul-André Calatayud, UMR EGCE (Evolution, Génome, Comportement, Ecologie), CNRS-IRD-Univ. Paris-Sud, IDEEV, Université Paris-Saclay, 91198 Gif-sur-Yvette Cedex, France. Email: Paul-Andre.Calatayud@egce.cnrs-gif.fr

## Introduction

The evolutionary success of phytophagous insects depends on their ability to utilise specific plants as food sources. Phytophagous insects are largely influenced by host-plant chemistry.<sup>1,2</sup> Each plant has a specific phytochemical profile consisting of both primary and secondary metabolites that form the basis of host selection and discrimination by an insect.<sup>3</sup> These chemicals inform the foraging insects about the suitability of a plant as food source<sup>1,2</sup> and determine an insect's food choice and its subsequent performance.<sup>4</sup> They are thus important in host-plant adaptation.<sup>1,5,6</sup>

Following the adaptation of a particular host-plant insects partly depend on the efficiency of their digestive physiology to use chemically diverse host plants as food source,<sup>7–10</sup> helping them for settlement on a particular plant. In most cases, this usually involves the induction of a cocktail of digestive and detoxifying gut enzymes that permit the exploitation of toxic plant allelochemicals encountered during foraging.<sup>11,12</sup> Digestive plasticity to plant chemical toxicants has been previously reported in a number of insect species as an adaptive mechanism to noxious chemical containing host plants. For example, a number of insect species including the grasshopper, *Melanoplus sanguinipes* (Fabricius),<sup>13</sup> the gypsy moth, *Lymantria dispar* L.,<sup>14</sup> the cotton bollworm, *Helicoverpa armigera* (Hübner),<sup>9,15</sup> and the beetle, *Trogoderma granarium* Everts (Coleoptera: Dermestidae)<sup>10</sup> have been reported to modify the activity of their gut enzymes in response to the chemical

composition of the food. Therefore, a complement of digestive enzymes quantitatively or qualitatively expressed in the larvae on insect feeding should reflect the type of chemicals present and exploited by an insect to start to feed and survive on a particular plant.

In sub-Saharan Africa, the oligophagous stem borer *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae) is an important pest of economically important food crops such as maize, *Zea mays* L. (Poaceae), and sorghum, *Sorghum bicolor* (L.) Moench (Poaceae).<sup>16</sup> The oligophagous feeding habit of *B. fusca* larvae can be associated with possible plastic biochemical, physiological, or evolutionary mechanisms that allow the insect to confront a variety of chemical complexities posed by diverse food plants. Similar studies on plastic responses to nutritive stresses have been reported for other insect species,<sup>12,17</sup> and they provide basic information on the mechanisms of host-plants' specialisation. Thus, knowledge of the assortments of digestive enzymes induced in larvae when they start to feed on different plants can aid in the understanding of the physiological mechanisms used by *B. fusca* larvae to choose a particular plant species characterised by diverse chemicals. In addition, this will help to understand the mechanisms used by the larvae that disarm secondary metabolites to enable them to adapt on a variety of plants.

This study focused on first instar *B. fusca* larvae, which among Lepidoptera is the critical stage for successful establishment on a host plant;<sup>4,8,18–20</sup> they are more discriminative and selective in their food choice than older larvae.<sup>8</sup> Among wild



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without

Poaceae plant species frequently found in *B. fusca* natural habitat,<sup>21</sup> with the exception of *Panicum deustum* Thunb (Poaceae) from which *B. fusca* larvae have never been recovered, all other plant species were reported to host *B. fusca* albeit at differing abundances<sup>21–24</sup> and suitability.<sup>25,26</sup> Thereby, wild sorghum, *Sorghum arundinaceum* (Desv.) Stapf., was the most suitable hosts followed by *Megathyrus maximus* (Jacq.) B.K. Simon & S.W.L. Jacobs, Napier grass *Pennisetum purpureum* Schumach, and *Arundo donax* L., whereas *P. deustum* and *Setaria megaphylla* (Steud.) Th. Dur. & Schinz have been shown to be completely unsuitable.<sup>25,26</sup> *Panicum deustum* and *S. megaphylla* are therefore considered as non-hosts. Among these 7 plant species, it was also found that during the first instar life of *B. fusca* (ie, about during the first 7 days after hatching), the relative growth rate (RGR) of *B. fusca* larvae was 0.13 mg/d on *S. arundinaceum* (compared with the cultivated crop, *Z. mays*, which was 0.15 mg/d), whereas they were 0.09 and 0.05 mg/d, respectively, on *P. purpureum* and *A. donax*; and only 0.06, 0.03, and 0.03 mg/d, respectively, on *M. maximus*, *P. deustum*, and *S. megaphylla*.<sup>26</sup> Moreover, very few first instar larvae (from 1% to 8%) were able to survive on *A. donax*, *M. maximus*, *P. deustum*, and *S. megaphylla*.<sup>26</sup>

In this context, we hypothesised that first instar *B. fusca* larvae can establish and then grow on a particular plant (to establish on it) as a result of induction of a complement of digestive enzymes that mediates host acceptance at first instars. To provide evidences for this hypothesis, we estimated larval physiological responses in relation to the differential assortments of digestive enzymes, by a semi-quantitative analysis, in the first instar larvae reared on host and non-host plants used in previous studies.<sup>25,26</sup>

## Materials and Methods

### Insects

The *B. fusca* larvae stemmed from laboratory-reared individuals maintained on the artificial diet of Onyango and Ochieng'-Odero<sup>27</sup> from the Animal Rearing and Containment Unit (ARCU) of the International Centre of Insect Physiology and Ecology (*icipe*, Nairobi, Kenya). Feral individuals collected from maize fields in western Kenya were added to the colony thrice a year to rejuvenate the laboratory culture. One-day-old neonates were used in the experiments.

### Plants

The maize cultivar 511 and 6 wild Poaceae species including wild sorghum *S. arundinaceum*, Napier grass *P. purpureum*, *A. donax*, *S. megaphylla*, *M. maximus*, and *P. deustum* were used. Maize was grown in 4-L pots (1 plant per pot) from seeds provided by Simlaw, Kenya Seeds Company, Nairobi. The other plant species were obtained from their natural habitats and propagated from tillers or stem cuttings in 4-L pots (1 plant per pot) in a greenhouse at *icipe*. The environmental

conditions for growth were approximately 31°C/17°C (day/night) with 12:12 h (L:D) photoperiod. Plants were watered 3 times weekly and once with a solution of calcium ammonium nitrate (26% N). All plants were 3 weeks old, when used in the experiments, with exception of *S. arundinaceum*, which, due to its slow growth, was 5 weeks old. Each plant was infested with approximately 100 first instar larvae and allowed to feed for 4 days. Previous study indicated a high mortality for longer feeding periods on plant species with low suitability and even no survival on *P. deustum* and *S. megaphylla*.<sup>26</sup>

### Enzyme induction in larvae on feeding on different plant species

A fast semi-quantitative analysis of 19 enzymatic activities present in first instar larvae previously fed on leaves of 7 plant species was performed using the API-ZYM system from the API® test kit of BioMérieux (Marcy l'Etoile, France), a test kit that can detect a multiplex of 19 specific enzyme activities present in a sample as described by Rahbé et al.<sup>28</sup> The API-ZYM system consists of a gallery of 20 micro-cups (Figure 1): 19 of each with a specific substrate for enzyme detection and 1 micro-cup empty (ie, with no substrate) as control. The kit was proposing to detect the following 19 enzyme activities: alkaline phosphatase (pH 8.5), esterase (C4), esterase lipase (C8), lipase (C14), leucine aminopeptidase, valine aminopeptidase, cystine aminopeptidase, trypsin (*N*-benzoyl-DL-arginine-2-naphtylamidase),  $\alpha$ -chymotrypsin (*N*-glutaryl-phenylalanine-2-naphtylamidase), acid phosphatase (pH 5.4), naphthol-AS-BI-phosphohydrolase,  $\alpha$ -galactosidase,  $\beta$ -galactosidase,  $\beta$ -glucuronidase,  $\alpha$ -glucosidase,  $\alpha$ -glucosidase, *N*-acetyl-beta-glucosaminidase,  $\alpha$ -mannosidase, and  $\alpha$ -fucosidase. These enzymes are commonly found in the guts of lepidopteran larvae.<sup>29</sup>

First, 4-day-old larvae were recovered from the infested plants. It was, however, not possible to isolate the midgut for enzyme analysis due to the tiny size of the recovered larvae. Thus, the entire larval body without the head (to prevent contamination by salivary enzymes) was used; it was assumed that the resulting larval homogenates contained mostly digestive enzymes as the fatty tissues at this developmental stage are still not developed.

From each plant species, 40 larvae were homogenised in 1300  $\mu$ L distilled water and centrifuged (18 000g, 10 min, +4°C). A 65- $\mu$ L aliquot of each of the resultant supernatant was then pipetted to each of the 20 porous plastic micro-cups of the API system (ie, 19 micro-cups dispersed with specific chromogenic substrate and 1 used as a control) and then incubated at 37°C for 4 hours. A similar protocol was used for the control experiment using larvae that were starved but water-fed for 4 days.

For each micro-cup, a coloured product whose intensity is directly proportional to the amount of substrate hydrolysed is formed following the specific enzyme-substrate reaction. For the control micro-cup, no coloration was formed. The



**Figure 1.** A picture showing the differential staining of the API-ZYM plate with ZYM A and ZYM B after 4 hours of incubation of larval homogenates with substrates dispersed on the porous plate. Each coloured well gives a rank score based on the intensity of colour formed.

approximate amount (in nmol) of the enzyme is given by comparison with a colour scale included in the kit, as compared with the control.

Enzyme activities in the tissue extracts were detected by the cleavage of a chromogenic substrate (naphthyl derivatives) dispersed dry on a porous plastic micro-cup (about 100  $\mu$ L) of the API-ZYM system. Enzymatic reactions were enhanced by application of an sodium dodecyl sulphate (SDS)-based acid Tris buffer (Zym A) as a drop in each well following plate incubation. The reactions were then visualised following the addition of a Fast Blue BB solution (Zym B) to each of the micro-cups. Each test experiment was repeated 3 times and visible colour changes of the medium in the micro-cups were considered positive. The enzyme activity in each micro-cup was visually determined by using a ranking expressed on a scale from 0 (no enzyme reaction) to 5 (very high enzyme concentration) corresponding to the intensity of the coloured product produced on comparison with the colour scale provided with the kit (from  $\leq 5$  to  $\geq 40$  nmol: 1 corresponded to 5 nmol, 2 to 10 nmol, 3 to 20 nmol, 4 to 30 nmol, 5 to 40 nmol and above of substrate released according to the manufacturer's specification; Figure 1). Rank values reflecting enzyme activity from water-fed larvae served as baseline. The final feeding score value was obtained by subtracting the rank values of this group from the rank values obtained for the experimental groups. Score values greater than 0 indicated induction of an appreciable amount of enzyme activities in all larval homogenates.

#### Statistical analysis

Rank values for enzyme activities were generated following the Kruskal-Wallis test and their means separated using Tukey-Kramer test (Proc GLM). Statistical analyses were done in R version 3.3.1.<sup>30</sup>

## Results

Of the 19 enzymes tested with the Api Zym kit, 14 enzymes' activities were evident in the larval homogenates (Table 1). Regardless of the plant species, the carbohydrate metabolising enzymes were more active than aminopeptidases and proteases. No alkaline phosphatase and lipase activity were detected, whereby esterases were only found in larvae that consumed leaves of *M. maximus*, *P. deustum*, and *S. megaphylla*.

The activities of the aminopeptidases and proteases were not varying significantly between plant species. By contrast, the activities of most of the carbohydrate metabolising enzymes varied significantly. This included  $\beta$ -galactosidase,  $\beta$ -glucuronidase,  $\alpha$ -glucosidase, and  $\beta$ -glucosidase. Highest activities of carbohydrates degrading enzymes were exhibited by larvae that consumed leaves of *Z. mays*, *S. arundinaceum*, and *P. purpureum*, whereby  $\beta$ -glucosidase activity was significantly highest for larvae fed only on *Z. mays* and *S. arundinaceum*.

## Discussion

Although quantitative enzymatic assays with zymographic detections are important to complement this study, this semi-quantitative analysis is a good first overview of the assortments of digestive enzymes induced in first instar larvae after their first feeding on a particular plant that might help the larvae to establish on it. Moreover, this assortment might be directly linked to the chemical composition of foliage consumed by larvae. The low performance of first instar larvae of *B. fusca* on *M. maximus*, *P. deustum*, and *S. megaphylla* reported in a previous study by Juma et al<sup>26</sup> may suggest the presence of antibiotic properties in these plants, possibly as a result of plant secondary metabolites or low nutritional quality, which cannot allow first instar larvae of *B. fusca* to establish. C4 esterase activity was only detected in homogenates of larvae fed on *M. maximus*, *P. deustum*, and *S. megaphylla*, which have been shown plant of low suitability.<sup>26</sup> Induction of esterases in insect midgut has

**Table 1.** Semi-quantitative analysis of enzyme activities in the homogenate of entire neonate larvae (with heads decapitated) reared for 4 days on different plant species. Enzyme activities (means  $\pm$  SE, n=5) correspond to the activities found in the crushed larval homogenate (40 larvae per replicate). Activities correspond to the release of 5, 10, and 20 nmol of substrate per 4 hours of incubation at 37°C (rate 0: activity not detected).

ENZYMES	ZEA MAYS	SORGHUM ARUNDINACEUM	PENNISETUM PURPUREUM	ARUNDO DONAX	MEGATHYRSUS MAXIMUS	PANICUM DEUSTUM	SETARIA MEGAPHYLLA
Alkaline phosphatase	–	–	–	–	–	–	–
Esterase (C4)	–	–	–	–	0.2 $\pm$ 0.2a	1.0a	0.2 $\pm$ 0.2a
Esterase lipase (C8)	–	–	–	–	0.2 $\pm$ 0.2	–	–
Lipase (C14)	–	–	–	–	–	–	–
Leucine aminopeptidase	–	–	–	–	–	–	–
Valine aminopeptidase	0.7 $\pm$ 0.2a	0.6 $\pm$ 0.2a	1.0a	0.4 $\pm$ 0.2a	–	–	0.6 $\pm$ 0.2a
Cystine aminopeptidase	–	0.2 $\pm$ 0.2a	–	–	0.2 $\pm$ 0.2a	0.2 $\pm$ 0.2a	–
Trypsin	–	–	–	–	0.5 $\pm$ 0.3	–	–
$\alpha$ -chymotrypsin	–	–	0.7 $\pm$ 0.2a	–	0.5 $\pm$ 0.3a	–	–
Acid phosphatase	–	–	–	–	–	–	–
Naphthol-AS-BI-phosphohydrolase	1.0b	0.6 $\pm$ 0.2ab	1.0b	1.0b	0.2 $\pm$ 0.2a	1.0b	0.8 $\pm$ 0.2ab
$\alpha$ -galactosidase	–	1.2 $\pm$ 0.5a	–	–	–	0.5 $\pm$ 0.3a	–
$\beta$ -galactosidase	1.2 $\pm$ 0.3b	1.2 $\pm$ 0.2b	1.7 $\pm$ 0.2b	0.8 $\pm$ 0.2a	0.5 $\pm$ 0.3a	–	0.2 $\pm$ 0.2a
$\beta$ -glucuronidase	1.2 $\pm$ 0.2a	2.0 $\pm$ 0.5b	2.0b	1.0a	1.0a	–	0.8 $\pm$ 0.2a
$\alpha$ -glucosidase	1.8 $\pm$ 0.2b	1.6 $\pm$ 0.2b	2.0b	1.2 $\pm$ 0.2a	–	0.5 $\pm$ 0.3a	1.0 $\pm$ 0.3a
$\beta$ -glucosidase	1.8 $\pm$ 0.5c	1.4 $\pm$ 0.2c	1.0b	0.4 $\pm$ 0.2a	0.7 $\pm$ 0.2ab	–	–
N-acetyl-beta-glucosaminidase	–	–	–	–	–	–	–
$\alpha$ -mannosidase	–	1.0 $\pm$ 0.3a	1.5 $\pm$ 0.3a	–	0.7 $\pm$ 0.2a	–	–
$\alpha$ -fucosidase	–	0.4 $\pm$ 0.2	–	–	–	–	–

Means within a line followed by different letters are significantly different at 5% level (Tukey-Kramer test).

been shown to be positively correlated with resistance to plant allelochemicals by several Lepidoptera species.<sup>31–37</sup> *Panicum* sp. and *Setaria* sp., for example, are known to possess beta-phenylethylamine alkaloids (such as *N*-methyltyramine) and a phenolic acid, setarin (4-allyloxycoumarin), respectively.<sup>38</sup>

The level of activities of glycosidases including  $\beta$ -galactosidase,  $\beta$ -glucuronidase,  $\alpha$ -glucosidase, and  $\beta$ -glucosidase was significantly induced in homogenates of larvae fed on maize, wild sorghum, and *P. purpureum*. These plants were previously demonstrated to support the best larval performance in the laboratory over the plant species used in this study.<sup>26</sup> This may be related to higher content of disaccharides and polysaccharides in young, developing leaves of these plants as compared with

the other plant species used as it has been observed for secondary metabolites in young maize leaf whorls by Bergvinson et al.<sup>39</sup> This indicates that these disaccharides and polysaccharides might mostly be hydrolysed by these enzymes into simple sugars for first instar larvae energy metabolism.

In addition,  $\beta$ -glucosidases, that were high in larvae fed on maize and sorghum, are also involved in the detoxification of a wide range of other plant-derived  $\beta$ -o-glycosyl containing allelochemicals such as the benzoxazinoids (DIMBOA and MBOA) present in maize<sup>40–42</sup> and dhurrin, the cyanogenic glycoside present in sorghum.<sup>43</sup> This suggests a possible physiological adaptation of *B. fusca* to toxic allelochemicals of these host plants as reported in other lepidoteran species including European corn

borer, *Ostrinia nubilalis* (Hübner) (Lepidoptera: Pyralidae)<sup>44,45</sup> and in *Spodoptera frugiperda* (J.E. Smith).<sup>42</sup> Similarly, specialised *Heliconius* caterpillars (Lepidoptera: Nymphalidae) are reported to efficiently convert sorghum-specific cyanogenic glycosides to soluble and harmless thiols<sup>29</sup> preventing the latter to release harmful cyanide and even allowing the caterpillars to utilise the toxic compounds as a nitrogen source.

Proteases including trypsin and chymotrypsin and aminopeptidases are the most predominant enzymes in the midgut of most lepidopteran insects.<sup>46–49</sup> Although third instar larvae of *B. fusca* were found to harbour digestive proteases,<sup>50</sup> the activity of proteases (aminopeptidases and serine proteases) in larval homogenate of first instar of *B. fusca* on all 7 plants was, however, lower than that of the carbohydrate-hydrolysing enzymes, suggesting that leaves of the plant species tested might be a poor source of proteins at least for first instar larvae of *B. fusca*, and therefore, the protein calorific value of the studied plants did not significantly induce the first instar larvae gut proteolytic activity. However, the level of free amino acids and soluble low-molecular-weight proteins in leaves might be sufficient for the development of young *B. fusca* larvae, and therefore, they did not induce gut extra proteolytic activities.

No lipase activity involved in fat digestion was found in first instar larvae of *B. fusca*. This indicates that *B. fusca*, at this initial stage of development, does not utilise fat for initial growth on plant.

In conclusion, our results indicated that first instar larvae of *B. fusca* generally relies on simple carbohydrates rather than proteins as food source. This might be linked to its specialisation to *Z. mays* and *S. arundinaceum*.<sup>22,23</sup> In fact, the advantage for host plant specialisation by herbivorous insects has been hypothesised to involve an increase in energetic efficiency.<sup>51</sup> Therefore, maize and wild sorghum should have a more suitable composition, level, or ratio of different carbohydrates required by *B. fusca* first instar larvae for energy unlike the other plant species considered in this study, thus allowing a better settlement of first instar of *B. fusca* on these plants than on the others.

## Acknowledgements

The authors thank Fritz Schulthess for his review of the manuscript.

## Author Contributions

The research was conducted by GJ under the supervision of PAC listed as co-author. GJ and PAC participated in defining the research methodology and all co-authors participated in writing/editing the paper.

## ORCID iD

Paul-André Calatayud  <https://orcid.org/0000-0002-9482-4646>

## REFERENCES

- Bernays EA, Chapman RF. *Host-Plant Selection by Phytophagous Insects*. New York, NY: Chapman and Hall; 1994.

- Schoonhoven LM, van Loon JJA, Dicke M. *Insect Plant Biology*. Oxford, UK: Oxford University Press; 2005.
- De Boer G, Hanson FE. Food plant selection and induction of feeding preference among host and non-host plants in larvae of tobacco hornworm *Manduca sexta*. *Entomol Exp Appl*. 1984;35:177–193.
- Barros E, Torres JB, Ruberson JR, Oliveira MD. Development of *Spodoptera frugiperda* on different hosts and damage to reproductive structures in cotton. *Entomol Exp Appl*. 2010;137:237–245.
- Foss LK, Rieske LK. Species-specific differences in oak foliage affect preference and performance of gypsy moth caterpillar. *Entomol Exp Appl*. 2003;108:87–93.
- Fischer DC, Kogan M, Paxton J. Effect of glyceollin, a soybean phytoalexin, on feeding by three phytophagous beetles (Coleoptera: Coccinellidae and Chrysomelidae): dose versus response. *Environ Entomol*. 1990;5:1278–1282.
- Brattsten LB. Bioengineering of crop plants and resistant biotype evolution in insect. *Arch Insect Biochem Physiol*. 1991;17:253–267.
- Zalucki MP, Clarke AR, Malcolm SB. Ecology and behavior of first instar larval Lepidoptera. *Annu Rev Entomol*. 2002;47:361–393.
- Wang Y, Cai QN, Zhang QW, Han Y. Effect of the secondary substances from wheat on the growth and digestive physiology of cotton bollworm *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Eur J Entomol*. 2006;103:255–258.
- Borzoui E, Naseri B, Namin FR. Different diets affecting biology and digestive physiology of the Khapra beetle, *Trogoderma granarium* Everts (Coleoptera: Dermestidae). *J Stored Prod Res*. 2015;62:1–7.
- Veenstra KH, Pashley DP, Ottea JA. Host-plant adaptation in fall armyworm host strains: comparison of food consumption, utilization, and detoxication enzyme activities. *Ann Entomol Soc Am*. 1995;88:80–91.
- Wouters FC, Blanchette B, Gershenzon J. Plant defense and herbivore counter-defense: benzoxazinoids and insect herbivores. *Phytochem Rev*. 2016;15:1127–1151.
- Hinks CF, Erlandson MA. The accumulation of haemolymph proteins and activity of digestive proteinases of grasshoppers (*Melanoplus sanguinipes*) fed wheat, oats or kochia. *J Insect Physiol*. 1994;41:425–433.
- Lazarevic J, Peric-Mataruga V. Nutritive stress effects on growth and digestive physiology of *Lymantria dispar* larvae. *Yugoslav Med Biochem*. 2003;22:53–59.
- Sarate PJ, Tamhane VA, Kotkar HM, et al. Developmental and digestive flexibilities in the midgut of a polyphagous pest, the cotton bollworm, *Helicoverpa armigera*. *J Insect Sci*. 2012;12:42.
- Kfir R, Overholt WA, Khan ZR, Polaszek A. Biology and management of economically important lepidopteran cereal stem borers in Africa. *Annu Rev Entomol*. 2002;47:701–731.
- Mello MO, Silva-Filho MC. Plant-insect interactions: an evolutionary arms race between two distinct defense mechanisms. *Braz J Plant Physiol*. 2002;14:71–81.
- Kaufmann T. Behavioural biology, feeding habits and ecology of three species of maize stemborers: *Eldana saccharina* (Lepidoptera: Pyralidae), *Sesamia calamistis* and *Busseola fusca* (Noctuidae) in Ibadan, Nigeria, West Africa. *J Georgia Entomol Soc*. 1983;18:259–272.
- Silva DM, da Bueno A, de F, et al. Biology and nutrition of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) fed on different food sources. *Sci Agric*. 2017;74:18–31.
- Scriber JM, Slansky FJ. The nutritional ecology of immature insects. *Annu Rev Entomol*. 1981;26:183–211.
- Calatayud P-A, Le Ru BP, van den Berg J, Schulthess F. Ecology of the African maize stalk borer, *Busseola fusca* (Lepidoptera: Noctuidae) with special reference to insect-plant interactions. *Insects*. 2014;5:539–563.
- Le Ru BP, Ong'amo GO, Moyal P, et al. Geographic distribution and host plant ranges of East African noctuid stem borers. *Ann Soc Entomol Fr*. 2006;42:353–361.
- Le Ru BP, Ong'amo GO, Moyal P, et al. Diversity of lepidopteran stemborers on monocotyledonous plants in eastern Africa and island of Madagascar and Zanzibar revisited. *Bull Entomol Res*. 2006;96:555–563.
- Ong'amo GO, Le Ru B, Dupas S, Moyal P, Calatayud P-A, Silvain J-F. Distribution, pest status and agro-climatic preferences of lepidopteran stemborers of maize in Kenya. *Ann Soc Entomol Fr*. 2006;42:171–177.
- Juma G, Thiongo M, Dutaur L, et al. Two sugar isomers influence host plant acceptance by a cereal caterpillar pest. *Bull Entomol Res*. 2013;103:20–28.
- Juma G, Ahuya PO, Ong'amo G, Le Ru B, Magoma G, Silvain J-F, Calatayud P-A. Influence of plant silicon in *Busseola fusca* (Lepidoptera: Noctuidae) larvae—Poaceae interaction. *Bull Entomol Res*. 2015;105:253–258.
- Onyango FO, Ochieng'-Odero JPR. Continuous rearing of the maize stem borer *Busseola fusca* on an artificial diet. *Entomol Exp Appl*. 1994;73:139–144.
- Rahbé Y, Sauvion N, Febvay G, Peumans WJ, Gatehouse AMR. Toxicity of lectins and processing of ingested proteins in the pea aphid *Acyrtosiphon pisum*. *Entomol Exp Appl*. 1995;76:143–155.
- Engler HS, Spencer KC, Gilbert LE. Insect metabolism-preventing cyanide release from leaves. *Nature*. 2000;406:144–145.
- R Development Core Team. *A Language and Environment for Statistical Computing R Foundation for Statistical Computing*. Vienna, Austria. <http://www.R-project.org/>. Up-dated 2013. Accessed February 2, 2015.

31. Ahmad S, Brattsen LB, Mullin CA, Yu SJ. Enzymes involved in the metabolism of plant allelochemicals. In: Brattsen LB, Ahmad S, eds. *Molecular Aspects of INSECT-Plant Associations*. New York, NY: Plenum Press; 1986:73–152.
32. Lindroth RL, Hemming JDC. Responses of the gypsy moth (Lepidoptera: Lymantriidae) to tremulacin, an aspen phenolic glycoside. *Environ Entomol*. 1990;19:842–847.
33. Lindroth RL, Bloomer MS. Biochemical ecology of the forest tent caterpillar: response to dietary proteins and phenolic glycosides. *Oecologia*. 1991;86:408–413.
34. Lindroth RL, Weisbrod AV. Genetic variation in response of the gypsy moth to aspen phenolic glycosides. *Biochem Syst Ecol*. 1991;19:97–103.
35. Lindroth RL, Jung SM, Feuker AM. Detoxification activity in the gypsy moth: effects of host CO<sub>2</sub> and NO<sub>3</sub>- availability. *J Chem Ecol*. 1993;19:357–367.
36. Hwang SY, Lindroth RL. Clonal variation in foliar chemistry of Aspen: effects on gypsy moths and forest tent caterpillars. *Oecologia*. 1997;111:99–108.
37. Ghumare SS, Mukherjee SN. Performance of *Spodoptera litura* (Fabricius) on different host plants: influence of nitrogen and total phenolics of plants and midgut esterase activity of the insect. *Indian J Exp Biol*. 2003;41:895–899.
38. Steglich W, Fugmann B, Lang-Fugman S. *ROMPP Encyclopedia, Natural Products*. Stuttgart, Germany; New York, NY: Georg Thieme Verlag; 2000.
39. Bergvinson DJ, Hamilton RI, Arnason JT. Leaf profile of maize resistance factors to European corn borer, *Ostrinia nubilalis*. *J Chem Ecol*. 1995;21:343–354.
40. Niemeyer HM. Hydroxamic acids derived from 2-hydroxy-2H-1,4-benzoxazin-3(4H)-one: key defense chemicals of cereals. *J Agric Food Chem*. 2009;57:1677–1696.
41. Niemeyer HM. Hydroxamic acids (4-hydroxy-1,4-benzoxazin-3-ones), defence chemicals in the Gramineae. *Phytochemistry*. 1988;27:3349–3358.
42. Wouters FC, Reichelt M, Glauser G, et al. Reglucosylation of the benzoxazinoid DIMBOA with inversion of stereochemical configuration is a detoxification strategy in lepidopteran herbivores. *Angew Chem Int Ed Engl*. 2014;53:11320–11324.
43. Conn EE. Cyanogenic glycosides. In: Bell EA, Charlwood BV, eds. *Secondary Plant Products*. Berlin, Germany; Heidelberg, Germany; New York, NY: Springer; 1980:461–492.
44. Campos F, Artkinson J, Arnason JT, et al. Toxicokinetics of 2, 4-dihydroxy-7-methoxy-1, 4-benzoxazin-3-one (DIMBOA) in the European corn borer, *Ostrinia nubilalis* (Hübner). *J Chem Ecol*. 1989;15:1989–2001.
45. Houseman JG, Campos F, Thie NMR, et al. Effect of the maize-derived compounds DIMBOA and MBOA on growth and digestive processes of European corn borer (Lepidoptera, Pyralidae). *J Econ Entomol*. 1992;85:669–674.
46. Terra WR, Ferreira C, Jordao BP, Dillon RJ. Digestive enzymes. In: Lehane MJ, Billingsley PF, eds. *Biology of the Insect Midgut*. London, England: Chapman and Hall; 1996:153–193.
47. Zhu YC, Zeng FR, Oppert B. Molecular cloning of trypsin-like cDNAs and comparison of proteinase activities in the salivary glands and gut of the tarnished plant bug *Lygus lineolaris* (Heteroptera: Miridae). *Insect Biochem Mol Biol*. 2003;33:889–899.
48. Stygar D, Dolezych B, Nakonieczny M, Migula P, Michalczyk K, Zaak M. Digestive enzymes activity in larvae of *Cameraria ohridella* (Lepidoptera: Gracillariidae). *C R Biol*. 2010;333:725–735.
49. Zibae A. Digestive enzymes of large cabbage white butterfly, *Pieris brassicae* L. (Lepidoptera: Pieridae) from developmental and site of activity perspectives. *Ital J Zool*. 2012;79:13–26.
50. George D, Ferry N, Back EJ, Gatehouse AM. Characterisation of midgut digestive proteases from the maize stem borer *Busseola fusca*. *Pest Manag Sci*. 2008;64:1151–1158.
51. Zibae A, Bandani AR, Kafil M, Ramzi S. Characterisation of  $\alpha$ -amylase in the midgut and the salivary glands of rice striped stem borer, *Chilo suppressalis* Walker (Lepidoptera: pyralidae). *J Asia-Pac Entomol*. 2008;11:201–205.