

# Three Decades of Antillean Manatee (Trichechus manatus manatus) Stranding Along the Brazilian Coast

Authors: Balensiefer, Deisi Cristiane, Attademo, Fernanda Loffler Niemeyer, Sousa, Gláucia Pereira, Freire, Augusto Carlos da Bôaviagem, da Cunha, Fábio Adônis Gouveia Carneiro, et al.

Source: Tropical Conservation Science, 10(1)

Published By: SAGE Publishing

# URL: https://doi.org/10.1177/1940082917728375

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Three Decades of Antillean Manatee (Trichechus manatus manatus) Stranding Along the Brazilian Coast

Tropical Conservation Science Volume 10: 1-9 © The Author(s) 2017 Reprints and permissions: sagepub.com/journalsPermissions.nav DOI: 10.1177/1940082917728375 journals.sagepub.com/home/trc

## **SAGE**

Deisi Cristiane Balensiefer<sup>1</sup>, Fernanda Loffler Niemeyer Attademo<sup>1</sup>, Gláucia Pereira Sousa<sup>1</sup>, Augusto Carlos da Bôaviagem Freire<sup>2</sup>, Fábio Adônis Gouveia Carneiro da Cunha<sup>1</sup>, Ana Emília Barboza Alencar<sup>2</sup>, Flávio José de Lima Silva<sup>2</sup>, and Fábia de Oliveira Luna<sup>1</sup>

### Abstract

Alive Antillean manatee stranding is one of the main challenges for the conservation of the species in Brazil. In the present study, data on Antillean manatee stranding recorded by the CMA/ICMBio's Wild Animals Rehabilitation Center from 1987 to 2015 were analyzed. The CMA and the REMANE's (Northeastern Aquatic Mammal Stranding Network's) institutions have rescued on average five Antillean manatees per year. Approximately 56% of the animals were alive when rescued, and 43.7% were found dead. The states of Ceará and Rio Grande do Norte have the highest records of stranding, rescue, and assistance of alive calves. Stranding of alive specimens is more frequent during the austral summer, especially in January, February, and March. In the North region, the main threat to the Antillean manatee population seems to be interaction with fishing activities. Studies on the causes of dependent calf stranding in northeast Brazil are urgently needed to promote the development of public policies for the conservation of the species. Brazil's manatee rehabilitation centers must undergo improvements to better assist stranded calves, which need special care to survive.

#### **Keywords**

newborn calves, anthropogenic threats, manatee, mortality, Brazil

# Introduction

Stranding of marine mammals occurs throughout the world and, although there is a lack of global statistical records, some countries have well-established programs and databases in response to these events (Aragones et al., 2010). The development of a database contributes to more refined studies of the species, yielding a greater success in preserving marine mammals (Moraes & Barreto, 2004). The institutions within the Northeastern Aquatic Mammal Stranding Network (REMANE, acronym in Portuguese) are the ones in charge of rescuing Antillean manatees (Trichechus manatus manatus) in Brazil. This network was developed by the government, research organizations, and academic institutions in the year 2000 to rescue marine mammals stranded in northeast Brazil. The National Center of Research and Conservation of Aquatic Mammals of the Chico Mendes Institute for Biodiversity Conservation (CMA/ICMBio,

acronym in Portuguese) is the government representative and network coordinator.

Antillean manatees inhabit warm coastal waters of the Caribbean Sea, Gulf of Mexico, and the western Atlantic Ocean (Lefebvre, Marmontel, Reid, Rathbun, & Domning, 2001). In Brazil, this subspecies occurs discontinuously along the coast from the state of Amapá to Alagoas. However, historically, the Antillean manatees

<sup>1</sup>Instituto Chico Mendes de Conservação da Biodiversidade, Itamaracá, Pernambuco, Brazil

<sup>2</sup>Universidade do Estado do Rio Grande do Norte, Natal, RN, Brazil Received 26 May 2017; Revised 13 July 2017; Accepted 31 July 2017

#### **Corresponding Author:**

Deisi Cristiane Balensiefer, Instituto Chico Mendes de Conservação da Biodiversidade, Estrada do Forte Orange, S/N, C.P 01, Itamaracá, Pernambuco 53900000, Brazil. Email: deisi.balensiefer@icmbio.gov.br



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons / tarbeter the terms of the Creative Commons / tarbeter terms of the work without further License (http://www.creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further License (http://www.creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further License (http://www.creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further terms of the terms of the terms of the terms of the terms of ter permission provided the original work is attributed as specified on the SAGE and Open Access pages (https://us.sagepub.com/en-us/nam/open-access-at-sage). Downloaded From: https://complete.bioone.org/journals/Tropical-Conservation-Science on 15 Jul 2025 Terms of Use: https://complete.bioone.org/terms-of-use

have inhabited areas further south, down to the state of Espírito Santo (Lima, Paludo, Soavinski, Silva, & Oliveira, 2011; Luna, Lima, Araújo, & Passavante, 2008). Currently, the Antillean manatee population is severely reduced in Brazil, being estimated at approximately 1,000 specimens, and is considered an endangered subspecies (Alves et al., 2015; Luna, Lima, et al., 2008; Self-Sullivan & Mignucci-Giannoni, 2008). The main threats to the Brazilian population of Antillean manatees are low genetic variability, change and loss of habitats, interaction with fisheries, stranding of calves in the northeast region, and hunting in the North region (Lima et al., 2011; Luna, Araújo, et al., 2008; Luna et al., 2012; Meireles, 2008).

Antillean manatee strandings have been recorded in Brazil since the 1980s (Meireles, 2008; Parente, Vergara-Parente, & Lima, 2004). The REMANE institution closest to the stranding site rescues the stranded animals, which, after a stabilization period, are allocated to the CMA/ICMBio. After receiving proper rehabilitation treatment at the CMA/ICMBio, the animals are released back to their natural habitat. In instances when a dead manatee is stranded, the institution rescuing it can keep the carcass. In the present study, we only evaluated the carcasses sent to the CMA/ICMBio. This study summarizes the CMA/ICMBio's records for Antillean manatee stranding and describes the spatial and temporal distribution of those strandings and differences in sex and length of manatees rescued more than 29 years.

## Methods

The CMA/ICMBio is a center specialized in Antillean manatee conservation and, until July 2013, it was the only institution in Brazil that kept this subspecies in captivity. The present analysis considered stranded manatees that were assisted or received by the CMA/ICMBio's Wild Animals Rehabilitation Center from 1987 to 2015. Calves born in captivity were not included in the analysis.

The area of assisted strandings extends from the state of Amapá (North Brazil) through the states of Maranhão, Ceará, Rio Grande do Norte, Paraíba, and Pernambuco to Alagoas (northeast Brazil). In the state of Amapá, the rainy and dry seasons occur from January to June and from July to December, respectively. The average annual rainfall is 2,500 mm, and the temperatures range from 27°C to 32°C (Ab'Saber, 1977). In the northeast region, due to its coastal configuration, the rainfall is irregular and varies spatially and temporally. In the eastern coast of the northeast region, the rainy season is between January and June, and the dry season occurs from July to December (Hastenrath, 2012). The average annual rainfall is 1,800 mm (da Silva, 2004) and, according to the National Weather Institute (Instituto Nacional *de Meteorologia*); the average annual temperature ranges from 20.7°C to 27.4°C. On the other hand, the northern coast has a semiarid climate with average annual temperatures from 23°C to 33°C. The rainy season is short, concentrated between February and May, while the dry season occurs from April to January, and the annual average rainfall is 537.5 mm (*Instituto de Desenvolvimento Sustentável e Meio Ambiente*, 1999).

The manatees were rescued, rehabilitated, or necropsied following the REMANE protocol (Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis, 2005). The specimens were rescued within the shortest period possible after stranding notifications issued by the network of collaborators in coastal communities. Depending on the animals' clinical condition and the environmental conditions of the stranding site, which includes the presence of other manatees in the area, the priority was to attempt immediate reintroduction, especially for calves, as described in the protocol. In cases of failure, the rescued animals were transported to the nearest RENAME institution and received initial care and stabilization. They were then transferred to the CMA/ ICMBio and received the necessary veterinary care. Animals deemed suitable for reintroduction were transferred to natural captivity for later release (Normande et al., 2015). Dead animals were necropsied, depending on the stage of carcass decomposition (Geraci & Lounsbury, 2005).

To verify the spatial distribution and seasonality of strandings, georeferenced records of manatee stranding from 1987 to 2015 were reviewed and organized by Brazilian state and date (month). The sex (male, female, and unknown) of stranded manatees (alive or dead) was checked as well as their length (cm), measured as a straight line from the tip of the nostril to the tip of the tail, which was compared using the Mann–Whitney test.

# Results

One hundred and forty-two Antillean manatees were translocated to the CMA throughout the 29 years of rescues (five manatees per year, on average). Of these, on average per year, three were stranded alive (56.3%) and two were found dead (43.7%). The male or female ratio was 1:1; however, it was not possible to determine the sex of 8% of the dead animals due to advanced decomposition of the carcasses (Code 4; Geraci & Lounsbury, 2005), making them impossible to necropsy.

The states of Rio Grande do Norte (n=40; 28%), Ceará (n=39; 27%), and Paraíba (n=29; 20%) had the highest number of strandings, followed by the states of Maranhão (n=18; 13%), Alagoas (n=9; 6%), Pernambuco (n=6; 4%), and Amapá (n=1; 1%). The highest number of strandings of alive Antillean manatees was recorded in eastern Ceará (n=34; 24%) and the northern coast of Rio Grande do Norte (n=26; 18%); Figure 1).



Figure 1. Antillean manatee stranding records and location by Brazilian state from 1987 to 2015.

The highest number of rescues was registered in 2004 and 2008 (n=11 manatees) with the same number of alive (n=4) and dead animals (n=7) recorded in both years. In 2008, four of the dead animals were found in a mangrove area of the state of Maranhão. The highest record of alive strandings during the studied period was in 2010 (n=9; Figure 2). Strandings of alive calves occurred in all months of the year but were more common during the austral summer, especially in January, February, and March. There were no records of stranding of dead calves during the winter. Strandings of dead adults were also registered in every month of the year with a higher incidence in February (Figure 3).

Of the 129 Antillean manatees measured, 75% were less than 220 cm long. The smallest specimen, at 67 cm long, was found dead in a mangrove area in the state of Maranhão in December 2009. Specimens rescued alive (n=77) were significantly smaller than those found dead (Mann–Whitney=503, df=1, p < .001; Table 1; Figure 4). Only two manatees rescued alive were not newborns according to the classification of Borges et al. (2012), their sizes being consistent with those of specimens older than 2 years old.

Calves rescued alive showed no signs of congenital diseases (hydrocephalus, eye and facial deformities, cardiac anomalies verified by auscultation) or other diseases diagnosed by clinical and laboratory tests. However, recently, one specimen is showing abnormal development likely related to congenital diseases, but some tests are needed to confirm the suspicion (F. Attademo, personal communication, September 2015). Commonly, the calves were found with remnants of the umbilical cord and folded tail, which are neonate characteristics. After being rescued, the animals underwent a health assessment, including complete blood count and serum chemistry. Their body condition score and body size were compared with those of healthy animals of the same age, and their body condition was always compatible with healthy neonates estimated with an average age of 0 to 3 days (database CMA/ICMBio).

# Discussion

### Alive Manatees

Strandings are invaluable sources of information on the biology and ecology of marine mammals and provide



Figure 2. Number of stranded Antillean manatees allocated to the CMA more than 29 years. Black bars = dead manatees; Gray bars = alive manatees.



Figure 3. Temporal distribution of records of manatee calves and adults stranded in Brazil. Black bars = dead manatees; Gray bars = alive manatees.

Table 1.	Length o	of Strand	ded Antillean	Manatees	Allocated	to the
CMA From	n <b>1987</b> t	to 2015.	Number of	Manatees =	= 129.	

	Alive		Dead		
	Female	Male	Female	Male	Unknown
Number	39	38	22	23	7
Minimum length (cm)	108	93	120	83	67
Maximum length (cm)	335	169	340	315	300
Mean length (cm) $\pm$ SD	$136\pm40$	$125 \pm 12$	$249 \pm 78$	$230\pm68$	$207\pm80$

important information about the animals' health (Borsa, 2006; Bossart, 2011). The stranding patterns of a particular species may be related to its distribution, migration, and reproduction (Perrin & Geracy, 2002). The data analyzed in the present study show that strandings are more common among calves; and stranding of alive manatees occurs in the northeast coast of Brazil, mainly in the states of Ceará and Rio Grande do Norte.

Studies have been cautioning about the stranding of newborn and immature calves in Brazil, indicating that this situation has been persistent over the years (Meireles, 2008; Parente et al., 2004). The eastern coast of the state of Ceará and the northern coast of the state of



Figure 4. Size of Antillean manatees rescued alive and dead in Brazil from 1987 to 2015.

Rio Grande do Norte have the highest rates of alive Antillean manatee stranding and rescue and are considered the most critical areas. Stranded calves are sent to rehabilitation centers and most are reintroduced into their natural habitat (76% success rate; Normande, Attademo, & Luna, 2016; Normande et al., 2015). However, when in captivity, calves miss a critical learning period and fail to learn how to find food, freshwater, and shelter on their own (Reep & Bonde, 2006), which may compromise their survival in the natural habitat after release.

Studies on the causes for calf stranding in Brazil are rare, but a significant anthropogenic impact has been verified in areas where stranding occurs, including the establishment of shrimp farms and lobster fishing in the northern coast of the Potiguar Basin, between the states of Rio Grande do Norte and Ceará (de Almeida, 2010). Some studies have also indicated that calf stranding is associated with the silting of rivers and degradation of coastal ecosystems, especially in relation to the estuarine complex and its recesses in the states of Ceará and Rio Grande do Norte (Lima et al., 2011; Parente et al., 2004). These anthropogenic disturbances limit the access of female manatees to inshore birthing sites; consequently, calves born in offshore waters may become separated from their mothers and strand along the beaches (Lima et al., 2011; Meireles, 2008). Adult manatees are able to free themselves from other environmental barriers, such as ocean currents and tidal changes (Hartman, 1979; Paludo & Langguth, 2002). For calves, however, the tidal fluctuations can cause stranding, since they lack the adults' agility and strength. For example, during low tide, a dependent calf was stranded alive in a silted area of the state of Paraíba. After 24 h searching for the

presence of others manatees in the surroundings, an adult female was found vocalizing close to the silted area. The calf was placed near the female and both vocalized substantially. Then, the female approached the calf and demonstrated parental care. This episode was considered a successful immediate reintroduction.

Adult Antillean manatees are able to move in shallow waters (1.2 m deep, on average) or along sandbanks and have been found resting in extremely shallow waters (0.4 m deep, on average; Paludo & Langguth, 2002). Therefore, it seems that adults are not disturbed by silting areas. As a matter of fact, in Brazil, between the beaches of Baía Formosa (state of Rio Grande do Norte) and Barra do Rio Camaratuba (state of Paraíba), adults were observed in shallow waters near the beach with their dorsum partially exposed.

Strandings of alive manatees were recorded through all the months of the studied years (1987–2015); however, the highest frequency was observed during the austral summer, which corresponds to the manatee reproductive season in northeast Brazil (Silva, Paludo, Oliveira, Lima, & Soavinski, 2011). The smallest manatee recorded in the present survey, which had been probably aborted (Ackerman, Wright, Bonde, Odell, & Banowetz, 1995; Mignucci-Giannoni et al., 2000) was found dead in a mangrove area in the state of Maranhão during the summer. Calf stranding was recorded in two occasions during or after the stranding of an adult specimen was recorded. However, it was not possible to verify if the calf stranding had its cause in the stranding of its mother. In some cases, the cause for the stranding was related to natural factors, such as diseases, congenital disorders or others, and, several times, to environmental factors. In the northern coast of the Potiguar Basin (state of Rio Grande do Norte), 12- to 15-knot winds can have an influence on strandings in the month of August to November (UERN, data not published). The seasonality of stranding records during the reproductive season demonstrates the need for maintaining rescue actions and for endeavoring to accomplish preventive actions. In many occasions, parturient females are also believed to remain at the stranding site in the days following the event. Therefore, an effort to locate these females for an average period of 72 h can be effective for the immediate reintroduction of the calf, when it is healthy. By bringing mother and offspring together again, the calf would be spared from remaining in captivity offsite.

In Brazil, there is no data showing the effects of stranding of alive calves on the size of the manatee population. In the last national conservation status assessment, *T. manatus* status downgraded from critically endangered to endangered. However, no study has proved that the population increased, since the existing surveys used different sampling methods (Alves et al., 2015; Luna, Lima, et al., 2008). The REMANE's

successful performance in the rescue of stranded calves has greatly contributed to the conservation of Antillean manatees in Brazil, considering that 76% of manatees were rehabilitated and reintroduced into their natural environment (Normande et al., 2015, 2016).

# Dead Manatees

The death of aquatic mammals due to entanglement in fishing nets has been frequently reported (Goldstein et al., 1999; Machado, Oliveira, & Montealegre-Quijano, 2015; Morizur, Berrow, Tregenza, Couperus, & Pouvreau, 1999; Ott et al., 2002; Read, 2008; Vidal, Waerebeek, & Findley, 1994; Yousuf et al., 2009). In northeast Brazil, these records have become priority in the last two decades, with the strengthening of the REMANE (Batista, Schiavetti, Santos, & Reis, 2012; Meirelles et al., 2009; Meirelles, Ribeiro, Siva, & Soares Filho 2010; Monteiro-Neto et al., 2000). Some small cetacean and sirenian species are incidentally captured in fishing nets because of their coastal habits (Adulyanukosol, Prasittipornkul, Man-Anansap, & Boukaew, 2009; Borgonia, Moreira, & Barreto, 2002; Cremer et al., 2013; Danilewicz et al., 2009; Dawson, 2006; Rodas-Trejo, Romero-Berny, & Estrada, 2008; Rojas-Bracho, Reeves, & Jaramillo-Legorreta, 2006; Rosas, Monteiro-Filho, & Oliveira, 2002). For the Antillean manatee, interaction with fisheries is also a cause of stranding and death of young and adult specimens in Brazil and in other countries (Castelblanco-Martinez et al., 2009; de Thoisy et al., 2003; Marsh & Lefebvre, 1994; Mignucci-Giannoni et al., 2000). Most of the specimens found dead in the states of Amapá and Maranhão were adults in advanced state of decomposition making it difficult to determine the cause of death at necropsy. However, cuts, bruises in the head, and tail injuries consistent with fishing nets were observed in three of those specimens. External postmortem examination indicated that the injuries were recent and interactions with fisheries may have led to death of animals by drowning. Gill nets (called malhadeira or malhão) are set perpendicularly to estuary channels in the state of Maranhão, obstructing the passage of manatees, which get entangled in the nets (Alvite, 2008). In the state of Ceará, manatees occur in coastal fishing areas (Alves et al., 2013), and the stranding and death of adults have been associated with the shrimp trawl activities (Meireles, 2008; Silva, Choi, & Carvalho, 2006). Collisions with motorboats may also cause edemas in the frontal part of the body and multiple cuts all over the body, as previously registered on the northeast coast of Brazil by Borges, Vergara-Parente, Alvite, Marcondes, and Lima (2007).

The present study did not aim to determine the cause of death of the reintroduced manatees (these specimens were considered in the group of calves stranded alive). However, death caused by anthropogenic activities, such as debris ingestion (Attademo et al., 2015) and injuries by firearms and sharp piercing objects (data not published because it is part of a criminal investigation) have already been described.

Cold stress syndrome is one of the causes of manatee (*T. m. latirostris*) and dugong (*Dugon dugong*) stranding in Florida and Queensland, respectively (Ackerman et al., 1995; Fertl et al., 2005; Owen, Flint, Limpus, Palmieri, & Mills, 2013). There are no records of manatee stranding due to this syndrome anywhere where *T. m. manatus* is distributed, since the average temperature in the area of species distribution is around  $24^{\circ}$ C throughout the year, which is not a stress factor for the population.

The number of dead Antillean manatees analyzed here may be underestimated because the CMA/ICMBio has not conducted a systematic monitoring during the studied years. Furthermore, manatees found dead in the states of Ceará and Rio Grande do Norte have not been sent to the Rehabilitation center since 2006; and in the states of Pará and Amapá, dead animals are sent to other institutions and local universities. Brazil has a Monitoring System of Marine Mammals (SIMMAM, in Portuguese), in which the records of strandings and sightings are registered. It is our belief that a greater commitment toward keeping that database updated would allow for more realistic estimates of strandings, which in turn would help in the conservation of the species.

# Implications for Conservation

Sirenians are K-strategists, have long lifespans, physically develop and reach sexual maturity slowly, produce small litters (often one offspring at a time), and provide long parental care for their few offspring (Reynolds & Marshall, 2012). The current scenario, associated with the species biological characteristics, raises the concern about the future of Antillean manatees in Brazil. Our hope is that the present study will stimulate researches into the causes of stranding of alive calves in the northeast region. It is also necessary to implement protection campaigns in fishing communities more often to prevent the death of manatees in the North and northeast regions. Finally, it is our hope that public policies for the conservation of the Antillean manatee continue for many years and include the conservation of their habitats as well, especially mangrove areas.

#### Acknowledgments

The authors thank L. N. Domingues for translating the manuscript and Ursula Migon for the revision and are grateful to anonymous reviewers, who provided valuable editorial comments on the manuscript. The authors also thank REMANE institutions, for their support and collaboration during the rescue of Antillean manatees.

#### **Declaration of Conflicting Interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

#### Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

#### References

- Ab'Saber, A. N. (1977). Os domínios morfoclimáticos da América do Sul: Primeira aproximação [South America morphoclimatic domains: First approach]. *Geomorfologia*, 52, 1–21.
- Ackerman, B. B., Wright, S. D., Bonde, R. K., Odell, D. K., & Banowetz, D. J. (1995). Trends and patterns in mortality of manatees in Florida, 1974–1992. In: T. J. O'Shea, B. B. Ackerman, & H. F. Percival (Eds.). *Population biology of the Florida manatee* (pp. 223–258). Washington, DC: National Biological Service Information and Technology.
- Adulyanukosol, K., Prasittipornkul, C., Man-Anansap, S., & Boukaew, P. (2009). Stranding records of dugong (Dugong dugon) in Thailand. In: N. Arai (ed.) Proceeding of 4th International symposium on Eastar 2000 and Asian Bio-logging Science (pp. 51–57). Phuket, TH: Kyoto University.
- Alves, M. D., Kinas, P. G., Marmontel, M., Borges, J. C. G., Costa, A. F., Schiel, N., & Arújo, M. E. (2015). First abundance estimate of the Antillean manatee (*Trichechus manatus manatus*) in Brazil by aerial survey. Journal of the Marine Biological Association of the United Kingdom, 1, 1–12.
- Alves, M. D. O., Schwamborn, R., Borges, J. C. G., Marmontel, M., Costa, A. F., Schettini, C. A. F., & Arújo, M. E. (2013). Aerial survey of manatees, dolphins and sea turtles off northeastern Brazil: Correlations with coastal features and human activities. *Biological Conservation*, 161, 91–100.
- Alvite, C. M. C. (2008). Indicadores populacionais e ecológicos de peixes-bois-marinhos (*Trichechus manatus manatus*) em duas áreas de manguezais e marismas no Maranhão (Dissertação, Mestrado em Sustentabilidade de Ecossistemas [Antillean manatee (*Trichechus manatus manatus*) ecological and population indicators in two mangrove and marsh areas of Maranhão. Thesis (Master's Degree in Ecosystem Sustainability]. São Luís, MA: Universidade Federal do Maranhão [São Luís, MA: Maranhão Federal University].
- Aragones, L. V., Roque, M. A. A., Flores, M. B., Encomienda, R. P., Laule, G. E., Espinos, B. G., ... Braun, R. C. (2010). The Philippine marine mammal strandings from 1998 to 2009: Animals in the Philippines in peril? *Aquatic Mammals*, 36(3): 219–233.
- Attademo, F. L. N., Balensiefer, D. C., Freire, A. C. B., Sousa, G. P., Cunha, F. A. G. C., & Luna, F. O. (2015). Debris ingestion by Antillean Manatee (*Trichechus manatus*). *Marine Pollution Bulletin*, 101(1): 284–287.
- Batista, R. L. G., Schiavetti, A., Santos, U. A., & Reis, M. S. S. (2012). Cetaceans registered on the coast of Ilhéus (Bahia), northeastern Brazil. *Biota Neotropica*, 12, 31–38.
- Borges, J. C. G., Freire, A. C. B., Attademo, F. L. N., Serrano, I. L., Anzolin, D. G., Carvalho, P. S. M., & Vergara-Parente, J. E. (2012). Growth pattern differences of captive born Antillean manatee (*Trichechus manatus*) calves and those rescued in the

Brazilian northeastern coast. *Journal of Zoo and Wildlife Medicine*, 43(3), 494–500.

- Borges, J. C. G., Vergara-Parente, J. E., Alvite, C. M. C., Marcondes, M. C. C., & Lima, R. P. (2007). Embarcações motorizadas: Uma ameaça aos peixes-boi marinhos (*Trichechus manatus*) no Brasil [Motorboat: a threat to Antillean manatees (*Trichechus manatus*) in Brazil]. *Biota Neotropica*, 7(3): 199–204.
- Borgonia, M. P., Moreira, P., & Barreto, A. S. (2002). Monitoramento da Morte Acidental de Pequenos Cetáceos no Litoral Centro-Norte de Santa Catarina, Brasil [Monitoring of Accidental Death of Small Cetaceans in the North-Central Coast of Santa Catarina, Brazil]. In *Reunión de trabajos de especialistas en mamíferos acuáticos de América del Sur, 10 y 4° Congreso de la Sociedad Latino Americana de Mamíferos Acuáticos* In *Collection of papers from South American aquatic mammal specialists, 10 and 4° Latin American Aquatic Mammal Society Congress*] (pp. 64–64). Valdívia, Chile: Universidad Austral de Chile.
- Borsa, P. (2006). Marine mammal strandings in the New Caledonia region, Southwest Pacific. *Comptes Rendus Biologies*, 329, 277–288.
- Bossart, G. D. (2011). Marine mammals as sentinel species for oceans and human health. *Veterinary Pathology*, 48(3): 676–690.
- Castelblanco-Martinez, D. N., Bermúdez-Romero, A. L., Gómez-Camelo, I. V., Rosas, F. C. W., Trujillo, F., & Zerda-Ordones, E. (2009). Seasonality of habitat use, mortality and reproduction of the Vulnerable Antillean manatee *Trichechus manatus manatus* in the Orinoco River, Colombia: Implications for conservation. *Oryx*, 43(2): 235–242.
- Cremer, M. J., Sartori, C. M., Holz, A. C., Schulze, B., Santos, N. Z., Alves, A. K. M., & Paitach, R. L. (2013). Franciscana strandings on the north coast of Santa Catarina State and insights into birth period. *Biotemas*, 26(4), 133–139.
- Danilewicz, D., Secchi, E. R., Ott, P. H., Moreno, I. B., Bassoi, M., & Borges-Martins, M. (2009). Habitat use patterns of franciscana dolphins (*Pontoporia blainvillei*) off southern Brazil in relation to water depth. *Journal of the Marine Biological Association of the United Kingdom*, 89(5): 943–949.
- da Silva, V. D. P. R. (2004). On climate variability in Northeast of Brazil. *Journal of Arid Environments*, 58(4): 575–596.
- Dawson, S. M. (2006). Incidental catch of Hector's dolphin in inshore gillnets. *Marine Mammal Science*, 7(3): 283–295.
- de Almeida, L. G. (2010). Caracterização das áreas de pesca artesanal de lagosta na Praia da Redonda, Icapuí, Ceará (Dissertação, Mestrado em Ciências Marinhas Tropicais) [Characterization of artisanal lobster fishery areas in Redonda beach, Icapuí, Ceará. Thesis (Master's Degree in Tropical Marine Science)]. Fortaleza, Brazil: Universidade Federal do Ceará [Ceará Federal University].
- de Thoisy, B., Spiegelberger, T., Rousseau, S., Talvy, G., Vogel, I., & Vié, J. C. (2003). Distribution, habitat, and conservation status of the West Indian manatee Trichechus manatus in French Guiana. *Oryx*, 37(4): 431–436.
- Fertl, D., Schiro, A. J., Regan, G. T., Beck, C. A., Adimey, N., Price-May, L.,...Crossland, R. (2005). Manatee occurrence in the northern Gulf of Mexico, West of Florida. *Gulf and Caribbean Research*, 17, 69–94.

- Geraci, J. R., & Lounsbury, V. J. (2005). *Marine mammals ashore: A field guide for strandings*. Baltimore, MD: National Aquarium in Baltimore.
- Goldstein, T., Johnson, S. P., Phillips, A.V., Hanni, K. D., Fauquier, D. A., & Gulland, F. M. D. (1999). Human-related injuries observed in live stranded pinnipeds along the central California coast 1986–1998. *Aquatic Mammals*, 25(1): 43–51.
- Hartman, D. S. (1979). Ecology and behavior of the Manatee, Trichechus manatus latirostris (Harlan) in Florida (Special Publication No. 5). Pittsburgh, PA: American Society of Mammalogists.
- Hastenrath, S. (2012). Exploring the climate problems of Brazil's Nordeste: A review. *Climatic Change*, *112*(2): 243–251.
- Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis. (2005). Protocolo de conduta para encalhes de mamíferos aquáticos/Rede de encalhe de mamíferos aquáticos do Nordeste [Conduct protocol for aquatic mammal stranding/ Northeastern Aquatic Mammal Stranding Network]. Recife, Brazil: Author.
- Instituto de Desenvolvimento Sustentável e Meio Ambiente (1999). Secretaria de Planejamento e Finanças. Instituto de Desenvolvimento Econômico e Meio Ambiente do Rio Grande do Norte – IDEMA [Department of Planning and Finances. Rio Grande do Norte Institute for Environment and Economic Development – IDEMA]. *Informativo Municipal*, 5, 1–14.
- Lefebvre, L. W., Marmontel, M., Reid, J. P., Rathbun, G. B., & Domning, D. P. (2001). Status and biogeography of the West Indian manatee. In: C. A. Woods, & F. E. Sergile (Eds.). *Biogeography of the West Indies: Patterns and perspectives* (pp. 425–474). Boca Raton, FL: CRC Press.
- Lima, R. P., Paludo, D., Soavinski, R. J., Silva, K. G., & Oliveira, E. M. A. (2011). Levantamento da distribuição, ocorrência e status de conservação do peixe-boi marinho (*Trichechus manatus*, Linnaeus, 1758) no litoral nordeste do Brasil [Survey on Antillean manatee (*Trichechus manatus*, Linnaeus, 1758) distribution, occurrence and conservation status in the Brazilian Northeast coast]. *Natural Resources*, 1, 41–57.
- Luna, F. O., Araújo, J. P., Lima, R. P., Pessanha, M. M., Soavinski, R. J., & Passavante, J. Z. O. (2008). Captura e utilização do peixe-boi marinho (*Trichechus manatus manatus*) no litoral norte do Brasil [Antillean manatee (*Trichechus manatus manatus*) capture and use in the Brazilian north coast]. *Biotemas*, 21, 115–123.
- Luna, F. O., Bonde, R. K., Attademo, F. L. N., Saunders, J. W., Meigs-Friend, G., Passavante, J. Z. O., & Hunter, M. E. (2012). Phylogeographic implications for release of critically endangered manatee calves rescued in northeast Brazil. *Aquatic Conservation Marine Freshwater Ecosystem*, 22, 665–672.
- Luna, F. O., Lima, R. P., Araújo, J. P., & Passavante, J. Z. O. (2008). Status de conservação do peixe-boi marinho (*Trichechus manatus manatus* Linnaeus, 1758) no Brasil [Antillean manatee (*Trichechus manatus manatus* Linnaeus, 1758) conservation status in Brazil]. *Revista Brasileira de Zoologia*, 10, 145–153.
- Machado, R., Oliveira, L. R., & Montealegre-Quijano, S. (2015). Incidental catch of South American sea lion in a pair trawl off southern Brazil. *Neotropical Biology and Conservation*, 10(1): 43–47.

- Marsh, H., & Lefebvre, L. W. (1994). Sirenian status and conservation efforts. Aquatic Mammals, 20(3): 155–170.
- Meireles, A. C. O. (2008). Mortality of the Antillean manatee, *Trichechus manatus manatus*, in Ceará State, northeastern Brazil. *Journal of the Marine Biological Association of the United Kingdom*, 88, 1133–1137.
- Meirelles, A. C. O., Monteiro-Neto, C., Martins, A. M. A., Costa, A. F., Barros, H. M. D. R., & Alves, M. D. O. (2009). Cetacean strandings on the coast of Ceará, northeastern Brazil (1992– 2005). Journal of the Marine Biological Association of the United Kingdom, 89(5): 1083–1090.
- Meirelles, A. C. O., Ribeiro, A. C., Siva, C. P. N., & Soares Filho, A. A. (2010). Records of Guiana dolphin, *Sotalia guianensis*, in the state of Ceará, Northeastern Brazil. *Latin American Journal* of Aquatic Mammals, 8(1–2): 97–102.
- Mignucci-Giannoni, A. A., Montoya-Ospina, R. A., Jiménez-Marrero, N. M., Rodríguez-López, M. A., Williams, E. H. Jr, & Bonde, R. K. (2000). Manatee mortality in Puerto Rico. *Environmental Management*, 25(2): 189–198.
- Monteiro-Neto, C., Alves, T. T., Jr., Ávila, F. J. C., Campos, A. A., Costa, A. F., Silva, C. P.N., & Furtado-Neto, M. A. A. (2000). Impact of fisheries on the tucuxi (*Sotalia fluviatilis*) and roughtoothed dolphin (*Steno bredanensis*) populations off Ceará state, northeastern Brazil. *Aquatic Mammals*, 26(1), 49–56.
- Moraes, C. G., & Barreto, A. S. (2004). Sistema de Monitoramento de Avistagens e Encalhes de Cetáceos na Costa Brasileira [Brazilian Coast Cetacean Sighting and Stranding Monitoring System]. In Reunión de trabajos de especialistas en mamíferos acuáticos de América del Sur, 11 y 5° Congreso de la sociedad latinoamericana de especialista en mamíferos acuáticos In [Collection of papers from South American aquatic mammal specialists, 11 and 5° Latin American Aquatic Mammal Specialists Society Congress] (pp. 83–83). Quito, Ecuador: Pontifica Universidad Católica del Ecuador.
- Morizur, Y., Berrow, S. D., Tregenza, N. J. C., Couperus, A. S., & Pouvreau, S. (1999). Incidental catches of marine-mammals in pelagic trawl fisheries of the northeast Atlantic. *Fisheries Research*, 41(3): 297–307.
- Normande, I. C., Attademo, F. L. N., & Luna, F. O. (2016). Antillean manatee release program in Brazil. In: P. S. Soorae (ed.) Global re-introduction perspectives: 2016. Case-studies from around the globe (pp. 162–165). Gland, Switzerland: IUCN/SSC Reintroduction Specialist Group/Abu Dhabi, UAE: Environment Agency-Abu Dhabi.
- Normande, I. C., Luna, F. O., Malhado, A. C. M., Borges, J. C. G., Viana, P. C., Jr Attademo, F. L. N., & Ladler, R. J. (2015). Eighteen years of Antillean manatee *Trichechus manatus manatus* releases in Brazil: Lessons learnt. *Oryx*, 49, 338–344.
- Ott, P. H., Secchi, E. R., Moreno, I. B., Danilewicz, D., Crespo, E. A., Bordino, P.,...Kinas, P. G. (2002). Report of the working group on fishery interactions. *Latin American Journal of Aquatic Mammals*, 1(1): 55–64.
- Owen, H. C., Flint, M., Limpus, C. J., Palmieri, C., & Mills, P. C. (2013). Evidence of sirenian cold stress syndrome in dugongs *Dugong dugon* from southeast Queensland, Australia. *Diseases* of Aquatic Organisms, 103, 1–7.
- Paludo, D., & Langguth, A. (2002). Use of space and temporal distribution of *Trichechus manatus manatus* Linnaeus in the region of Sagi, Rio Grande do Norte State, Brazil (Sirenia, Trichechidae). *Revista Brasileira de Zoologia*, 19, 205–215.

- Parente, C. L., Vergara-Parente, J. E., & Lima, R. P. (2004). Strandings of Antillean manatees, *Trichechus manatus manatus*, in northeastern Brazil. *Latin American Journal of Aquatic Mammals*, 3, 69–75.
- Perrin, W. F., & Geraci, J. R. (2002). Stranding. In: W. F. Perrin, B. Würsig, & J. G. M. Thewissen (Eds.). *The encyclopedia of marine mammals* (pp. 1192–1197). San Diego, CA: Academic Press.
- Read, A. J. (2008). The looming crisis: Interactions between marine mammals and fisheries. *Journal of Mammalogy*, 89(3): 541–548.
- Reep, R. L., & Bonde, R. K. (2006). The Florida manatee. Biology and conservation. Gainesville, FL: University Press of Florida.
- Reynolds, J. E. III, & Marshall, C. D. (2012). Vulnerability of sirenians. In: E. M. Hines, J. E. Reynolds III, L. V. Aragones, A. A. Mignucci-Giannoni, & M. Marmontel (Eds.). Sirenian conservation: Issues and strategies in developing countries (pp. 12–19). Gainesville, FL: University Press of Florida.
- Rodas-Trejo, J., Romero-Berny, E. I., & Estrada, A. (2008). Distribution and conservation of the West Indian manatee (*Trichechus manatus manatus*) in the Catazajá wetlands of northeast Chiapas, México. *Tropical Conservation Science*, 1, 321–333.
- Rojas-Bracho, L., Reeves, R. R., & Jaramillo-Legorreta, A. (2006). Conservation of the vaquita *Phocoena sinus*. *Mammal Review*, 36(3): 179–216.
- Rosas, F. C. W., Monteiro-Filho, E. L. A., & Oliveira, M. R. (2002). Incidental catches of Franciscana (*Pontoporia blainvillei*) on the southern coast of São Paulo State and the coast of

- Self-Sullivan, C., & Mignucci-Giannoni, A. (2008). Trichechus manatus ssp. manatus. The IUCN Red List of Threatened Species. Version 2015.2. Retrieved from http://www.iucnred list.org
- Silva, C. P. N., Choi, K. F., & Carvalho, V. L. (2006). Relatório de encalhe de dois espécimes de peixe-boi marinho (*Trichechus manatus manatus*) mortos, ocorridos no município de Aracati-CE. Associação de Pesquisa e Preservação de Ecossistemas Aquáticos – AQUASIS (Report) [Report on the stranding of two specimens of dead Antillean manatees (*Trichechus manatus manatus*), occurring in the city of Aracati-CE. Aquatic Ecosystem Preservation and Research Association – AQUASIS (Report)]. Ceará, Brazil.
- Silva, K. G., Paludo, D., Oliveira, E. M. A., Lima, R. P., & Soavinski, R. J. (2011). Distribuição e ocorrência do peixe-boi marinho (*Trichechus manatus*) no estuário do rio Mamanguape, Paraíba, Brasil [Distribution and occurrence of manatee (*Trichechus manatus*) in the Mamanguape River estuary, Paraíba, Brazil]. *Natural Resources*, 1, 5–14.
- Vidal, O., Waerebeek, K. V., & Findley, L. T. (1994). Cetaceans and gillnet fisheries in Mexico, Central America and the Wider Caribbean: A preliminary review. *Report of the International Whale Commission*, 15, 221–233.
- Yousuf, K. S. S. M., Anoop, A. K., Anoop, B., Afsal, V. V., Vivekanandan, E., Kumarran, R. P., ... Jayasankar, P. (2009). Observations on incidental catch of cetaceans in three landing centres along the Indian coast. *Marine Biodiversity Records*, 2, e64.