

Bioremediation of Fluorophenols by Glycosylation with Immobilized Marine Microalga Amphidinium Crassum

Authors: Shimoda, Kei, and Hamada, Hiroki

Source: Environmental Health Insights, 4(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/EHI.S5392>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

Bioremediation of Fluorophenols by Glycosylation with Immobilized Marine Microalga *Amphidinium Crassum*

Kei Shimoda¹ and Hiroki Hamada²

¹Department of Chemistry, Faculty of Medicine, Oita University, 1-1 Hasama-machi, Oita 879-5593, Japan. ²Department of Life Science, Faculty of Science, Okayama University of Science, 1-1 Ridai-cho, Kita-ku, Okayama 700-0005, Japan. Corresponding author email: hamada@dls.ous.ac.jp

Abstract: Fluorophenols are used as agrochemicals and released into environment as pollutants. Cultured marine microalga *Amphidinium crassum* (*Gymnodinium*) glucosylated 2-fluorophenol (**1**), 3-fluorophenol (**2**), and 4-fluorophenol (**3**) to the corresponding β -D-glucosides, ie, 2-fluorophenyl β -D-glucoside (**4**, 60 μ g/g cells), 3-fluorophenyl β -D-glucoside (**5**, 20 μ g/g cells), and 4-fluorophenyl β -D-glucoside (**6**, 40 μ g/g cells). On the other hand, 2-, 3-, and 4-fluorophenols were efficiently converted by immobilized *A. crassum* in sodium alginate gel to give their β -D-glucosides in higher yields (**4**: 140 μ g/g cells; **5**: 60 μ g/g cells; **6**: 100 μ g/g cells). In repetitive batch use, the immobilized cells of *A. crassum* maintained the potential for the glucosylation of the substrate fluorophenol after 5 times of usage.

Keywords: glycosylation, fluorophenol; β -glycoside, *Amphidinium crassum*, immobilized marine microalga

Environmental Health Insights 2010:4 87–91

doi: [10.4137/EHI.S5392](https://doi.org/10.4137/EHI.S5392)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.



Introduction

Fluorophenolic compounds are used to agrochemicals and pharmaceuticals, and cause serious environmental contamination.¹ Number of fluorine-containing agricultural chemicals has grown from 4% to approximately 9% of all agrochemicals and has increased in number faster than non-fluorinated agrochemicals over the past 15 years.¹ These compounds are primarily used as herbicides (48%), insecticides (23%), and fungicides (18%), and their residues are released as pollutants into rivers and seas. Many studies on biological degradation of halophenols by microorganisms have been reported.^{1–9} However, halophenols containing stable carbon-halogen bond in their structures have been described to be much more resistant to microbial degradation than unsubstituted analogs.¹

Cell cultures are ideal systems for studying biotransformations such as glycosylation reactions.¹⁰ Some freshwater microalga can convert exogenous toxic compounds into glycosides.¹¹ From the viewpoint of seawater pollution control, glycosylation of halophenols by marine microalga is of importance. This study focused on the metabolism of mono-fluorophenols by cultured and immobilized marine microalga of *Amphidinium crassum*.

Material and Methods

General

Substrates, 2-, 3-, and 4-fluorophenols, were purchased from Aldrich Chemical Co. The ¹H and ¹³C NMR, H-H COSY, C-H COSY, and HMBC spectra were recorded using a Varian XL-400 spectrometer in CD₃OD solution. Chemical shifts were expressed in δ (ppm) referring to tetramethylsilane. FABMS spectra were measured using a JEOL MStation JMS-700 spectrometer. *A. crassum* cells (5 g) were cultivated in a synthetic seawater (500 ml) for 2 weeks at 20 °C with constant aeration by air (1l/min) in 1 l flasks under illumination (1000lx). Constituents of synthetic seawater are summarized in Table 1.

Biotransformation of fluorophenols by *A. crassum*

Cultured *A. crassum* cells were harvested by centrifugation at 1000 g for 15 min and washed twice by adding 100 ml of synthetic seawater followed by centrifugation (1000 g for 15 min). To the 500 ml

Table 1. The constituents of synthetic seawater.

Constituents ^a	Conc. (g/l)	Constituents ^a	Conc. (g/l)
NaCl	20.7	AlCl ₃ ·6H ₂ O	8 × 10 ⁻⁶
MnCl ₂ ·4H ₂ O	1 × 10 ⁻⁶	FeCl ₃ ·6H ₂ O	5 × 10 ⁻⁶
MgCl ₂ ·6H ₂ O	9.5	Na ₂ WO ₄ ·2H ₂ O	2 × 10 ⁻⁷
CaCl ₂ ·6H ₂ O	1.3	(NH ₄) ₆ MO ₇ O ₂₄	2 × 10 ⁻⁵
Na ₂ SO ₄	3.5	Na ₂ SiO ₃	5 × 10 ⁻²
KCl	0.6	NaNO ₃	0.2
NaHCO ₃	0.2	Na ₂ HO ₄	1 × 10 ⁻²
KBr	9 × 10 ⁻²	EDTA-2 Na	9 × 10 ⁻⁴
Na ₂ B ₄ O ₇ ·10H ₂ O	3 × 10 ⁻²	Vitamin B ₁₂	2 × 10 ⁻⁶
SrCl ₂	1 × 10 ⁻²	Thiamine·HCl	8 × 10 ⁻⁵
NaF	3 × 10 ⁻³	Biotin	1 × 10 ⁻⁶
LiCl	1 × 10 ⁻³	EDTA-Fe	3 × 10 ⁻³
KI	7 × 10 ⁻⁵	H ₂ NC(CH ₃ OH) ₃	5 × 10 ⁻³
CoCl ₂ ·6H ₂ O	2 × 10 ⁻⁷		

Note: ^aConstituents are dissolved in distilled water.

flask containing 9 g of cultured *A. crassum* cells and 300 ml of a synthetic seawater was added 0.2 mmol of substrate. Cultures were incubated at 20 °C with aerobic shaking for five days under illumination (1000lx). After the incubation period, the cells and synthetic seawater were separated by centrifugation at 1000 g for 15 min. Synthetic seawater was extracted with ethylacetate and then *n*-butanol. Cells were extracted (three times) by homogenization with methanol, and the methanol fraction was concentrated and partitioned between water and ethylacetate. Ethylacetate fractions were analyzed by HPLC, combined, and concentrated. This water fraction and *n*-butanol fraction extracted from medium were analyzed by HPLC, combined, evaporated, and re-dissolved in water. This water fraction was applied to a Diaion HP-20 column and the column was washed with water followed by elution with methanol. Methanol eluate was subjected to preparative HPLC to give glycosylation products. Yield of the products was determined on the basis of the peak area from HPLC. Product yields in μ g/g cells (dry weight) was calculated based on the starting cell concentration. Control free suspended cells, which had not been treated with substrates, were subjected to the same extraction procedures and HPLC analyses as planktonic biotransformation experiments, and no products or substrates were detected in the cells and in the medium despite careful HPLC analyses.

Spectral data of products are as follows.

2-Fluorophenyl β -D-glucoside: FAB-MS m/z : 297 $[M+Na]^+$; 1H NMR (400 MHz, CD_3OD , δ in ppm): 3.41–3.89 (6H, m, H-2'-H-6'), 4.97 (1H, d, $J = 8.0$ Hz, H-1'), 6.96–7.01 (1H, m, H-4), 7.04–7.09 (2H, m, H-3, H-6), 7.25–7.30 (1H, dd, $J = 1.6$ Hz, $J = 8.4$ Hz, H-5); ^{13}C NMR (100 MHz, CD_3OD , δ in ppm): 62.3 (C-6'), 71.1 (C-4'), 74.6 (C-2'), 77.8 (C-3'), 78.0 (C-5'), 102.5 (C-1'), 117.0 (1C, d, $J_{F-3C} = 18.9$ Hz, C-3), 119.1 (C-6), 123.8 (1C, d, $J_{F-4C} = 6.6$ Hz, C-4), 125.4 (1C, d, $J_{F-5C} = 4.2$ Hz, C-5), 146.3 (1C, d, $J_{F-1C} = 10.8$ Hz, C-1), 153.9 (1C, d, $J_{F-2C} = 244.2$ Hz, C-2).

3-Fluorophenyl β -D-glucoside; FAB-MS m/z : 297 $[M+Na]^+$; 1H NMR (400 MHz, CD_3OD , δ in ppm): 3.39–3.90 (6H, m, H-2'-H-6'), 4.92 (1H, d, $J = 7.2$ Hz, H-1'), 6.74 (1H, m, H-4), 6.85 (1H, m, H-2), 6.90 (1H, dd, $J = 2.2$ Hz, $J = 8.0$ Hz, H-6), 7.27 (1H, m, H-5); ^{13}C NMR (100 MHz, CD_3OD , δ in ppm): 62.3 (C-6'), 71.2 (C-4'), 74.7 (C-2'), 77.7 (C-3'), 78.0 (C-5'), 102.1 (C-1'), 105.2 (1C, d, $J_{F-4C} = 25.5$ Hz, C-4), 109.7 (1C, d, $J_{F-2C} = 20.6$ Hz, C-2), 113.4 (1C, d, $J_{F-6C} = 2.4$ Hz, C-6), 131.3 (1C, d, $J_{F-5C} = 9.9$ Hz, C-5), 160.1 (1C, d, $J_{F-1C} = 10.5$ Hz, C-1), 164.5 (1C, d, $J_{F-3C} = 241.7$ Hz, C-3).

4-Fluorophenyl β -D-glucoside; FAB-MS m/z : 297 $[M + Na]^+$; 1H NMR (400 MHz, CD_3OD , δ in ppm): 3.35–3.90 (6H, m, H-2'-H-6'), 4.85 (1H, d, $J = 7.2$ Hz, H-1'), 6.95–7.01 (2H, m, H-3, H-5), 7.08–7.12 (2H, m, H-2, H-6); ^{13}C NMR (100 MHz, CD_3OD , δ in ppm): 62.4 (C-6'), 71.2 (C-4'), 74.7 (C-2'), 77.8 (C-3'), 78.0 (C-5'), 102.9 (C-1'), 116.4 (2C, d, $J_{F-3,5C} = 23.1$ Hz, C-3, C-5), 119.3 (2C, d, $J_{F-2,6C} = 8.3$ Hz, C-2, C-6), 155.2 (1C, d, $J_{F-1C} = 2.5$ Hz, C-1), 159.4 (1C, d, $J_{F-4C} = 237.6$ Hz, C-4).

Biotransformation of fluorophenols by immobilized *A. crassum* cells in sodium alginate gel

Sodium alginate (2%) was suspended in water (50 ml), which was autoclaved at 120 °C for 30 min. Cultured *A. crassum* cells (9 g) were added to this solution and the mixture was stirred for 2h until it became homogeneous. Suspension was added drop-wise from a dropping funnel with a glass tube into a 5% $CaCl_2$ solution (1l) with stirring to form pieces




Figure 1. Biotransformation of 2-fluorophenol (1) to 2-fluorophenyl β -D-glucoside (4) by cultured and immobilized cells of *A. crassum*.

of spherical sodium alginate gel with 5 mm diameter immediately. After washing with a synthetic seawater, immobilized *A. crassum* cells were used for biotransformation experiments.

To the immobilized *A. crassum* cells which included 9 g of *A. crassum* cells with 250 ml of the synthetic seawater in a 500 ml flask was added 0.2 mmol of substrate. Effect of sodium alginate concentration on the glycosylation of 2-fluorophenol (1) was examined by using the cells immobilized with sodium alginate at final concentrations of 1, 2, 3, 4, and 5%. Flask was incubated at 20 °C with aerobic shaking for five days under illumination (1000lx). Isolation of products was carried out by the similar procedures to planktonic biotransformation experiment described above.

Results and Discussion

Biotransformation product was isolated from the cultured cells of *A. crassum*, which had been treated with 2-fluorophenol (1) for 48 h. Potentially glucosylated product (4) was obtained, and no additional conversion




Figure 2. Effects of sodium alginate concentration on the glucosylation activity of the immobilized cells of *A. crassum* against 2-fluorophenol (1). Concentration (µg/g cells) of 2-fluorophenyl β -D-glucoside (4) (♦) is plotted.

Table 2. The production of **4** by immobilized *A. crassum*.



Batch	Conc. of products (μg/g cells) ^a
1	147
2	143
3	123
4	117
5	90

Note: ^aResults are expressed as average of three times experiments.

products were observed in spite of the careful HPLC analyses. Chemical structure of the product was determined on the basis of their FABMS, ¹H and ¹³C NMR, H-H COSY, C-H COSY, and HMBC spectra as 2-fluorophenyl β-D-glucoside (60 μg/g cells) (Fig. 1). Other microalga, ie, *Chattonella* sp. and *Heterosigma* sp., showed no glycosylation action against **1**.

In addition, 2-fluorophenol (**1**) was subjected to the biotransformation by immobilized *A. crassum* cells with sodium alginate. Fig. 2 shows that the glucosylation activity of immobilized cells was enhanced at 2% sodium alginate concentration. Immobilized *A. crassum* cells, which had been prepared at 2% sodium alginate concentration, were used for biotransformation experiments. After 48 h incubation of 2-fluorophenol (**1**), 2-fluorophenyl β-D-glucoside (140 μg/g cells) was obtained in higher yield in comparison with the case of the biotransformation using planktonic cells. Repetitive use of the immobilized cells was carried out. Yields of the product **4** obtained after each batch reaction were shown in Table 2. Experiments were repeated three times, and average is shown in Table 2. In repetitive batch use, the immobilized cells of *A. crassum* maintained the potential for the glucosylation of **1** (100 μg/g cells) after 5 times of usage.

Next, *A. crassum* cells were incubated with 3-fluorophenol (**2**) for 48h in the same biotransformation conditions as 2-fluorophenol (**1**). Glucosylation

**Figure 3.** Biotransformation of 3-fluorophenol (**2**) to 3-fluorophenyl β-D-glucoside (**5**) by cultured and immobilized cells of *A. crassum*.**Figure 4.** Biotransformation of 4-fluorophenol (**3**) to 4-fluorophenyl β-D-glucoside (**6**) by cultured and immobilized cells of *A. crassum*.

product was isolated from the cell cultures of *A. crassum* by preparative HPLC and was identified as 3-fluorophenyl β-D-glucoside (**5**, 20 μg/g cells) by spectroscopic methods (Fig. 3). Use of the immobilized *A. crassum* cells, which had been prepared at 2% sodium alginate concentration, improved the conversion yield of the product to 60 μg/g cells. When the substrate 4-fluorophenol (**3**) was used for the biotransformation experiment, 4-fluorophenyl β-D-glucoside (**6**) was obtained in 40 μg/g cells (Fig. 4). Incubation of immobilized *A. crassum* cells prepared at 2% sodium alginate concentration with 4-fluorophenol (**3**) afforded 4-fluorophenyl β-D-glucoside (**6**) in 100 μg/g cells.

This study demonstrated that the cultured cells of *A. crassum* can convert 2-, 3-, and 4-fluorophenols into the corresponding β-D-glucosides. Highest yield was found in the case of the biotransformation of 2-fluorophenol with *ortho*-phenolic hydroxyl group. Results of this experiment revealed that cultured cells of *A. crassum* can incorporate the environmentally toxic compounds such as 2-, 3-, and 4-fluorophenol and biotransform into β-D-glucosides. Yield of β-D-glucosides produced from fluorophenols by immobilized *A. crassum* cells in sodium alginate gel was improved in comparison with the case of biotransformation by planktonic *A. crassum* cells. Immobilized cells of *A. crassum* would be useful to detox fluorophenols to stable glycosides. This procedure is a simple operation and is environmentally friendly. Further studies on the enzymes which catalyze the glucosylation of fluorophenols are now in progress.

Acknowledgement

This work was supported by grant from The Salt Science Research Foundation, No. 1007.

Disclosure

This manuscript has been read and approved by all authors. This paper is unique and is not under



consideration by any other publication and has not been published elsewhere. The authors and peer reviewers of this paper report no conflicts of interest. The authors confirm that they have permission to reproduce any copyrighted material.

References

1. Key BD, Howell RD, Criddle CS. Fluorinated organics in the biosphere. *Environ Sci Technol*. 1997;31:2445–54.
2. Haggblom MM. Microbial breakdown of halogenated aromatic pesticides and related compounds. *FEMS Microbiol Rev*. 1992;9:29–71.
3. Kiyohara H, Hatta T, Ogawa Y, Kakuda T, Yokoyama H, Takizawa N. Isolation of *Pseudomonas pickettii* strains that degrade 2,4,6-trichlorophenol and their dechlorination of chlorophenols. *Appl Environ Microbiol*. 1992;58:1276–83.
4. Hofrichter M, Bublitz F, Fritsche W. Unspecific degradation of halogenated phenols by the soil fungus *Penicillium frequentans* Bi 7/2. *J Basic Microbiol*. 1994;34:163–72.
5. Bae HS, Lee JM, Lee ST. Biodegradation of 4-chlorophenol via a hydroquinone pathway by *Arthrobacter ureafaciens* CPR706. *FEMS Microbiol Lett*. 1996;145:125–9.
6. Hollender J, Hopp J, Dott W. Degradation of 4-chlorophenol via the meta cleavage pathway by *Comamonas testosteroni* JH5. *Appl Environ Microbiol*. 1997;63:4567–72.
7. Boersma MG, Dinarieva TY, Middelhoven WJ, et al. ^{19}F nuclear magnetic resonance as a tool to investigate microbial degradation of fluorophenols to fluorocatechols and fluoromuconates. *Appl Environ Microbiol*. 1998;64:1256–63.
8. Bondar VS, Boersma MG, van Berkel WJ, et al. Preferential oxidative dehalogenation upon conversion of 2-halophenols by *Rhodococcus opacus* 1G. *FEMS Microbiol Lett*. 1999;181:73–82.
9. Boersma MG, Solyanikova IP, van Berkel WJ, Vervoort J, Golovleva LA, Rietjens IM. ^{19}F NMR metabolomics for the elucidation of microbial degradation pathways of fluorophenols. *J Ind Microbiol Biotechnol*. 2001;26:22–34.
10. Ishihara K, Hamada H, Hirata T, Nakajima N. Biotransformation using plant cultured cells. *J Mol Catal B Enz*. 2003;23:145–70.
11. Nakajima N, Teramoto T, Kasai F, et al. Glycosylation of bisphenol A by freshwater microalgae. *Chemosphere*. 2007;69:934–41.

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>