

Isolation, Characterization and Quantity Determination of Aristolochic Acids, Toxic Compounds in Aristolochia bracteolata L.

Authors: Abdelgadir, Abdelgadir A., Ahmed, Elhadi M., and Eltohami, Mahgoub Sharif

Source: Environmental Health Insights, 5(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/EHI.S6292>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Isolation, Characterization and Quantity Determination of Aristolochic Acids, Toxic Compounds in *Aristolochia bracteolata* L.

Abdelgadir A. Abdelgadir¹, Elhadi M. Ahmed¹ and Mahgoub Sharif Eltohami²

¹Department of Chemistry and Pharmacognosy, Faculty of Pharmacy, University of Gezira, Sudan. ²Department of Pharmacognosy, Faculty of Pharmacy, Omdurman Islamic University, Sudan.

Corresponding author email: yahadi54@yahoo.com

Abstract:

Background: Aristolochic Acids (AAs) are major components of plants in *Aristolochia* and have been found to be nephrotoxic, carcinogenic and mutagenic. Herein reported are the isolation, identification and quantity determination methods of Aristolochic Acid-I (AA-I) and Aristolochic Acid-II (AA-II) toxic compounds of *Aristolochia bracteolata* indigenous to Central Sudan and medicinally used in diverse biological functions including analgesic and diuretic effects, treatment of tumors, malaria and/or fevers.

Methods and results: AAs mixture was extracted with methanol from the defatted material of *Aristolochia bracteolata* whole plant at room temperature and was isolated from the aqueous methanol extract by chloroform. Moreover, Silica-gel column chromatography and Preparative Thin Layer Chromatography (PTLC) using chloroform/methanol gradient mixtures were used to isolate AAs mixtures as a yellow crystalline solid. A preliminary detection of AAs was made by Thin Layer Chromatography (silica-gel, chloroform: methanol (6:1)). The R_f value of the acids mixture was 0.43–0.46. The presence of AAs in plant sample was confirmed by High Performance Liquid Chromatography/Ultraviolet (HPLC/UV) analysis using 1% acetic acid and methanol (40:60) as mobile phase and maximum absorption wave length of 250 nm. Quantitative determination of AA-II (49.03 g/kg) and AA-I (12.98 g/kg) was also achieved by HPLC/UV.

Recommendation: It is recommended that the use of *Aristolochia bracteolata* as a medicinal plant should be extremely limited or strictly prohibited. The chromatograms obtained in this study can serve as fingerprints to identify AAs in plant samples.

Keywords: Aristolochic acid-I, Aristolochic acid-II, *Aristolochia*, *Aristolochia bracteolata*

Environmental Health Insights 2011:5 1–8

doi: [10.4137/EHI.S6292](https://doi.org/10.4137/EHI.S6292)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.

Introduction

The genus *Aristolochia* grows over a wide area of different climatic zones and consists of about 500 species.¹ Various *Aristolochia* species have been used in herbal medicines since antiquity in obstetrics and in treatment of snakebite,² festering wounds, and tumors, and they remain in use particularly in Chinese herbal medicine.^{3,4} *Aristolochia bracteolata* is known as “worm killer” due to supposed antihelminthic activity and trypanocidal effect.⁵ *Aristolochia bracteolata* also possess a potent antiallergic activity⁶ and has pronounced antibacterial and antifungal activities.⁷

In Sudan, *Aristolochia* is traditionally used as an analgesic, antiscorpion, and antisnake. It is also used in the treatment of tumors, malaria and for fevers.⁸

Aristolochic acids (Fig. 1) are compounds of nitrophenanthrene carboxylic acids that occur naturally in plants of the family Aristolochiaceae, primarily in the genera *Aristolochia* and *Asarum*^{9,10} as well as in butterflies that feed on such plants.¹¹ Botanical products from plants containing aristolochic acids are used in traditional folk medicines, particularly in Chinese herbal medicine, especially those used as a part of weight-loss regimens.⁴

Aristolochic acids are known to be toxic and a rodent carcinogen, in addition to their carcinogenicity, aristolochic acids are also highly nephrotoxic agents.^{1,10–12} In the present work *Aristolochia bracteolata* grown in Sudan was investigated for the presence of Aristolochic acids.

Materials and Methods


Plant material

The plant *Aristolochia bracteolata* was collected during December 2009 from the Southern Gezira scheme, Sudan. The plant was taxonomically identified in the Department of Chemistry and Pharmacognosy, University of Gezira, and a voucher specimen was deposited.

Methods


Extraction of plant material

The shade dried whole plant sample (200 g) of *Aristolochia bracteolata* were milled in the form of a coarse powder. The powdered sample was defatted by maceration at room temperature using petroleum ether (40–60 °C) for 6 hours to yield dark green oily


 $C_{17}H_{11}NO_7$

Rel. mol. mass: 341.27

Aristolochic acid I (8-methoxy-6-nitrophenanthro [3, 4-*d*][1, 3] dioxole-5-carboxylic acid).


 $C_{16}H_9NO_6$

Rel. mol. mass: 311.25

Aristolochic acid II (6-nitrophenanthro [3, 4-*d*]-1, 3-dioxole-5-carboxylic acid).

Figure 1. Chemical structure of aristolochic acid I and aristolochic acid II.

residue after evaporation of the solvent. As described in previous studies,^{13,14} and with some modifications (defatting before methanol extraction), the dried defatted residue was macerated in methanol at room temperature for 72 hours and filtered. Solvent was evaporated to dryness under reduced pressure using a rotary evaporator at 60 °C to give a dark green methanol extract.

Liquid-liquid extraction

Twenty grams of *Aristolochia bracteolata* methanol extract was partitioned between water (450 ml) and chloroform (800 ml). Chloroform layer was separated and dried by rotary evaporator at 40 °C to give chloroform extract.

Column chromatography

Chloroform extract of *Aristolochia bracteolata* (4.0 g) was chromatographed on a silica-gel column. For column fractionation the chloroform extract was dissolved in 20 ml chloroform and applied to the top of a column (8.0 cm ID) backed with 160 g silica-gel of a mesh size of 60–120.

Elution was commenced with gradient chloroform/methanol mixture (8:1). Polarity of the solvent was increased in a gradient of 8:1, 4:1, 1:1, 0:1 and finally the column was washed with 100 ml of water. 200 ml of eluents were used and fractions of 50 ml were collected and thin layer chromatography (TLC) monitored. Similar fractions were combined to give three fractions 1–7 (a), 8–14 (b) and 15–17 (c). Fraction (b) was further subjected to column chromatography (3.5 cm ID) backed with 40 g silica-gel of mesh size 60–120, and eluted with a chloroform/methanol mixture of increasing polarity as: 8:1, 4:1, 1:1, 0:1. Fractions of 25 ml were collected and TLC monitored using chloroform/methanol mixture (6:1) as mobile phase (Fig. 2).

Thin layer chromatography (TLC)

TLC glass backed plates were prepared using silica-gel G type 60 (s d Fine-Chem Limited, Mumbai, India). A slurry of silica-gel in water was applied by means of a spreader to glass plates, previously




Figure 2. Isolated Aristolochic acids (left), silica-gel fluorescence TLC aluminum sheet, chloroform/methanol (6:1) mobile phase (under UV at 365 nm), compare to standard Aristolochic acids (right).

cleaned with methanol. A layer 0.25 mm thick was used routinely. Plates were activated by heating at 100–110 °C for an hour. TLC aluminum sheets 20 × 20 cm, 1 mm thick, silica-gel 60 F₂₅₄ (Merk, Germany) and Cellulose plastic papers 20 × 20 cm, 1 mm thick (Macherey-Nagel, Germany) were also used. Different mobile phases chloroform/methanol (6:1);¹⁵ chloroform/methanol/acetic acid (65:20:5);¹⁶ benzene/acetone (14:1),¹⁷ chloroform/methanol/water (60:40:10)¹⁸ were checked for the separation of aristolochic acids.

Chromatographic responses (Table 1) were obtained and compared to those of Aristolochic acids reference sample Aristolochic acid mixture of AA-I and AA-II (Aldrich Chemical Company, USA). Day light, ultraviolet light (UV), 0.5% Diphenylamine in 60% H₂SO₄ (DPA/H₂SO₄) and Dragendurff's reagents were used as locating agents to disclose the aristolochic acids on TLC and preparative TLC chromatograms (Fig. 2).

Preparative thin layer chromatography (PTLC)

The dried material obtained from the combined column chromatography fractions which are rich in aristolochic acids (Fraction b) was dissolved in 10 ml chloroform/methanol mixture (6:1) and subjected to preparative thin layer chromatography (PTLC) using 20 × 20 cm cellulose papers for TLC with chloroform/methanol/water (6:4:1) as mobile phase and TLC aluminum sheets: (20 × 20 cm), silica-gel 60 F₂₅₄ with chloroform/methanol (6:1) as mobile phase. The disclosed bands of aristolochic acids were then scraped off, dissolved in chloroform/methanol mixture (6:1) and filtered. The solvent was removed by evaporation and the obtained isolate containing aristolochic acids was subjected to further PTLC for purification using silica-gel fluorescent TLC aluminum sheets as described above (Figs. 3 and 4). Scraped Aristolochic acids were dissolved and filtered. Solvent was removed to yield a yellow crystalline aristolochic acids mixture.

C18-High-performance liquid chromatography (HPLC)

The separation and quantification of aristolochic acids was achieved on HPLC chromatograph, (Waters® 2996, USA); C18 250 × 4.6 mm column with UV detector. A mixture of 1% acetic acid and methanol

Table 1. Observed chromatographic responses of Aristolochic acids on TLC chromatograms.

Type of adsorbent	Solvent system	Rf value	Appearance of spots/bands		
			Day light	UV at 365 nm	0.5% DPA/60% H ₂ SO ₄
Silica-gel 60 F ₂₅₄ , precoated aluminum sheets, 20 × 20 cm (Merk, Germany)	CHCl ₃ /MeOH (6:1)	0.43	Yellow fluorescence	Black spot	Pale grey spot
Silica-gel G glass plates (s d Fine-Chem., Limited, Mumbai, India)	CHCl ₃ /MeOH (6:1)	0.46	Yellow fluorescence	Faint brown spot	Pale grey spot
Cellulose, precoated plastic sheets, 20 × 20 cm (Macherey-Nagel, Germany)	CHCl ₃ /MeOH/Water (6:4:1)	0.83	Yellow fluorescence	Faint brown spot	Pale grey spot

(40:60) utilized as mobile phase, flow rate 1.0 ml/min, pressure 3015 psr, temperature 20–25 °C, UV detection 250 nm, Helium degassing, and injection volume 20 µL.

Preparation of standard Aristolochic acids, isolated Aristolochic acids and methanol extract of *Aristolochia bracteolata*

Fifteen milligrams of Aristolochic acids mixture of AA-I and AA-II (Aldrich Chemical Company, USA), fifteen mg of isolated aristolochic acids of *Aristolochia bracteolata* and one g of methanol extract of *Aristolochia bracteolata* were weighed separately using digital balance, each was placed in a small beaker, transferred into volumetric flask (50 ml), washed by methanol-HPLC grade and adjusted to 50 ml, filtered and sonicated for 10 minutes. 20 µL were separately

injected in the HPLC system. Identification was made by comparing the samples retention times with those of the standard reference (Figs. 5–7).

Results and Discussion

To isolate and identify Aristolochic acids (AAs) in *Aristolochia bracteolata* L., the whole plant (200 g) was defatted to yield dark green oily residue (5.35%). As described in previous studies^{13,14} and with some modifications (defatting before methanol extraction, further n-butanol was not used in successive partitioning), the defatted plant material produced on drying 16.5% crude dark green extract on subsequent methanol maceration. On column chromatography of chloroform fraction obtained from the aqueous methanol extract, fraction (b) eluted by 4:1 and 1:1 solvent mixture was found to be rich in Aristolochic acids as




Figure 3. Aristolochic acids PTLC chromatogram, silica-gel fluorescence TLC aluminum sheet, chloroform/methanol (6:1) mobile phase, (day light).





Figure 4. Aristolochic acids PTLC chromatogram, silica-gel fluorescence TLC aluminum sheet, chloroform/methanol (6:1) mobile phase, (under UV at 365 nm).



	Peak Name	RT	Area	% Area	Height
1	Peak1	2.402	26796089	32.59	1292511
2	Peak2	3.029			
3	Peak3	3.281	9595634	11.67	330078
4		4.041	3932822	4.78	134660
5		4.741	1810100	2.20	56422
6		5.331	963237	1.17	49228
7		5.837	3009459	3.66	69800
8	Peak4	6.933			
9	Peak5	7.359	4581045	5.57	119377
10	Peak6	10.482	632544	0.77	17826
11	AA-II	18.426	24433508	29.72	289625
12	AA-I	27.897	6469776	7.87	63686

Figure 5. HPLC chromatogram of methanol extract sample of *Aristolochia bracteolata*.


Condition: column, C18 (Varian® microsorb-MV 300-8 C18 250 × 4.6 mm), mobile phase, 1% acetic acid/methanol (40:60), temperature 20–25 °C, flow rate 1.0 ml/min, UV detection 250 nm.

revealed by TLC monitoring. Column chromatography fractionation of fraction (b) with the same gradient mobile phase and TLC monitoring showed that Aristolochic acids were obtained as major spots in the first ten fractions. Sixteen milligrams of a yellow powder of Aristolochic acids mixture (Fig. 2) was produced thereafter by preparative thin layer chromatography.

It was noted in previous studies that many attempts have been made to assay active components in Aristolochiaceae by various methods including Thin Layer Chromatography, High Performance Liquid Chromatography-Ultraviolet. However, most of these

methods focus on analysis of AA-I and AA-II in samples.^{11,14,15,19,20}

In the present study TLC and PTLC chromatograms clearly demonstrated the presence of Aristolochic acids. Results also indicated that the obtained AAs produced varying chromatographic responses with different chromatographic solvent systems and/or locating agents (Table 1). Aristolochic acids generally acquired black color on fluorescent silica-gel in UV-light at 365 nm while they fluoresce yellow in day light. Rf values ranged 0.43–0.46 (chloroform/methanol (6:1); silica-gel) and 0.83 (chloroform/methanol/water (6:4:1); cellulose).



	Peak Name	RT	Area	% Area	Height
1	Peak1	2.351	74044	0.85	5365
2	Peak2	2.633			
3	Peak3	3.067	22897	0.26	1983
4		3.209	40940	0.47	2449
5		3.583	751552	8.59	2666
6	Peak4	4.265			
7	Peak5	5.127			
8	Peak6	7.951			
9	Peak7	10.094			
10		18.839	5480168	62.62	71021
11		27.992	2381515	27.21	26055


Figure 6. HPLC chromatogram of isolated AAs sample of *Aristolochia bracteolata*.

Condition: column, C18 (Varian® microsorb-MV 300-8 C18 250 × 4.6 mm), mobile phase, 1% acetic acid/methanol (40:60), temperature 20–25 °C, flow rate 1.0 ml/min, UV detection 250.

For separation and quantification of Aristolochic acids by HPLC earlier reports mainly used a mobile phase of methanol and water^{21–24} or acetonitrile and water^{21,25,26} and the optimum chromatographic conditions were investigated by varying the content of methanol and water.

In this study, and for the first time in Sudan, the presence of Aristolochic acids in *Aristolochia bracteolata* methanol extract was confirmed and quantified by High Performance Liquid Chromatography as shown in Figures 5–7 using a mixture of 1% acetic acid and methanol (40:60) as mobile phase. Aristolochic acid II (AA-II) and Aristolochic acid I (AA-I) were detected as

major peaks of components at a retention time of about 18.4 min and 27.9 min respectively and exactly corresponding to data profile exhibited by the Aristolochic acid reference sample. HPLC data also showed that AA-II was represented in a higher calculated amount of 49.03 g/kg compared to AA-I (12.98 g/kg) in *Aristolochia bracteolata* L. whole plant while in literature, other species of *Aristolochia* had recorded far more lower amounts of AAs such as *Aristolochia debilis* that contained 0.18 g/kg of AA-II and 1.01 g/kg of AA-I; *Aristolochia manshuriensi* contained 1.0 g/kg of AA-II and 8.82 g/kg of AA-I; *Aristolochia fangchi* also contained 0.22 g/kg of AA-II and 2.22 g/kg of AA-I.^{1,11,27–32}



	Peak Name	RT	Area	% Area	Height
1	Peak1	2.408	133058	0.28	8675
2	Peak2	3.029	15311	0.03	1354
3	Peak3	3.361	38589	0.08	2092
4	Peak4	6.933	15507	0.03	1391
5	Peak5	7.503	407054	0.85	14153
6	Peak6	10.946	156466	0.33	4078
7	AA-II	19.078	21782025	45.33	281529
8	AA-I	27.877	25503012	53.07	286251

Figure 7. HPLC chromatogram of AAs standard sample.

Condition: column, C18 (Varian® microsorb-MV 300-8 C18 250 × 4.6 mm), mobile phase, 1% acetic acid/methanol (40:60), temperature 20–25 °C, flow rate 1.0 ml/min, UV detection 250 nm.

It is worth noting that, in the reported data AA-I obtained from different *Aristolochia* species was found to be in a higher quantity than AA-II, contrary to results obtained in this study. This may be attributed to climatic or biological variations which could affect metabolic pathways of interconversion of Aristolochic acids and analogues.

In conclusion, the presence of Aristolochic acids toxic compounds in *Aristolochia bracteolata* L. methanol extract was evident through the multi-component method including Column Chromatography (CC), Thin Layer Chromatography (TLC), Preparative Thin Layer Chromatography (PTLC) and had been confirmed and quantified by High Performance Liquid Chromatography (HPLC).

Thus it is recommended that the use of *Aristolochia bracteolata* as a medicinal plant should be extremely limited or strictly prohibited. Moreover, the chromatograms obtained in this study can serve as fingerprints to identify plants suspected to contain or adulterated with Aristolochic acids.

Disclosure

This manuscript has been read and approved by all authors. This paper is unique and is not under consideration by any other publication and has not been published elsewhere. The authors and peer reviewers of this paper report no conflicts of interest. The authors confirm that they have permission to reproduce any copyrighted material.



References

- IARC (International Agency for Research on Cancer). Some Traditional Herbal Medicines, Some Mycotoxins, Naphthalene and Styrene. *IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans*. 2002;82:69–128. Lyon, France.
- Meenatchisundaram S, Prajish Parameswari G, Subbraj T, Michael A. Studies on antivenom activity of *Andrographis paniculata* and *Aristolochia indica* plant extracts against *Echis carinatus* venom. *The Internet Journal of Toxicology*. 2009;6(1):1559–3916.
- Balachandran P, Wei F, Lin R, Khan I, Pasco D. Structure activity relationships of aristolochic acid analogues: Toxicity in cultured renal epithelial cells. *Journal of Kidney International*. 2005;67:1797–805.
- NTP (National toxicological program), United States department of health and human services. Final report on carcinogens background document for aristolochic acids. *Research Triangle Park, NC 27709*. 2008:1–12.
- Samia HAR, Elmalki KH, Khalid HS. Therapeutic Effect of *Aristolochia bracteolata* Extract Against Experimental *Trypanosoma evansi* Infection. *International Journal of Tropical Medicine*. 2006;1(4):170–2.
- Chitme H, Malipatil M, Chandrashekar V, Prashand P. Antiallergic activity of *Aristolochia bracteolata* Lank in animal model. *Indian Journal of Experimental Biology*. 2010;48:46–5.
- Kavitha D, Nirmaladevi R. Assessment of *Aristolochia bracteolata* leaf extracts for its biotherapeutic potential. *African Journal of Biotechnology*. 2007;8(17):4242–4.
- El-Tahir A, Satti G, Khalid S. Antiplasmodial Activity of Selected Sudanese Medicinal Plants with Emphasis on *Acacia nilotica*. *Phytotherapy Research*. 1999;13:474–8.
- Huang C, Tseng M, Lin J. Analyzing Aristolochic Acids in Chinese Herbal Preparations Using LC/MS/MS. *Journal of Food and Drug Analysis*. 2005; 13:125–31.
- Hsieh S, Huang M, Lin B, Chang H. Determination of aristolochic acid in Chinese herbal medicine by capillary electrophoresis with laser-induced fluorescence detection. *Journal of Chromatography*. 2006;1105:127–34.
- Yuan J, Liu Q, Wei G, Tang F, Ding L, Yao S. Characterization and determination of six aristolochic acids and three aristolactams in medicinal plants and their preparations by high-performance liquid chromatography-photodiode array detection/electrospray ionization mass spectrometry. *Rapid Communication in Mass Spectrometry*. 2007;21:2332–42.
- Jou J, Li CP, Schelonka E, Lin C, Wu T. Analysis of the Analogues of Aristolochic Acid and Aristolactam in the Plant of *Aristolochia* Genus by HPLC. *Journal of Food and Drug Analysis*. 2004;12:40–5.
- Wu T, Chan Y, Leu Y. The constituent of the Root and Stem of *Aristolochia cucubifolia* Hayata and Their Biological Activity. *Chemical and Pharmaceutical Bulletin*. 2000;48(7):1006–9.
- Stacheno E, Andres O, Armando M, Rene M. Determination of Volatile and Semi-volatile Secondary Metabolites, and Aristolochic Acids in *Aristolochia Ringens* Vahl. *Journal of Chromatographic Science*. 2009;47(9):817–21.
- Watanabe K, Miyakado M, Iwai T, Izumi K, Yanagi K. Isolation of Aristolochic Acid and Aristolic Acid from *Cocculus trilobus* DC as Potent Seed Germination Inhibitor. *Agricultural and Biological Chemistry*. 1988;52(4):1079–82.
- Ioset JR, Raelison GE, Hostettmann K. Detection of aristolochic acid in Chinese phytomedicines and dietary supplements used as slimming regimens. *Food and Chemical Toxicology*. 2003;41(1):29–36.
- Almahy HA, Khalid HE. Chemical Examination of the Leaves *Nerium oleander*. *International Journal of Tropical Medicine*. 2006;1(2):58–62.
- Seto T, Hamano T, Shioda H, Kamimura H. Analysis of Aristolochic acid and in Kampo Medicine Preparations. *Journal of Health and Science*. 2002;48(5):412–7.
- Chinese Pharmacopoeia Committee. The Chinese Pharmacopoeia. *Chemical Industry Press*: 25, 35, 37, 101, Beijing; 2005.
- Sathornviriyapong S, Pichansoonthon C, Tiasakul R, Tiyaworanant S, Reutrakul V. Botanical Origin Identification of Krai-Krue Herbal Plant. *Kasetsart Journal (Natural Science)*. 2007;41:420–432.
- Sun Q, Wu Y, Jia L. Quantitative Determination of Aristolochic Acid in *Asarum heterotropoides* Fr. Schmidt var. *mandshuricum* (Maxim.) Kitag. *Asian Journal of Traditional Medicines*. 2001:5–13.
- Xie ZM, Li SX, Liao HC. Determination of aristolochic acid A in herba asari by HPLC. *Central South Pharmacy*. 2003;1(3):165–7.
- Jiang X, Wang ZM, You LS, et al. Determination of aristolochic acid in Radix Aristolochiae and Herba Asari by RP-HPLC. *China Journal of Chinese Materia Medica*. 2004;29(5):408–10.
- Guan WQ, Li XM, Xiao JS. Quantitative determination of aristolochic acid in herba asari and its preparations by RPHPLC. *Chinese Remedies and Clinics*. 2005;5(4):283–5.
- Lu J. Studies on the aristolochic acid related crude traditional Chinese medicine. *Chinese Material Medica Standard*. 2002;3(2):49–50, 61.
- Geng LJ, Li BG. Determination of aristolochic acid A in herba asari and its preparation. *Chemical Analysis and Measure*. 2005;14(2):25–7.
- Hashimoto K, Higuchi M, Makino B, et al. Quantitative analysis of aristolochic acids, toxic compounds, contained in some medicinal plants. *Journal of Ethnopharmacology*. 1999;64:185–9.
- Ong ES, Woo SO. Determination of aristolochic acids in medicinal plants (Chinese) prepared medicine using capillary zone electrophoresis. *Electrophoresis*. 2001;22:2236–41.
- Lee TY, Wu ML, Deng JF, Hwang DF. High-performance liquid chromatographic determination for aristolochic acid in medicinal plants and slimming products. *Journal Chromatography B, Analytical Technologies in the Biomedical and Life Sciences*. 2001;766(1):169–74.
- Zhai ZD, Luo XP, Shi YP. Separation and determination of aristolochic acids in herbal medicines by microemulsion electrokinetic chromatography. *Analytica Chimica Acta*. 2006;561(1–2):119–25.
- Zhang C, Wang X, Shang M, et al. Simultaneous determination of five aristolochic acids and two aristolactams in *Aristolochia* plants by high-performance liquid chromatography. *Biomedical Chromatography*. 2006b; 20(4):309–18.
- Wu KM, Farrelly JG, Upton R, Chen J. Complexities of the herbal nomenclature system in traditional Chinese medicine (TCM): lessons learned from the misuse of *Aristolochia*-related species and the importance of the pharmaceutical name during botanical drug product development. *Phytomedicine*. 2007a;14(4):273–9.

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>