

Allium Achaium Boiss. (Alliaceae), a Critical Species of Greek Flora

Authors: Bogdanovic', Sandro, Brullo, Cristian, Brullo, Salvatore, Galdo, Gianpietro Giusso del, Maria Musarella, Carmelo, et al.

Source: Candollea, 66(1) : 57-64

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2011v661a3>

BioOne Complete (complete.bioone.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Allium achaicum Boiss. (Alliaceae), a critical species of Greek flora

Sandro Bogdanović, Cristian Brullo, Salvatore Brullo, Gianpietro Giusso del Galdo,
Carmelo Maria Musarella & Cristina Salmeri

Abstract

BOGDANOVIC, S., C. BRULLO, S. BRULLO, G. GIUSSO DEL GALDO, C. M. MUSARELLA & C. SALMERI (2011). Allium achaicum Boiss. (Alliaceae), a critical species of Greek flora. *Candollea* 66: 57-64. In English, English and French abstracts.

Allium achaicum Boiss. (*Alliaceae*), a critical and misappreciated species of Greek flora is investigated from a taxonomic point of view and illustrated. It belongs to sect. *Codonoprasum* Rchb. and shows close relationships with the taxa of the *Allium stamineum* Boiss. group. Karyology, ecology and distribution of that species are examined.

Key-words

ALLIACEAE – Allium – Greece – Taxonomy – Karyology

Résumé

BOGDANOVIC, S., C. BRULLO, S. BRULLO, G. GIUSSO DEL GALDO, C. M. MUSARELLA & C. SALMERI (2011). Allium achaicum Boiss. (Alliaceae), une espèce critique de la flore grecque. *Candollea* 66: 57-64. En anglais, résumés anglais et français.

Allium achaicum Boiss. (*Alliaceae*), une espèce critique et mal connue de la flore grecque, est étudiée du point de vue taxonomique et illustrée. Elle appartient à la sect. *Codonoprasum* Rchb., et montre une remarquable affinité avec les taxons du groupe de *Allium stamineum* Boiss. La caryologie, l'écologie et la distribution de cette espèce sont aussi examinées.

Addresses of the authors: SB: University of Zagreb, Faculty of Science, Department of Botany and Botanical Garden, Marulićev trg 9a, 10000 Zagreb, Croatia.
Email: sandro@botanic.hr

CB, SB, GGG, CMM: Università degli Studi di Catania, Dipartimento di Botanica, Via A. Longo 19, 95125 Catania, Italia.

CS: Università di Palermo, Dipartimento di Scienze Botaniche, Via Archirafi 38, 90123 Palermo, Italy.

Submitted on July 9, 2009. Accepted on February 17, 2011.

Edited by P. Bungener

Introduction

The aim of this work is to clarify the taxonomical position of *Allium achaicum* Boiss., a critical species of Greek flora. It was described by BOISSIER (1882) based on specimens collected by Orphanides in Mt Klokos (North Peloponnisos) and treated as closely related to *A. paniculatum* L. Besides, the same author included within *A. achaicum* a population from Mt. Parnassos describing it as var. *parnassicum* Boiss. Afterwards, this species was treated by HALÁCSY (1904), HAYEK (1924-1933) and STEARN (1978, 1980, 1981) as synonym of *A. frigidum* Boiss. & Heldr., while ANDERSSON (1991), TZANOUDAKIS & TAN (2000) and TAN & IATROU (2001) keep it at a specific rank.

Morphological investigations, carried out both on dried and living material coming from the *locus classicus* and many other Greek localities, highlighted that the true *A. achaicum* is a well differentiated species, belonging to the *A. stamineum* Boiss. group. Therefore, the taxonomic treatment of *A. achaicum* as synonym of *A. frigidum* must be rejected. According to BRULLO & al. (2001), *A. frigidum* is characterized by spathe valves shorter than the pendulous inflorescence, stamens included in the perigon, with interstaminal teeth and white filaments, ovary ellipsoid and capsule subglobose, while *A. achaicum* has spathe valves longer than the umbel, which is erect, stamens exserted from the perigon, without interstaminal teeth, and filaments purplish above, ovary subglobose and capsule globose.

Several morphological characters allow *A. achaicum* to be well distinguished from *A. parnassicum* (Boiss.) Halácsy, which is indeed closely related to *A. frigidum* (cf. BRULLO & al., 2001).

Due to its geographical distribution, *A. achaicum* occurs only on some mountains of Peloponnisos, therefore the records of ANDERSSON (1991) and TAN & IATROU (2001) for Sterea Ellas should be considered a misinterpretation. In fact, based on herbarium and field investigations, this species is a rare orophyte found within the thorny, cushion-like plant communities of North and Central Peloponnisos.

Materials and methods

Our investigation was based on living material of *A. achaicum* collected in various mountains of Peloponnisos (Mt. Klokos, Mt. Parnon, Mt. Panachaiko, Mt. Chelmos, Mt. Erimanthos and Mt. Menalo), and cultivated in the Botanic Garden of Catania. In addition, herbarium specimens of *A. achaicum*, deposited in many European herbaria (BM, CAT, G, K, WU), were examined for the morphological study. The karyological analyses were made on mitotic plates from root-tip cells of cultivated bulbs coming from above mentioned localities, pre-treated with 0.3% colchicine water solution, fixed in ethanol-acetic (3:1) and stained according to the Feulgen technique. Metaphases observations and chromosome

measurements were made using the image analysis systems IKAROS 4.6 (Metasystem) and AXIOVISION 5.1 (ZEISS). Karyotyping was done with the software Cromolab[©] 1.1. (BRULLO, 2002-2003) specialized in recognizing chromosome pairs, ordering them by size and classifying by morphology, and assembling the karyotype formula based on the centromere position (LEVAN & al., 1964; TZANOUDAKIS, 1983). Eleven mitotic plates from six individuals were used for determining the karyotype parameters. To estimate the karyotype asymmetry, different numerical parameters were calculated, such as categories of STEBBINS (1971), REC and SYI (GREILHUBER & SPETA, 1976), TF% (HUZIWARA, 1962), A1 and A2 (ROMERO ZARCO, 1986), DI (LAVANIA & SRIVASTAVA, 1992), CV_{CI}, CV_{CL} and AI index (PASZKO, 2006). The leaf anatomy was studied on the living material, sectioned using the freezing microtome.

Results

Allium achaicum Boiss., Fl. Orient. 5: 259. 1882 (Fig. 1, 2).

Lectotypus (designated by ANDERSSON, 1991): **GREECE**.

PELOPONNISOS: In monte Clocos Achajae prope Bostitzam loco dicto Pente Vryses, 4000 m, 9-21.VII.1855, *Orphanides* 427 (G-BOIS!; iso-: G!, K!, WU!).

Bulb ovoid, 12-25 × 10-15 mm, with outer tunics fibrous, dark-brown, splitting in thin parallel fibres, the inner ones pale-brown to whitish. *Stem* 10-30 cm tall, cylindrical, 2-5 mm in diameter, glabrous, erect, covered for $\frac{1}{2}$ - $\frac{2}{3}$ of its length by leaf sheaths. *Leaves* 4, 7-15 cm long, 1.5-3 mm wide, compact, subequal or shorter than stem, green-glaucous, glabrous. *Spathe* valves unequal, persistent, the larger one longer than the umbel, 3-8 cm long, 7-nerved, the smaller one shorter to longer than the umbel, 2-3.5 cm long, 5-nerved. *Umbel* lax and expanded, 3-5 cm in diameter, 9-45 flowered, with unequal pedicels, 10-20-(25) mm long. *Perigon* campanulate; tepals equal, white-yellowish, tinged with pink, pruinose, elliptical, smooth, rounded at the apex, 4.5-5 mm long and 1.8-2 mm wide, midrib green-purplish. *Stamens* exserted from perigon, with simple filaments, subequal, purplish above and white below, 4-6 mm long, below connate into an anulus 0.5 mm high. *Anthers* yellow, elliptical, 1.5-1.6 mm long, 0.8-0.9 mm wide, rounded at the apex. *Ovary* ovoid, green, lightly tuberculate above, 2-2.2 mm long, 1.8-2 mm wide, slightly contracted below. *Style* white, 2-2.5 mm long. *Capsule* trivalved, globose, lightly flattened, 2.8-3 × 3.3-3.8 mm.

Distribution, ecology and conservation status. – *Allium achaicum* occurs in several mountains of Peloponnisos (Mt. Klokos, Mt. Parnon, Mt. Panachaiko, Mt. Chelmos, Mt. Erimanthos and Mt. Menalo), where it grows at an altitude of ca. 1200-2000 m (Fig. 3). It is normally quite rare, chiefly localizing on rocky places colonized by thorny cushion-like




Fig. 1. – *Allium achaicum* Boiss. A. Habit; B. Spathe valves; C. Flower; D. Open perigon; E. Anther; F. Ovary; G. Capsule.

[Brullo, Giusso & Musarella s.n., CAT] [Drawn by Salvatore Brullo]



Fig. 2. – Inflorescence of *Allium achaicum* Boiss.

[Photo: S. Brullo]



Fig. 3. – Geographical distribution of *Allium achaicum* Boiss.

plant communities. Floristically, this vegetation is characterized by the dominance of numerous pulvinate orophytes, most of them endemic (*Astragalus rumelicus* subsp. *taygeticus* (Sirj.) Podlech, *Astragalus angustifolius* Lam. subsp. *angustifolius*, *Marrubium cylindricum* Boiss. & Heldr, *Cerastium candidissimum* Correns, *Galium thymifolium* Boiss. & Heldr, *Pterocephalus perennis* Coult. subsp. *perennis*, *Erodium chrysanthum* DC., *Asperula boissieri* Boiss., *Centaurea pichleri* Boiss., etc.). The bioclimate affecting the growing site of *Allium achaicum* prevalently coincides with the supramediterranean belt, only marginally with the mesomediterranean one, while the substrate is represented by Mesozoic limestones and dolomites.

At present, *A. achaicum* can be considered as a threatened species on the few populations with low number of individuals. Therefore, this species should be added to the ‘Red List of Threatened Species’ as vulnerable (VU), and following the criteria adopted by IUCN (2001, 2003, 2006), it should be included in the category VU and D1.

Karyology. – *Allium achaicum* is a tetraploid species with a somatic chromosome number $2n = 32$ (Fig. 4A); both the somatic number and the basic one ($x = 8$) are quite common



Fig. 4. – *Allium achaicum* Boiss. A. Metaphasic plate ($2n = 32$); B. Karyogram.

within the *Allium* sect. *Codonoprasum* Rchb. All investigated specimens show a very regular and homogeneous complement with an autoploid arrangement. On the whole, chromosomes assemble in 8 sets of four homologues which are nearly all metacentric. Only one set tends towards the submetacentric type (arm ratio exceeding 1.30) and chromosomes are indicated as «msm» according to TZANODAKIS (1983). Short satellites are well evident only in Mt. Klokos populations, occurring on the short arm of one metacentric set (Fig. 4B). The karyotype formula can be resumed as $2n = 4x = 32: 24m + 4m^{sat} + 4msm$. The absolute chromosome length varies from $11 \pm 1.95 \mu\text{m}$ for the longest chromosome, to $6.56 \pm 0.99 \mu\text{m}$ for the shortest one, while the relative length ranges from $3.86 \pm 0.37\%$ to $2.20 \pm 0.13\%$. Many different indices were also calculated to evaluate the symmetry degree of the chromosome complement; all values remarked a high homogeneity in the karyotype of *A. achaicum* (Table 1).

Leaf anatomy. – The leaf cross section of *Allium achaicum* has a subpentagonal outline, with rounded edges. The epidermis is covered by a well developed cuticle. The several stomata are distributed along the whole leaf perimeter. The palisade tissue is regular, compact, and two-layered, with the cells of

the abaxial face bigger than those ones of adaxial face. The spongy tissue is compact with small cells, slightly larger in the middle part. Several secretory canals occur in the outermost part of the spongy tissue. The vascular bundles are 20, of which 12 abaxial (6 are bigger), and 8 adaxial (all small) (Fig. 5).

Taxonomic discussion. – According to the literature data (HALÁCSY, 1904; HAYEK, 1924-1933; STEARN, 1980), *A. achaicum* is a misappreciated species usually treated as synonym of *A. frigidum*. This is probably due to both its infrequency and very scattered distribution and the lack of in-depth morphological investigations of this species. Moreover, it must be highlighted that in the most important European herbaria, specimens belonging to the typical population of this species are rather rare. For *A. frigidum*, its ecological requirements are similar to those of *A. achaicum*, as testified by their usual co-occurrence in the same stands, but remarkable morphological differences as mentioned above and in Table 2 distinguish these species quite well. Also the chromosome number differs, since *A. frigidum* is usually diploid or rarely triploid ($2n = 16, 24$), as quoted by TZANODAKIS (1983) and BRULLO & al. (2001). Taxonomically, *A. achaicum* is instead closely related to *A. stamineum* s.l., which is widely distributed in the eastern




Fig. 5. – *Allium achaicum* Boiss. Leaf cross section.

Table 1. – *Allium ochrarium* Boiss. Measures and classification of chromosomes.

Chrom.	Absolute length [µm]		Relative length [%]		Ratio I/s	
	Long arm Mean ± S.D.	Short arm Mean ± S.D.	Total length Mean ± S.D.	Long arm Mean ± S.D.	Short arm Mean ± S.D.	Total length Mean ± S.D.
1	5.79 ± 0.99	5.3 ± 0.99	11.09 ± 1.95	2.02 ± 0.15	1.84 ± 0.08	3.86 ± 0.21
2	5.68 ± 1.14	5.18 ± 0.93	10.85 ± 2	1.97 ± 0.18	1.97 ± 0.18	3.77 ± 0.18
3	5.35 ± 0.83	4.83 ± 0.81	10.18 ± 1.6	1.87 ± 0.1	1.68 ± 0.02	3.54 ± 0.09
4	5.21 ± 0.85	4.85 ± 0.78	10.06 ± 1.6	1.81 ± 0.08	1.69 ± 0.04	3.5 ± 0.06
5	6.4 ± 0.78	4.74 ± 1.31	11.14 ± 2	2.24 ± 13	3.87 ± 0.16	3.87 ± 0.16
6	6.16 ± 0.69	4.59 ± 1.37	10.75 ± 1.94	2.16 ± 0.15	1.58 ± 0.25	3.74 ± 0.19
7	6.09 ± 0.87	4.36 ± 1.12	10.45 ± 1.98	2.13 ± 0.1	1.51 ± 0.18	3.63 ± 0.24
8	5.9 ± 0.96	4.39 ± 1.34	10.29 ± 2.1	2.06 ± 0.14	1.51 ± 0.26	3.57 ± 0.31
9	5.25 ± 0.81	4.39 ± 0.53	9.64 ± 1.24	1.83 ± 0.1	1.54 ± 0.13	3.37 ± 0.12
10	4.99 ± 0.8	4.38 ± 0.73	9.36 ± 1.51	1.74 ± 0.04	1.52 ± 0.04	3.26 ± 0.01
11	4.76 ± 0.8	4.46 ± 0.64	9.23 ± 1.44	1.66 ± 0.03	1.56 ± 0.05	3.21 ± 0.03
12	4.73 ± 0.78	4.3 ± 0.68	9.03 ± 1.42	1.64 ± 0.05	1.5 ± 0.07	3.14 ± 0.03
13	4.91 ± 0.94	4.61 ± 0.82	9.53 ± 1.75	1.7 ± 0.07	1.6 ± 0.05	3.31 ± 0.12
14	4.73 ± 0.89	4.51 ± 0.83	9.24 ± 1.71	1.64 ± 0.1	1.57 ± 0.07	3.21 ± 0.17
15	4.85 ± 0.95	4.3 ± 0.86	9.14 ± 1.77	1.68 ± 0.11	1.49 ± 0.13	3.17 ± 0.2
16	4.8 ± 0.92	4.25 ± 0.92	9.05 ± 1.83	1.67 ± 0.1	1.47 ± 0.1	3.14 ± 0.22
17	4.83 ± 0.83	4.21 ± 0.55	9.04 ± 1.35	1.68 ± 0.11	1.47 ± 0.08	3.15 ± 0.15
18	4.85 ± 0.79	4.06 ± 0.66	8.91 ± 1.39	1.69 ± 0.15	1.41 ± 0.03	3.11 ± 0.15
19	4.6 ± 0.49	3.91 ± 0.49	8.51 ± 0.94	1.62 ± 0.15	1.37 ± 0.07	2.98 ± 0.2
20	4.53 ± 0.54	3.73 ± 0.35	8.25 ± 0.84	1.59 ± 0.14	1.31 ± 0.09	2.89 ± 0.21
21	4.65 ± 0.76	4.06 ± 0.8	9.15 ± 1.6	1.62 ± 0.07	1.41 ± 0.07	3.18 ± 0.19
22	4.55 ± 0.84	3.9 ± 0.87	8.88 ± 1.59	1.58 ± 0.07	1.35 ± 0.1	3.09 ± 0.17
23	4.39 ± 0.64	4.03 ± 1.01	8.83 ± 1.76	1.53 ± 0.04	1.39 ± 0.19	3.07 ± 0.31
24	4.33 ± 0.77	3.63 ± 0.66	8.2 ± 1.3	1.5 ± 0.03	1.26 ± 0.1	2.86 ± 0.04
25	4.59 ± 0.92	3.54 ± 0.68	8.13 ± 1.54	1.59 ± 0.14	1.23 ± 0.05	2.82 ± 0.14
26	4.35 ± 0.75	3.54 ± 0.71	7.89 ± 1.45	1.51 ± 0.06	1.23 ± 0.05	2.74 ± 0.08
27	4.21 ± 0.74	3.44 ± 0.65	7.65 ± 1.37	1.47 ± 0.08	1.19 ± 0.06	2.66 ± 0.08
28	4.2 ± 0.81	3.26 ± 0.63	7.46 ± 1.36	1.46 ± 0.12	1.13 ± 0.08	2.59 ± 0.08
29	3.86 ± 0.77	3.39 ± 0.67	7.25 ± 1.43	1.34 ± 0.14	1.18 ± 0.07	2.52 ± 0.21
30	3.61 ± 0.48	3.26 ± 0.61	6.88 ± 1.07	1.27 ± 0.13	1.13 ± 0.08	2.4 ± 0.19
31	3.69 ± 0.69	3.09 ± 0.38	6.78 ± 1.06	1.28 ± 0.11	1.08 ± 0.06	2.36 ± 0.16
32	3.56 ± 0.52	3 ± 0.51	6.56 ± 0.99	1.25 ± 0.13	1.04 ± 0.02	2.29 ± 0.13

Complement length (2n): 287.35 ± 46.88 µm
 Symmetry indices: Stebbins' Cat.: 1A; Al: 0.81; CV_{Cl}: 17.01; CV_{Ci}: 4.77; A1: 0.15; A2: 0.17; REC: 95.70; SYi: 85.16; TF%: 45.75

Table 2. – Comparative scheme of the main features between *Allium achaicum* Boiss., *A. guicciardii* Heldr. and *A. frigidum* Boiss. & Heldr.

	<i>A. achaicum</i>	<i>A. guicciardii</i>	<i>A. frigidum</i>
Bulb size (mm)	10-25 × 10-15	10-18 × 8-12	5-14 × 5-12
Outer tunics	fibrous	coriaceous	coriaceous
Stem length (cm)	10-30	10-50	[6]-8-22
Leaf number	4	4-6	3
Leaf max. length (cm)	7-15	25	3-7
Leaf width (mm)	1.5-3	1-2	0.6-0.8
Leaf sheathing the stem	1/2-2/3	2/3	1/3-1/2
Nerves of larger spathe	7	7	5-8
Nerves of smaller spathe	5	5	3-5
Larger spathe valve length [cm]	3-8	6-15	1.2-2.8
Smaller spathe valve length [cm]	2-3.5	4-9	1-1.8
Pedicel length (mm)	10-20(-25)	10-50	5-15
Inflorescence feature	erect	erect	pendulous
Inflorescence flowers	9-45	50-70	5-20(-25)
Perigon colour	white-yellowish, pruinose	yellowish-green, pruinose	white-pinkish
Tepal shape	elliptical	elliptical	elliptical to oblong-elliptical
Tepal apex	rounded	rounded	obtuse-mucronate
Tepal length [mm]	4.5-5	4.5-5	5-6
Tepals width [mm]	1.8-2	2-2.3	1.5-2.2
Stamen filament colour	purplish above	white	white
Stamen filament size	unequal, exserted	equal, exserted	unequal, included
Stamen filament length [mm]	4-6	6.5-7	1.8-2.6
Length of filament anulus [mm]	0.5	0.4-0.6	1, with interstaminal teeth
Anther size [mm]	1.5-1.6 × 0.8-0.9	1.5 × 1	1-1.2 × 0.8-0.9
Anther colour	yellow	yellow	white-yellowish
Ovary shape	ovoid	subglobose to ovoid	ovoid-ellipsoid
Ovary size [mm]	2-2.2 × 1.8-2	1.5-2 × 1.8-2	3-3.5 × 1.8-2
Style length [mm]	2-2.5	4-5	0.5-1
Capsule shape	globose	subglobose	subglobose
Capsule size [mm]	2.8-3 × 3.3-3.8	3.5-5 × 3.5-5	4.8-5 × 4-5
Chromosome number [2n]	32	32	16, 24

Mediterranean area (BRULLO & al., 2007). Among the taxa belonging to this group, *A. achaicum*, for its tetraploid chromosome complement and some morphological characters, such as the habit, inflorescence and flower, seems to be more closely related to *A. guicciardii* Heldr. occurring in C-N Greece (BRULLO & al., 2007). Nevertheless, *A. guicciardii* is a thermophilous species, chiefly found on rocky habitats, more or less disturbed by human activities, and usually localized in the thermo-mediterranean belt. In addition, *A. guicciardii* morphologically differs from *A. achaicum* in having coriaceous bulb tunics, stems covered up to 2/3 of its length by leaf sheaths, inflorescence 50-70-flowered, tepals unequal, yellowish-green, stamens with filaments 6.5-7 mm long, ovary shorter with style 4-5 mm long, capsule larger (Table 2). Another important difference regards the karyotype structure, as *A. guicciardii* is a tetraploid taxon with a diploid arrangement of its chromosomes (BRULLO & al., 2007).

Specimens examined. – GREECE: Vostitza, VII.1861, *Helreich s.n.* (G-BOIS); in monte Klokos Achaia prope Vostitza, rariss. 4000 m, 21.VII.1855, *Orphanides* 427 (BM, G); Peloponneso, Mt. Panachaiko presso Prof. Ilias, 1500 m, 24.VI.2006, Brullo, Giusso & Musarella s.n. (CAT); Peloponneso, Mt. Erimanthos, versante nord, ca. 1300 m, 25.VI.2006, Brullo, Giusso & Musarella s.n. (CAT); Peloponneso, Mt. Parnon, a Prof. Ilias presso Agriani, ca. 1400 m, 29.VI.2006, Brullo, Giusso & Musarella s.n. (CAT); Peloponneso, Mt. Menalon, ca. 1600 m, 1.VII.2006, Brullo, Giusso & Musarella s.n. (CAT); Peloponneso, Mt. Chelmos a Pouliou Vrisi, ca. 2000 m, 2.VII.2006, Brullo, Giusso & Musarella s.n. (CAT); Peloponneso, Mt. Klokos presso la cima, ca. 1700 m, 4.VII.2006, Brullo, Giusso & Musarella s.n. (CAT); Peloponneso, Mt. Klokos, ca. 1600 m, 30.VI.2007, Brullo, Giusso & Musarella s.n. (CAT).

Acknowledgements

The authors wish to thank the curators of BM, CAT, G, K, WU herbaria for the facilities during the examination of the collections. In addition, the M.I.U.R. (Ministero dell’Istruzione, dell’Università e della Ricerca) is gratefully acknowledged for its financial support.

References

- ANDERSSON, I. A. (1991). Allium L. In: STRID, A. & K. TAN (ed.), *Mountain Fl. Greece* 2: 701-714.
- BOISSIER, P. E. (1882). *Fl. orientalis* 5. Basileae, Genevae & Lugduni.
- BRULLO, F. (2002-2003). *CromoLab*©. Version 1.1. Dipartimento di Botanica, Catania.
- BRULLO, S., A. GUGLIELMO, P. PAVONE & C. SALMERI (2001). Cytotaxonomical notes on some rare endemic species of Allium (Alliaceae) from Greece. *Caryologia* 54: 37-57.
- BRULLO, S., A. GUGLIELMO, P. PAVONE & C. SALMERI (2007). Cytotaxonomic consideration on Allium stamineum Boiss. group (Alliaceae). *Bocconeia* 21: 325-343.
- GREILHUBER, J. & F. SPETA (1976). C-banded karyotypes in the Scilla hohenackeri Group, S. persica and Puschkinia (Liliaceae). *Pl. Syst. Evol.* 126: 149-188.
- HALÁCSY, E. von (1904). *Conspectus Flora Graecae* 3. Lipsiae.
- HAYEK, A. von (1924-1933). Prodromus Flora Peninsulae Balcanicae. *Repert. Spec. Nov. Regni Veg. Beih.* 30.
- HUZIWARA, Y. (1962). Karyotype analysis in some genera of Compositae. VIII. Further studies on the chromosomes of Aster. *Amer. J. Bot.* 49: 116-119.
- IUCN (2001). *IUCN Red List Categories and Criteria, Version 3.1*. IUCN.
- IUCN (2003). *Guidelines for Application of IUCN Red List Criteria at Regional Levels: Version 3.0*. IUCN.
- IUCN (2006). *Guidelines for using IUCN Red List Categories and Criteria, Version 6.1*. IUCN.
- LAVANIA, U. C. & S. SRIVASTAVA (1992). A simple parameter of dispersion index that serves as adjunct to karyotype asymmetry. *J. Biosci.* 17: 179-182.
- LEVAN, A., K. FREDA & A. A. SANDBERG (1964). Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201-220.
- PASZKO, B. (2006). A critical review and a new proposal of karyotype asymmetry indices. *Pl. Syst. Evol.* 258: 39-48.
- ROMERO ZARCO, C. (1986). A new method for estimating karyotype asymmetry. *Taxon* 35: 526-530.
- STEARNS, W. T. (1978). European species of Allium and allied genera of Alliaceae: a synonymic enumeration. *Ann. Mus. Goulandris* 4: 83-198.
- STEARNS, W. T. (1980). Allium L. In: TUTIN, T. G. & al. (ed.), *Fl. Eur.* 5: 49-69.
- STEARNS, W. T. (1981). The genus Allium in the Balkan Peninsula. *Bot. Jahrb. Syst.* 102: 201-213.
- STEBBINS, G. L. (1971). *Chromosomal evolution in higher plants*. Edward Arnold.
- TAN, K. & G. IATROU (2001). *Endemic plants of Greece. The Peloponnese*. Gads Forlag.
- TZANOUDAKIS, D. (1983). Karyotypes of ten taxa of Allium section Scorodon from Greece. *Caryologia* 36: 259-284.
- TZANOUDAKIS, D. & K. TAN (2000). Allium samothracicum Tzanoudakis, Strid & Kit Tan, a new species from the North Aegean area, Greece. *Portugalae Acta Biol., Sér. B, Sist.* 19: 355-360.