



A taxonomic revision of *Melanoxerus* (Rubiaceae), with descriptions of three new species of trees from Madagascar

Author: Kainulainen, Kent

Source: Candollea, 76(1) : 105-116

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2021v761a11>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A taxonomic revision of *Melanoxerus* (Rubiaceae), with descriptions of three new species of trees from Madagascar

Kent Kainulainen

Abstract

KAINULAINEN, K. (2021). A taxonomic revision of *Melanoxerus* (Rubiaceae), with descriptions of three new species of trees from Madagascar. *Candollea* 76: 105–116. In English, English and French abstracts. DOI: <http://dx.doi.org/10.15553/c2021v761a11>

This paper provides a taxonomic revision of *Melanoxerus* Kainul. & Bremer (Rubiaceae) – a genus of deciduous trees with eye-catching flowers and fruits that is endemic to Madagascar. Descriptions of three new species, *Melanoxerus antsiranensis* Kainul., *Melanoxerus atropurpureus* Kainul., and *Melanoxerus maritimus* Kainul. are presented along with distribution maps and a species identification key. The species distributions generally reflect the ecoregions of Madagascar, with *Melanoxerus antsiranensis* being found in the dry deciduous forests of the north; *Melanoxerus atropurpureus* in the inland dry deciduous forests of the west; *Melanoxerus maritimus* in dry deciduous forest on coastal sands; and *Melanoxerus suavissimus* (Homolle ex Cavaco) Kainul. & B. Bremer in the dry spiny thicket and succulent woodlands of the southwest.

Résumé

KAINULAINEN, K. (2021). Révision taxonomique du genre *Melanoxerus* (Rubiaceae), avec la description de trois nouvelles espèces d'arbres de Madagascar. *Candollea* 76: 105–116. En anglais, résumés anglais et français. DOI: <http://dx.doi.org/10.15553/c2021v761a11>

Cet article propose une révision taxonomique de *Melanoxerus* Kainul. & Bremer (Rubiaceae), un genre d'arbres à feuilles caduques avec des fleurs et des fruits attrayants qui est endémique de Madagascar. La description de trois nouvelles espèces, *Melanoxerus antsiranensis* Kainul., *Melanoxerus atropurpureus* Kainul. et *Melanoxerus maritimus* Kainul. est présentée accompagnée de cartes de répartition et d'une clé d'identification des espèces. La répartition des espèces reflète généralement les écorégions de Madagascar: *Melanoxerus antsiranensis* pousse dans les forêts sèches de feuillus du nord; *Melanoxerus atropurpureus* dans les forêts de feuillus sèches de l'intérieur des terres de l'ouest; *Melanoxerus maritimus* dans la forêt sèche décidue sur sables côtiers; et *Melanoxerus suavissimus* (Homolle ex Cavaco) Kainul. & B. Bremer dans le fourré épineux sec et les forêts succulentes du sud-ouest.

Keywords

RUBIACEAE – Gardenieae – Euclinia – Melanoxerus – Madagascar – Taxonomy – New species

Address of the author:

Gothenburg Botanical Garden, Carl Skottbergs Gata 22A, 41319 Gothenburg, Sweden. E-mail: kent.kainulainen@vgregion.se

Submitted on August 20, 2020. Accepted on February 8, 2021.

First published online on March 22, 2021.

ISSN: 0373-2967 – Online ISSN: 2235-3658 – *Candollea* 76(1): 105–116 (2021)

© CONSERVATOIRE ET JARDIN BOTANIQUES DE GENÈVE 2021

Introduction

The genus *Melanoxerus* Kainul. & B. Bremer is part of the tribe *Gardenieae* in the flowering plant family *Rubiaceae*. It was segregated from *Euclinia* Salisb. based on the results of the molecular phylogenetic analyses by KAINULAINEN & BREMER (2014). Whereas *Euclinia* is restricted to central Africa, *Melanoxerus* is endemic to Madagascar, where it is widely distributed in the subarid to dry southern and western regions but also occurs in the dry to subhumid north. *Melanoxerus* is a genus of deciduous or semi-deciduous trees with smooth flaky bark and sympodial plagiotropic branches with terminally clustered leaves and scarious stipules covering the terminal buds. The flowers are showy and very fragrant, and the large fleshy fruits are lenticellate with conspicuous patterns. The genus can be further diagnosed in that leaves and flowers turn black when drying. It is readily distinct from the other genera of tribe *Gardenieae* in Madagascar, i.e., *Catunaregam* Wolf, *Gardenia* Ellis, and *Hyperacanthus* E. Mey. ex Bridson. *Melanoxerus* lacks the opposite-decussate spines of *Catunaregam*, the sheathing persistent stipules and resinous exudate of *Gardenia*, and the flowers and fruits are terminal, whereas in *Hyperacanthus* these appear axillary (SCHATZ, 2001; MADAGASCAR CATALOGUE, 2021).

A taxonomic revision of *Melanoxerus* is warranted. *Melanoxerus suavisissimus* (Homolle ex Cavaco) Kainul. & B. Bremer is currently the only recognized species in the genus. It was originally described by CAVACO (1967) as *Gardenia suavisissima* Homolle ex Cavaco, but shortly thereafter transferred to *Euclinia* by LEROY (1974). In the protologue, CAVACO (1967) lists several paratypes from western Madagascar that do not correspond to the same species as the holotype from Ambovombe in southernmost Madagascar. The populations from northern Madagascar are also morphologically distinct. Since plants of *Melanoxerus* are deciduous and often flower before the leaves flush, specimens collected at different times of the year can be difficult to match up, and this may in part explain how the new species presented in this paper have escaped taxonomic attention in the past despite their eye-catching flowers and fruits. In this study I recognize a total number of four species in Madagascar, with *M. antsiranensis* Kainul., *M. atropurpureus* Kainul., and *M. maritimus* Kainul. being newly described.

The molecular phylogenetic study by KAINULAINEN & BREMER (2014) included samples from three of the four species recognized here, and their phylogenetic reconstruction from plastid DNA data indicates that *Melanoxerus atropurpureus* (as “sp. 2”) and *M. suavisissimus* are more closely related than either are to *M. antsiranensis* (as “sp. 1”). *Melanoxerus maritimus* was not included in that study. KAINULAINEN et al. (2017) estimated that the *Melanoxerus* clade diverged from an African ancestor around 9.4 Ma (with a 95 % highest posterior density interval of 6.6–12.4 Ma), and that the crown age,

and consequently the minimum age of the dispersal of this lineage from Africa to Madagascar, is 4.2 Ma (95 % HPD: 1.9–6.9 Ma).

Taxonomic treatment

Melanoxerus Kainul. & B. Bremer in Taxon 63: 828. 2014.

Typus: *Melanoxerus suavisissimus* (Homolle ex Cavaco) Kainul. & B. Bremer (= *Gardenia suavisissima* Homolle ex Cavaco. = *Euclinia suavisissima* (Homolle ex Cavaco) J.-F. Leroy).

Deciduous or semi-deciduous *shrubs* or *trees* up to 26 m tall. Sympodial plagiotropic branching, usually with much reduced distal shoot internodes, terminally clustered leaves, and scarious stipules covering the terminal buds. *Leaves* opposite, petiolate, glossy, papyraceous to subcoriaceous, cordiform to obovate, drying dark brown to black. *Stipules* triangular, caducous. *Flowers* bisexual. *Inflorescences* terminal, with few or solitary flowers. *Calyces* shortly tubular; lobes narrowly triangular, caducous. *Corollas* funnellform or campanulate to broadly urceolate; lobes contorted in bud, overlapping to the left, spreading horizontally at anthesis. *Styles* club-shaped, ribbed, with secondary pollen presentation in the distal end. *Pollen* dispersed in tetrads. *Fruits* fleshy-indehiscent, globose to ovoid, with a thick mesocarp and an exocarp covered with conspicuous lenticels. *Seeds* flattened, irregularly angled, horizontally inserted and immersed in soft pulp.

Distribution, habitat and ecology. – *Melanoxerus* is endemic to Madagascar and distributed across most of the southern and western regions (i.e. Anosy, Androy, Atsimo-Andrefana, Menabe, Melaky, Boeny, and Sofia Regions), as well as in northernmost Madagascar in the DIANA and SAVA Regions. *Melanoxerus* occurs in vegetation types ranging from subarid thickets to subhumid semi-deciduous forests, from sea level to about 800 m in elevation, and on various substrates from coastal sands to laterite, including skeletal soils in karstic limestone formations (tsingy). It is absent from the Central Highlands and the humid east coast. Although there are exceptions, in general the species distributions reflect the ecoregions of Madagascar (cf. OLSON et al., 2001), with *M. antsiranensis* being found in the dry deciduous forests of the north; *M. atropurpureus* in the inland dry deciduous forests of the west; *M. maritimus* in dry deciduous forest on coastal sands; and *M. suavisissimus* in the dry spiny thicket and succulent woodlands of the southwest (Fig. 1).

Little is known about the ecology of this genus, and no information on pollination or seed dispersal has been recorded for the specimens that I have examined. The flowers are large and ± fleshy, with funnellform to broadly urceolate corollas that are externally white (sometimes tinged purple or pale green)

and internally purple or white with green, red, or purple markings. Pollen is usually deposited on the upper part of the style (secondary pollen presentation), which is club-shaped with longitudinal ridges. The flowers are strongly fragrant with a sweet smell like jasmine.

The fruits are large, indehiscent “berries” with a fleshy pulp and are most likely an adaptation to endozoochory. The seeds are presumably dispersed by lemurs, but it is possible that seed dispersal, at least in part, has relied on the recently extinct Malagasy megafauna, as has been suggested for *Adansonia* L. (BAUM, 1995) – a genus which also has large indehiscent fruits, and a distribution in Madagascar that is similar to that of *Melanoxerus*. The leaves of *Melanoxerus* appear to be attractive to herbivores because in the specimens seen in this study the leaves are usually damaged. Although not documented, insects are probably responsible for part of the damages, although slugs (pers. obs.) and lemurs of the genus *Propithecus* (information on the label of Meyers & Boltz 110) have been observed feeding on leaves of *M. antsiranensis*.

Key to the species of *Melanoxerus*

1. Leaves broadly lanceolate to obovate; flowers solitary or not; corolla tube campanulate to broadly urceolate; corolla throat glabrous; mature fruits with lenticels 1–2.5 mm in diam. 2
- 1a. Leaves cordiform or ovate to lanceolate; flowers solitary; corolla tube funnelform; corolla throat pubescent; mature fruits with lenticels 0.1–1 mm in diam. 3
2. Leaves obovate, 7–18 × 4.5–8.5 cm; corolla throat white with red, purple or violet (and sometimes also green) spots 1. *M. antsiranensis*
- 2a. Leaves broadly lanceolate, 10–30 × 6.5–14.5 cm; corolla throat uniformly purple to violet ... 2. *M. atropurpureus*
3. Leaves broadly ovate to cordiform; leaf length/width ratio 4:3 to 1:1; corolla lobes rounded to indented 3. *M. maritimus*
- 3a. Leaves ovate to lanceolate; leaf length/width ratio 3:1 to 3:2; corolla lobes acute 4. *M. suavissimus*

Taxonomy

1. *Melanoxerus antsiranensis* Kainul., **sp. nov.** (Fig. 2).
– *Euclinia* sp. 1 in KAINULAINEN & BREMER (2014: 820).

Holotypus: MADAGASCAR. **Reg. DIANA [Prov. Antsiranana]:** Distr. Ambilobe, Mahamasina, Réserve Spéciale d'Ankarana, 12°57'15"S 49°07'39"E, 100 m, 10.II.2003, fr., Bardot-Vaucoulon et al. 1406 (P [P00455719]!; iso-: K [K000551041]!, MO!, TAN).

Melanoxerus antsiranensis Kainul. differs from *M. suavissimus* (Homolle ex Cavaco) Kainul. & B. Bremer by

its pauciflorous inflorescences (vs. solitary flowers), campanulate (vs. funnelform) corollas, and obovate (vs. ovate to lanceolate) leaves. It most resembles *M. atropurpureus* Kainul., from which it differs in that the corolla throat is white with purple and sometimes green spots (vs. uniformly purple), and in its smaller (7–18 × 4.5–8.5 cm vs. 10–30 × 6.5–14.5 cm) and obovate (vs. broadly lanceolate) leaves.

Shrubs or trees up to 18 m tall and 40 cm dbh. Bark beige or whitish pale grey and smooth. Stipules 9–14.5 × 4–7 mm, triangular, interpetiolar, caducous. *Leaves* deciduous, petioles 0.7–6 cm long, adaxially canaliculate; leaf blades papyraceous, obovate, 7–18 × 4.5–8.5 cm; bases cuneate to attenuate, apices (rounded) obtuse to apiculate or shortly acuminate; adaxial surface: dark green when fresh, drying dark brown to black, smooth and glossy, glabrous; 8–12 pairs of conspicuous, ± pinnate secondary veins, tertiary venation reticulate; midribs prominent, whitish green when fresh, ± the same colour as the leaf when dry; abaxial surface green when fresh, drying dark brown to blackish, glabrous except for tuft domatia that are often present in the vein axils. *Flowers* solitary or few (< 10) in a terminal, sessile, congested and umbel-like inflorescence; pedicels up to 30 mm long; hypanthium narrowly urceolate, 5–8 × 3–4 mm, glabrous. *Calyces* green; calyx tubes 0.5–2 mm long, externally and internally glabrous; calyx lobes 4.3–11 × 0.8–3.5 mm, narrowly triangular, spreading or recurved. *Corollas* fleshy, campanulate; corolla tubes 1.5–2.6 cm long, externally (yellowish) white, sometimes with purple and pale green streaks, glabrous, internally white with red, purple or violet (and sometimes also pale green) spots, glabrous or with a few trichomes (0.1–0.5 mm long) at the very base; corolla lobes white (sometimes with purple spots on the external side), 1–4.5 × 0.8–2.5 cm, ovate to deltate, rounded to acute, spreading horizontally at anthesis; stamens subsessile (filaments c. 2 mm long), attached 10–12 mm below corolla sinus and 3 mm from anther base; anthers 11–18 × 1.5–2 mm, linear, included or shortly exserted (to 2 mm above corolla sinus). *Styles* simple, clavate, with shallow longitudinal grooves, 14–20 mm long, glabrous, included or shortly exserted; stigma shortly bifid (lobes 0.5–3.5 mm long). *Fruits* (globose) ellipsoid, 3–7.6 × 2.7–6 cm, smooth, glabrous, olive to brown, with numerous beige to pale grey lenticels (1–2 mm in diam.), fleshy-indehiscent with yellowish to white pulp; calyx tube sometimes persistent but calyx lobes caducous. *Seeds* flattened, irregularly angled, 10–15 mm in diam., c. 2 mm thick, smooth, ± translucent, yellowish brown when fresh, drying black.

Vernacular names. – “Mamoampoma” (*Rakotondrafara et al.* 505); “Mantalana” (*Service Forestier* 8329); “Voavandrikala” (*Manjakahery & Sola* 83).




Fig. 1. – Distribution of *Melanoxerus antsiranensis* Kainul. (yellow circles), *M. atropurpureus* Kainul. (pink circles), *M. maritimus* Kainul. (triangles), and *M. suavis* (Homolle ex Cavaco) Kainul. & B. Bremer (stars) in Madagascar. [mapped on the bioclimatic zones of Madagascar (after CORNET, 1974; see SCHATZ, 2000)]

Distribution, habitat and ecology. – *Melanoxerus antsiranensis* is only known from the DIANA and SAVA Regions in Antsiranana Province where it is widely distributed (Fig. 1). The new species grows in littoral forests and in dry deciduous forests from sea level to c. 700 m in elevation on Montagne d'Ambre.

Phenology. – Flowering material has been collected from September to December and fruiting material from September to July.

Conservation status. – *Melanoxerus antsiranensis* is known from numerous localities in northern Madagascar. Its distribution includes five protected areas, i.e. Ankarana, Loky-Manambato, Montagne d'Ambre, and Montagne des Français. *Melanoxerus antsiranensis* is therefore assigned a preliminary conservation status of "Least Concern" [LC] according the IUCN Red list Categories and Criteria (IUCN, 2012).

Notes. – The specimen *Razafitsalama* & *Ludovic* 16 is unusual in having distinctly velutinous leaves and is tentatively included here. The label on *Meyers* & *Boltz* 110 reports that the lemur *Propithecus tattersalli* eats the young leaves of this species. *Melanoxerus antsiranensis* flowers at the same time or just before the new leaves flush. It resembles *M. atropurpureus* in having pauciflorous inflorescences and flowers with campanulate corollas. However, the phylogenetic analysis of KAINULAINEN & BREMER (2014) indicates that they are not sister species, a result that suggests that these shared characters may be plesiomorphic rather than synapomorphic.

Melanoxerus antsiranensis is distinguished from *M. atropurpureus* in its smaller leaves (7–18 × 4.5–8.5 cm vs. 10–30 × 6.5–14.5 cm) that are obovate rather than broadly lanceolate. Furthermore, the corolla throat is white with red to violet spots (vs. uniformly dark purple in *M. atropurpureus*).

Selected additional material examined. – **MADAGASCAR. Reg. DIANA [Prov. Antsiranana]:** Masorolava, Mahagaga, Antsoroby forest, 12°42'37"S 48°58'11"E, 53 m, 27.IX.2007, fr., *Andriamihajarivo et al.* 1434 (MO, P); Ankarana SR, 2 km W of Mahamasina village, 12°57'26"S 49°07'46"E, 132 m, 21–30.III.1995, fr., *Andrianantoanina* & *Bezara* 792 (BR, G, K, MO); Montagne d'Ambre NP, c. 8 km E of Bobakilandy village, 12°37'37"S 49°06'26"E, 533 m, 5.XII.1995, fl., *Andrianantoanina* & *Bezara* 895 (BR, G, K, MO); Andrafiomena, forest near Anjahankely, 12°54'23"S 49°19'30"E, 450 m, 6.I.2011, fr., *Burivalova* ZB 176 (G, K); Ankarana, 12°50'47"S 49°06'18"E, 82 m, 14.I.2002, fr., *De Block et al.* 1226 (BR, MO, P, TAN, UPS, WAG); Ankarana, 4–9.III.1951, fr., *Humbert* & *Capuron* 25493 (BR, K); *ibid. loco*, 12°54'42"S 49°06'43"E, 240–260 m, 22.V.1993, fr., *Jongkind* & *Rapanarivo* 979 (K, MO); Montagne des Français, 12°19'25.5"S 49°20'11"E, 230–250 m, 5.III.2016, ster., *Kainulainen et al.* 370 (TAN); Anosy à 4 km au NE du village d'Ambolobozokely [Nosy Voanio], 12°26'49"S 49°33'07"E, 22 m, 30.III.2007, fr., *Rakotondrafara et al.* 505 (BR, MO, P); Andranovondronina, Andakorobe, village le plus proche Antsaravy, 12°08'51"S 49°20'30"E, 19 m, 17.III.2006, fr., *Rakotondrafara et al.* 474 (CNARP, K, MO, P, TAN); Montagne d'Ambre, 12°34'59"S 49°05'59"E, 683 m, 5.II.2012, fr., *Ramandimbimanana* & *Randimbison* SDR 372 (BR, G, MO); Sahafary SF, 12°35'51"S 49°26'31"E, 250 m, 7.XI.2006, fl., *Ranaivojaona et al.* 1551 (BR, MO, K, P, TAN); Nosy Voanio, 4 km E of Ambolobozokely, 12°26'01"S 49°33'00"E, 15 m, 15.II.2006, fr., *Randianaivo et al.* 1364 (BR, MO); Mosorolava, Ampombiantambo, Antsoroby forest, 12°42'13"S 48°58'12"E, 75 m, 23.IX.2007, fl. & fr., *Ratovoson et al.* 1332 (CNARP, G, MO, P, TAN); Ankarana, 11–14.I.2002, fr., *Razafimandimbison* & *Andrianantoanina* 454 (BR, MO, UPS); Ankarongana, Irodo, N of Analamerana forest, Ampandriantsira, 12°39'42"S 49°32'37"E, 50–123 m, 14.II.2001, fr., *Razafitsalama* & *Ludovic* 16 (MO, P); Montagne des Français, 250 m, 18.II.1954, fr., *Service Forestier* 8329 (P, TEF); Ankarefo, 18.V.1954, fr., *Service Forestier* 10003 (P); Diego-Suarez, Bemaskabe forest, 200–500 m, 17.X.1927, fr., *Ursch* 231 (P). **Reg. SAVA [Prov. Antsiranana]:** Distr. Vohemar, Daraina, Bekaraoka south, 13°11'15"S 49°42'E, 9.IV.2008, fr., *Bremer et al.* 5113 (MO, S, TAN); Hillside next to Daraina, 13°08'S 49°27'E, 10.IV.2008, fr., *Bremer et al.* 5131 (MO, S, TAN); Daraina, Bobankora forest, 13°12'22"S 49°46'27"E, 200 m, 2.II.2003, fr., *Gautier et al.* LG 4195 (G, K, MO); Daraina, Bekaraoka forest, 13°03'90"S 49°42'35"E, 420 m, 22.XI.2006, fl., *Gautier et al.* LG 4911 (G); Nosibe, Anjiabe, Anaborano, 13°04'18"S 49°54'32"E, 33 m, 11.XII.2004, fr., *Manjakabery* & *Sola* 83 (MO); Mantamena, 7 km NW of Daraina, 13°08'S 49°42'E, 112–330 m, IV.1990, *Meyers* & *Boltz* 110 (MO); Nosibe, Anjiabe, Analabe forest, 13°04'43"S 49°54'04"E, 11.V.2004, fr., *Rabehevitra et al.* 915 (MO, P); Tsarabaria, Manakana, Ambondrombe forest, 13°42'58"S 50°05'37"E, 3 m, 24.X.2002, fr., *Rabenantoandro et al.* 1019




Fig. 2. – *Melanoxerus antsiranensis* Kainul. **A.** Habit; **B.** Fruit and smooth bark of a trunk (c. 20 cm dbh); **C.** Fruiting branch; **D–F.** Flowering branch with young leaves; **G.** Immature fruits; **H.** Cross-section of a fruit showing the seeds embedded in whitish pulp. [A–B, H: Bremer et al. 5113; C, G: Ramandimbimanana & Randimbiarison SDR 372; D–E: Ratovoson 1332; F: Randrianaivo 1433] [Photos: A–B, H: K. Kainulainen; C, G: S.D. Ramandimbimanana; D–E: F. Ratovoson; F: R. Randrianaivo]

(MO); Nosibe, Anjiabe, Anaborano forest, near lake Sahaka, 13°04'42"S 49°54'13"E, 25 m, 2.XI.2002, fl., *Rabenantoandro et al.* 1091 (MO, TEF); *ibid. loco*, 13°04'43"S 49°54'04"E, 10 m, 23.II.2003, fr., *Rabenantoandro et al.* 1272 (MO, TEF); Mosorolava forest, 13°05'44"S 49°54'43"E, 23 m, 5.X.2013, fl., *Rakotovoao* 6482 (MO); Fanambana, Ambohimahazo forest, 13°35'44"S 50°00'04"E, 60–100 m, 30.III.2001, fr., *Ranaivojaona et al.* 355 (BR, MO, P); Solanampilana forest 35 km N of Daraina, 13°05'42"S 49°34'57"E, 137 m, 9.XII.2006, fl. & fr., *Randrianaivo et al.* 1433 (G, MO, P, TAN); Daraina, Ambilondomba forest, 13°09'39"S 49°38'47"E, 390 m, 31.I.2004, fr., *Ranirison et al.* PR 367 (G); Vohimarina, Fanambana, Antsatoby, 13°36'55"S 49°59'55"E, 77 m, 8.VII.2007, fr., *Rasoafaranaivo* 239 (MO); Analabe forest, near Anaborano village and Sahaka lake, 13°04'59"S 49°54'26"E, 10.VII.2003, fr., *Razakamalala* 519 (MO, P, TEF); Tsarabaria, Manakana, E of Ambondrobe village, 13°43'08"S 50°05'50"E, 13.III.2004, fr., *Razakamalala* RZK 1022 (K, MO, P, TEF).

2. *Melanoxerus atropurpureus* Kainul., sp. nov. (Fig. 3).

– *Euclinia* sp. 2 in KAINULAINEN & BREMER (2014: 820).

Holotypus: MADAGASCAR. **Reg. Melaky [Prov. Mahajanga]:** Distr. Maintirano, Beanka, partie S, Sarodrano, 18°03'27"S 44°30'50"E, 359 m, 26.XI.2011, fl., *Gautier, Tabinarivony & Bolliger* LG 5744 (G [G00340493]!; iso-: BR!, K [K000918390]!, MO!, P!, TEF).

Melanoxerus atropurpureus Kainul. is similar to *M. antsiranensis* Kainul. in its pauciflorous inflorescences and flowers with campanulate corollas, but it differs in having larger leaves (10–30 × 6.5–14.5 cm vs. 7–18 × 4.5–8.5 cm) that are broadly lanceolate (vs. obovate), and by the uniformly purple corolla throat (vs. whitish corolla throat with red, purple or violet and sometimes also green spots).

Shrubs or *trees* up to 26 m tall and 25 cm dbh. Bark grey and smooth, flaky. Stipules 2.5–13 × 3–5 mm, triangular, interpetiolar, caducous. *Leaves* deciduous, petioles 2–7.5 cm long, adaxially canaliculate; leaf blades papyraceous, broadly lanceolate, 10–30 × 6.5–14.5 cm; bases cuneate to rounded, sometimes oblique; apices acute to acuminate; adaxial surface dark green when fresh, drying dark brown to black, smooth and glossy, sparsely pubescent along venation; 8–12 pairs of conspicuous, ± pinnate secondary veins, tertiary venation reticulate; midrib prominent, pale green when fresh, ± the same colour as the leaf when dry; abaxial surface pale green when fresh, drying dark brown to blackish, glabrous except the pubescent veins (trichomes c. 0.1 mm); tuft domatia often present in the vein axils. *Flowers* solitary or few (< 10), sessile, congested and umbel-like *inflorescence*; pedicels up to 15 mm long; hypanthium narrowly urceolate, 8–15 × 5–8 mm, glabrous. *Calyces* green; calyx tubes 2–3.5 mm long, externally and internally glabrous; calyx lobes 3–6.5 × 0.5–3.5 mm, narrowly triangular and recurved; corollas fleshy, campanulate to broadly urceolate; corolla tubes 1.8–4.8 cm long, externally white with purple and pale green streaks, glabrous, internally dark purple to violet, glabrous or with a sparse indument

(trichomes 0.4–1 mm) in the lowermost (c. 0.5 cm) part of the tube. *Corollas* lobes white, 1.5–3 × 1.2–2.5 cm, ovate, rounded to acute, spreading horizontally at anthesis; stamens subsessile (filaments 1–4 mm long), attached 25–35 mm below corolla sinus and 4–7 mm from anther base; anthers 22–27 × 2–3 mm, linear, included (tips 5–6 mm below corolla sinus). *Styles* simple, clavate, with shallow longitudinal grooves, 28–33 mm long, glabrous, included; stigma shortly bifid (lobes 1–1.5 mm long). *Fruits* (globose) ellipsoid, 3.5–7 × 3–6 cm, smooth, glabrous, olive to brown or maroon with numerous beige to pale grey lenticels (1–2 mm in diam.), fleshy-indehiscent with yellowish to white pulp; calyx tube sometimes persistent, but lobes caducous. *Seeds* flattened, irregularly angled, 8–13 mm in diam., c. 2 mm thick, smooth, ± translucent, pale yellow when fresh, drying black.

Etymology. – The specific epithet *atropurpureus* refers to the dark purple colour of the interior of the corolla tube in this species.

Vernacular names. – “Andrivitsy” (*Réserves Naturelles* 7791); “Hazondanitra” (*Service Forestier* 4953); “Voafotaka” (*Labat & Deroin* 2280); “Voangorindambo” (*Service Forestier* 29775).

Distribution, habitat and ecology. – *Melanoxerus atropurpureus* occurs in the Boeny, Melaky, and Sofia Regions (Mahajanga Province) and in the southernmost part of the DIANA Region (Antsiranana Province) (Fig. 1), where it grows in dry to humid, deciduous or evergreen forests from sea level to 450 m in elevation.

Phenology. – Flowering material has been collected from October to November and fruiting material from December to July.

Conservation status. – *Melanoxerus atropurpureus* has a wide distribution in western Madagascar and is known from several protected areas including Ankarafantsika, Bamaraha, and Namoroka. It can therefore be assigned a preliminary conservation status of “Least Concern” [LC] according the IUCN Red list Categories and Criteria (IUCN, 2012). However, it is not a commonly collected species, and more information is needed about its population size.

Notes. – The specimen *Perrier de la Bathie* 1155A from the Ankara plateau is in poor shape but probably belongs to this species. In this species the exocarp of the fruit is often almost completely hidden by lenticels (Fig. 3F).

Melanoxerus atropurpureus is the tallest growing species in the genus. It mostly resembles *M. antsiranensis* but has larger leaves (10–30 × 6.5–14.5 cm vs. 7–18 × 4.5–8.5 cm) that are broadly lanceolate rather than obovate. Furthermore, the



Fig. 3. – *Melanoxerus atropurpureus* Kainul. A–D. Corolla; E. Cross-section of a fruit showing the seeds embedded in whitish pulp; F. Fruiting branch. [A–D: Gautier et al. LG 5744; E: Hanitrarivo et al. HRM 146] [Photos: A–D: L. Gautier; E: M. Hanitrarivo; F: L. Rajaovelona, Ankarafantsika (Boeny), 17.III.2020]

corolla throat is uniformly dark purple (white with red to violet spots in *M. antsiranensis*).

Selected additional material examined. – **MADAGASCAR. Reg. Boeny [Prov. Mahajanga]:** Distr. Marovoay, Ankarafantsika NP, Ampijoroa forest station, 16°18'52"S 46°49'04"E, 190 m, 24.I.2000, fr., *Davis et al.* 2500 (K, TAN); *ibid. loco*, 16°20'S 46°51'E, 200 m, 14.IV.1984, *Dorr* 3048 (BR, K, P, MO); *ibid. loco*, 16°19'S 46°49'E, 70–100 m, 7.IV.1988, *Gentry & Schatz* 62050 (MO); Plateau d'Ankara, X.1900, fl., *Perrier de la Bâthie* 1155A (P); Manongarivo (Ambongo), XI.1903, fl., *Perrier de la Bâthie* 1640 (P); *ibid. loco*, X.1905, fl., *Perrier de la Bâthie* 4103 (P); *ibid. loco*, I.1904, fr., *Perrier de la Bâthie* 6944 (P); Ankarafantsika, 16°20'05"S 46°47'06"E, 3–9.II.1997, fr., *Rabevohitra* 3027 (TEF); *ibid. loco*, 16°18'S 46°49'E, 22.IV.1994, fr., *Rajemisa* 19 (MO, P, TAN); Distr. Mahajanga II, Mariarano, Ankatsabe forest, 15°28'30"S 46°41'24"E, 16 m, 20.IV.2007, fr., *Rakotoarivelo* 94 (MO, P, TAN); Soalala, 15.VII.1977, fr., *Rakotozafy* 1849 (TAN); Distr. Soalala, Andranomavo, Vilanandro, Tsingy Namoroka, Ambatomay forest, 16°28'36"S 45°20'52"E, 153 m, 9.IX.2012, fr., *Ravelonarivo et al.* 4523 (MO, TAN); Andranomavo, 19.XI.1955, fl., *Réserves Naturelles* 7791 (P, TEF); Ankarafantsika, s.d., fr., *Service Forestier* 19 (P); Tsingy Namoroka, 4.IV.1933, fr., *Service Forestier* 36 (P); *sine loco*, 4.IV.1933, fr., *Service Forestier* 54 (BR, K, P); Ankarafantsika, Ampijoroa, 15.V.1952, fr., *Service Forestier* 4953 (P, TEF); *ibid. loco*, 16.VI.1954, fr., *Service Forestier* 10204 (P); Ambato-Boeny, Andranovato, Ankijabe, 20.I.1959, fr., *Service Forestier* 19261 (TEF); Ambongo, Soalala (baie de Baly), 22.XI.1965, fl., *Service Forestier* 24244 (BR, P, TEF); by lake Ampijoroa, 28.II.1980, fr., *Service Forestier* 29775 (TEF). **Reg. DIANA [Prov. Antsiranana]:** Distr. Ambanja, Ampasindava, Andranomatavy forest, 13°40'04"S 47°59'21"E, 295 m, 25.XI.2009, fl., fr., *Madiomanana et al.* MAD 266 (G, K, MO, P, TEF); along Andranomatavy river, 13°40'39.19"S 47°58'30.24"E, 260 m, 2.VII.2012, fr., *Tabinarivony & Rasoanaivo* TAJ 699 (G). **Reg. Melaky [Prov. Mahajanga]:** Beanka, Ambinda-Nord, 17°55'51"S 44°28'39"E, 216 m, 5.II.2012, fr., *Hanitrivoo et al.* HRM 146 (G, S); Ambodirina, 1 km E of Ambinda (RN 9, near Antsalova), 18°38'S 44°42'E, 100–200 m, 3.XII.1992, fr., *Labat & Deroin* 2280 (K, P, TAN); Beanka, Sarodrano, 18°02'55"S 44°31'13"E, 446 m, 11.XII.2011, fl., *Nusbaumer et al.* LN 3073 (G, K); Antsalova, 20.III.1993, fr., *Villiers et al.* 4851 (MO, P); Belitsaka, Ankilimanarivo, Beanka forest, 18°03'37"S 44°31'37"E, 319 m, 16.IV.2011, fr., *Razakamalala & Germain* 6382 (MO, S). **Reg. Sofia [Prov. Mahajanga]:** entre Antsohihy et Befandriana-Nord, X.1962, fl., *Bosser* 16697 (P, TAN); Distr. Antsohihy, Anjimangirana, Analananbe, Andrafiborizina, 15°09'19"S 47°44'22"E, 200 m, 30.V.2000, fr., *Rakotonasolo* RNF 177 (BR, K, P, TAN).

3. *Melanoxerus maritimus* Kainul., sp. nov. (Fig. 4).

Holotypus: MADAGASCAR. **Reg. Melaky [Prov. Mahajanga]:** Distr. Maintirano, Lac Mandrozo, 17°32'43"S 44°02'26"E, 18 m, 23.XI.2015, fl. & fr., *Gautier, Luino & Rasolonjatovo* LG 6196 (G [G00419027]); iso-: MO!, P!, S!, TEF).

Melanoxerus maritimus Kainul. differs from *M. suavissimus* (*Homolle ex Cavaco*) Kainul. & B. Bremer in its rounder leaves (leaf length/width ratio 4:3–1:1 vs. 3:1–3:2), that have a truncate to cordate leaf base (vs. cuneate to attenuate), and corolla lobes that are rounded to indented (vs. acute).

Shrubs or trees up to 10 m tall and 28 cm dbh, with mostly sympodial branching. Bark pale grey to pale brown, smooth, flaky. Stipules 3.5–4.7 × 3.6–4.5 mm, broadly triangular, interpetiolar, caducous. *Leaves* semi-deciduous, petioles 2.8–5.2 cm long, adaxially canaliculate; leaf blades

papyraceous to subcoriaceous, cordiform to broadly ovate, 7.5–12.7 × 5–10.5 cm; bases cordate to truncate; apices acute; adaxial surface dark green when fresh, drying dark brown to black, smooth and very glossy, glabrous except for the lower part of the midrib (trichomes 0.1–0.5 mm long); 6–9 pairs of conspicuous, ± pinnate secondary veins, tertiary venation reticulate; midrib prominent, pale green when fresh, ± the same colour as the leaf when dry; abaxial surface green when fresh, drying dark brown to blackish, glossy, glabrous, but with tuft domatia often present in the vein axils. *Flowers* solitary, terminal, very fragrant; pedicels up to 14 mm long; hypanthium narrowly urceolate, 5–18 × 3–4 mm, glabrous; calyces green; calyx tubes 1–4.5 mm long, externally and internally glabrous; calyx lobes 3.4–6.2 × 1.2–2 mm, narrowly triangular and recurved; corollas fleshy, funnelform, straight; corolla tubes 2–3.7 cm long, externally pale yellow with a greenish base, glabrous, internally white with red spots and lines, finely pubescent (trichomes c. 0.5 mm long). *Corollas* lobes white, 2–2.7 × 1.6–3 cm, ovate to orbicular, rounded to truncate or indented, spreading horizontally at anthesis; stamens subsessile (filaments 2–4 mm long), attached c. 10 mm below corolla sinus and c. 4 mm from anther base; anthers c. 15 × 2 mm, linear, exerted for 1–3 mm. *Styles* simple, clavate, with shallow longitudinal grooves, 30–35 mm long, glabrous, exerted for 5–9.5 mm; stigma shortly bifid (lobes c. 3 mm long). *Fruits* ellipsoid, 2.5–5.1 × 1.6–3.7 cm, smooth, glabrous, (colour of mature fruit unknown) with numerous beige lenticels (0.1–1 mm in diam.), fleshy-indehiscent; calyx tube persistent but lobes caducous. *Seeds* not seen.

Etymology. – This species grows near the sea and that is what the specific epithet *maritimus* refers to.

Vernacular names. – “Ndraniana” (*Service Forestier* 10585); “Tselitselika” (*Bertrand* 16).

Distribution, habitat and ecology. – *Melanoxerus maritimus* is known from sea level to 30 m in elevation in coastal areas of the Boeny, Melaky, and Sofia Regions (Mahajanga Province) (Fig. 1). It grows in low canopy, dry deciduous forests on dunes and on edges of salt flats or mangroves.

Phenology. – Flowering material has been collected from June to November, and fruiting material from July to February.

Conservation status. – *Melanoxerus maritimus* is known from less than ten locations, (three of which are within the recently protected areas of Antrema, Bombetoka Belem-boka, and Mandrozo), with an estimated area of occupancy (AOO) of less than 2000 km². It appears restricted to littoral dry forest. This habitat is impacted for firewood or charcoal production (also noted on label of *De Block & Rakotonasolo*



Fig. 4. – *Melanoxerus maritimus* Kainul. A. Habit; B. Flowering branch; C. Flower bud; D–E. Open flowers; F. Transverse section of the corolla and stigma; G–H. Immature fruits. [A–C, F–G: Gautier et al. LG 6196; D: Randriatsivry 207; E: Labat et al. 3534; H: Labat et al. 3535] [Photos: A–C, F–G: L. Gautier; D: M. Randriatsivry; E, H: J.-N. Labat]

806). It is projected that the number of locations and the number of mature individuals continue to decline, and this species can therefore be assigned a preliminary conservation status of “Vulnerable” [VUB2ab(iii)] according to IUCN Red list Categories and Criteria (IUCN, 2012). However, more information is needed about the population size of this species.

Notes. – *Melanoxerus maritimus* has solitary flowers with a funnellform corolla and pubescent corolla throat, and in these characters it most resembles *M. suavissimus*, but it is recognized by its $\pm$ cordiform leaves that are almost as wide as long (leaf length/width ratio 4:3–1:1 vs. 3:1–3:2 in *M. suavissimus*), and by having corolla lobes that are rounded to indented rather than acute. The leaves of *M. maritimus* are glossy below and very glossy above. They remain glossy when dried and almost appear varnished. It is notable that none of the specimens examined in this study is lacking leaves, and that specimens collected during the dry season still retain some leaves. This species is therefore probably best considered semi-deciduous as opposed to deciduous.

Selected additional material examined. – **MADAGASCAR. Reg. Boeny [Prov. Mahajanga]:** [Distr. Mahajanga I] Antananamasadza [probably Antanimasaja, now part of Mahajanga town], 1912, *Afzelius* 266 (P, S); Katsepy, c. 10 m, 19.II.1950, ster., *Bertrand* 16 (P); Distr. Mahajanga II, Antsanitia, Belamonty forest, 15°34'19"S 46°25'04"E, 9 m, 14.II.1999, fr., *De Block & Rakotonasolo* 806 (BR, MO, TAN); Katsepy, 15°45'56"S 46°14'21"E, 0 m, 22.VII.1995, fl. & fr., *Edmondson et al.* 95–17 (MO, P); *ibid. loco*, 8.IX.1912, fl., *Kaudern s.n.* (S); *ibid. loco*, 15°42'35"S 46°12'03"E, 10.XI.2005, fl., *Labat et al.* 3534 (K, P); *ibid. loco*, 10.XI.2005, fr., *Labat et al.* 3535 (G, K, MO, P); Antsanitia, 15°34'31"S 46°25'7"E, 30 m, 7.I.2000, fr., *Rakotonasolo RNF* 132 (K, TAN); Forest station E of Antsanitia village, 15°34'31"S 46°24'26"E, 16 m, 15.II.2013, fr., *Ralimanana et al.* 1803 (BR, K, MO, P, TAN); Distr. Mahajanga II, Ampasimariny, 9 m, 23.VIII.1954, fl. & fr., *Service Forestier* 10585 (MO, P, TEF). **Reg. Melaky [Prov. Mahajanga]:** Region de Tambohorano, 23.VI.1930, fl., *Decary* 8083 (P); Ambongo, 1841, fr., *Pervillé* 673 (K, P). **Reg. Sofia [Prov. Mahajanga]:** SW of Antangena, 15°28'S 47°09'E, 0 m, 25.X.1987, fl. & fr., *Bisset M12* (K); Distr. Analalava, Antonibe, 15°01'44"S 47°13'17"E, 19.VIII.2007, fl. & fr., *Randriatsivory* 207 (MO, P, TAN).

4. *Melanoxerus suavissimus* (Homolle ex Cavaco) Kainul. & B. Bremer (Fig. 5).

= *Gardenia suavissima* Homolle ex Cavaco in *Adansonia* ser. 2, 7: 177. 1967. = *Euclinia suavissima* (Homolle ex Cavaco) J.-F. Leroy in *Adansonia* ser. 2, 14: 52. 1974.

Holotypus: MADAGASCAR. Reg. Androy [Prov. Toliara]: Ambovombe, 27.X.1924, *Decary* 3357 (P [P00852562]).

Shrubs or **trees** up to 13 m tall and 40 cm dbh, with mostly sympodial branching. Bark (mottled) whitish or pale grey to pale brown, smooth, flaky. Stipules 3–10 $\times$ 2–4 mm, triangular, interpetiolar, caducous. **Leaves** deciduous, petioles 0.5–5 cm long, adaxially canaliculate; leaf blades papyraceous, ovate to lanceolate, 6.5–16 $\times$ 2–9(–11) cm; bases (rounded) cuneate to attenuate (often asymmetrically); apices acute to acuminate;

adaxial surface dark green when fresh, drying dark brown to black, smooth, matte or glossy, $\pm$ glabrous to (velvety) pubescent in particular along midrib and venation; (6–)8–10(–12) pairs of conspicuous, $\pm$ pinnate secondary veins, tertiary venation reticulate; midribs prominent, pale green when fresh, $\pm$ the same colour as the leaf when dry; abaxial surface pale green when fresh, drying dark brown to blackish, matte or glossy, $\pm$ glabrous or with pubescent midrib and venation, tuft domatia often present in the vein axils. **Flowers** solitary, terminal, very fragrant; pedicels up to 9 mm long; hypanthium narrowly urceolate, 4.5–12 $\times$ 2.5–5.5 mm, glabrous; calyces green; calyx tubes 1–5.5 mm long, externally and internally glabrous; calyx lobes 4.5–8 $\times$ 1–3 mm, narrowly ovate, acuminate, recurved; corollas fleshy, funnellform, straight; corolla tubes 2–4.5 cm long, externally white to pale yellow often with a greenish base, glabrous, internally white with pink, red, or purple spots, finely pubescent (trichomes 0.5–1.5 mm long). **Corollas** lobes white (sometimes pale yellow on the external side), 2–4.5 $\times$ 1–3 cm, ovate, acute, spreading horizontally at anthesis; stamens subsessile (filaments c. 1 mm long), attached 10–14 mm below corolla sinus and 6–10 mm from anther base; anthers 22–26 $\times$ 1.5–2 mm, linear, exerted for c. 5 mm. **Styles** simple, clavate, with shallow longitudinal grooves, to c. 5 cm long, glabrous, usually shortly exerted for c. 5 mm; stigma shortly bifid (lobes c. 5 mm long). **Fruits** globose to ellipsoid, 3–5 $\times$ 2.5–3.5 cm, smooth, glabrous, olive to grey brown or dark purple to black with numerous beige to pale grey lenticels (0.1–1 mm in diam.), fleshy-indehiscent; calyx tube usually persistent, but lobes caducous. **Seeds** flattened, irregularly angled, 7–10 mm in diam., c. 2 mm thick, smooth, yellowish brown when fresh, drying black.

Vernacular names. – “Asomitsoy” (*Rakotonasolo et al.* RNF 932); “Hazomfotaka” (*Réserves Naturelles* 6762); “Kaliona” (*Koechlin s.n.*); “Mahabiboala” (*Service Forestier* 29830); “Vala” (*Service Forestier* 5797); “Voafotaka” (*Decary* 4556); “Volatsiva” (*Service Forestier* 4136).

Distribution, habitat and ecology. – *Melanoxerus suavissimus* is widespread in subarid vegetation in the Androy, Anosy, Atsimo-Andrefana, and Menabe Regions (Toliara Province), and occurs as far north as the Manambolo river in the southernmost part of the Melaky Region (Mahajanga Province) (Fig. 1). It is found from sea level to c. 800 m in elevation.

Phenology. – Flowering specimens have been collected in (April) July to December, and fruiting specimens throughout the year.

Conservation status. – *Melanoxerus suavissimus* is widespread in south-western Madagascar and has been assigned an IUCN conservation status of “Least Concern” [LC] (IUCN, 2021).



Fig. 5. – *Melanoxerus suavisissimus* (Homolle ex Cavaco) Kainul. & B. Bremer. **A.** Habit and habitat; **B.** Top view of the flower; **C.** Flower; **D.** Flowering branch; **E–F.** Immature fruits. [**A–B:** Ranaivojaona et al. 1456; **D:** Phillipson 2413; **E–F:** Andriamihajarivo et al. 1814] [Photos: **A–B, D–E:** P. Phillipson; **C:** S. Landrein, Ifaty (Toliara), 5.X.2006; **F:** K. Kainulainen]

Notes. – Since CAVACO's (1967) original description of *Melanoxerus suavisissimus* (as *Gardenia suavisissima*) included specimens of both *M. atropurpureus* and *M. maritimus*, an amended description is provided.

This is the most widespread and the most commonly collected species. However, few collections have been made in from May to August (none from June), and presumably this species is usually leafless during that time. *Melanoxerus suavisissimus* usually flowers at the same time or just before the new leaves flush (cf. Fig. 5A, D). Apparently, the juice of the fruits turns black and is used for making tattoos (SCHATZ, 2001). Information about the use of *Melanoxerus* by local

people is otherwise scant, although according to the label information on the specimen Ranaivojaona et al. 248, the fruits are edible.

Selected additional material examined. – MADAGASCAR. **Reg. Androy [Prov. Toliara]:** Besakoa, 24°56'36"S 45°27'59"E, 227 m, 4.III.2003, fr., Andriamahay 497 (K); Imanombo, XI.1952, fr., Bosser 3787 (P); Beraketa, X.1956, fl., Bosser 10237 (P); Ambovombe, XII.1959, fl. & fr., Bosser 13353 (P, TAN); Beraketa, on road to Isoanala, 10 km S of Soanala, 646 m, 19.XII.2004, fr., Rakotonasolo et al. RNF 932 (K, TAN); Antanimora, 30.VII.1926, fr., Decary 4556 (P); Behara, 22.X.1954, fl., Réserves Naturelles 6762 (P); Ampandrandava, 700–800 m, X.1942, fl., Seyrig 122 (P); Beloha, 19.II.1949, fr., Service Forestier 454 (P, TEF); Imanombo-Ifotaka (rte), 26.IX.1952, fr., Service Forestier 5797 (P, TEF). **Reg. Anosy [Prov. Toliara]:** Distr. Amboasary Sud, Marotsiraka,

Atsonjo Analamainty, Ankotsy, 24°17'44"S 46°07'43"E, 453 m, 10.XI.2008, fr., *Andriamihajarivo et al.* 1527 (G, MO); Anadabolava, 200–250 m, XII.1933, fr., *Humbert* 12450 (P); Andohahela, Ihazofotsy, 24°50'S 46°32'E, 100 m, 16.IV.1996, fr., *Laba* 75 (BR, G, K, MO); Fort Dauphin, Analamatahitra, 31.I.1952, fr., *Service Forestier* 4463 (P, TEF); Fort Dauphin, Anteviala, 10.IV.1960, fr., *Service Forestier* 10789 (P, TEF). **Reg. Atsimo-Andrefana [Prov. Toliara]:** Distr. Sakaraha, forêt d'Analaraty, 22°49'14"S 44°17'16"E, 520 m, 20.XII.2010, fr., *Andriamihajarivo et al.* 1814 (MO, P, TAN); Zombitse, lieu dit Poakafo, 22°46'21"S 44°40'25"E, 540 m, 8.IV.2006, fr., *Andriamihajarivo et al.* 905 (MO, TAN); Befandriana-Sud, 150 m, 9.IX.1961, fl. & fr., *Appert* 203 (MO, Z); along rte #10 between Betioky and Ejeda, 175–225 m, 14.II.1975, fr., *Croat* 31241 (MO, TAN); St. Augustin, 23°28'58"S 43°45'58"E, 0 m, 6.I.1999, fr., *De Block et al.* 561 (BR, K, MO, P, TAN); road from Betioky to Ampanihy, c. 23 km after Ejeda, 24°31'17"S 44°37'25"E, 244 m, 6.II.2007, fr., *De Block et al.* 2323 (BR, MO, P, S, TAN, WAG); entre Tongobory et aux environs d'Ambovombe, 117–214 m, 20.I.2004, fr., *Hong-Wa et al.* 178 (G, MO); forêt de Besomaty entre le Fiherenana et l'Isahaina (Mangoky), 750–800 m, X.1933, fl., *Humbert* 11250 (P); Manombo, Fiherenamasay, 18.XI.1967, fl., *Koechlin s.n.* (P); St. Augustin, 23°27'S 43°46'E, 10 m, 1.II.1990, fr., *Labat & Du Puy* 2045 (K, MO, P); vallée du Fiharenana, 0–200 m, 9.XI.1960, fl. & fr., *Leandri* 3778 (P); lac Tsimanampetsotsa, 0–100 m, 23.XI.1960, fr., *Leandri & Saboureau* 4451 (BR, K, P); 20–30 km N of Tulear on the road to Morombe, 23°12'S 43°37'E, 5–10 m, 11.XI.1989, fl. & fr., *Miller & Keating* 4513 (K, MO, P, TAN); rte d'Ankazoabo, II.1967, fr., *Morat* 2455 (P, TAN); vallée de Fiherenana, X.1924, fl., *Perrier de la Bathie* 16653 (P); Beza Mahafaly Reserve near Betioky, 23°40'S 44°35'E, 160 m, 17.V.1987, fr., *Phillipson* 1812 (MO); *ibid. loco*, 23°40'S 44°38'E, 150 m, 20.X.1987, fl., *Phillipson* 2413 (K, MO, P, TAN); *ibid. loco*, 23°41'S 44°39'E, 160 m, 13.XI.1987, fr., *Phillipson* 2558 (K, MO, P, TAN); Itampolo, Montagne de Haktokaliotse, 24°24'26"S 43°51'01"E, 100 m, 1.IX.1998, fr., *Rakotomalaza & Messmer* 1653 (G, K); Ambohimahavelona, Sept Lacs, Andranolahy, 23°31'32"S 44°09'27"E, 100 m, 15.IX.1998, fl., *Rakotomalaza & Messmer* 1725 (G, K); Manombo, near PK 32, 23°03'00"S 43°41'42"E, 9.XII.2004, fr., *Rakotonasolo et al.* RNF 880 (BR, K, MO, TAN); Basibasy, Mikea forest, 22°10'13"S 43°31'30"E, 100 m, 31.I.2000, fr., *Ranaivojaona et al.* 248 (G, K, MO); Belalanda, Ranobe, 22°56'49"S 43°41'08"E, 135 m, 15.III.2006, fr., *Ranaivojaona et al.* 1425 (P); *ibid. loco*, 22°58'04"S 43°40'22"E, 108 m, 5.IX.2006, fl., *Ranaivojaona et al.* 1456 (G, K, MO, P); Sakaraha, Mahaboboka, Ambinanintelo, Forêt de Satra, 22°48'10"S 44°17'54"E, 470 m, 23.III.2013, fr., *Randrianarivony et al.* 570 (MO, P); Zombitse-Vohibasia NP, Isako, 22°41'S 44°50'E, 4.XII.2003, ster., *Razafimandimbison & Bremer* 496 (MO, TAN, UPS); Beroroha, Antsoa, Androtsy forest, 21°05'43"S 45°05'48"E, 254 m, 9.I.2011, fr., *Razakamalala* 5851 (MO); Imonty, Ankazoabo, Sakaraha, 27.XI.1951, fr., *Service Forestier* 4136 (P, TAN, TEF); Antseva, Besifake, 500 m, 11.II.1955, fr., *Service Forestier* 12815 (P, MO, TEF); Sarodrano, XII.1961, fl., *Service Forestier* 20830 (BR, K, MO, P, TEF). **Reg. Melaky [Prov. Mahajanga]:** Tsingy de Bemaraha, 19°09'S 44°49'E, 50 m, 10.XII.1996, fr., *Jongkind et al.* 3482 (BR, K, L, MO, TAN). **Reg. Menabe [Prov. Toliara]:** village avant la Réserve d'Andranomena, 17.X.1999, fl., *Allorge* 2428 (P); on the road to Belo, c. 68.5 km NE of Morondava, 19°55'06"S 44°36'00"E, 5 m, 22.II.2000, fr., *Davis et al.* 2587 (BR, K, MO, TAN); Kirindy Mitea NP, 20°53'14"S 43°55'22"E, 20 m, 21.III.2011, fr., *Razafimandimbison & Kainulainen* 1205 (MO, S); N de Tsimafana env. 2 km, 25.V.1980, fr., *Service Forestier* 29830 (TEF).

Acknowledgements

I thank the curators of G, K, MO, P, S, TAN, TEF, and UPS for providing access to their collections, and Laurent Gautier, Mitia Hanitrarivo, the late Jean-Noël Labat, Sven Landrein, Peter Phillipson, Landy Rita Rajaovelona, Solotiana D. Raman-dimbimanana, Richard Randrianaivo, Monique Randriatsivory,

Fidisoa Ratovoson, Désiré Ravelonarivo, and A. Jacques Tahirivony for providing photographs. Conservatoire et Jardin botaniques de Genève, Muséum national d'Histoire naturelle, and Missouri Botanical Garden have also made photographs available. I also thank Petra De Block, Laurent Gautier, Sylvain Razafimandimbison, and Martin Callmänder for helpful comments to improve the manuscript.

References

- BAUM, D. (1995). A systematic revision of *Adansonia* (Bombacaceae). *Ann. Missouri Bot. Gard.* 82: 440–471.
- CAVACO, A. (1967). Un nouveau *Gardenia* (Rubiaceae) de Madagascar. *Adansonia* ser. 2, 7: 177–179.
- CORNET, A. (1974). Essai de cartographie bioclimatique à Madagascar. *Notice Explicative ORSTOM* 55.
- IUCN (2012). *IUCN Red List Categories and Criteria: Version 3.1*. Ed. 2. IUCN Species Survival Commission, Gland and Cambridge.
- IUCN (2021). *The Red List of threatened species*. [https://www.iucn-redlist.org]
- KAINULAINEN, K. & B. BREMER (2014). Phylogeny of *Euclinia* and allied genera of *Gardenieae* (Rubiaceae), and description of *Melanoxerus*, an endemic genus of Madagascar. *Taxon* 63: 819–830. DOI: https://doi.org/10.12705/634.2
- KAINULAINEN, K., S.G. RAZAFIMANDIMBISON, N. WIKSTRÖM & B. BREMER (2017). Island hopping, long-distance dispersals and species radiations in the Western Indian Ocean: historic biogeography of the *Coffeae* alliance (Rubiaceae). *J. Biogeogr.* 44: 1966–1979. DOI: https://doi.org/10.1111/jbi.12981
- LEROY, J.-F. (1974). Recherches sur les Rubiacées de Madagascar: les genres *Mantalania* et *Pseudomantalania* (Gardéniees). *Adansonia* ser. 2, 14: 29–52.
- MADAGASCAR CATALOGUE (2021). *Catalogue of the plants of Madagascar*. Missouri Botanical Garden, St. Louis & Antananarivo. [http://www.tropicos.org/Project/Madagascar]
- OLSON, D.M., E. DINERSTEIN, E.D. WIKRAMANAYAKE, N.D. BURGESS, G.V.N. POWELL, E.C. UNDERWOOD, J.A. D'AMICO, I. ITOUA, H.E. STRAND, J.C. MORRISON, C.J. LOUCKS, T.F. ALLNUTT, T.H. RICKETTS, Y. KURA, J.F. LAMOREUX, W.W. WETTENGEL, P. HEDAO & K.R. KASSEM (2001). Terrestrial ecoregions of the world: a new map of life on Earth. *BioScience* 51: 933–938.
- SCHATZ, G.E. (2000). Endemism in the Malagasy tree flora. In: LOURENÇO, W.R. & S.M. GOODMAN (ed.), *Diversity and Endemism in Madagascar*: 1–9. Mémoires de la Société de Biogéographie, Paris.
- SCHATZ, G.E. (2001). *Generic tree flora of Madagascar*. Royal Botanic Gardens, Kew.