

Use of geometric morphometrics to distinguish trapdoor spider morphotypes (Mygalomorphae: Anamidae: Proshermacha): a useful tool for mygalomorph taxonomy

Authors: Sagastume-Espinoza, Kevin O., Simmons, Leigh W., and Harvey, Mark S.

Source: The Journal of Arachnology, 52(1) : 31-40

Published By: American Arachnological Society

URL: <https://doi.org/10.1636/JoA-S-22-033>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Use of geometric morphometrics to distinguish trapdoor spider morphotypes (Mygalomorphae: Anamidae: *Proshermacha*): a useful tool for mygalomorph taxonomy

Kevin O. Sagastume-Espinoza^{1,2,3,5}, Leigh W. Simmons¹ and Mark S. Harvey^{3,4}: ¹Centre for Evolutionary Biology, School of Biological Sciences, The University of Western Australia, Crawley, WA 6009, Australia; E-mail: sagastume.espinozak@gmail.com; ²Laboratorios de Zoología, Escuela de Biología, Universidad Nacional Autónoma de Honduras (UNAH), Tegucigalpa, Honduras; ³Collections and Research, Western Australian Museum, 49 Kew Street, Welshpool, WA 6106, Australia; ⁴School of Biological Sciences, The University of Western Australia, Crawley, WA 6009, Australia; ⁵Bennelongia Environmental Consultants, Jolimont, WA 6014.

Abstract. Taxonomic studies have evolved greatly since their early stages and new techniques have been incorporated to improve species descriptions. Those involving the comparison of traits, either quantitatively or qualitatively, can be difficult because the identification of a species must rely on the experience of the observer and errors can occur when cryptic species are involved. Molecular methods have been used to fill these gaps, but morphological methods are still needed to match the recognized molecular species with an adequate taxonomic description. Focusing on the trapdoor spider genus *Proshermacha* Simon, 1908, we provide a case study using Geometric Morphometrics (GM) techniques to identify morphological divergence between species found in the south-western Australia region. We used GM to identify morphological divergence from museum-preserved specimens by examining shape variation of sexual characters from 39 male specimens from five different localities on a single mountain range. Variation in the shape of both the palpal bulb and tibia provided strong evidence to distinguish two morphotypes, while metatarsus shape showed fewer between-locality differences. Our results illustrate the utility of GM methods, when applied to a few taxonomically-informative structures, as a quantitative species delimitation tool for taxonomic studies.

Keywords: Araneae, morphology, Stirling Range, biodiversity hotspot, taxonomy
<https://doi.org/10.1636/JoA-S-22-033>

Taxonomy is a key discipline for describing and understanding biodiversity and is critical for documenting and inventorying undescribed species before they reach extinction due to climate change or habitat loss (Wheeler 2018, 2020). Even though taxonomy has been recently recognized as extremely important (Bond et al. 2022), it has been regarded by some as less relevant and with little intellectual content, resulting in what has been called “taxonomy in crisis” (Agnarsson & Kunter 2007). It is true that many taxonomic groups, such as mygalomorph spiders (e.g., Rix et al. 2017; Harvey et al. 2018; Opatova et al. 2020) often remain poorly resolved due to the occurrence of multiple cryptic species that offer little morphological variation that can be used by taxonomists in species description. In this study, we show how multivariate statistical analysis of trait shape can assist in species delimitation in taxonomic studies of mygalomorph spiders.

The genus *Proshermacha* Simon, 1908 (Mygalomorphae: Anamidae) was recently resurrected (Harvey et al. 2018) to include nine species (World Spider Catalog 2021) from southern Australia (Main 1982; Raven 2000). Members of the genus are relatively gracile mygalomorph spiders that construct silk-lined burrows with an open entrance (Harvey et al. 2018). Molecular data indicate the existence of numerous undescribed species within its distributional range (Harvey et al. 2018). One locality of interest for the genus is the Stirling Range, a mountain formation estimated to have originated ~1.2 billion years ago (Rasmussen et al. 2002). The Range is home to ancient endemic invertebrate species including land snails, onychophorans, assassin spiders, and trapdoors spiders (Rix et al. 2015), all of which are considered relictual groups that can be traced back to the Gondwanan super-continent (Cooper et al. 2011). It has been suggested that biodiversity within the Range has been driven by multiple mechanisms of diversification, such as the remnants of relictual fauna, vicariant isolation, or *in situ* speciation (Rix et al. 2015).

Several studies have explored the biogeography and evolution of Stirling Range fauna. For instance, five species of the millipede

genus *Atelomastix* Attems, 1911 (Diplopoda: Spirostreptida: Iulomorphidae) (Edwards & Harvey 2010), three species of the genus *Bertmainius* Harvey, Main, Rix & Cooper, 2015 (Araneae: Migidae) (Harvey et al. 2015), and four species of the trapdoor spider genus *Cataxia* Rainbow, 1914 (Araneae: Idiopidae) (Rix et al. 2017) have been identified and are known to be endemic to the Range. In each of these genera, individual species are found on different peaks within the range, supporting the idea that the Stirling Range acted as a natural refuge for relict invertebrates and, given its isolated sky-island nature, the peaks represent an important driver of allopatric speciation for species with limited-dispersal (Main 1993). Although little is known regarding the habitat requirements, biogeography, and speciation patterns of the genus *Proshermacha*, like all mygalomorph spiders they are likely to be dispersal-limited (Buzatto et al. 2021), and we might therefore expect to find a similar pattern of vicariant isolation in the Stirling Range.

Extracting both qualitative and quantitative data from morphological characters of mygalomorph spiders can often be difficult. Morphological traits might suggest geographically isolated populations that represent a single, undifferentiated species, which when analyzed through molecular methods result in distinct molecular species (Bond & Stockman 2008). Molecular data has without doubt contributed to an increasing appreciation of the extent of biodiversity (Franzini et al. 2013; Castalanelli et al. 2014; Teixeira Jr. et al. 2016; Hupalo et al. 2020) by recognizing molecular divergence when there are seemingly few morphological differences between taxa (Bond et al. 2001), and by solving delimitation of species complexes (Bond & Stockman 2008) or differentiating between cryptic species (Leavitt et al. 2015). Molecular data have helped develop modern-day systematic classifications of mygalomorph spiders (e.g., Bond et al. 2012; Opatova et al. 2020; Harvey et al. 2018, 2020) that provide a better understanding of the evolutionary relationships amongst groups and lineages.




Figure 1.—Map of morphotype distributions and sampling localities. ○ represents Morph1 and ☆ represents Morph2. Notice sympatry of morphotypes at Mondurup, Mt. Magog (separated altitudinally), and Toolbrunup.

Taxonomy has improved greatly since its initial species descriptions based on comparative morphology, with new and complementary methods being developed for delimiting species boundaries (Dayrat 2005; Padial et al. 2010). Although molecular sequencing provides more reliable, reproducible, universal, and scalable data (Sharma et al. 2021), both natural history and morphological data are crucial, and should be used in combination to support species hypotheses derived from molecular data (Giribet 2015; Muster & Michalik 2020). Traditional and modern approaches should be combined into an integrative taxonomy (Bond et al. 2022). The role of morphology becomes even more evident for those species groups that have not been revised and are known only from museum material which can not be analyzed with molecular methods due to inadequate preservation (Derkarabetian et al. 2019). In the field of arachnology, most molecular studies work in tandem with qualitative morphological descriptions (with some exceptions; see Bond 2012), but few incorporate multivariate statistical methods during data analysis to support morphological differences between molecularly distinct species (Wilson et al. 2021).

Geometric morphometrics (GM) is a technique that relies on landmarks placed over specific anatomical structures from which quantitative shape variation can be obtained (Bookstein 1991; Rohlf & Marcus 1993). Soon after its development (Bookstein 1991), GM was adopted by numerous biologists and has improved morphological studies by allowing multivariate analysis to be conducted to capture shape variation (Adams et al. 2004). GM has been used widely in taxonomic groups such as vertebrates, reinforcing species delimitations based on multilocus data in milksnakes (Ruane 2015), to maximize shape coverage in 2D models of varanid lizards (Openshaw et al.

2017), and even to explore optimal sample sizes and sampling error when utilizing GM using horse teeth as a model (Cardini 2014). Within arachnids GM has been used to compare phylogenetic and morphological divergences, reveal sexual dimorphism in shape allometry, quantify intraspecific variation, and to delimit species within a species-complex (Crews & Hedin 2006; Fernández-Montraveta & Marugán-Lobón 2017; Torres et al. 2018; Wilson et al. 2021). Like many fields, GM has grown rapidly with new techniques becoming available such as the use of semi-landmarks that allow capturing curvature data (Adams et al. 2013), the development of specialized programs for landmark acquisition and manipulation (Rohlf 2013, 2017, 2021), and specific statistical packages to analyze landmark data (Adams et al. 2016). The use of GM provides taxonomists a quantitative tool with which to test for morphological differences among species (Wilson et al. 2021). Although there is the added cost of greater effort and identification time, quantitative shape analysis may be the only means available when discrete morphological features are unavailable. We had access to numerous alcohol-preserved specimens belonging to the genus *Proshermacha* that were collected during the early 1990s. These samples provided an ideal study group for applying quantitative geometric morphometric analysis to test for distinct morphotypes and their distributions through the Stirling Range.

METHODS

Study site.—Samples used in this study came from five different localities in the Stirling Range National Park in southwest Western Australia (Fig. 1). Mondurup and Talyuberlup Peaks, in




Figure 2.—Visually distinct types of bulbs, ventral view (A–C); tibia, prolateral side of leg (E–F); metatarsus, prolateral side of leg (H–I); and detail of landmark placement (D, G, and J). Arrow in B points to the flanged process at the tip of the embolus. Scale bar represents 1mm for all images.

the western and southern parts of the range, respectively, are dominated by thicket plant formations in the higher areas, and mallee and woodlands at lower elevations. Mount Magog and Toolbrunup, in the northern and eastern parts of the range, reach higher elevations, and both are dominated by a combination of thicket and mallee (Keighery 1993). Two Mile Lake is located on the lower elevations at the eastern edge of the park and is

dominated by open landscapes with remnants of bushes and short mallee (Keighery 1993).

Material examined.—Samples were collected in pitfall traps with ethyl-glycol (70–30 mix) as preservation liquid and stored in 70% ethanol in the Western Australian Museum (WAM). Given the lack of female samples across all localities, only adult males were used for this study. A total of 39 specimens from an undescribed

Table 1.—Landmark placements and their contribution to shape variation for each structure. Type I are fixed landmarks, Type II are sliding semi-landmarks (Bookstein 1991, Zelditch et al. 2004).

BULB				TIBIA				METATARSUS			
Landmark	Type	Description	Contribution	Landmark	Type	Description	Contribution	Landmark	Type	Description	Contribution
LM1	I	left origin of genital bulb	1.55%	LM1	I	left distal origin	2.67%	LM1	I	left proximal origin	1.13%
LM2	II	middle point between left origin of genital bulb and left upper limit of seminal duct	1.79%	LM2	II	left middle point between distal origin and origin of tibial apophysis	1.48%	LM2	II		1.09%
LM3	II	left upper limit of seminal duct	1.87%	LM3	II	origin of tibial apophysis	2.87%	LM3	II		1.18%
LM4	II	left middle section of seminal duct	1.18%	LM4	II	origin of tibial spur	16.94%	LM4	II	evenly distributed points along external face	1.35%
LM5	II	left lower limit of seminal duct	0.46%	LM5	II	right-faced middle point on tibial spur	20.37%	LM5	II		2.11%
LM6	II	middle point between left lower limit of seminal duct and left end of genital bulb	0.30%	LM6	I	tip of tibial spur	0.23%	LM6	II		2.37%
LM7	I	left end of genital bulb	0.33%	LM7	II	left-faced middle point of tibial spur	18.76%	LM7	II		1.76%
LM8	II	evenly distributed points along left-side of embolus	0.56%	LM8	II	end of tibial spur	12.84%	LM8	I	left distal origin	2.94%
LM9	II		1.30%	LM9	II	midpoint of apophysis between end of spur and end of apophysis	2.81%	LM9	II	middle point between left and right distal origins	11.07%
LM10	II		3.85%	LM10	II	end of tibial apophysis	2.18%	LM10	I	right distal origin	6.65%
LM11	II		8.10%	LM11	II	evenly distributed points on left proximal side	1.40%	LM11	II	evenly distributed points along internal face	2.66%
LM12	II		6.00%	LM12	II		1.19%	LM12	II		2.64%
LM13	II		6.73%	LM13	I	left proximal origin	0.93%	LM13	II		2.59%
LM14	I	tip of embolus	7.79%	LM14	II	middle point between left and right proximal origins	2.52%	LM14	II	origin of metatarsal depression	5.71%
LM15	II	evenly distributed points along right-side of embolus	23.29%	LM15	I	right proximal origin	2.08%	LM15	II	evenly distributed points along metatarsal depression	13.62%
LM16	II		13.92%	LM16	II	evenly distributed points on right proximal side	1.06%	LM16	II		12.99%
LM17	II		9.27%	LM17	II		0.62%	LM17	II		9.43%
LM18	II		4.01%	LM18	II		0.74%	LM18	I	right proximal origin	9.37%
LM19	II		1.26%	LM19	II		0.96%	LM19	II	middle point between left and right proximal origins	9.32%
LM20	II		0.64%	LM20	II		0.92%				
LM21	I	right end of genital bulb	0.36%	LM21	I	right distal origin	1.56%				
LM22	II	right middle point between right lower limit of seminal duct and right end of genital bulb	0.22%	LM22	II	middle point between left and right distal origins	4.87%				
LM23	II	right lower limit of seminal duct	0.34%								
LM24	II	right middle section of seminal duct	0.52%								
LM25	II	right upper limit of seminal duct	0.71%								
LM26	II	middle point between right origin of genital bulb and right upper limit of seminal duct	1.02%								
LM27	I	right origin of genital bulb	1.25%								
LM28	II	middle point between left and right origins of genital bulb	1.39%								

Table 2.—Percentage of variance in trait shape explained by each Relative Warp (RW). Only those RW corresponding to >95% of trait shape variance per structure are included.

Structure	RW	Individual %	Cumulative %
Palpal bulb	1	71.09	71.09
	2	9.5	80.59
	3	6.27	86.87
	4	3.93	90.80
	5	2.39	93.19
	6	1.83	95.02
Tibia	1	61.84	61.84
	2	14.43	76.27
	3	8.8	85.07
	4	5.49	90.55
	5	3.48	94.04
	6	2.04	96.08
Metatarsus	1	60.48	60.48
	2	17.25	77.73
	3	8.07	85.8
	4	3.53	89.33
	5	2.52	91.86
	6	1.93	93.79
	7	1.47	95.26

species were analyzed, coming from the localities as follows: Mondurup $n = 5$; Mt. Magog Low $n = 4$; Mt. Magog High $n = 4$; Talyuberlup $n = 5$; Toolbrunup $n = 15$; Two Mile Lake $n = 6$. Across all specimens, the right palpal bulb was separated from the cymbium (Figs. 2A–C), and the left leg I was detached from the body for capturing images of the tibia (Figs. 2E,F) and metatarsus (Figs. 2H, I) to allow for accurate landmark placement. Structures were fully submerged in 70% ethanol on a petri dish and held in place by colorless hair styling gel when capturing images, orienting the bulb to capture the ventral side and the retrolateral side for leg I.

Image acquisition was conducted with the Leica Application Suite v4.6 using the multi-focus setting, visualized through a Leica DFC 500 digital camera attached to a Leica MZ16A microscope. Image scaling and landmark digitalization were performed using tpsDig2 v2.3.1 (Rohlf 2017) and following a counterclockwise direction for each anatomical structure. Landmarks consisted of Type I fixed and Type II sliding semi-landmarks and followed the typology of Bookstein (1991) and Zelditch et al. (2004) selection criteria and are described in Table 1 (see Fig. 2D, G, and J for landmark placement).

Morphometrics and statistical analysis.—Landmark coordinates were aligned and superimposed by Generalized Procrustes Analysis (GPA) with the software tpsRelw (Rohlf 2013) which provided data including centroid size (CS), partial warps (PW), and relative warps (RW). Measurements of cephalothorax length were obtained using the linear measurement tool in the Leica Application Suite v4.6 and were used as the standard measure of body size (*Size*) for each sample. Repeatability analyses were conducted for each of the three traits (bulb, tibia, and metatarsus) to determine if the selected landmarks provided repeatable estimates of trait size and shape. Landmarks were placed on images from 10 randomly selected individuals (2 per site), on 3 different occasions. Relative warps and centroid sizes were extracted and the repeatability analyses were conducted as Gaussian data type with 1000 bootstrap iterations using the rptR package (Stoffel et al. 2017).




Figure 3.—Scatterplots showing positive and negative values for RW1 and RW2. Notice the clear formation of two distinct clusters for (A) bulb and (B) tibia, but not so clear differences for (C) metatarsus.

Multivariate analysis of variance (MANOVA) was conducted using those RWs that represented 95% of the cumulative shape variance of each structure. *Locality* was used as the explanatory variable, and both *Size* and *CS* were used as covariates to control for body size and trait size respectively during analysis. Pairwise comparisons between populations were made using only RW1 for each structure as it accounted for most of the shape variation (> 60%). All values were set to 0.05 significance and were conducted in RStudio (RStudio 2019) using R v.4.0.5 (2021).

RESULTS

Shape analysis.—Repeatability analysis showed that landmark placement provided repeatable estimates for *bulb* centroid size ($P < 0.001$, $R = 0.98$ [CI = 0.941, 0.993]) and shape RW1 ($P < 0.001$, $R = 0.96$ [CI = 0.876, 0.986]), *tibia* centroid size




Figure 4.—Boxplots for quantitative values of RW1 per locality for (A) bulb, (B) tibia, (C) metatarsus, and (D) size (measured as cephalothorax length).

($P < 0.001$, $R = 0.98$ [CI = 0.955, 0.996]) and shape RW1 ($P < 0.001$, $R = 0.988$ [CI = 0.957, 0.995]), and *metatarsus* centroid size ($P < 0.001$, $R = 0.99$ [CI = 0.978, 0.998]) and shape RW1 ($P < 0.001$, $R = 0.99$ [CI = 0.979, 0.998]). For the three structures, RW1 explained most of the trait shape variance (Table 2). Visualisation of RW1 and RW2 revealed the existence of two clusters of specimens for *bulb* (Fig. 3A) and *tibia* (Fig. 3B), each of which contained specimens from multiple localities, but no clear clusters could be observed for *metatarsus* (Fig. 3C).

Statistical analysis and data visualization.—MANOVA revealed significant differences in mean shape for all structures across localities: *bulb* ($P < 0.001$, $F_{30,160} = 3.48$), *tibia* ($P < 0.001$, $F_{30,160} = 3.75$), and *metatarsus* ($P < 0.001$, $F_{35,155} = 3.02$). Data from RW1 showed differences between localities for *bulb* (Fig. 4A), *tibia* (Fig. 4B), and *size* (Fig. 4D), but no clear differences could be observed with *metatarsus* RW1 (Fig. 4C). Shape differences corresponding to extremes in RW1 were visualized using thin-plate spline deformation grids (Fig. 5). For the *bulb*, positive RW1 values represent a long, elongated structure without a flanged tip (Fig. 5A), while negative RW1 values represent a shorter, flanged structure (Fig. 5C). For *tibia*, positive RW1 values represent a narrower structure (Fig. 5D), and negative RW1 values represent a wider structure (Fig. 5F). For *metatarsus*, positive RW1 values represent a slim structure with a short metatarsal depression (Fig. 5G), and negative RW1 shows a broad structure with a long metatarsal depression (Fig. 5I), though variation in shape along this axis was not statistically significant.

Locality comparisons.—Samples from Mt. Magog Low, Mondurup, and Two Mile Lake show positive RW1 values for *bulb*, negative RW1 values for *tibia*, and positive RW1 values for *metatarsus*. Those from Mt. Magog High, Talyuberlup, and Toolbrunup show

the opposite, with negative RW1 values for *bulb*, positive RW1 values for *tibia*, and negative RW1 values for *metatarsus* (Fig. 5). *Bulb* data (Table 3A) suggests there is no difference between samples from Mt. Magog High, Talyuberlup, and Toolbrunup, and shows a strong difference between Mt. Magog Low & Mt. Magog High, Talyuberlup, and Toolbrunup. Two Mile Lake samples differ from Mt. Magog High, Mt. Magog Low, Talyuberlup, and Toolbrunup, while Mondurup did not differ significantly from any other locality.

Similar results were found with *tibia* data (Table 3B), with the exception that no difference was found between Mt. Magog Low and Two Mile Lake. *Metatarsus* showed almost no differences across all localities, with differences only detected between Two Mile Lake samples and those from Talyuberlup and Toolbrunup (Table 3C).

DISCUSSION

Morphotype recognition and distribution.—Based on the two statistically significant traits, *bulb* and *tibia*, two distinct morphotypes could be recognized amongst the five localities. *Morph1* is characterized by having a long, elongated bulb (Fig. 2A), a wide tibia (Fig. 2F), and was found on individuals from Mt. Magog Low, Mondurup, and Two Mile Lake (Fig. 1). *Morph2* has a short, flanged bulb (Fig. 2B), a narrow tibia (Fig. 2E), and was found on individuals from Mt. Magog High, Talyuberlup, and Toolbrunup (Fig. 1).

Biogeographical patterns could not be accurately estimated given the limited sample size across localities. Considering the overlapping distributions of both morphotypes, and the fact that there is no apparent altitudinal separation between them, the morphotypes are expected to show some level of sympatry. This contradicts what has




Figure 5.—Thin-plate spline deformation grids showing extreme positive RW values for (A) bulb, (D) tibia, and (G) metatarsus; consensus values for (B) bulb, (E) tibia, and (H) metatarsus; and extreme negative RW values for (C) bulb, (F) tibia, and (I) metatarsus. Consensus deformation grids (B, E, H) show landmark placement at each structure (see Table 1 for descriptions).

been found for other dispersal-limited invertebrate species within the Stirling Range (Edwards & Harvey 2010; Harvey et al. 2015; Rix et al. 2017). It is possible that even though both morphotypes of *Proshermacha* can be found on the same mountain, they inhabit isolated areas with little to no overlap between them and are separated by unknown ecological constraints. Further sampling efforts should be conducted to address this topic and provide a better understanding of the morphotype biogeography within the Stirling Range.

Trait effectiveness for morphotype recognition.—Out of the three structures, *bulb* and *tibia* showed more evident differences between localities, while *metatarsus* provided no statistical

separation and does not appear to differentiate localities clearly (Fig. 4). Copulatory structures in spiders have been widely used in spider systematics to differentiate between morphologically similar species (Huber 2004). Interestingly, not only do primary copulatory organs evolve rapidly but also secondary copulatory structures such as the tibial spurs are considered useful when differentiating between closely related species (Simmons 2014). Indeed, genital morphology is one of the best traits when it comes to understanding species-level divergence (Eberhard 2010), and the rapid evolution of the copulatory organs is the most likely reason why our results suggest stronger evidence for recognizing two distinct morphotypes when using bulb and tibia (tibial spur). Both structures are known to aid directly in mating by affecting insemination and grasping female fangs/legs/coxae during copula, respectively (Foeelix 2011; Pérez-Miles & Perafán 2017). In contrast, little is known about how the metatarsus might influence mating in spiders.

An initial visual examination of samples might suggest there to be three different bulb types among the available samples (Figs. 2A–C). However, our data analysis recognized only two morphotypes (Figs. 2A,B). Through a qualitative approach, the individual in Fig. 2C may appear visually distinct, but under a quantitative analysis of shape, it appears to represent a size variation of Morph1. Further study including molecular data will provide information to determine whether there is a substantial genetic divergence between morphotypes to be considered different valid species. Additional studies including female samples need to be made, and given female genitalia can diverge as rapidly as males (Simmons & Fitzpatrick 2019), it is expected that morphological examination of female samples should reveal a similar pattern of divergence across localities.

Geometric morphometrics as a taxonomical tool.—With further methodological advances, different lines of evidence can be integrated onto species delimitation (Agnarsson & Kunter 2007), combining quantitative data along with descriptive characters. Here we provide an example where GM appears as a useful tool for arachnid taxonomy. In mygalomorphs, contradicting results have been found when contrasting morphological and molecular data. Some studies have distinguished between different morphological and molecular lineages within cryptic species (Wilson et al. 2021), while others have found no morphological differences despite clear molecular divergence (Bond & Stockman 2008). Bond & Stockman (2008) found no morphological differences when using male mating claspers (leg I) in a group of trapdoor spiders, and as extensive as their quantitative measures were, our results suggest that the assessment of other reproductive structures might yet reveal differences between species.

It has been suggested that morphological data have become accessory to molecular phylogenetics when documenting speciation (Bond & Stockman 2008; Satler et al. 2013; Sharma et al. 2021). However, morphological data should remain part of an integrative approach for species descriptions, providing one of many methods contributing to species delimitations and part of the transformation in taxonomy to a collaborative, integrative information science (Bond et al. 2022). An approach like GM can provide statistical support in the recognition of morphological divergence among samples and could become a fundamental statistical approach in the arachnologists' toolkit.

Limitations of Geometric Morphometrics.—Landmark placement and orientation can become complicated tasks when dealing with intricate morphological structures, and it increases in complexity with the addition of semi-landmarks for analyzing outlines or

Table 3.—Pairwise comparisons between localities using RW1 for each structure. Bold values represent significant differences.

A. BULB					
	Mt. Magog H	Mt. Magog L	Mondurup	Talyuberlup	Toolbrunup
Mt. Magog L	0.0062	-	-	-	-
Mondurup	0.1855	0.7821	-	-	-
Talyuberlup	1.000	0.0049	0.1220	-	-
Toolbrunup	1.000	<0.001	0.1774	1.000	-
Two Mile Lake	0.0038	0.0383	1.000	0.0013	0.0026
B. TIBIA					
	Mt. Magog H	Mt. Magog L	Mondurup	Talyuberlup	Toolbrunup
Mt. Magog L	0.0014	-	-	-	-
Mondurup	0.1375	1.000	-	-	-
Talyuberlup	1.000	0.0011	0.1679	-	-
Toolbrunup	1.000	<0.001	0.1113	1.000	-
Two Mile Lake	<0.001	1.000	1.000	<0.001	<0.001
C. METATARSUS					
	Mt. Magog H	Mt. Magog L	Mondurup	Talyuberlup	Toolbrunup
Mt. Magog L	1.000	-	-	-	-
Mondurup	1.000	1.000	-	-	-
Talyuberlup	0.5995	0.5051	0.4189	-	-
Toolbrunup	1.000	0.6581	0.7670	1.000	-
Two Mile Lake	1.000	1.000	1.000	<0.001	0.007

curvature, such as in the palpal bulb. Consistent orientation of the structures is crucial for the landmarks to effectively capture any shape variation, and this can sometimes become a complicated mission when dealing with small structures such as spider genitalia.

When studying closely related populations, a sample size of several dozen individuals has been suggested (Cardini et al. 2015) and limitations on sample size can limit the power of shape analysis. Confidence intervals and variance estimates, as measures of uncertainty, are negatively correlated with sample size, where smaller samples exhibit larger confidence intervals and variance estimates. Not only statistical results, but also the visualization of shape variation might also be affected by small sample size, resulting in estimates of mean shapes not being truly representative of true biological variation (Cardini & Elton 2007). There is a possibility that the “third” bulb type identified from the initial visual examination (Fig. 2C) might represent a distinct morphotype but was not recognized in our statistical analysis because the program failed to fully eliminate size variation from a bimodal distribution of male size due to limited sample size, even after a Generalized Procrustes Analysis was conducted. However, the capacity of our data to differentiate between sympatric individuals suggests that this was not the case and that the specimens matching Fig. 2C morphology are a variation of Morph1. Nonetheless, further sampling should be conducted on the different Stirling Range populations to corroborate these findings.

Evaluating a single plane of view (ventral) from 2D images alone might be a limiting factor in the identification of morphological divergence between populations, and alternate planes of view could provide additional information. Other studies have compared the effectiveness of 3D over 2D images in shape analysis, suggesting that 2D images may underestimate shape variation, resulting in erroneous conclusions (Cardini 2014). This is likely to be especially problematic for complex structures with greater variation (Buser et al. 2018). Nevertheless, this is not always the case, and for some structures a traditional 2D approach can be effective (McWhinnie & Parsons 2019).

In conclusion, even though GM can provide quantitative evidence for improving taxonomic studies it should not be implemented alone, but rather used as a template to which other morphological datasets such as MRI (Ziegler et al. 2011) or μ CT scans (Semple et al. 2018) can be added. Our findings demonstrate that 2D GM can identify morphological divergence between sympatric populations, and as others have shown (Crews & Hedin 2006; Franzini et al. 2013; Wilson et al. 2021) they can be paired with molecular methods to resolve species hypotheses with quantitative morphological support.

ACKNOWLEDGMENTS

This research formed part of KSE master's degree which received funding through the *Hondufuturo-PCB 2019* and *Programa Becas 20/20* scholarships, for which the author is greatly thankful. Many of the specimens used in the study were kindly collected by Sarah Barrett (Department of Biodiversity, Conservation, and Attractions). The authors thank Peter Michalik, Jeremy D. Wilson, and two anonymous reviewers for their comments which significantly improved the manuscript.

LITERATURE CITED

- Adams D, Collyer M, Kaliontzopoulou A, Sherratt E. 2016. geomorph: Software for geometric morphometric analyses. R package, version 3.0.2. Online at cran.r-project.org/web/packages/geomorph/index
- Adams D, Rohlf J, Slice D. 2004. Geometric morphometrics: Ten years of progress following the 'revolution'. *Italian Journal of Zoology* 71:5–16.
- Adams D, Rohlf J, Slice D. 2013. A field comes of age: geometric morphometrics in the 21st century. *Hystrix, the Italian Journal of Mammalogy* 24:7–14.
- Agarsson I, Kunter M. 2007. Taxonomy in a changing world: seeking solutions for a science in crisis. *Systematic Biology* 56:531–539.

- Bond JE. 2012. Phylogenetic treatment and taxonomic revision of the trapdoor spider genus *Aptostichus* Simon (Aranea, Mygalomorphae, Euctenizidae). *ZooKeys* 252:1–209.
- Bond JE, Stockman AK. 2008. An integrative method for delimitating cohesion species: Finding the population-species interface in a group of Californian trapdoor spiders with extreme genetic divergence and geographic structuring. *Systematic Biology* 57:628–646.
- Bond JE, Godwin RL, Colby JD, Newton LG, Zahnle XJ, Agnarsson I, et al. 2022. Improving taxonomic practices and enhancing its extensibility- An example from Araneology. *Diversity* 14:5.
- Bond JE, Hedin MC, Ramirez MG, Opell BD. 2001. Deep molecular divergence in the absence of morphological and ecological change in the Californian coastal dune endemic trapdoor spider *Aptostichus simus*. *Molecular Ecology* 10:899–910.
- Bond JE, Hendrixson BE, Hamilton CA, Hedin MC. 2012. A reconsideration of the classification of the spider infraorder Mygalomorphae (Arachnida: Araneae) based on three nuclear genes and morphology. *PloS ONE* 7: e38753.
- Bookstein F. 1991. Morphometric Tools for Landmark Data: Geometry and Biology. Cambridge University Press.
- Buser T, Sidlauskas B, Summers A. 2018. 2D or Not 2D? Testing the utility of 2D vs. 3D landmark data in geometric morphometrics of the Sculpin Subfamily Oligocottinae (Pisces; Cottoidea). *The Anatomical Record* 301:806–818.
- Buzatto BA, Haessler L, Tamang N. 2021. Trapped indoors? Long-distance dispersal in mygalomorph spiders and its effect on species ranges. *Journal of Comparative Physiology A* 207:279–292.
- Cardini A. 2014. Missing the third dimension in geometric morphometrics: how to assess if 2D images really are a good proxy for 3D structures? *Hystrix, the Italian Journal of Mammalogy* 25:73–81.
- Cardini A, Elton S. 2007. Sample size and sampling error in geometric morphometric studies of size and shape. *Zoomorphology* 126:121–134.
- Cardini A, Seetah K, Barker G. 2015. How many specimens do I need? Sampling error in geometric morphometrics: testing the sensitivity of means and variances in simple randomized selection experiments. *Zoomorphology* 134:149–163.
- Castalanelli MA, Teale R, Rix MG, Kennington WJ, Harvey MS. 2014. Barcoding of mygalomorph spiders (Araneae: Mygalomorphae) in the Pilbara bioregion of Western Australia reveals a highly diverse biota. *Invertebrate Systematics* 28:375–385.
- Cooper SJB, Harvey MS, Saint KM, Main BY. 2011. Deep phylogeographic structuring of populations of the trapdoor spider *Moggridgea tingle* (Migidae) from southwestern Australia: evidence for long-term refugia within refugia. *Molecular Ecology* 20:3219–3236.
- Crews SC, Hedin MC. 2006. Studies in morphological and molecular phylogenetic divergence in spiders (Araneae: *Homalonychus*) from the American southwest, including divergence along the Baja California Peninsula. *Molecular Phylogenetics and Evolution* 38:470–487.
- Dayrat B. 2005. Towards integrative taxonomy. *Biological Journal of the Linnean Society* 85:407–417.
- Derkarabetian S, Benavides LR, Giribet G. 2019. Sequence capture phylogenomics of historical ethanol-preserved museum specimens: Unlocking the rest of the vault. *Molecular Ecology Resources* 19:1–14.
- Eberhard WG. 2010. Evolution of genitalia: theories, evidence, and new directions. *Genetica* 138:5–18.
- Edwards KL, Harvey MS. 2010. A review of the Australian millipede genus *Atelomastix* (Diplopoda: Spirostrepida: Iulomorphidae). *Zootaxa* 2371:1–63.
- Fernández-Montraveta C, Marugán-Lobón J. 2017. Geometric morphometrics reveals sex-differential shape allometry in a spider. *PeerJ* 5: e3617.
- Foelix R. 2011. Biology of Spiders. Third edition. Oxford University Press.
- Franzini PZ, Dippenaar-Schoeman AS, Yessoufou K, Van der Bank FH. 2013. Combined analysis of genetic and morphological data indicate more than one species of *Cyrtophora* (Araneae: Araneidae) in South Africa. *International Journal of Modern Biological Research* 1:21–34.
- Giribet G. 2015. Morphology should not be forgotten in the era of genomics - a phylogenetic perspective. *Zoologischer Anzeiger* 256: 96–103.
- Harvey MS, Hillyer MJ, Main BY, Moulds TA, Raven RJ, Rix MG, et al. 2018. Phylogenetic relationship of the Australasian open-holed trapdoor spiders (Araneae: Mygalomorphae: Nemesiidae: Anamirinae): multi-locus molecular analysis resolve the generic classification of highly diverse fauna. *Zoological Journal of the Linnean Society* 184: 407–452.
- Harvey MS, Main BY, Rix MG, Cooper SJB. 2015. Refugia within refugia: in situ speciation and conservation of threatened *Bertmainius* (Araneae: Migidae), a new genus of relictual trapdoor spider endemic to the mesic zone of south-western Australia. *Invertebrate Systematics* 29:511–553.
- Harvey MS, Rix MG, Hillyer MJ, Huey JA. 2020. The systematics and phylogenetic position of the troglobitic Australian spider genus *Trogloplura* (Araneae: Mygalomorphae), with new classification for Anamiridae. *Invertebrate Systematics* 34:799–822.
- Huber BA. 2004. The significance of copulatory structures in spider systematics. Pp. 89–100. In *Biosemiotek: Praktische Anwendung und Konsequenzen für die Einzelwissenschaften*. VWB, Berlin.
- Hupalo K, Karaouzas I, Mamos T, Grabowski M. 2020. Molecular data suggests multiple origins and diversification times of freshwater gammarids on the Aegean archipelago. *Nature Scientific Reports* 10: 19813.
- Keighery G. 1993. Plant communities. Pp. 43–54. In *Mountains of Mystery: a Natural History of the Stirling Range*. (Thomson C, Hall G, Friend G (eds)). Department of Conservation and Land Management.
- Leavitt DH, Starrett J, Westphal MF, Hedin MC. 2015. Multilocus sequence data reveal dozens of putative cryptic species in a radiation of endemic Californian mygalomorph spiders (Araneae, Mygalomorphae, Nemesiidae). *Molecular Phylogenetics and Evolution* 91:56–67.
- Main BY. 1982. Further studies on the systematics of Australian Diplurinae (Araneae: Mygalomorphae, Dipluridae): the taxonomic status of *Prothermacha* Simon and *Chenistonia tepperi* Hogg. *Australian Entomological Magazine* 8:83–88.
- Main BY. 1993. Spiders and other invertebrates. Pp. 117–125. In *Mountains of Mystery: a Natural History of the Stirling Range*. (Thomson C, Hall G, Friend G (eds)). Department of Conservation and Land Management.
- McWhinnie K, Parsons K. 2019. Shaping up? A direct comparison between 2D and low-cost 3D shape analysis using African cichlid mandibles. *Environmental Biology of Fishes* 102:927–938.
- Muster C, Michalik P. 2020. Cryptic diversity in ant-mimic *Micraria* spiders (Araneae, Gnaphosidae) and a tribute to early naturalists. *Zoologica Scripta*, 49:197–209.
- Opatova V, Hamilton CA, Hedin MC, Montes de Oca L, Král J, et al. 2020. Phylogenetic systematics and evolution of the spider infraorder Mygalomorphae using genomic scale data. *Systematic Biology* 69: 671–707.
- Openshaw G, D'Amore D, Vidal-García M, Keogh J. 2017. Combining geometric morphometric analyses of multiple 2D observation views improves interpretation of evolutionary allometry and shape diversification in monitor lizard (*Varanus*) crania. *Biological Journal of the Linnean Society* 120:539–552.
- Padial JM, Miralles A, De la Riva I, Vences M. 2010. The integrative future of taxonomy. *Frontiers in Zoology* 7:16.
- Pérez-Miles F, Perafán C. 2017. Behavior and biology of Mygalomorphae. Pp. 29–34. In *Behaviour and Ecology of Spiders, Contributions from the Neotropical Region*. (Viera C, Gonzaga MO (eds.)) Springer.
- R Development Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. Online at <https://www.R-project.org>

- Rasmussen B, Bengston S, Fletcher I, McNaughton N. 2002. Discoidal impressions and trace-like fossils more than 1200 million years old. *Science* 296:1112–1115.
- Raven RJ. 2000. Taxonomica Araneae I: Barychelidae, Theraphosidae, Nemesiidae and Dipluridae (Araneae). *Memoirs of the Queensland Museum* 45:569–575.
- Rix MG, Bain K, Main BY, Raven RJ, Austin AD, Cooper SJB, et al. 2017. Systematics of the spiny trapdoor spider of the genus *Cataxia* (Mygalomorphae: Idiopidae) from southwestern Australia: documenting a threatened fauna in a sky-island landscape. *Journal of Arachnology* 45:395–423.
- Rix MG, Edwards DL, Bryne M, Harvey MS, Joseph L, Roberts JD. 2015. Biogeography and speciation of terrestrial fauna in the south-western Australian biodiversity hotspot. *Biological Reviews* 90:762–793.
- Rohlf J. 2013. Relative warps. Ecology & Evolution, SUNY at Stony Brook.
- Rohlf J. 2017. tpsDIG2 Version 2.31. *Ecology & Evolution & Anthropology*, Stony Brook University.
- Rohlf J. 2021. tpsUtil. *Ecology & Evolution and Anthropology*, Stony Brook University.
- Rohlf J, Marcus L. 1993. A Revolution in Morphometrics. *Trends in Ecology and Evolution* 8:129–132.
- RStudio T. 2019. RStudio: Integrated Development for R. RStudio Inc., Boston, MA, <http://www.rstudio.com>.
- Ruane, S. 2015. Using geometric morphometrics for integrative taxonomy: an examination of head shapes of milksnakes (genus *Lampropeltis*). *Zoological Journal of the Linnean Society* 174:394–413.
- Satler JD, Carstens BC, Hedin MC. 2013. Multilocus species delimitation in a complex of morphologically conserved trapdoor spiders (Mygalomorphae, Antrodiaetidae, *Aliatypus*). *Systematics Biology* 62:805–823.
- Semple T, Peakall R, Tatarnic N. 2018. A comprehensive and user-friendly framework for 3D-data visualization in invertebrates and other organisms. *Journal of Morphology* 280:223–231.
- Sharma PP, Ballesteros JA, Santibáñez-López CE. 2021. What is an “Arachnid”? Consensus, consilience, and confirmation bias in the Phylogenetics of Chelicerata. *Diversity*, 13: 568. <https://doi.org/10.3390/d13110568>.
- Simmons LW 2014. Sexual selection and genital evolution. *Austral Entomology* 53:1–17.
- Simmons LW, Fitzpatrick JL. 2019. Female genitalia can evolve more rapidly and divergently than male genitalia. *Nature Communications* 10:1–8.
- Simon E. 1908. Araneae. Ire partie. Pp. 359–446. In Die Fauna Südwest-Australiens. Ergebnisse der Hamburger südwest-australischen Forschungsreise 1905. (Michaelson W, Hartmeyer R. (eds.)) Gustav Fischer, Jena.
- Stoffel MA, Nakagawa S, Schielzeth H. 2017. rptR: Repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution* 8:1639–1644.
- Teixeira M, Jr. Prates I, Nisa C, Silva-Martins N, Strüssmann C, Rodrigues M. 2016. Molecular data reveal spatial and temporal patterns of diversification and a cryptic new species of lowland *Stenocercus* Duméril & Bibron, 1837 (Squamata: Tropiduridae). *Molecular Phylogenetics and Evolution* 94:410–423.
- Torres RA, Atencia PL, Liria J. 2018. Morphometric variation in *Phrynus barbadensis* (Pocock, 1893) (Amblypygi: Phrinidae) from Colombia. *Revista de la Sociedad Entomológica Argentina* 77:18–23.
- Wheeler Q. 2018. Blank canvas: the case for descriptive taxonomy. *Integrative and Comparative Biology* 58:1118–1121.
- Wheeler Q. 2020. A taxonomic renaissance in three acts. *Megataxa* 1:4–8.
- Wilson JD, Zapata LV, Barone ML, Cotoras DD, Poy D, Ramírez MJ. 2021. Geometric morphometrics reveal sister species in sympatry and a cline in genital morphology in a ghost spider genus. *Zoologica Scripta* 50:1–15.
- World Spider Catalog. 2021. World Spider Catalog. Version 22.5. Natural History Museum Bern. Online at wsc.nmbe.ch, accessed on September 28 2022. doi:10.24436/2
- Zelditch M, Swiderski D, Sheets D, Fink W. 2004. Geometric Morphometrics for Biologists. Elsevier Academic Press.
- Ziegler A, Kunth M, Mueller S, Bock C, Pohmann R, Schröder L, et al. 2011. Application of magnetic resonance imaging in zoology. *Zoomorphology* 130:227–254.

Manuscript received 29 May 2022, revised 9 December 2022, accepted 11 December 2022.