

An Updated Key to the Genera of Caligidae (Copepoda: Siphonostomatoida)

Authors: Bernot, James P., Boxshall, Geoffrey A., Kihara, Terue Cristina, and Arbizu, Pedro Martínez

Source: Journal of Parasitology, 111(1): 10-25

Published By: American Society of Parasitologists

URL: https://doi.org/10.1645/24-97

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Published 20 January 2025

Contents and archives available through www.bioone.org or www.jstor.org



Journal of Parasitology

DOI: 10.1645/24-97

journal homepage: www.journalofparasitology.org

AN UPDATED KEY TO THE GENERA OF CALIGIDAE (COPEPODA: SIPHONOSTOMATOIDA)

James P. Bernot^{1,2}, Geoffrey A. Boxshall³, Terue Cristina Kihara⁴, and Pedro Martínez Arbizu⁵

¹ Department of Ecology and Evolutionary Biology, University of Connecticut, 75 N Eagleville Road, Unit 3043, Storrs, Connecticut 06269. ² Department of Invertebrate Zoology, Smithsonian National Museum of Natural History, 10th Street and Constitution Avenue NW, Washington, D.C. 20560.

³ Department of Life Sciences, Natural History Museum, London SW7 5BD, United Kingdom.

⁴ INES—Integrated Environmental Solutions GmbH, German Center for Marine Biodiversity Research, Senckenberg am Meer, 26382, Wilhelmshaven, Germany.

⁵ German Center for Marine Biodiversity Research, Senckenberg am Meer, 26382, Wilhelmshaven, Germany.

Correspondence should be sent to James P. Bernot (https://orcid.org/0000-0002-1769-8631) at: james.bernot@uconn.edu

KEY WORDS	ABSTRACT
Caligidae Sea lice Caligus Midias Markevichus Parasitology Confocal laser scanning microscopy Taxonomy	Members of the copepod family Caligidae are some of the most common parasites of marine fishes. There are 503 recognized species divided into about 30 genera, with 75% of species belonging to the 2 largest genera, <i>Caligus</i> and <i>Lepeophtheirus</i> . More than 30 caligid species are known to cause serious pathologic changes as ectoparasites in marine teleost aquaculture. This study was undertaken to provide a key to the valid genera of Caligidae, to examine their morphology in new detail with confocal laser scanning microscopy, and to review uncertainties concerning boundaries between several genera. There have been several substantial changes to the taxonomy of Caligidae over the last decade: <i>Metacaligus, Sciaenophilus,</i> and <i>Sinocaligus</i> have been synonymized with <i>Caligus</i> and the validity of <i>Midias</i> has been questioned. Here, we formally propose that <i>Midias</i> and <i>Markevichus</i> are junior subjective synonyms of <i>Caligus</i> . In total, we recognize 27 valid genera; provide a key to these genera; present confocal laser scanning micrographs for 25 of the genera, many of which have never been imaged before; and highlight helpful diagnostic features used in the key. We also discuss some concerns regarding the generic boundaries separating <i>Belizia, Caritus, Parapetalus, Parechetus, Pseudechetus,</i> and <i>Synestius</i> from <i>Caligus</i> and those separating <i>Anuretes, Pseudanuretes</i> , and <i>Mappates</i> from <i>Lepeophtheirus</i> .

Sea lice, members of the family Caligidae Burmeister, 1835, are ectoparasitic copepods found on most groups of marine fishes. Adult caligids typically inhabit the body surface, buccal cavity, or gills of their hosts. These parasites are abundant in natural ecosystems and can cause substantial pathogenesis in aquaculture (Mustafa et al., 2000). Caligids are of major economic significance because several species have emerged as serious pests in marine finfish aquaculture in coastal waters around the world, including cold temperate and subtropical environments (Johnson et al., 2004). These copepods are perhaps most well-known as problematic pathogens in salmon farms, where they are responsible for over \$1 billion in losses each year (Boxaspen et al., 2022). It is not uncommon for dozens or even hundreds of individuals to be found on a single fish, feeding on mucus, tissue, and blood and causing substantial pathologic changes (Johnson et al., 1996, 2004; Bravo, 2003; Treasurer and Bravo, 2011; Kodama et al., 2021; Rodger et al., 2022).

Caligids are characterized by the possession of a shield-shaped cephalothorax incorporating the first to third pedigerous somites

(i.e., leg-bearing body segments). The cephalothorax typically forms a suction cup, enhanced by a marginal membrane, closed off anteriorly by the paired frontal plates, and delimited posteriorly by the broad, apron-like third swimming legs (Ohtsuka et al., 2021). This sucker is used for host attachment, and the suction force can be rapidly generated and released for attachment or free movement as needed (Ohtsuka et al., 2021). However, there are several interesting and rather extreme deviations from this body plan, which are imaged here with confocal laser scanning microscopy (CLSM) and revealed in new detail.

The most recent key to the genera of Caligidae is from the detailed monographic study of Dojiri and Ho (2013), who considered 31 genera to be valid, but many recent taxonomic changes have occurred since. Dojiri and Ho (2013) synonymized many genera and noted that boundaries between several others were poorly defined. Since then, there have been several revisions, and the genera *Metacaligus* Thomsen, 1949, *Sciaenophilus* van Beneden, 1852, and *Sinocaligus* Shen, 1957 have all been synonymized



with *Caligus* Müller, 1785 (Özak et al., 2017, 2024; Boxshall and Barton, 2023). Although Dojiri and Ho (2013) treated *Midias* Wilson, 1911 as valid, we agree with Kabata (1979) and others in regarding *Midias* as a synonym of *Caligus*. We also formally recognize *Markevichus* Özdikmen, 2008 as a junior subjective synonym of *Caligus*. In total, we recognize 27 genera as valid, which collectively include 503 species. Over 75% of caligid species are members of the 2 largest genera, *Caligus* (276 species) and *Lepeophtheirus* von Nordmann, 1832 (124 species), followed by *Anuretes* Heller, 1865 with 21 species (Walter and Boxshall, 2024). The remaining 24 genera have fewer than 10 species each, and 10 genera are monotypic.

Here, we present a new key to genera of Caligidae supported by images generated with CLSM and macrophotography for nearly all valid genera. The goals of this study were to facilitate the identification of the 27 genera, to highlight the morphological diversity of Caligidae, and to discuss limitations in data supporting several taxa. Molecular phylogenetic analyses are needed to establish more stable and robust generic boundaries, but because appropriate specimens for sequencing are not available for most genera and species, a more formal systematic revision awaits the collection of suitable material for sequencing.

MATERIALS AND METHODS

Specimens of 25 of the 27 valid caligid genera were imaged; only Arrama Dojiri and Cressey, 1991 and Dartevellia Brian, 1939 were not examined. The following specimens were imaged with CLSM: Abasia platyrostris Pillai, 1963 (NHMUK 2017.173), Alanlewisia fallolunulus (Lewis, 1967) (NHMUK 2010.658), Alebion gracilis Wilson, 1905 (NHMUK 2010.974), Anchicaligus nautili Stebbing, 1900 (NHMUK 2010.979), Anuretes heckelii (Krøyer, 1863) (NHMUK 1979.592), Belizia brevicauda Cressey, 1990 (USNM 241670), Caligodes laciniatus (Krøyer, 1863) (NHMUK 2017.192), Caligus diaphanus von Nordmann, 1832 (NHMUK 1975.180), Caritus serratus Cressey, 1967 (USNM 120354), Echetus typicus Krøyer, 1864 (USNM 107877), Euryphorus brachypterus (Gerstaecker, 1853) (NHMUK 2014.637), Gloiopotes huttoni (Thomson, 1890) (from a "sailfish," locality unknown), Hermilius longicornis Bassett-Smith, 1898 (NHMUK 2017.334), Kabataella indica Prabha and Pillai, 1983 (USNM 268272), Lepeophtheirus pectoralis (Müller, 1776) (NHMUK 1975.482), Mappates plataxus Rangnekar, 1958 (NHMUK 2017.346), Paralebion elongatus Wilson, 1911 (NHMUK 1994.716), Parapetalus occidentalis Wilson, 1908 (NHMUK 1984.13), Parechetus carangis (Bassett-Smith, 1898) (NHMUK 1999.628), Pseudanuretes papernai Kabata and Deets, 1988 (USNM 231871), Pseudechetus fimbriatus Prabha and Pillai, 1979 (USNM 274345), Pupulina flores van Beneden (1892) (USNM 1087132), Synestius caliginus Steenstrup and Lütken, 1861 (NHMUK 1994.1212), and Tuxophorus caligodes Wilson, 1908 (NHMUK 1954.9.20.7).

For CLSM, specimens were stained overnight in a saturated solution of Congo Red in 100% ethanol, rinsed in changes of distilled water until no stain could be seen diffusing, and prepared as temporary mounts in a 50% solution of glycerin and distilled water on a glass slide under a coverslip. For thicker specimens, the coverslip was suspended above the specimen with pieces of paraffin or, for the largest specimens, with rubber washers glued to a slide to create a deep well. Specimens were examined with a Leica TCS SP5 equipped with a Leica DM5000 B upright microscope and the Leica Application Suite Advanced Fluorescence software LAS AF 2.2.1 (Leica, Wetzlar, Germany). We used a 561 nm excitation wavelength from a DPSS 10 mW 561 nm laser set at 80% power and collected the emitted fluorescence in 2 channels: 570–630 nm artificially colored green and 630–715 nm artificially colored red. A series of image stacks was collected, and the final images were obtained by maximum projection of the overlaid channels using the same Leica software or ImageJ v2.14. For some specimens, multiple stacks or multiple fields of view were needed to produce high-resolution images of the entire specimen. Multiple stacks were combined using the concatenate tool in ImageJ or manually using Adobe Photoshop v25.0, and multiple fields of view were combined using the ImageJ pairwise stitching plugin (Preibisch et al., 2009) or with Adobe Photoshop.

RESULTS

Systematics Caligidae Burmeister, 1835

Valid genera: Abasia Wilson, 1908, Alanlewisia Boxshall, 2008, Alebion Krøyer, 1863, Anchicaligus Stebbing, 1900, Anuretes Heller, 1865, Arrama Dojiri and Cressey, 1991, Avitocaligus Boxshall and Justine, 2005, Belizia Cressey, 1990, Caligodes Heller, 1865, Caligus Müller, 1785, Caritus Cressey, 1967, Dartevellia Brian, 1939, Echetus Krøyer, 1863, Euryphorus Milne Edwards, 1840, Gloiopotes Steenstrup and Lütken, 1861, Hermilius Heller, 1865, Kabataella Prabha and Pillai, 1983, Lepeophtheirus von Nordmann, 1832, Mappates Rangnekar, 1958, Paralebion Wilson, 1911, Parapetalus Steenstrup and Lütken, 1861, Parechetus Pillai, 1962, Pseudanuretes Yamaguti, 1936, Pseudechetus Prabha and Pillai, 1979, Pupulina van Beneden, 1892, Synestius Steenstrup and Lütken, 1861, and Tuxophorus Wilson, 1908.

Remarks *Midias* Wilson, 1911 is a junior subjective synonym of *Caligus* Müller, 1785

Before the landmark monograph of Kabata (1979), the genus Midias was classified in the family Euryphoridae. However, Kabata (1979) noted that Midias could not be validly retained in the Euryphoridae because it lacks the diagnostic dorsal plates on the fourth pedigerous somite. He stated that "except for the shape of the abdomen and the rather prominent fifth legs, it could become a typical member of the genus Caligus" (Kabata, 1979, p. 202). Dojiri and Ho (2013) considered the appendages of Midias nearly identical to those of Caligus but upheld the validity of Midias within the Caligidae (now incorporating the genera previously placed in the Euryphoridae) based on 4 characteristics: (1) the bulbous expansion of the first abdominal somite of the female, (2) the setiferous conical projection representing leg 5 of the female, (3) 2 large claw-like outer spines on the first exopodal segment of leg 3, and (4) the spinulose posterolateral processes on the anal somite. Of these characters, only the presence of spinulose posterolateral processes on the anal somite is unique. Even in that case, similar processes are present in Caligus formicoides Redkar, Rangnekar and Murti, 1949, but they are unornamented. As

detailed below, none of these characters is unique and we consider that they are likely autapomorphies.

Similar lateral expansions of the first abdominal somite of the female are found in several *Caligus* species, including *Caligus alepicolus* Boxshall, 2018, *C. formicoides, Caligus grandiabdominalis* Yamaguti, 1954, *Caligus kurochkini* Kazachenko, 1975, *Caligus parapetalopsis* Hameed and Pillai, 1973, and *Caligus seriolicolus* Boxshall, 2018. These species represent at least 3 different species groups within the genus: *bonito*-group (*C. grandiabdominalis*), *diaphanus*-group (*C. formicoides*), and *confusus*-group (*C. alepicolus, C. kurochkini, C. parapetalopsis*, and *C. seriolicolus*) (Boxshall, 2018), and we infer that such lateral expansions of the abdomen have arisen independently on several occasions within the Caligidae.

A projecting conical fifth leg is also found in several other species of *Caligus*, including: *Caligus cordiventris* Shiino, 1952, *Caligus parvilatus* Kim, 1998, *Caligus praecinctorius* Hayes, Justine and Boxshall, 2012, *Caligus sclerotinosus* Roubal, Armitage and Rohde, 1983, and *Caligus tylosuri* (Rangnekar, 1956). The possession of large conical fifth legs was once used to distinguish members of the genus *Dentigryps* Wilson, 1913 from *Lepeophtheirus*, but as noted by Dojiri and Ho (2013, p. 378), "the taxonomic value of the length of any appendage is highly questionable" and the generic boundary between *Dentigryps* and *Lepeophtheirus* was somewhat arbitrary. Dojiri and Ho (2013) agreed with Hewitt (1971) that *Dentigryps* should be treated as a junior synonym of *Lepeophtheirus*. We consider using the fifth leg to distinguish between *Midias* and *Caligus* to be equally suspect.

The character "presence of two large claw-like outer spines on the first exopodal segment of leg 3" (Dojiri and Ho, 2013, p. 275) requires clarification. Examination of *Midias lobodes* specimens in the collections of the NHM (NHMUK 2010.623-628) reveals that only the larger distal spine is a true armature element articulated at its base. The smaller proximal structure is a spinous outgrowth of the segment itself. Such spinous projections, often ornamented with strips of membrane, are present at the outer distal corner of the first exopodal segment of leg 3 in several *Caligus* species, including *Caligus mortis* Kensley, 1970, *Caligus pageti* Russell, 1925, *C. parvilatus*, and *Caligus saucius* Dojiri, 1989, although they do not curve across the surface of the segment.

In summary, the character states identified by Dojiri and Ho (2013) as diagnostic for Midias (when found in combination) are scattered through the morphospace occupied by Caligus. Given the variability exhibited within *Caligus*, the largest genus in the entire Copepoda, we consider that none of these characters is robust as a generic-level discriminant. We therefore follow Kabata (1979) and Boxshall and Halsey (2004) in treating Midias as a junior subjective synonym of Caligus. Furthermore, we consider that its type species, *Midias lobodes*, is probably a member of the Caligus confusus-group within Caligus because it shares most of the character states diagnostic for the species group, namely, leg 4 is 3-segmented and armed with I, I, III spines; leg 3 bears a rosette and rib on the apron, and the true outer spine on the exopod segment 1 is robust and strongly recurved; and the maxillule bears a small accessory tine. In addition, the distal antennulary segment is quite long, the plumose posterior setae on leg 1 are short, and the outer margin of leg 2 endopod segment 1 is strongly convex, which we note here are all additional states commonly exhibited by members of the *confusus*-group.

Markevichus Özdikmen, 2008 is a junior subjective synonym of *Caligus*

Markevichus is the replacement name proposed for the genus Caligopsis Markevitsch, 1940, established to accommodate a single new species, Caligopsis ponticus Markevitsch, 1940. As noted by Dojiri and Ho (2013), Markevitsch (1940) distinguished his new genus primarily on the absence of leg 4, and the genus currently remains monotypic (Walter and Boxshall, 2024). Kabata (1979) suggested that Caligus apodus (Brian, 1924) (as Pseudocaligus apodus) belongs to Markevichus (as Caligopsis) because this genus lacked leg 4. However, there is still uncertainty regarding the state of leg 4 in this species. A redescription of C. apodus (as P. apodus) from Tunisia included the presence of vestiges of leg 4 in both sexes (Ben-Hassine, 1983). Subsequently, Özak et al. (2013) re-examined 2 syntype females of P. apodus in the collections of the Museum National d'Histoire Naturelle in Paris and found that 1 female lacked any trace of leg 4 while the other carried a minute rod-like vestige, possibly representing leg 4, on 1 side of the body. Özak et al. (2013) also reported that newly collected material of both sexes of C. apodus from Turkish waters carried an indistinctly segmented leg 4 in the male, armed with a total of 4 setae, but lacked any obvious trace of leg 4 in the female. It seems possible that the state of the vestigial leg 4 is variable in C. apodus, especially in the female. The genus Pseudocaligus has been recognized as a junior subjective synonym of Caligus based on both morphological (Dojiri and Ho, 2013) and molecular (Freeman et al., 2013) evidence.

Dojiri and Ho (2013, p. 374) raised the key question concerning the validity of this genus, namely, "Even if leg 4 is absent, can *Caligopsis* be considered valid based solely on the absence of leg 4, a characteristic of questionable value?" They then noted that species formerly placed in *Pseudocaligus* form a gradation of progressive reduction of leg 4. This series culminates in the loss of leg 4, as described for *Caligopsis ponticus* by Markevitsch (1940).

Although Dojiri and Ho (2013) clearly articulated the argument against continuing to treat *Markevichus* as valid and they dealt with it in their chapter "Miscellaneous Genera" (which comprised the other genera they considered invalid), they did not formally place *Markevichus* in synonymy with *Caligus*. Here we propose to transfer *Caligopsis ponticus* Markevitsch, 1940 to *Caligus* as *Caligus ponticus* (Markevitsch, 1940) n. comb. and thus treat *Markevichus* as a junior subjective synonym of *Caligus*. We also consider *C. ponticus* (Markevitsch, 1940) n. comb. to be a nomen dubium because it is inadequately described and the types have been lost (Titar, pers. comm. cited by Dojiri and Ho, 2013).

Metacaligus Thomsen, 1949 is a junior subjective synonym of *Caligus*

Metacaligus was originally established by Thomsen (1949) as a subgenus of Caligus to accommodate a new species, Caligus (Metacaligus) uruguayensis Thomsen, 1949. It was raised to generic status by Ho and Bashirullah (1977). Until the recent paper by Özak et al. (2024), 5 species were treated as valid (Walter and Boxshall, 2024), namely, Metacaligus trichiuri (Krøyer, 1863) (as M. uruguayensis (Thomsen, 1949)), Metacaligus rufus (Wilson, 1908), Metacaligus unguidentatus (Rangnekar and Murti, 1950) (as Metacaligus hilsae (Shen, 1957)), Metacaligus latus Ho and Lin, 2002, and Metacaligus yucatanensis SuárezMorales, Kim and Escamilla, 2012. Dojiri and Ho (2013) treated *Metacaligus* as a valid genus and highlighted 4 diagnostic character states, the combination of which could be used to separate this genus from *Caligus*. These states were (1) the absence of a sternal furca; (2) the lack of accessory processes on spines 2 and 3 on the distal margin of the second exopodal segment of leg 1; (3) the 3 plumose setae on the posterior margin of the same segment are short (i.e., shorter than the segment); and (4) the presence of only 1 outer spine on the third exopodal segment of leg 2. Dojiri and Ho (2013) emphasized that all these character states can be found individually in particular species of *Caligus*, but the full combination is found in only species of *Metacaligus*.

Özak et al. (2024) considered the list of characters provided by Dojiri and Ho (2013) to support the validity of Metacaligus not to be robust. The sternal furca has been lost or extremely reduced in at least 10 species of Caligus, including members of several different species groups: for example, Caligus enormis (Caligus productus-group), Caligus vucatanensis (Caligus undulatus-group), and Caligus afurcatus. The reduced length of the 3 plumose setae on the posterior margin of the distal exopodal segment of leg 1 is found in about 20 other species of *Caligus*; it is typical of members of the C. confusus-group and is found in some other species such as Caligus paulini Nuñes-Ruivo and Fourmanoir, 1956, a member of the Caligus bonito-group (Ismail et al., 2024). The absence of accessory processes from spines 2 and 3 on the distal exopodal segment of leg 1 is found in over 40 species of *Caligus*, is typical of the C. diaphanus-group, and is found in some members of other species groups. The loss of the proximal outer spine of the third exopodal segment of leg 2 is rare in Caligus; however, it is lacking in Caligus chinglonglini Ohtsuka and Boxshall, 2019 and is reduced in size in members of the Caligus pseudorhombigroup and C. undulatus-group (Ohtsuka and Boxshall, 2019; Ohtsuka et al., 2020; Özak et al., 2024). In addition, Özak et al. (2024) demonstrated numerous similarities between M. yucatanensis and their 2 new species Caligus izmiriensis Özak, Kurt, Kamanli, Akbulut, Yanar and Boxshall, 2024, and Caligus sarosi Özak, Kurt, Kamanli, Akbulut, Yanar and Boxshall, 2024. These similarities include the loss of the outer margin spine on the second exopodal segment of leg 2, which is shared by C. sarosi and M. yucatanensis. In light of the numerous shared character states, Ozak et al. (2024) transferred Metacaligus yucatanensis to Caligus as C. yucatanensis and placed it in the C. undulatus-group established by Ohtsuka et al. (2020). Özak et al. (2024) also proposed to treat Metacaligus as a synonym of Caligus and recognized the 4 remaining species previously assigned to Metacaligus as a new species group, the Caligus trichiuri-group. We agree with that proposal.

Caligodes megacephalus Wilson, 1905 is a junior subjective synonym of *Caligodes laciniatus* (Krøyer, 1863)

Dojiri and Ho (2013) noted that these 2 species are very similar, and the main feature distinguishing them is the presence of a posteromedian process on the genital complex in *C. megacephalus*. We examined the holotype of *C. megacephalus* (USNM 6103) and confirmed that, as suggested by Dojiri and Ho (2013), this process was a misinterpretation by Wilson; the suspected process is simply a series of folds in the cuticle on the dorsal surface of the genital complex that intersect posteriorly (as drawn by Pillai, 1961, fig. 15A) rather than an extension of the body (as drawn by Wilson, 1905, fig. 187). The maxillipeds were missing on the type specimen, but we agree with Dojiri and Ho (2013) that the maxillipeds are similar in these 2 species, especially given the more detailed redescription by Pillai (1961). All other appendages were the same between these 2 species, including the unusual setation patterns on legs 1, 2, and 4. We therefore propose to treat *C. megacephalus* as a junior subjective synonym of *C. laciniatus*.

Key to Caligidae genera

1a.	Dorsal shield of cephalothorax typical, dorso-ven-
	trally flattened
	(e.g., Fig 1B-J, L, O-X)
1b.	Lateral margins of cephalothorax folded ventrally 22
	(e.g., Fig 2A-G)
2a.	Leg 1 endopod 2-segmented; Leg 3 endopod 3-seg-
	mented
	(Fig. 3A, B)
2b.	Leg 1 endopod unsegmented with reduced or vesti-
	gial setal elements: Leg 3 endopod with $<$ 3 segments 3
	(Fig. 3C, D)
39	Leg 1 endopod armed with 2 setae \pm setal vestige:
Ju.	Characteristic rounded lunule like projections
	extending enteriorly from frontal plates near base of
	extending anteriorry nom nomai plates near base of
	antennules Alanlewisia
21	(Fig. 1B, 4D)
30.	Leg I endopod with vestigial setae only or unarmed;
4	With true lunules or lacking lunules
4a.	Lunules present
41	(Fig. 4A-C)
4b.	Lunules absent
Sa.	Genital complex with long posterior processes
	(extending almost to, or beyond, caudal rami);
	Abdomen without plate-like lateral expansions
	(Fig. 5B-E)
5b.	Genital complex with plate-like lateral expansions
	extending posteriorly (at least one-third the length of
	the genital complex); Abdomen with plate-like lateral
	expansions 7
	(Fig. 5G, F)
5c.	Genital complex without long posterior processes;
	Abdomen without plate-like lateral expansions
6a.	Total of 6 processes present (4 on genital complex
	and 2 on abdomen) Pseudechetus
	(Fig. 5C, D)
6b.	Total of 4 processes present (all on genital complex).
	<i>Synestius</i>
	(Fig. 5E)
6c.	Total of 2 processes present (both on genital com-
	plex)
	(Fig. 5B)
7a.	With elongate "neck" present between cephalothorax
	and genital complex ("neck" longer than cephalotho-
	rax) Parochotus
	(Fig. 1S)
7h	Without elongate "neck" Paranetalus
, 0.	(Fig. 1R)



Figure 1. Caligid genera, dorsal view, confocal laser scanning microscopy (except F). (A) Abasia platyrostris. (B) Alanlewisia fallolunulus. (C) Alebion gracilis. (D) Anchicaligus nautili. (E) Anuretes heckelii. (F) Avitocaligus assurgericola (macrophotography, holotype MNHN Cp-2185). (G) Belizia brevicauda. (H) Caligodes laciniatus. (I) Caligus diaphanus. (J) Echetus typicus. (K) Euryphorus brachypterus; arrowheads indicate dorsal plates on fourth pediger. (L) Gloiopotes huttoni. (M) Hermilius longicornis. (N) Kabataella indica. (O) Lepeophtheirus pectoralis. (P) Mappates plataxus. (Q) Paralebion elongatus. (R) Parapetalus occidentalis. (S) Parechetus carangis. (T) Pseudanuretes papernai. (U) Pseudechetus fimbriatus. (V) Pupulina flores. (W) Synestius caliginus. (X) Tuxophorus caligodes; arrowheads indicate dorsal plates on fourth pediger. Color version available online.

- 8a. Fourth pedigerous somite with dorsal aliform plates. (Fig. 1X) 9a. Extremely elongate neck present (derived from fourth pedigerous somite and anterior part of genital complex)..... Echetus (Fig. 1J) 9b. Lacking extremely elongate neck 10 10a. Thoracic zone of cephalothorax extending posteriorly, completely concealing fourth pedigerous somite Belizia (Fig. 6F. G) 10b. Thoracic zone of cephalothorax not concealing fourth pedigerous somite 11 11a. Leg 3 endopod lacking setae Caritus 11b. Leg 3 endopod with 6 setae (or rarely 2–5 setae)...... 12 12a. Nauplius eye with enlarged conical lenses.... Anchicaligus (Fig. 1D) 12b. Nauplius eye with small lenses present, or (rarely), if lenses enlarged, lenses extending laterally Caligus (Fig. 1I) 13a. Abdomen expanded into large paired, flattened processes extending posteriorly, incorporating caudal rami Dartevellia 13b. Abdomen not expanded into flattened processes extending posteriorly; caudal rami (typically) distinct. 14 14a. Genital complex bearing long digitiform processes extending posteriorly Paralebion (Fig. 5A) 14b. Genital complex without digitiform posterior pro-15a. Dorsal cephalothoracic shield with main transverse supporting rib located posteriorly (almost at tips of lateral thoracic zones of cephalic shield); Posterior process of maxillule forming large rounded lobe Mappates (Fig. 6D) 15b. Dorsal cephalothoracic shield supported by Hshaped ribs with main transverse rib near mid-level of lateral thoracic zones of cephalic shield; Posterior process of maxillule tapering conical or absent...... 16 16a. First endopodal segment of leg 3 with inner plumose seta Lepeophtheirus (Fig. 10, 6C) 16b. First endopodal segment of leg 3 lacking inner plumose seta 17 17a. Antenna of female with accessory claw..... Pseudanuretes (Fig. 6B) 17b. Antenna of female without accessory claw Anuretes (Fig. 6A) 18a. Leg 4 biramous 19 18b. Leg 4 uniramous (lacking endopod) 20 19a. Fourth pedigerous somite lacking flattened dorsal plates; Genital complex with 2 pairs of large leaf-like plates overlying abdomen; Pair of similar, large leaf-
- like plates on abdomen Avitocaligus (Fig. 1F)

(Fig. 1K)

- 20a. Leg 4 reduced, 1-segmented; Some outer spines on exopods of legs modified, fan-like...... *Alebion* (Fig. 1C, 3B)

- 21b. Fourth pedigerous somite without dorsal plates; Leg 1 distal exopod segment with spines 2 and 3 at tip of segment simple; Leg 3 endopod segment 2 unmodified, lacking plate-like extension *Pupulina* (Fig. 1V)
- 22a. Leg 3 reduced to minute setiferous lobe, lacking intercoxal plate *Arrama*

- 23b. Antenna terminal segment lacking accessory claw; Leg 1 distal exopod segment bearing spines shorter than length of segment; Leg 1 distal exopod segment with reduced or vestigial setae on posterior margin 24
- 24a. Leg 4 with 4 segments Kabataella (Fig. 2F-H)

DISCUSSION

Here, we review questionable evidence supporting the validity of some caligid genera and speculate on potential evolutionary relationships based on new morphological interpretations. Boundaries between many genera are poorly defined and have often been further eroded by the discovery of species exhibiting intermediate combinations of character states. The few morphological features in caligids that have been examined in a molecular phylogenetic context have often been found to be homoplasious. For example, the segmentation of leg 4 was historically used to distinguish the genera *Markevichus, Pseudocaligus,* and *Pseudolepeophtheirus,* which possess a highly reduced leg 4 or lack leg 4 altogether. Although variability in the segmentation of leg 4 remains one of the most useful features for identifying species, molecular phylogenetic studies have revealed that leg 4 has been reduced to a single segment at least 3 times independently



Figure 2. Confocal laser scanning microscopy of caligid genera with ventrally folded cephalothorax (*Arrama* not shown). *Hermilius longicornis*: (A) dorsal, (B) ventral, (C) lateral. *Abasia platyrostris*: (D) dorsal, (E) ventral. *Kabataella indica*: (F) dorsal, (G) ventral, (H) leg 4 ventral with segments numbered. Color version available online.

(Freeman et al., 2013), providing support for existing synonymies of *Markevichus* and *Pseudocaligus* with *Caligus* and *Pseudolepeophtheirus* with *Lepeophtheirus*.

The presence or absence of the sternal furca is also patchy throughout the Caligidae, and we consider it very likely that this structure also exhibits a high degree of homoplasy. This situation is analogous to that in the genus *Dissonus* Wilson, 1906 (family Dissonidae), the species of which may possess a typical sternal furca or a modified sternal stylet or may lack the sternal furca altogether

(Boxshall et al., 2008). Given that what little we know about morphological evolution in caligids suggests there are many instances of convergent evolution, in the interest of taxonomic stability we have avoided making large taxonomic changes until we have evidence from molecular phylogenetic analysis to support or refute clades. We caution that taxonomic changes in this diverse and globally important group should not be made without robust support.

Perhaps the most problematic set of generic distinctions relates to *Anuretes*, *Pseudanuretes*, *Mappates*, and *Lepeophtheirus*. These



Figure 3. Confocal laser scanning microscopy for comparison of legs 1 and 3. (A) Leg 1 with 2-segmented endopod; *Alebion gracilis*, arrowheads indicate segments. (B) Leg 3 with 3-segmented endopod in *A. gracilis* with segments numbered, arrowhead indicates 1 of many modified spines characteristic of *Alebion*. (C) Leg 1 with unsegmented endopod in *Caligus hirsutus*, endopod indicated with arrowhead. (D) Leg 3 with 2-segmented endopod in *Caligus epidemicus* with segments numbered. Color version available online.

genera are traditionally distinguished from each other primarily by the extent of the elongation of the free margin of the thoracic zone of the cephalothorax in the female; however, there is significant overlap in this feature between the genera (e.g., Fig. 6A–D). Dojiri and Ho (2013) included a detailed discussion of this history and evidence, or lack thereof, supporting their separation. These genera already have a complex taxonomic history given that *Anuretes* has already been synonymized with *Lepeophtheirus* and then resurrected (Ho and Dojiri, 1977; Dojiri and Ho, 2013) and that 3 of the 9 currently accepted species of *Pseudanuretes* (Walter and Boxshall, 2024) were considered as questionable by Dojiri and Ho (2013) because their inclusion would require broadening of the generic boundaries. Indeed, it is challenging to key out these genera because virtually every feature used to distinguish *Anuretes* is also found in some species of *Lepeophtheirus* or *Pseudanuretes*. Because there are no molecular data for any species of *Anuretes*, *Pseudanuretes*, or *Mappates*, we have elected to retain these 3 genera until a molecular phylogenetic analysis can better address the boundaries separating them.

Ten genera appear closely related to *Caligus* based on morphology: *Anchicaligus, Belizia, Caligodes, Caritus, Echetus, Parapetalus, Parechetus, Pseudechetus, Synestius,* and *Tuxophorus.* Despite the



Figure 4. Confocal laser scanning microscopy of lunules and lunule-like structures in the Caligidae. (A) *Caligus schlegeli* exhibits relatively large lunules. (B) *Caligus confusus* exhibits typical lunules. (C) *Caligus furcisetifer* has the smallest lunules in *Caligus*. (D) *Alanlewisia fallolunulus* female possesses lunule-like structures that are not true lunules. Color version available online.

occasional dramatic elongations or lateral extensions of body somites (e.g., "necks," waists, and foliaceous processes on the abdomen or genital complex), the appendage structure across these genera is remarkably conserved. They share the same segmentation on all appendages except for leg 4 (there are also rare fusions of the terminal 2 exopodal segments of leg 3 in a few species of *Caligus*) and share the same unique evolutionary novelty—the paired lunules on the frontal plate of the dorsal cephalothoracic shield (Kaji et al., 2012). Essentially all these genera are erected based on a difference in body somites relative to *Caligus*. The following 6 genera are monotypic and based on only autapomorphies relative to *Caligus: Anchicaligus, Belizia, Echetus, Parechetus, Pseudechetus*, and *Synestius*. The robustness of all 10 of these genera awaits to be tested in a molecular phylogenetic context. Regardless of their validity, these taxa are very interesting from an evolutionary perspective, and resolution of their phylogenetic relationships will give insights into the evolution of some remarkable body structures (e.g., "necks," foliaceous processes, and enlarged lenses in the nauplius eye), attachment methods and sites (e.g., specialization for attachment to the operculum or to the wall of the buccal cavity by *Echetus, Parapetalus, Parechetus,* and *Pseudechetus,* with a mesoparasitic style in *Echetus*), and host associations (e.g., the colonization of *Nautilus* Linnaeus, 1758 by *Anchicaligus*). We suspect that the divergent morphology of at least some of these taxa is not the result of a unique phylogenetic history but that they are terminal branches of larger clades within



Figure 5. Confocal laser scanning microscopy of posterior processes on caligid genera. (A) Paralebion elongatus. (B) Caligodes laciniatus. (C) Pseudechetus fimbriatus. (D) Pseudechetus fimbriatus ventral. (E) Synestius caliginus. (F) Parapetalus occidentalis. (G) Parechetus carangis. Abbreviations: GC, processes on genital complex; Abd, processes on abdomen. Color version available online.



Figure 6. Confocal laser scanning microscopy of caligid genera with dorsal expansion of cephalothoracic shield partially covering pediger 4. (A) Anuretes heckelii. (B) Pseudanuretes papernai. (C) Lepeophtheirus atypicus. (D) Mappates plataxus. (E) Caligus epidemicus. (F) Belizia brevicauda. (G) Belizia brevicauda, ventral view. Color version available online.

Caligus, and their divergent morphology may be linked to changes in hosts and/or attachment sites.

The distinction between *Belizia* and *Caligus* is similar to that separating *Anuretes*, *Pseudanuretes*, and *Mappates* from *Lepeoph*-*theirus* in that *Belizia* is distinguished primarily by the extension of the posterior margin of the thoracic zone of the cephalothorax so that it overlaps and conceals the fourth pedigerous somite in dorsal view (Fig. 6F). Cressey (1990) and Dojiri and Ho (2013) further distinguished *Belizia* based on the presence of only 4 setae on the caudal ramus, the sternal furca being reduced to 2 sclerotized knobs (Fig. 6G), the presence of 4 setae on the terminal endopodal segment of leg 3, and the velum not distinctly

separated from the first endopodal segment of leg 3. However, we do not find these derived differences particularly robust. The sternal furca is reduced or absent in at least 10 species of *Caligus*. Although most species of *Caligus* have 6 setae on the terminal endopodal segment of leg 3, several species have 5, *Caligus longipes* (Moon and Kim, 2012) and *C. laciniatus* have 4, and *Caligus uniartus* (Ho, Kim, Cruz-Lacierda and Nagasawa, 2004) has 3 (Ho et al., 2004; Moon and Kim, 2012). The velum of leg 3 is not much different from that of most species of *Caligus* except that it is a bit smaller; this is not so surprising given that *Belizia*, at 1.6 mm body length, is 1 of the smallest caligids and has a small leg 3. We suspect molecular phylogenetic analyses will show that

Belizia is a morphologically unusual species of *Caligus*, but we retain it as a valid genus at present.

Caritus also closely resembles Caligus. Caritus serratus, the type species, was redescribed from type material by Dojiri and Ho (2013) and was distinguished from Caligus by the absence of a posterior process on the proximal antennal segment, the absence of the postantennal process, the absence of the sternal furca, the possession of a mandible with an elongate terminal section, the possession of modified exopodal spines on leg 2, the absence of setae on the terminal endopodal segment of leg 3, and the relatively large lobe-like leg 5. This latter character is not particularly robust because several species of *Caligus* and *Lepeoph*theirus have an enlarged lobe-like leg 5, as discussed in detail in the remarks on Midias, above. Furthermore, numerous species of genera such as Abasia and Caligus lack a posterior process on the proximal antennal segment, a postantennal process, or sternal furca and occasionally lack 2 of these 3 features. In fact, the antennal posterior process is very reduced in *Echetus typicus*, and this species also lacks a postantennal process and sternal furca. Although no species of Caligus lacks all setae on the leg 3 terminal endopodal segment, several species have fewer than 6, and C. uniartus has only 3. Caritus tolii Rangnekar, 1984 is relevant to this discussion. Although poorly described, it appears to have the typical complement of 6 setae on the terminal endopodal segment of leg 3, and it appears to lack the large lobe-like leg 5 of the type species (Rangnekar, 1984), further eroding the distinction between Caritus and Caligus. If C. tolii truly lacks the posterior process on the proximal segment of the antenna and lacks the postantennal process, the sternal furca, and the lobe-like leg 5, that would leave the genus distinguished by only modified spines on the exopodal segments of leg 2. We consider that Caritus tolii should be placed in *Caligus*, as proposed by Morales-Serna et al. (2024). Indeed, based on these morphological interpretations, we infer that the 2 species of *Caritus* are not sister taxa and that both are nested within *Caligus*, but this hypothesis requires testing with molecular data.

Interestingly, *Midias* (=*Caligus*), *Parapetalus*, *Parechetus*, Pseudechetus, and Synestius share a suite of characters not just with Caligus but specifically with the Caligus confusus-group. With the addition of C. lobodes here, there are now 25 species in the *confusus*-group (Boxshall, 2018), and the group is defined by the following set of characters: antenna with posterior process on proximal segment (process typically spatulate); accessory tine present on postantennal process; maxillulary posterior process bifid; apron of leg 3 with raised cuticular rib and circular array of large denticles (also known as a rosette); outer margin spine on first exopodal segment of leg 3 stout and strongly recurved; leg 4 with 4 segments and with terminal 3 segments bearing spines I, I, III (Boxshall, 2018). In addition, the distal segment of the antennule is nearly always elongated, the setae on the posterior margin of the exopod of leg 1 are usually somewhat reduced in size, and members of the *confusus*-group typically parasitize species of the Carangidae. Although species assigned to the *confusus*-group typically possess this entire suite of character states, some species lack 1 or more of them.

We find it remarkable that many of these *confusus*-group character states are shared with *Parapetalus*, *Parechetus*, *Pseudechetus*, and *Synestius*. These genera are distinguished by superficially obvious differences involving processes and expansions on the abdomen or genital complex (Fig. 5C–G). We suspect that these gross differences in body form, currently used as generic discriminants, may not be robust with respect to phylogeny, especially given the larger suite of shared characters detailed below. We have synonymized *Midias* with *Caligus* here but have maintained the other genera until their phylogenetic relationships can be more formally evaluated with molecular data.

Of these 4 *confusus*-like genera, *Pseudechetus* most clearly resembles the *C. confusus*-group. The sole species, *Pseudechetus fimbriatus*, is distinctive because it possesses an elongated waist (also referred to as a neck) between the fourth pedigerous somite and the genital complex and has 6 elongate posterior processes: 4 on the genital complex and 2 on the abdomen (Fig. 5D). Despite these differences in the overall appearance of the body, *P. fimbriatus* possesses all of the *confusus*-group characters listed above, is known from only carangid hosts, and further resembles *C. confusus* itself in that it possesses a postantennal process that has 2 accessory processes (a posterior tine and an additional anterior-lateral tine) rather than the typical single accessory process (Fig. 5D).

Parechetus is easily identified by its possession of an elongated waist and single pair of long processes on the genital complex and abdomen (Fig. 5G), but it also possesses many of the *confusus*-group features. As redescribed by Pillai (1985), the type species *P. carangis* has a bifid postantennal process, a leg 3 protopod with a small rosette, a leg 3 exopod with a stout recurved spine, and a leg 4 typical of the *confusus*-group. It also possesses a long antennule distal segment and has reduced setae on the exopod of leg 1 (in this case, reduced to a single seta), plus it is known from only the carangid *Carangoides ferdau*. Its possession of the remaining *confusus*-group characters cannot be confirmed because Pillai (1985) did not figure the maxillule or the rib on the apron of leg 3.

The status of *Parechetus constrictus* Kirtisinghe, 1964 is uncertain. It is currently treated as a synonym of *P. carangis* (Walter and Boxshall, 2024), but Pillai (1985) did not formally propose a synonymy. He suggested only that the constricted neck in Kirtisinghe's *P. constrictus* might be an artifact, and if so they would likely be synonyms (Pillai, 1985). Dojiri and Ho (2013) also considered that these species were likely synonyms. However, the fact that the NMNH specimen of *P. fimbriatus* imaged here (Fig. 5C) has a constriction in just the same area of the "neck" indicates that this character might not be an artifact. Pillai (1985) specifically stated that he could not synonymize the 2 taxa because he had not looked at Kirtisinghe's types, so both should be treated as valid until further review.

The single species of *Synestius, S. caliginus,* possesses 4 elongate posterior processes on the genital complex (Fig. 5E) but also exhibits many *confusus*-group features. The distal segment of the antennule is elongate; the postantennal process has an accessory process; leg 1 has reduced setae on the posterior margin of the exopod; leg 3 bears a rosette, rib, and robustly recurved exopodal spine; and leg 4 is of the typical form (Dojiri and Ho, 2013). Beyond the 4 large posterior processes on the genital complex, *S. caliginus* differs from typical *confusus*-group species in several notable ways: the postantennal process has an accessory process that is a anterio-lateral knob rather than the typical posteriorly directed pointed process; the maxillulary process is simple rather than bifid and is less elongate; and it is known to parasitize hosts belonging to the families Stromateidae and Lutjanidae but not Carangidae (Walter and Boxshall, 2024).

The situation in *Parapetalus* is more complicated than that of the previous 3 genera because almost since its inception it has comprised a heterogenous group of species loosely united by degrees of expansion of the lateral margins of the abdomen. In the last decade, 7 species previously assigned to this genus have been transferred to Caligus (Boxshall, 2018; Boxshall and Barton, 2023; Walter and Boxshall, 2024). The 4 remaining species are Parapetalus dewani Pillai and Hameed, 1981, Parapetalus longipennatus Rangnekar, 1956, Parapetalus occidentalis, and the type species Parapetalus orientalis Steenstrup and Lütken, 1861. Each possesses a different number of confusus-group characters. Para*petalus longipennatus* exhibits the most: it has an antennule with an elongate distal segment; an antenna with a broadening posterior process; a post-antennal process bearing an accessory process; a bifid maxillulary process; a leg 3 bearing a rib, rosette, and exopod armed with a robust recurved spine; reduced setae on leg 1; and is so far known only to parasitize carangid hosts (Pillai, 1985; Walter and Boxshall, 2024). Parapetalus dewani exhibits the next largest suite of confusus-group characters: long antennular distal segment; postantennal process with an accessory tine (although the maxillulary process is not bifid); leg 3 bearing a rosette and robust recurved exopodal spine; and reduced setae on leg 1 (Pillai and Hameed, 1981); it is unclear from available descriptions whether this species possesses a rib on the apron of leg 3. Parapetalus dewani is known from only Rachycentron canadum (Rachvcentridae), and although this host is not in the Carangidae it is a member of the order Carangiformes.

Parapetalus occidentalis and P. orientalis share a more reduced suite of *confusus*-group features. They lack an accessory process on the postantennal process, and they have a simple rather than bifid maxillulary process, but they are reminiscent of the confusus-group in possessing an elongate distal antennulary segment. reduced setae on the exopod of leg 1, a rib on leg 3, a strongly recurved or oblique spine on the exopod of leg 3, and a leg 4 typical of the group (Ho and Lin, 2004; Dojiri and Ho, 2013). Parapetalus occidentalis parasitizes R. canadum, and P. orientalis parasitizes carangids. Given the suite of shared characters, it is tempting to transfer the remaining species of Parapetalus to Caligus. However, in the interest of taxonomic stability, until the diagnostic utility of the various *confusus*-group characters can be evaluated in a phylogenetic context or the evolutionary relationships of these species can be directly tested, we have elected to retain these species in *Parapetalus* given their unusual body plans and mosaic of character states.

The variety of posterior foliaceous and digitiform processes among species of Parapetalus, Parechetus, Pseudechetus, and Syn*estius* (Fig. 5C–G) raises the question: what is their function? Parasites in general, and indeed many caligids, tend to have enlarged reproductive systems that increase their egg production capacity, but the processes on the genital complex of these genera are very flattened and do not appear to contain reproductive tissue. These posteriorly directed processes may serve to protect egg strings, perhaps from the activity of cleaner fish, which has been suggested for similar posterior processes in other groups of copepods such as the Lernanthropidae (Boxshall et al., 2020). All of these genera are relatively large caligids, so the expansions and processes may contribute additional surface area for respiration, but species of Pupulina and Gloiopotes and even Caligus curtus are substantially larger and lack such processes. These 4 genera have modified attachment sites relative to most caligids and live on the interior of the operculum or in the buccal cavity rather than on the body surface or gills. These regions may be more sheltered from shearing forces, which is particularly relevant because these copepods tend to parasitize rapidly swimming fishes. Under reduced shearing forces, the bodies of these copepods may have a broader adaptive landscape. Lastly, it is interesting to speculate that there may be a developmental element to this process. Many body parts in the confusus-group have additional processes or extensions: the posterior process of the maxillule and the postantennal process both tend to have accessory tines, and the rib on the apron of leg 3 often has a bifid tip. In Pseudechetus, the sternal furca is also bifid. Could there be a signaling pathway that leads to the development of additional axes of growth such as those listed above, and if so might this same process be involved in forming the foliaceous processes on the genital complex and abdomen in Parapetalus, Parechetus, Pseudechetus, and Synestius?

ACKNOWLEDGMENTS

We are grateful to the staff of the Natural History Museum, London and the Smithsonian National Museum of Natural History for loans of specimens used for this study. This material is based in part upon work supported by the NSF Postdoctoral Research Fellowships in Biology Program under grant 2010898 to JPB. This is publication number 72 which uses data from the Senckenberg am Meer Confocal Laser Scanning Microscope Facility.

LITERATURE CITED

- BASSETT-SMITH, P. W. 1898. Some new or rare parasitic copepods found on fish in the Indo-Tropic region. Annals and Magazine of Natural History 2: 357–372.
- BEN-HASSINE, O. K. 1983. Les Copépodes parasites de Poissons Mugilidae en Mediterranée occidentale (côtes françaises et tunisiennes). Ph.D. thesis. Université des Sciences et Techniques du Languedoc, Montpellier, France, 452 p.
- BOXASPEN, K. K., Ø. KARLSEN, T. SVÅSAND, AND L. ASPLIN. 2022. Impacts of sea lice. *In* Sea Lice Biology and Control, J. Treasurer, I. Bricknell, and J. Bron (eds.). Kings, Great Easton, Essex, U.K., p. 531–547.
- BOXSHALL, G. 2018. The sea lice (Copepoda: Caligidae) of Moreton Bay (Queensland, Australia), with descriptions of thirteen new species. Zootaxa 4398: 1–172.
- BOXSHALL, G. A. 2008. A new genus of sea louse (Copepoda: Siphonostomatoida: Caligidae) parasitic on the bluespine unicornfish (*Naso unicornis*). Folia Parasitologica 55: 231–240.
- BOXSHALL, G. A., AND D. P. BARTON. 2023. Caligid sea lice (Copepoda: Caligidae) from golden snapper *Lutjanus johnii* (Bloch) in Australian waters, with the recognition of *Sinocali*gus Shen, 1957 as a junior synonym of *Caligus* Müller, 1785. Systematic Parasitology 100: 487–504.
- BOXSHALL, G. A., J. P. BERNOT, D. P. BARTON, B. K. DIGGLES, Q. RUSSELL, T. ATKINSON-COYLE, AND K. S. HUTSON. 2020. Parasitic copepods of the family Lernanthropidae Kabata, 1979 (Copepoda: Siphonostomatoida) from Australian fishes, with descriptions of seven new species. Zootaxa 4736: 1–103.
- BOXSHALL, G. A., AND S. H. HALSEY. 2004. An Introduction to Copepod Diversity. Ray Society, London, U.K., 2000 p.

- BOXSHALL, G. A., AND J. L. JUSTINE. 2005. A new genus of parasitic copepod (Siphonostomatoida: Caligidae) from the razorback scabbardsh, *Assurger anzac* (Trichiuridae) off New Caledonia. Folia Parasitologica 52: 349–358. doi:10.14411/fp. 2005.048.
- BOXSHALL, G. A., C. L. LIN, J. S. HO, S. OHTSUKA, B. A. VENMATHI MARAN, AND J. L. JUSTINE. 2008. A revision of the family Dissonidae Kurtz, 1924 (Copepoda: Siphonostomatoida). Systematic Parasitology 70: 81–106.
- BRAVO, S. 2003. Sea lice in Chilean salmon farms. Bulletin of the European Association of Fish Pathologists 23: 197–200.
- BRIAN, A. 1924. Copepoda. Copépodes commensaux et parasites des cotes mauritaniennes. Parasitologica Mauritanica. Matériaux pour la Faune Parasitologique en Mauritanie 1: 1–66.
- BRIAN, A. 1939. Copépodes parasites recueillis par ME Dartevelle à l'embouchure du fleuve Congo. Revue de Zoologie et de Botanique Africaines 32: 176–198.
- BURMEISTER, H. 1835. Beschreibung einiger neuen oder weniger bekannten Schmarotzerkrebse, nebst allgemeinen Betrachtungen uber die Gruppe, welche sie angehoren. Acta Verhandlungen der Leopoldinisch-Carolinischen Akademie der Naturforscher 17: 269–336.
- CRESSEY, R. 1990. Belizia brevicauda, a new genus and species of caligid copepod from the western Caribbean Sea. Systematic Parasitology 15: 151–154.
- CRESSEY, R. F. 1967. *Caritus*, a new genus of caligoid copepod, with a key to the genera of Caliginae. Proceedings of the United States National Museum. 123: 1–8.
- DOJIRI, M. 1989. Two species of *Caligus* (Copepoda: Siphonostomatoida) parasitic on fishes from southern Africa. Journal of Natural History 23: 263–274.
- DOJIRI, M., AND R. F. CRESSEY. 1991. Arrama, new genus (Siphonostomatoida: Caligidae), with two new species, copepods parasitic on Australian fishes. Journal of Crustacean Biology 11: 594–606.
- DOJIRI, M., AND J.-S. Ho. 2013. Systematics of the Caligidae, Copepods Parasitic on Marine Fishes. Brill, Leiden, The Netherlands, 461 p.
- FREEMAN, M. A., H. ANSHARY, AND K. OGAWA. 2013. Multiple gene analyses of caligid copepods indicate that the reduction of a thoracic appendage in *Pseudocaligus* represents convergent evolution. Parasites & Vectors 6: 336. doi:10.1186/1756-3305-6-336.
- GERSTAECKER, A. 1853. Über eine neue und eine weniger gekannte Siphonostomen-Gattung. Archiv für Naturgeschichte 19: 58–70.
- HAMEED, S. M., AND N. K. PILLAI. 1973. Description of a new species of *Caligus* (Crustacea: Copepoda) from Kerala. Zoological Anzeiger 191: 114–118.
- HAYES, P., J. L. JUSTINE, AND G. A. BOXSHALL. 2012. The genus *Caligus* Müller, 1785 (Copepoda: Siphonostomatoida): Two new species from reef associated fishes in New Caledonia, and some nomenclatural problems resolved. Zootaxa 3534: 21–39.
- HELLER, C. 1865. Crustaceen. In Reise der österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von Wüllerstorf-Urbair. Zoologie 2: 1–280.
- HEWITT, G. C. 1971. Species of *Lepeophtheirus* (Copepoda, Caligidae) recorded from the ocean sunfish (*Mola mola*) and their implications for the caligid genus *Dentigryps*. Journal of the Fisheries Research Board of Canada 28: 323–334.

- Ho, J.-S., AND A. K. M. BASHIRULLAH. 1977. Two species of caligid copepods (Crustacea) parasitic on marine fishes of Venezuela, with discussion of *Metacaligus* Thomsen, 1949. Journal of Natural History 11: 703–714.
- Ho, J.-S., AND M. DOJIRI. 1977. Parasitic copepods on the fishes of the Great Barrier Reef, Australia. Part II. Caligoida: *Dissonus, Lepeophtheirus*, and *Dentigryps*. Publications of the Seto Marine Biological Laboratory 24: 77–97.
- Ho, J.-S., I. H. KIM, E. R. CRUZ-LACIERDA, AND K. NAGASAWA. 2004. Sea lice (Copepoda, Caligidae) parasitic on marine cultured and wild fishes of the Philippines. Journal of the Fisheries Society of Taiwan 31: 235–249.
- Ho, J.-S. AND C. L. LIN. 2002. New species of *Metacaligus* (Caligidae, Copepoda) parasitic on the cutlassfish (*Trichiurus lepturus*) of Taiwan, with a cladistic analysis of the family Caligidae. Zoological Sciences 19: 1363–1376.
- Ho, J.-S., AND C.-L. LIN. 2004. Sea Lice of Taiwan: Copepoda, Siphonostomatoida, Caligidae. Sueichan Press, Keelung, Taiwan, 388 p.
- ISMAIL, N., Y. NISHIDA, S. OHTSUKA, G. BOXSHALL, AND J. P. BERNOT. 2024. First record of *Caligus dussumieri* Rangnekar, 1957 (Copepoda, Siphonostomatoida, Caligidae) from Malaysia, with notes on caligids found from Malaysia and on hostspecificity of caligids on lutjanid fishes. Biodiversity Data Journal 12: e116598. doi:10.3897/BDJ.12.e116598.
- JOHNSON, S. C., R. B. BLAYLOCK, J. ELPHICK, AND K. D. HYATT. 1996. Disease induced by the sea louse (*Lepeophtheirus salmonis*) (Copepoda: Caligidae) in wild sockeye salmon (*Oncorhynchus nerka*) stocks of Alberni Inlet, British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 53: 2888–2897.
- JOHNSON, S. C., S. BRAVO, K. NAGASAWA, Z. KABATA, J. HWANG, J.-S. HO, AND C. T. SHIH. 2004. A review of the impact of parasitic copepods on marine aquaculture. Zoological Studies 43: 229–243.
- KABATA, Z. 1979. Parasitic Copepods of British Fishes. The Ray Society, London, U.K., 468 p.
- KABATA, Z. AND G. B. DEETS. 1988. Pseudoanuretes papernai sp. nov. (Copepoda: Caligidae) with comments on the genus Pseudanuretes Yamaguti, 1936. Canadian Journal of Zoology 66: 680–684.
- KAJI, T., B. A. VENMATHI MARAN, Y. KONDOH, S. OHTSUKA, G. A. BOXSHALL, AND A. TSUKAGOSHI. 2012. The lunule of caligid copepods: An evolutionarily novel structure. Evolution & Development 14: 465–475.
- KAZACHENKO, V. N. 1975. New species of the genus *Caligus* Müller, 1785 (Copepoda parasitica, Caligidae) from *Usacar*anx georgianus (Cuvier et Valenciennes) from the Great Australian Gulf. Parazitologiia 9: 425–431.
- KENSLEY, B. 1970. A new species of *Caligus* from South West Africa (Copepoda, Caligidae). Crustaceana 18: 167–172.
- KIM, I.-H. 1998. Caligus parvilatus n. nov., a replaced name of Caligus latus Kim, 1995 (Crustacea, Copepoda, Caligidae), with description of male. Animal Systematics, Evolution and Diversity 14: 117–120.
- KIRTISINGHE, P. 1964. A review of the parasitic copepods of fish recorded from Ceylon, with descriptions of additional forms. Bulletin of the Fisheries Research Station, Ceylon 17: 45–132.
- KODAMA, T., K. IKEDA, O. ARAKAWA, Y. KONDO, M. ASAKAWA, K. KAWATSU, AND S. OHTSUKA. 2021. Evidence of accumulation of

tetrodotoxin (TTX) in tissues and body parts of ectoparasitic copepods via their feeding on mucus of TTX-bearing puffer-fish. Toxicon 204: 37–43.

- KRØYER, H. 1863. Bidrag til Kundskab om Snyltekrebsene. Naturhistorisk Tidsskrift Series 3: 75–320.
- KRØYER, H. 1864. Bidrag til Kundskab om Snyltekrebsene. Naturhistorisk Tidsskrift Series 3: 321–426.
- LEWIS, A. G. 1967. Copepod crustaceans parasitic on teleost fishes of the Hawaiian islands. Proceedings of the United States National Museum 121: 1–204.
- MARKEVITSCH, A. P. 1940. Novi vidi parazitichnykh Copepoda. Dopovidi Akademiyi Nauk, Ukrayin'skoi RSR 11: 13–21.
- MILNE EDWARDS, H. 1840. Histoire Naturelle des Crustacés, Comprenant l'Anatomie, la Physiologie et la Classification de ces Animaux. Encyclopédique, Volume III. Roret, Paris, France, 638 p.
- MOON, S. Y., AND I. H. KIM. 2012. Sea lice (Copepoda, Siphonostomatoida, Caligidae) new to Korea, including three new species. Journal of Species Research 1: 175–217.
- MORALES-SERNA, F. N., B. YÁÑEZ-RIVERA, AND S. GÓMEZ. 2024. Caligus selenecola sp. nov. (Siphonostomatoida, Caligidae) parasitic on the hairfin lookdown Selene brevoortii (Gill) from the Gulf of California, Mexico, with some comments on Caritus tolii Rangnekar, 1984. ZooKeys 1209: 43–68. doi:10.3897/ zookeys.1209.120812.
- MÜLLER, O. F. 1776. Zoologiae Danicae prodromus, seu Animalium Daniae et Norvegiae indigenarum: characteres, nomina, et synonyma imprimis popularium. Hallagerii, Copenhagen, Denmark, 282 p.
- MÜLLER, O. F. 1785. Entomostraca seu Insecta Testacea quae ni aquis Daniae et Norvegia reperit. F. W. Thiele, Copenhagen, Denmark, 134 p.
- MUSTAFA, A., C. MACWILLIAMS, N. FERNANDEZ, K. MATCHETT, G. A. CONBOY, AND J. F. BURKA. 2000. Effects of sea lice (*Lep-eophtheirus salmonis* Kröyer, 1837) infestation on macrophage functions in Atlantic salmon (*Salmo salar* L.). Fish & Shellfish Immunology 10: 47–59.
- NUÑES-RUIVO, L., AND P. FOURMANOIR. 1956. Copépodes parasites de poissons de Madagascar. I. Mémoires de l'Institut Scientifique de Madagascar Series A 10: 69–80.
- OHTSUKA, S., AND G. A. BOXSHALL. 2019. Two new species of the genus *Caligus* (Crustacea, Copepoda, Siphonostomatoida) from the Sea of Japan, with a note on the establishment of a new species group. ZooKeys 893: 91–113. doi:10.3897/ zookeys.893.46923.
- OHTSUKA, S., M. NAWATA, Y. NISHIDA, M. NITTA, K. HIRANO, K. ADACHI, Y. KONDO, B. A. V. MARAN, AND E. SUÁREZ-MORALES. 2020. Discovery of the fish host of the 'planktonic' caligid *Caligus undulatus* Shen & Li, 1959 (Crustacea: Copepoda: Siphonostomatoida). Biodiversity Data Journal 8: e52271: doi:10.3897/BDJ.8.e52271.
- OHTSUKA, S., Y. NISHIDA, K. HIRANO, T. FUJI, T. KAJI, Y. KONDO, S. KOMEDA, S. TASUMI, K. KOIKE, AND G. A. BOXSHALL. 2021. The cephalothoracic sucker of sea lice (Crustacea: Copepoda: Caligidae): The functional importance of cuticular membrane ultrastructure. Arthropod Structure & Development 62: 1–24.
- ÖZAK, A. A., I. DEMIRKALE, G. A. BOXSHALL, AND M. ETYEMEZ. 2013. Parasitic copepods of the common sole, *Solea solea* (L.), from the eastern Mediterranean coast of Turkey. Systematic Parasitology 86: 173–185.

- ÖZAK, A. A., T. T. KURT, S. A. KAMANLI, G. AKBULUT, A. YANAR, AND G. A. BOXSHALL. 2024. Two new species of planktonic *Caligus* O.F. Müller 1785 (Copepoda: Caligidae) from Türkiye with an updated review and checklist of planktonic caligids. Zootaxa 5437: 301–335. doi:10.11646/zootaxa. 5437.3.
- ÖZAK, A. A., A. YANAR, AND G. A. BOXSHALL. 2017. The discovery of *Caligus macrurus* Heller, 1865 (Copepoda: Caligidae) in the Mediterranean Sea, and the recognition of *Sciaenophilus* van Beneden, 1852 as a junior synonym of *Caligus* Müller, 1785. Systematic Parasitology 94: 97–109.
- ÖZDIKMEN, H. 2008. Nomenclatural changes for nine crustacean genera (Crustacea: Copepoda). Munis Entomology and Zoology 3: 265–274.
- PILLAI, N. K. 1961. Copepods parasitic on South Indian fishes. Part I. Caligidae. Bulletin of the Research Institute, University of Kerala 8: 87–130.
- PILLAI, N. K. 1962. A revision of the genera *Parapetalus* Steenstrup & Lütken and *Pseudopetalus* nov. Crustaceana 3: 285–303.
- PILLAI, N. K. 1963. Observations on the genus *Abasia* (Copepoda) with the description of a new species. Crustaceana 5: 1–9.
- PILLAI, N. K. 1985. The Fauna of India: Copepod Parasites of Marine Fishes. White Lotus Press, Calcutta, India, 900 p.
- PILLAI, N. K., AND M. S. HAMEED. 1981. Description of a new species of *Parapetalus* (Caligidae: Copepoda) from Kerala. Journal of the Marine Biological Association of India 23: 77–81.
- PRABHA, C., AND N. K. PILLAI. 1979. *Pseudechetus fimbriatus* gen. et sp. nov., a caligid copepod from Kerala coastal waters. Parasitology 79: 425–429.
- PRABHA, C., AND N. K. PILLAI. 1983. Additions to the copepods parasitic on the marine fishes of India. 2. *Kabataella indica* gen. et sp. nov. (Caligoida). Records of the Zoological Survey of India 81: 1–5.
- PREIBISCH, S., S. SAALFELD, AND P. TOMANCAK. 2009. Globally optimal stitching of tiled 3D microscopic image acquisitions. Bioinformatics 25: 1463–1465. doi:10.1093/bioinformatics/ btp184.
- RANGNEKAR, M. P. 1956. Parasitic copepods from the marine fishes of Bombay. Journal of the University of Bombay 24: 42–65.
- RANGNEKAR, M. P. 1958. *Mappates plataxus*, gen. et sp. nov., a copepod parasitic on the fish, *Platax teira* (Forsk.). Records of the Zoological Survey of India 53: 303–308.
- RANGNEKAR, M. P. 1984. Parasitic copepods from marine fishes of Bombay. Crustaceana(Suppl. 7): 344–351.
- RANGNEKAR, P. G. AND N. N. MURTI. 1950. A new caligid copepod parasitic on the fish Clupea toli. Journal of the University of Bombay (B) 18: 21–28.
- REDKAR, M. V., P. G. RANGNEKAR, AND N. N. MURTI. 1949. Four new species of parasitic copepods from the marine fishes of Bombay. Journal of the University of Bombay 18: 36–50.
- RODGER, H., A. HERRERO, AND C. MATTHEWS. 2022. Pathologies associated with sea lice. *In* Sea Lice Biology and Control, J. Treasurer, I. Bricknell, and J. Bron (eds.). 5M Books Ltd., Great Easton, U.K., p. 134–149.
- ROUBAL, F. R., J. ARMITAGE, AND K. ROHDE. 1983. Taxonomy of metazoan ectoparasites of snapper, *Chrysophrys autratus* (family Sparidae), from southern Australia, eastern Australia

and New Zealand. Australian Journal of Zoology Supplementary Series 31: 1–68.

- RUSSELL, F. S. 1925. A new species of Caligus from Egypt, *Caligus pageti* sp. n. Annals and Magazine of Natural History Series 9: 611–618.
- SHEN, C. J. 1957. Parasitic copepods from fishes of China. II. Caligoida, Caligidae (1). Acta Zoologica Sinica 9: 351–377.
- SHIINO, S. M. 1952. Copepods parasitic on Japanese fishes. I. On the species of *Caligus* and *Lepeophtheirus*. Report of the Faculty of Fisheries, Prefectural University of Mie 1: 79–113.
- STEBBING, T. R. R. 1900. On the Crustacea brought by Dr Wiley from the South Seas. *In* Zoological Results Based on Material from New Britain, New Guinea, Loyalty Islands and Elsewhere Collected During the Years 1895, 1896 and 1897. Part V, A. Willey (ed.). Cambridge University Press, Cambridge, U.K., p. 64–74.
- STEENSTRUP, J. J. S., AND C. F. LÜTKEN. 1861. Bidrag til Kundskab om det aabne Havs Snyltekrebs og Lernaeer samt om nogle andre nye eller hidtil kun ufuldstaendigt kjendte parasitiske Copepoder. Kongelige Danske Videnskabernes Selskabs Skrifter, Naturhistorisk og Mathematisk Afdeling 5: 341–432. doi:10.5962/bhl.title.59539.
- SUÁREZ-MORALES, E., I. H. KIM, AND J. B. ESCAMILLA. 2012. On some caligids (Copepoda: Caligidae) from plankton of a coastal lagoon in the Gulf of Mexico with a description of a new species of *Metacaligus*. Zoological Studies 51: 804–818.
- THOMSON, G. M. 1890. Parasitic Copepoda of New Zealand, with descriptions of new species. Transactions of the New Zealand Institute 22: 353–376.
- THOMSEN, R. 1949. Copepods parasites de los peces marinos del Uruguay. Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 3: 1–41.
- TREASURER, J. W., AND S. BRAVO. 2011. The spatial distribution patterns of *Caligus rogercresseyi* and *C. elongatus* on Atlantic salmon hosts (*Salmo salar*). Aquaculture 320: 154–158.

- VAN BENEDEN, P. J. 1852. Note sur quelques parasites d'un poisson rare sur nos côtes (le Maigre d'Europe, *Sciaena aquila* Cuv). Bulletin de l'Academie royale de Belgique 19: 98–109.
- VAN BENEDEN, P. J. 1892. Quelques nouveaux Caligidés de la côte d'Afrique et de l'archipel des Açores. Bulletin de l'Academie royale de Belgique 3: 241–262.
- VON NORDMANN, A. 1832. Mikrographische Beiträge Zur Naturgeschichte der Wirbellosen Thiere. G. Reimer, Berlin, Germany, 150 p.
- WALTER, T. C., AND G. BOXSHALL. 2024. World of Copepods Database. Available at: https://www.marinespecies.org/ copepoda. Accessed 31 January 2024. doi:10.14284/356.
- WILSON, C. B. 1905. New species of parasitic copepods from the Massachusetts coast. Proceedings of the Biological Society of Washington 18: 127–131.
- WILSON, C. B. 1906. Report on some parasitic Copepoda collected by Professor Herdman, at Ceylon, in 1902. *In* Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar, W. A. Herdman (ed.). The Royal Society, London, U. K., p. 189–210.
- WILSON, C. B. 1908. North American parasitic copepods: New genera and species of Caliginae. Proceedings of the United States National Museum 33: 593–627. doi:10.5479/si.00963801.33-1580.593.
- WILSON, C. B. 1911. North American parasitic copepods: Descriptions of new genera and species. Proceedings of the United States National Museum 39: 263–400.
- WILSON, C. B. 1913. Crustacean parasites of West Indian fishes and land crabs, with descriptions of new genera and species. Proceedings of the United States National Museum 44: 189– 227. doi:10.5479/si.00963801.44-1950.189.
- YAMAGUTI, S. 1936. Parasitic copepods from fishes of Japan. Part 2. Caligoida I. S. Yamaguti, Kyoto Imperial University, Kyoto, Japan, p. 1–22.
- YAMAGUTI, S. 1954. Parasitic copepods from fishes of Celebes and Borneo. Publications of the Seto Marine Biological Laboratory 3: 375–398.