



**First Report of *Geoica utricularia* (Hemiptera: Aphididae) Population on Parasitic Broomrape *Orobanche foetida***

Authors: Boukhris-Bouhachem, Sonia, Youssef, S. Ben, and Kharrat, M.

Source: Florida Entomologist, 94(2) : 343-344

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0230>

---

BioOne Complete ([complete.bioone.org](https://complete.bioone.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## FIRST REPORT OF *GEOICA UTRICULARIA* (HEMIPTERA: APHIDIDAE) POPULATION ON PARASITIC BROOMRAPE *OROBANCHE FOETIDA*

SONIA BOUKHRIS-BOUHACHEM<sup>1</sup>, S. BEN YOUSSEF<sup>2</sup> AND M. KHARRAT<sup>3</sup>

<sup>1</sup>Plant Protection Laboratory, National Institute of Agricultural Research of Tunisia

<sup>2</sup>Animal and Forage Laboratory, National Institute of Agricultural Research of Tunisia

<sup>3</sup>Field Crop Laboratory, National Institute of Agricultural Research of Tunisia INRAT, Rue Hédi Karray, 2049, Tunisia

*Geoica utricularia* (Passerini) is known as a species that alternates between galls on *Pistacia* and the roots of Gramineae, and occasionally Cyperaceae (Blackman & Eastop 1985). Its distribution has been recorded as Morocco, Europe, the Middle East, Central Asia and North America (Blackman & Eastop 1985). In 2009, we observed severe attacks of *G. utriculariae* on wheat and barley in Manouba and Cap-Bon regions in Northeast Tunisia. Previously, *G. utriculariae* had been captured in suction traps in the same region (Boukhris-Bouhachem et al. 2007).

Very few insects have been reported to feed on *Orobanche* (Orobanchaceae). The case of *Phytomyza orobanchia* Kalt. (Diptera: Agromyzidae) has been reported (Linke 1990; Klein & Kroschel 2002), and Zermane et al. (2001) found *Smicronyx cyaneus* Gyll. (Coleoptera: Curculionidae) on *Orobanche foetida* Poir. in Tunisia. Holman (2009) reported *Smynturodes betae* WestWood (Hemiptera: Aphididae) on 3 *Orobanche* species (*O. aegyptiaca*, *O. crenata* and *O. variegata*).

Populations of aphids were found on the subterranean parts, and feeding on spikes of two broomrape plants, *Orobanche foetida*, attached to faba bean (*Vicia faba* L.) in May 2010 in a field located at Ariana (36°47' 57"N, 10°10'32"E). Samples of aphids were collected and prepared for microscopic examination in Canada balsam. They were identified using the Remaudière & Seco Fernandez (1990) and Blackman & Eastop (2000) keys, and also submitted for confirmation to Jon H. Martin at the Natural History Museum in London, UK. Aphid specimens in the colony were white cream, lightly dusted with wax, and broadly oval (Fig. 1). The aphids were identified as *G. utricularia* (Fig. 2) (determination confirmed by J. H. Martin) and compared favorably with specimens in the museum collection from Morocco and elsewhere. Voucher specimens were deposited at the British museum.



Fig. 1. Apterous viviparae *Geoica utricularia*.



Fig. 2. Colony of *G. utricularia* feeding on *O. foetida*.

To our knowledge, no associations of this aphid species have been reported previously on *Orobanche foetida*, so even if the species identification is suspect due to the unusual host range, it represents a host range expansion for the genus. Broomrape infestations are increasing in different regions of Tunisia (Kharrat et al. 2004). Herefore, the association between *G. utriculariae* and *Orobanche* has not been reported, and it is quite curious because the aphids normally feed on Gramineae except when forming galls on *Pistacia*. A supplementary study will be conducted to determine if aphids can reduce seed production of *Orobanche*.

#### SUMMARY

A new association between aphids (Aphididae) and *Orobanche foetida* was observed in Tunisia. The aphids were identified as *Geoica utricularia* (Passerini). The feeding of *G. utricularia* on *Orobanche* is illustrated.

#### REFERENCES CITED

- BLACKMAN, R. L., AND EASTOP, V. F. 2000. Aphids on the World's Crops: an Identification and Information Guide. Wiley and Sons, Chichester, UK 475 pp.
- BOUKHRIS-BOUHACHEM, S., SOUISSI, R., TURPEAU, E., ROUZE-JOUAN, J., FAHEM, M., BEN BRAHIM, N., AND HULLE, M. 2007. Aphid (Hemiptera, Aphidoidea) diversity in Tunisia in relation to seed production. Ann. Soc. Entomol. France 43: 311-318.
- HOLMAN, J. 2009. Host Plant Catalog of Aphids, Palearctic Region. Academy of Science of Czech Republic. Springer, 1216 pp.
- KHARRAT, M., AND SOUISSI, T. 2004. Research on *Orobanche foetida* and *O. crenata* in Tunisia, pp. 106-110 In R. Dahan and M. El-Mourid [eds.], Integrated Management of Orobanche in Food Legumes in the Near East and North Africa. Proc. Expert Consultation on IPM for Orobanche in Food Legume Systems Near East and North Africa, Rabat, Morocco. ICARDA/INRA/FAO, ICARDA, Aleppo, Syria.
- KLEIN, O., AND KROSCHER, J. 2002. Biological control of *Orobanche* spp. with *Phytomyza orobanchia*, a review. Biocontrol 47(3): 245-277.
- LINKE, K.-H., VORLAENDER, C., AND SAXENA, M. C. 1990. Occurrence and impact of *Phytomyza orobanchia* [Diptera: Agromyzidae] on *Orobanche crenata* [Orobanchaceae] in Syria. Biocontrol 35(4): 633-639.
- REMAUDIÈRE, G., AND SECO FERNANDEZ, MA. V. 1990. Claves para ayudar al reconocimiento de alados de pulgones trapeados en la region mediterranea (Hom. Aphidoidea). Universidad De León, León, 2V, 205 pp.
- ZERMANE, N., KROSCHER, J., SOUISSI, T., AND KHARRAT, M. 2001. First report of *Smicronyx cyaneus* Gyll. on *Orobanche foetida* Poiret: investigations in Tunisia. Haustorium (Parasitic Plants Newsletter), 39: 5-6 (Research Note).