

Diversity of Frugivorous Flies (Diptera: Tephritidae and Lonchaeidae) and their Relationship with Host Plants (Angiospermae) in Environments of South Pantanal Region, Brazil

Authors: Nicácio, José, and Uchôa, Manoel A.

Source: Florida Entomologist, 94(3) : 443-466

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0309>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DIVERSITY OF FRUGIVOROUS FLIES (DIPTERA: TEPHRITIDAE AND LONCHAEIDAE) AND THEIR RELATIONSHIP WITH HOST PLANTS (ANGIOSPERMAE) IN ENVIRONMENTS OF SOUTH PANTANAL REGION, BRAZIL

JOSÉ NICÁCIO AND MANOEL A. UCHÔA

Laboratório de Insetos Frugívoros, Caixa Postal 241, Faculdade de Ciências Biológicas e Ambientais (FCBA), Universidade Federal da Grande Dourados (UFGD), Rod. Dourados Itahum, Km 12, CEP 79804-970, Cidade Universitária, Dourados, MS, Brazil

E-mails: <uchoa.manoel@gmail.com>; <jose.nicacio1@uol.com.br>

ABSTRACT

Tephritidae and Lonchaeidae (Tephritoidea) are major pests of horticultural crops worldwide. Knowledge of the interactions between these flies and their host plants is needed for rational methods of population control. In the South Pantanal and adjacent areas in Brazil we sampled fruits from 92 plant species (22 orders and 36 families) in natural environments. Fifty-three species of plants were infested, and 39 not infested. Some aspects of the biology and patterns of species diversity, abundance, frequency, pupation period for males and conspecific females, and the interactions among species of frugivorous flies and their host plants were quantified. Twenty-six species of flies from 4 genera and 2 families were reared from the fruits: 17 species of Tephritidae (16 *Anastrepha* spp. and *Ceratitis capitata*), and 9 species of Lonchaeidae, comprising 8 species of *Neosilba* McAlpine, and 1 species of *Lonchaea* Fallén (Lonchaeidae). The lonchaeids infested 48 species of fruits, and the tephritids 30 species. The most polyphagous species were *Neosilba zadolicha* McAlpine & Steyskal (36 hosts), *Neosilba pendula* (Bezzi) (18 hosts), and *Neosilba inesperata* Strikis & Prado (14 hosts) (Lonchaeidae); and *Anastrepha sororcula* Zucchi (10 hosts), *Ceratitis capitata* (Wiedemann) (10 hosts), *Anastrepha obliqua* (Macquart) (7 hosts), and *Anastrepha zenildae* (6 hosts) (Tephritidae). All the males of Tephritidae (16 *Anastrepha* spp. and *C. capitata*) emerged before their conspecific females. Conversely, all females of the 8 species of *Neosilba* emerged in advance of their conspecific males.

Key Words: biodiversity, Cerrado, frugivory, fruit infestation, Neotropics, taxonomy

RESUMEN

Tephritidae y Lonchaeidae (Tephritoidea) son las principales plagas de la horticultura en todo el mundo. El conocimiento de las interacciones entre estas moscas y sus plantas huéspedes es necesario para el uso de los métodos racionales de control de sus poblaciones. En el Pantanal Sur y zonas adyacentes en Brasil fueron tomadas muestras de frutos de 92 especies de plantas (22 órdenes y 36 familias) en ambientes naturales, y 26 especies frugívoras de Tephritoidea fueron creadas a partir de sus plantas hospederas (17 especies de Tephritidae y nueve especies de Lonchaeidae). Cincuenta y tres especies de plantas estaban infestadas y 39 no estaban infestadas. Fueron investigados algunos patrones biológicos y fue cuantificada la diversidad de especies, abundancia, frecuencia, período de pupación de machos y hembras de las especies, y las interacciones entre las especies de moscas y sus plantas hospederas. Veinte y seis especies de moscas de 4 géneros y 2 familias fueron criados de los frutos: diecisiete especies de Tephritidae (16 *Anastrepha* spp. y *Ceratitis capitata*), y nueve especies de Lonchaeidae, siendo 8 especies de *Neosilba* McAlpine y una de *Lonchaea* Fallén (Lonchaeidae). El lonqueidos colonizaron 48 especies de frutas, y los tefritidos colonizaron 30 especies. Las especies más polífaga fueron: *Neosilba zadolicha* McAlpine & Steyskal (36 hospederos), *Neosilba pendula* (Bezzi) (18 hospederos), y *Neosilba inesperata* Strikis & Prado (14 hospederos) (Lonchaeidae); *Anastrepha sororcula* Zucchi (10 hospederos), *Ceratitis capitata* (Wiedemann) (10 hospederos), y *Anastrepha obliqua* (Macquart) (7 hospederos) (Tephritidae). Los machos de Tephritidae emergieron de las pupas antes que las hembras (16 *Anastrepha* spp. y *C. capitata*). Mientras, todas las hembras de las 8 especies de *Neosilba* emergieron antes que sus machos co-específicos.

Translation provided by the authors.

The polyphagous and oligophagous species of Tephritidae (fruit flies) and Lonchaeidae (lance flies) (Tephritoidea) are the main pests of horticulture in the Neotropical Region. They are the cause of quarantine constraints imposed by many countries to curtail their entry, and they significantly influence commerce inside and outside of the countries producing fruits and vegetables around the world (Uchôa et al. 2002; Uchôa & Nicácio 2010). On the other hand, many stenophagous and monophagous tephritoid species are important as potential biological control agents for invading species of plants (Norrbon & McAlpine 1997), in the decomposition of fruits and cycling of nutrients in the food web, and as alternative hosts for the multiplication of parasitoids, which may reduce the populations of pest flies in orchards.

The Pantanal and Cerrado biomes host important components of Neotropical biodiversity, and the insects are among the taxa most poorly known in both environments (Myers et al. 2000; Junk et al. 2006). Frugivorous Tephritoidea (fruit flies and lance flies) are biodiverse in all Brazilian regions, but their interactions with their host plants are poorly studied, and surveys of the Cerrado and Pantanal biomes are scarce.

The determination of the host plant status for species of fruit flies on commercially important fruits is a controversial topic that frequently causes commercial disputes between countries around the world, and knowledge about the interactions among fruit flies and their host plants is missing for decades. One of the reasons for this delay is a methodological problem (Aluja & Mangan 2008). As pointed out by Aluja & Mangan (2008), first it is necessary to determine if a plant is a natural host, a conditional host (non-natural infestation that occurs when only one species of fruit is offered to a fruit fly species with no option of choice in laboratory conditions), or a non-host (when even in the laboratory under forced conditions the species of fruit fly does not oviposit in the only species of fruit offered as a resource). Some host plant records for fruit flies may be not trustworthy, because inappropriate methodology was applied to study fruit fly and plant interactions.

Aluja & Mangan (2008) classify the species of fruit flies into four types, according to their ability to colonize host plants: the species that infest fruits of different families/orders of plants are polyphagous; those that develop in fruits of the same family are oligophagous; species that infest fruits of a single genus of plants are stenophagous, and finally, monophagous species of fruit flies colonize fruits of only a single species of plant.

The most species rich genera of frugivorous Tephritoidea in Brazil are *Anastrepha* Schiner (Tephritidae) and *Neosilba* McAlpine (Lon-

chaeidae), with 251 (Uramoto et al. 2008; Norrbom & Korytkowski 2009; Canal 2010; Uramoto & Zucchi 2010; Norrbom & Korytkowski 2011; Norrbom & Uchôa 2011), and 17 described species (Strikis & Prado 2005, 2009; Strikis & Lereña 2009), respectively.

No host plant is known for 51 of the 112 species of *Anastrepha* reported from Brazil, (Zucchi 2008; Garcia et al. 2008; Uramoto & Zucchi 2010; Uchôa & Nicácio 2010; Norrbom & Uchôa 2011). In the genus *Anastrepha*, some species are of economic importance in the American tropics and subtropics (Calkins & Malavasi 1995) and about half of the species of *Neosilba* have been associated with fruits and vegetables in the Neotropics.

Knowledge of trophic interactions between frugivorous Tephritoidea and their host plants is important to support strategies for integrated management of fruit fly pests (polyphagous or oligophagous), and for the conservation of stenophagous and monophagous species in their natural environments.

The aims of this study were (1) identify the diversity, abundance and frequency of species of frugivorous Tephritidae and Lonchaeidae (Tephritoidea) in the Pantanal and surrounding areas, (2) study the occurrence and larval viability of the species of fruit flies and lance flies in different naturally infested host fruits in the Pantanal and adjacent areas, and (3) quantify the mean period of pupation (from the exit of larva from the host fruit until adult emergence) for both males and females of the different species of Tephritidae and Lonchaeidae.

MATERIALS AND METHODS

Characterization of Climate, Relief and Vegetation

The climate in the state of Mato Grosso do Sul is basically tropical and influenced by cold fronts during the winter (Jun-Sep), when the environment is drier and rainfall vary from 30-60 mm / month. Monthly mean temperature vary from 20-34°C, and the mean annual precipitation ranges from 1300 to 1700 mm. The altitude ranges from 100 to 1000 m above sea level, and vegetation types include savanna, seasonal semideciduous, deciduous, and sub-mountain. In the plateaus near the foothills there are dense shrub-herbaceous strata (1.5-2.0 m high), an intermediate continuous arboreal strata forming a canopy about 10 m above the ground with some trees of 15-25 m (Radam Brasil 1982; Galati et al. 2006).

The biome Pantanal (16-20°S/55-58°W, an environment that resembles the Florida Everglades), is the largest periodically flooded plain in the world, with an area of approximately 160,000 km² (140,000 km² in Brazilian territory, 15,000 km² in Bolivia and 5000 km² in Paraguay). It is recognized by the United Nations Educational,

Scientific and Cultural Organization (UNESCO) as a natural heritage of mankind, biological reserve of the biosphere, and included among the most fragile and threatened biomes of the world (Junk et al. 2006). The Pantanal is connected with the Serra de Maracajú and Cerrado.

The Serra de Maracajú is a plateau that divides the state of Mato Grosso do Sul into 2 distinct biomes: the Cerrado and the Pantanal. To the east of the capital (Campo Grande), Cerrado environments predominate, with soil composed mainly of sandstones, and to the west of Campo Grande, the Pantanal begins (Radam Brasil 1982).

The Cerrado biome, an environment that resembles an African savanna, is located mainly in Central Brazil and occupies approximately 2 million km² (about 25% of the Brazilian territory). Vegetation varies, but is characterized by its great diversity and high rates of endemism (about 2% of global diversity of plants). Among 26 global biodiversity-hotspots listed for preservation of habitat, it is listed as the sixth most important (Myers et al. 2000; Carvalho et al. 2009).

The original vegetation of the Cerrado biome occupied 1,783,200 km², but currently only 356,630 km² remain natural, which represent 20% of the original area. Of this, only 22,000 km² (6.2%) are located in protected areas, such as National Parks or private reserves. About 10,000 plant species are cataloged in the Cerrado, but for animals only 4 groups of vertebrates are well known (mammals, birds, reptiles and amphibians: 27,298 species). The diversity of invertebrates, totaling more than 95% of living species of animals and represented mainly by insects, is poorly documented (Myers et al. 2000).

Sampling of Plants

A total of 184 trips was made to sample the native fruits of South Pantanal, Cerrado and Serra de Maracajú. Fruits of 92 plant species from 22 orders and 36 families were evaluated for the occurrence of frugivorous tephritoid larvae.

The samples of fruits were taken during the seasons of fruit maturation for angiosperms in the South Pantanal Region (Pott & Pott 1994). Fruits were harvested directly from trees, or occasionally fruits recently fallen to the soil were collected, between Apr 1998 and Aug 2000, in 11 municipalities in the 3 different environments (Pantanal, Cerrado, and Sierra de Maracajú) of the state of Mato Grosso do Sul. The fruits of *Ximena americana* L. (Olacaceae) also were collected in Oct 2003 and Oct 2004 in the Pantanal.

Sampled Environments

Pantanal: Miranda (20°20'36"S/56°23'49"W, altitude 121 m); Aquidauana: Fazenda Aguapé

(20°6'18"S/55°58'11"W, 144 m), Fazenda Rio Negro (19°34'6"S/56°14'44"W, 107 m) and Aquidauana City - Lagoa dos Padres (20°28'46"S/55°48'35"W, 141 m); Corumbá: Passo do Lontra (19°35'32"S/57°32'34"W, 89 m) and Fazenda São Domingos (19°31'24"S/57°2'22"W, 90 m).

Cerrado: Aquidauana (20°28'36"S/55°47'15"W, altitude 151 m); Anastácio (20°29'1"S/55°49'48"W, 148 m) and Fazenda Nossa Senhora Aparecida (20°33'2"S/55°52'45"W, 173 m), Dois Irmãos do Buriti (20°40'47"S/55°17'46"W, 318 m), Terenos (20°27'2"S/55°5'3"W, 263 m), Campo Grande (20°26'34"S/54°38'47"W, 581 m); Nioaque (21°8'7"S/55°49'48"W, 213 m) and Rio Negro (19°26'58"S/54°59'13"W, 261 m).

Serra de Maracajú: Aquidauana - Fazenda Rio Doce: (20°21'52"S/55°36'35"W, altitude 478 m) and Morro do Paxixi (20°28'1"S/55°39'40"W, 192 m), Rochedo (19°57'11"S/54°53'33"W, 257 m), Miranda (20°20'36"S/56°23'49"W, 121 m) and Bodoquena (20°5'19"S/56°46'54"W, 130 m).

Collection of Larvae of Frugivorous Flies

Fruit samples were transported to the Laboratório de Controle Biológico de Insetos, Departamento de Biociências, Universidade Federal de Mato Grosso do Sul (UFMS), Campus de Aquidauana, where fruits were counted and weighed, then held to obtain associated insects according to methodology by Uchôa & Zucchi (1999). Photoperiod was 12h:12h (L:D). The temperature and relative humidity ranged from 15 to 40°C and from 40 to 90%.

Voucher specimens of plants were deposited in the Central Herbarium of the Universidade Federal de Mato Grosso do Sul (UFMS) in Campo Grande-MS, and in the Herbarium of the Departamento de Botânica, Universidade de São Paulo (USP) in São Paulo-SP, Brazil.

Exiting third instar tephritoids (L3) were counted and separated in transparent acrylic containers (300 mL) with sterile sand moistened with distilled water until adult emergence. Each container was labeled to identify the host fruit species and sample date, place, and municipality of collection, fly family, and date of L3 leaving the fruits. The larvae of Lonchaeidae and Tephritidae were differentiated by the shape of the posterior spiracles, body color, and morphology of the mandibles. The posterior spiracles of tephritid larvae (*Anastrepha* and *Ceratitis capitata* (Wiedemann)) are yellowish, inconspicuous, and lacking a sclerotized peritreme. The cephalopharyngeal region is shorter, compared to that of lonchaeids. The Lonchaeidae larvae have the posterior spiracles blackish, with a sclerotized peritreme, and more noticeable than those of tephritids, and the body of the lonchaeids is more filiform in shape, whitish, with the cephalopharyngeal region more elongated. Furthermore, the mouth hooks of the

Lonchaeidae larvae are darker and more visible than those of tephritids.

The adult flies after emergence, were kept alive for 2 or 3 d to allow full development of color patterns of the body and of the wings of each species. After that, they were killed by exposure to ethyl acetate, and preserved in vials with 70% ethanol.

Identification of Host Plants and Frugivorous Tephritoidea

The plants were identified by the botanists Ubirazilda Maria Rezende from the Herbarium of the Universidade Federal de Mato Grosso do Sul (UFMS) at Campo Grande-MS, and José Rubens Pirani from the Herbarium of the Departamento de Botânica, Universidade de São Paulo (USP, São Paulo), Brazil.

The species of *Anastrepha* and *C. capitata* were identified in the Laboratório de Insetos Frugívoros, Faculdade de Ciências Biológicas e Ambientais (FCBA), Universidade Federal da Grande Dourados (UFGD), by M. A. Uchôa using keys and species description (Lima 1934; Stone 1942; Foote 1980; Norrbom & Korytkowski 2009).

The species of *Neosilba* were identified mainly based on the morphology of the male genitalia (McAlpine & Steyskal 1982) by Pedro Carlos Strikis, UNICAMP, Campinas-SP, Brazil. Voucher specimens of Tephritidae are deposited in the Entomological Collection of the Museu da Biodiversidade (MuBio), FCBA-UFGD, Dourados-MS, Brazil, and voucher specimens of *Neosilba* (Lonchaeidae) are in the Zoological Collection of Universidade Estadual de Campinas-UNICAMP, Campinas-SP, Brazil.

Definition of Variables, Statistical Analysis and Methods

Statistical analysis was based on 5 qualitative variables: species, orders and families of plants, species and environments of emerged frugivorous flies. Five quantitative variables were evaluated: number of emerged flies, pupal period of the species of fly, number of third instar larvae (L3) / the number of fruit and number of L3 / the mass of fruit (g).

The fruit infestation levels were measured by 2 indexes: average number of third instars (= pre-pupal larva or L3) emerged / fruit, and average number of L3 / mass (g) of fruit. The first index was categorized into 3 intervals of classes: (1) low infestation (<0.1 larvae / fruit); (2) mean infestation (0.1 to 1.0 larvae / fruit); and (3) high infestation (>1.0 larvae / fruit).

The pupal period is defined as the time between the exit of the pre-pupal larva (L3) from the fruit until the emergence of the adult. Four class intervals were established for the average of

pupal period: 1 (6-10 d); 2 (10-14 d); 3 (14-18 d); and 4, over 18 d. The averages of pupal period were compared for 13 species of *Anastrepha*, *C. capitata* and 6 species of *Neosilba*. The statistical analysis included only those species that occurred in at least 3 fruit samples.

The viability of the pre-pupal larva (L3) was calculated by the equation: %V = No. of EA / L3 $\times$ 100, where: %V = Percent of viability), No. of EA = number of emerged adults, and L3 = Total number of pre-pupal larva; and the quotient was multiplied by 100.

The tests used were chi-square of Pearson (χ^2) to test the hypothesis of independence between 2 variables (Dawson & Trapp 2003), and the Kruskal-Wallis Test to compare each 2 variables (Pestana & Gageiro 2008).

We considered $\alpha < 1\%$ (highly significant difference), $1\% < \alpha < 5\%$ (significant difference), and $5\% < \alpha < 10\%$ (marginally significant difference).

Analysis of variance (ANOVA) was used to compare means when assumptions of homogeneity occurred (Levene's Test), and normality (Kolmogorov-Smirnov Test) for dependent variables (Maroco 2007).

RESULTS

From the total of 92 species of fruits sampled, 53 were infested by tephritoid larvae, from which 40 species had not been surveyed previously for Tephritoidea infestation (Uchôa et al. 2002; Zucchi 2007; Uramoto et al. 2008; Garcia et al. 2008; Uchôa & Nicácio 2010) (Table 1). A total of 89,593 fruit were collected weighting 1,141.55 kg, from which 28,714 third-instars (L3) exited.

In all the species of host fruit (except from *Duguetia furfuraceae* Saint Hilaire) only pulp feeding frugivorous flies were obtained. The Tephritidae comprised 22,505 pre-pupal larva (L3) and 13,443 adults, and the Lonchaeidae: 6,209 larvae (L3) and 3,770 adults (Table 1).

Thirty-nine species of plants (Aristolochiaceae [Aristolochiales], Bromeliaceae [Bromeliales], Cactaceae [Cactales]; Cucurbitaceae [Cucurbitales], Erythroxylaceae [Gentianales], Bombacaceae and Malvaceae [Malvales], Smilacaceae [Liliflorae], Meliaceae [Rutales], Sapindaceae [Sapindales], and Verbenaceae [Verbenales] were not infested by the tephritoid larvae (Uchôa & Nicácio 2010).

In this series of samples, 26 species of Neotropical Tephritoidea (16 species of *Anastrepha*, *C. capitata*, Tephritidae, and 9 species of Lonchaeidae) were reared from their host fruits (Table 2).

Tephritidae. The tephritids infested 30 species of fruits in 17 families and 13 orders of Angiospermae. Seventeen species emerged (16 *Anastrepha* and *C. capitata*), and a single species emerged from 5 hosts. The highest species richness of Te-

TABLE 1. HOST PLANTS (ANGIOSPERMAE) AND INFESTATION INDICES BY FRUGIVOROUS LARVAE OF FRUIT FLIES AND LANCE FLIES (DIPTERA: TEPHRITOIDEA) IN THE SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004).

Order	Sampled Plants			Obtained Tephritoidea						
	Fruit Tree Species			Larvae		Tephritidae		Lonchaeidae		
	Families	Infested Fruit Tree Species	Not Infested	Total of Sampled fruits	Total Biomass of Fruits (Kg)	Number of 3rd Instars (L3)	Number of L3 Larvae	Number of Adults	Number of L3Larvae	Number of Adults
22	36	53	39	89,593	1,141.55	28,714	22,505	13,443	6,209	3,770

phritidae occurred in the mountainous environment, where 10 species were recovered (Table 2).

The most ubiquitous species was *Anastrepha sororcula* Zucchi, which occurred in all 3 environments, followed by *Anastrepha zenildae* Zucchi in mountainous and cerrado areas. At the other extreme, *Anastrepha undosa* Stone was characterized as a monophagous species restricted to the swamp environment of the Pantanal (Table 2).

The host fruits *Terminalia catappa* L., *Psidium kennedyanum* Morong and *Ximenia americana* L. had the highest abundance of larvae. The most frequent species were *A. sororcula*, *Anastrepha striata* Schiner, and *A. zenildae*, which occurred in 26, 16, and 15 samples of fruits, respectively (Table 2).

The highest infestation rates (L3/fruit) occurred in *Andira cuyabensis* Benthams, followed by *Terminalia catappa* and *Ximenia americana*, with 9.59, 9.26, and 4.75 larvae per fruit, respectively. However, taking into account the rate of infestation number of pre-pupal larva by fruit mass (L3/g of fruit), the highest rates occurred in *T. catappa* (0.4 L/g), followed by *Andira cuyabensis* (0.33 L3 /g), and *Psidium kennedyanum* (0.31 L3/g) (Table 2).

The percentage of viability for the third instar (L3) was highest in the hosts *Sorocea sprucei saxicola* (Baillon) J. F. Macbride (95.83%), *Licania tomentosa* (94.29%), and *Andira cuyabensis* (94.0%) (Table 2).

The more generalist species of Tephritidae were *A. sororcula*, *C. capitata*, *Anastrepha obliqua* (Macquart), and *A. zenildae*, which infested 10, 10, 7, and 6 hosts, respectively (Table 2).

Lonchaeidae. Nine lonchaeid species (8 *Neosilba* spp. and 1 *Lonchaea* sp.) occurred in fruit of 48 species of plants from 24 families and 15 orders. From 2 hosts (*Syagrus flexuosa* (Martius) Beccari, and *Eugenia pitanga* (O. Berg) Kiaerskou, larvae exited but no adults emerged. Only species of *Neosilba* were obtained from 21 species of plants, and the highest species richness of this genus occurred in the Cerrado environment (S = 8) (Table 3).

The highest rates of fruit infestation by the species of *Neosilba* occurred in the hosts *Duguetia furfuracea* Saint Hilaire (10.78 L3/fruit and 0.22 L3/g of fruit), *Annona crassiflora* Martius (4.97 L3/fruit and 0.01 L3/g of fruit), and *Citrus sinensis* (L.) Osbeck (1.36 L3/fruit and 0.01 L3/g of fruit) (Table 3).

The highest viability (% of adult emergence from L3 larvae) for the Lonchaeidae occurred in *Andira cuyabensis* Benthams (100%), *Physalis angulata* L. (99.55%), *Licania tomentosa* (96.30%), and *Alibertia edulis* (Richard) A. Richard (93.75%) (Table 3).

The most abundant, frequent, and ubiquitous species of Lonchaeidae were *Neosilba zadolicha* McAlpine & Steyskal, *Neosilba pendula* (Bezzi),

TABLE 2. TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES			SPECIES OF FRUIT FLIES																INDICES / RATES								
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha	A. alveatoides	A. distincta	A. fraterculus	A. hastata	A. leptozona	A. macrura	A. obliqua	A. nr. pickeli	A. rheediae	A. serpentina	A. sororcula	A. striata	A. turpiniae	A. undosa	A. zenilidae	A. zeryni	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3
Arecales																												
Areaceae																												
Syagrus flexuosa (Martius)																												
Beccari	p-n	620	4,200	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Allogoptera leucocalyx (Drude)	c-n	264	2,350	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Celastrales																												
Hippocrateaceae																												
Cheiloclinium cognatum (Miers)	c-n	161	1,471.4	131	31	—	6	—	—	—	25	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.81	0.09	23.66
AC. Smith																												
Salacia elliptica (Martius)	s-n	374	16,792.5	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
G. Don																												
Ebenales																												
Ebenaceae																												
Diospyros hispida A. de Candolle	s/c-n	593	18,565.4	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Ericales																												
Sapotaceae																												

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES				SPECIES OF FRUIT FLIES																	INDICES / RATES							
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zeryni Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3	
Chrysophyllum gonocarpum (Martius & Eichler) Engler	s-n	521	22,821.9	6	5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3	2	0.01	0	83.33
	c-n	1,578	14,650	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Chrysophyllum soboliferum Rizzini	c-n	752	17,552	1,070	825	—	389	—	—	—	—	—	—	—	—	—	—	—	—	—	—	436	—	—	—	—	1.42	0.06	77.10
Pouteria glomerata (Miquel) Radlkofer	c-n	1,111	21,954	694	156	1	64	—	—	—	—	—	—	—	—	—	81	—	—	—	—	—	—	1	3	0.62	0.03	22.48	
Pouteria rami-flora (Martius) Radlkofer	p-n	1,600	34,593.5	616	185	15	90	—	—	—	—	64	—	—	—	—	16	—	—	—	—	—	—	—	—	—	0.39	0.02	30.03
Pouteria torta (Martius) Radlkofer	c/s-n	66	1,915	633	595	—	335	—	—	—	—	—	—	—	—	—	—	—	—	5	—	255	—	—	—	—	9.59	0.33	94.00
Fabales Fabaceae 'Andira cuyabensis Bentham	s/c-n	880	1,100	10	5	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.01	0.01	50.00
Flacourtiaceae Banara arguta Briquet	s-n	2,009	1,970	6	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	0	0	16.67
Casearia sylvestris Swartz	c-n																												

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRTIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES			SPECIES OF FRUIT FLIES																INDICES / RATES										
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zernyi Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3		
Mimosaceae																														
Inga laurina (Swartz)																														
Willdenow	c/p/s-n	1,042	15,192	49	21		1		3																5	12	0.05	0	42.86	
Gentianales																														
Apocynaceae																														
Hancornia																														
speciosa Gomez	c-n	395	8,769.5	4	2																				2		0.01	0	50.00	
Loganiaceae																														
Strychnos																														
pseudoquina																														
Saint-Hilaire	c/s-n	1,335	17,160	0	0																									
Rubiaceae																														
Alibertia edulis																														
(A. Richard) A. Richard	c/p-n	304	19,306.5	0	0																									
Genipa ameri-																														
cana L.	c/p-n	741	82,773	0	0																									
Tocoyena																														
formosa																														
(Chamisso & Schlechtendal)																														
K. Schumann	c/p-n	162	4,670	0	0																									

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES					SPECIES OF FRUIT FLIES																	INDICES / RATES						
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zernyi Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3	
Geraniales																													
Euphorbiaceae																													
Manihot sp.	c-n	396	6,350	12	11	—	7	—	—	—	—	—	—	—	4	—	—	—	—	—	—	—	—	—	—	—	0.03	0	91.67
Miller																													
Guttiferales																													
Clusiaceae																													
Rheedia																													
brasiliensis																													
Planchon &																													
Triana	p-n	460	1,311.5	174	49	3	33	—	—	—	—	—	—	—	—	13	—	—	—	—	—	—	—	—	—	—	0.38	0.13	28.16
Magnoliales																													
Annonaceae																													
Annona																													
crassiflora																													
Martius	s-n	29	24,413.4	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Duguetia																													
furfuracea (Saint-																													
Hilarie) Bentham	c-n	184	9,049	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Lauraceae																													
Persea americana																													
Miller	c-e Mex.	31	21,100	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Myrtales																													
Combretaceae																													

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES					SPECIES OF FRUIT FLIES																	INDICES / RATES						
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zeryni Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3	
Buchenavia sp. Echler Terminalia catappa L.	c-n	300	1,722	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	c/p-eMal.	987	23,121	9,139	5,344	—	9	—	—	—	—	—	—	—	—	—	—	2	—	—	—	3	—	2,681	2,649	9.26	0.4	58.47	
Melastomataceae Mouriri elliptica Martius	c-n	378	5,697.4	931	576	41	273	—	—	—	—	—	—	—	—	—	—	2	—	—	—	259	—	1	—	2.46	0.16	61.87	
Myrtaceae Syzygium jam- bos (L.) Alston Syzygium cumini (L.) Skeels Eugenia pitanga (O. Berg) Kiaerskou Psidium cattle- ianum Sabine Psidium guajava L.	c/p-n	1,030	4,034.4	3	2	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	0	0	66.67	—	
	c-n	6,651	16,607	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	p-n	105	72	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	c-n	524	12,745	541	326	—	163	—	—	—	—	—	—	—	—	—	—	—	154	9	—	—	—	—	—	1.03	0.04	60.26	
c/s-n	399	17,672.5	774	562	1	145	—	—	—	—	—	—	—	4	—	—	—	106	13	10	—	—	—	134	149	1.94	0.04	72.61	
Psidium kennedyanum Morong	c/s-n	2,694	11,821.9	3,648	2,667	220	1,265	—	—	5	—	—	—	2	—	—	—	1,023	151	1	—	—	—	—	—	1.35	0.31	73.11	
Rosales Chrysobalaceae																													

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES			SPECIES OF FRUIT FLIES																	INDICES / RATES							
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zernyi Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3
Licania tomentosa (Bentham) Fritsch	c-n	916	36,899.7	210	198	—	8	—	—	—	—	—	—	—	—	—	—	10	—	—	—	3	—	91	86	0.23	0.01	94.29
Rutales Malpighiaceae																												
Byrsonina orbignyana A. Jussieu	c/p-n	5,192	8,887	3	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	0	0	33.33
Rutaceae Citrus jambhiri Lush	c/p/s-n	79	11,113	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Citrus sinensis (L.) Osbeck	c/p-n	33	4,411	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Santalales Olacaceae																												
Schoepfia sp.	s-n	1,601	3,715	15	8	—	1	—	—	—	—	—	1	—	—	—	—	1	—	—	—	—	5	—	—	0.01	0	53.33
Ximenia americana L.	p-n	646	14,761	3,069	1,396	465	428	503	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.75	0.21	45.49
Sapindales Anacardiaceae																												

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES				SPECIES OF FRUIT FLIES																	INDICES / RATES								
	Host Environments	N° of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zernyi Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3		
Anacardium humile	c-n	715	21,335.8	70	40	—	25	—	—	—	—	—	—	15	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0	57.14	
Saint-Hilaire		—	—	—	—	—	—	—	—	—	—	—	—	—	5	—	—	—	1	—	—	—	—	—	—	—	0.18	0	37.93	
Anacardium othonianum	c-n	157	9,000	29	11	—	5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Rizzani	c-n	111	23,060	320	188	—	66	—	—	—	—	—	—	91	—	—	—	—	—	—	—	—	—	—	11	20	2.88	0.01	58.75	
Mangifera indica		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
L.	c-e (Ind)	246	21,159	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Spondias dulcis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Parkinson	c/s-n	120	1,024	260	165	6	72	—	—	—	—	—	—	87	—	—	—	—	—	—	—	—	—	—	—	—	2.17	0.25	63.46	
'Spondias lutea		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
L.	p-n	468	6,880.8	12	10	—	7	—	—	—	—	—	—	3	—	—	—	—	—	—	—	—	—	—	—	—	0.03	0	83.33	
Spondias purpureae L.		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tubiflorae	c-n	1,073	3,661.5	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Convolvulaceae		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Operculina alata	c-n	7,254	5,302.7	1	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	0	0	100.00	
(Hamilton)		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Urban	c-n	616	693.7	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Solanaceae		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Solanaceae	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Physalis		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
angulata L.	c-n	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		

TABLE 2. (CONTINUED) TAXONS OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). ENVIRONMENT LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL, AND ORIGIN OF THE SPECIES OF PLANTS: N = NATIVE, E = EXOTIC.

SAMPLES					SPECIES OF FRUIT FLIES															INDICES / RATES								
PLANT TAXA	Host Environments	N° Of fruits	Mass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Anastrepha spp. ♂ ♀	A. alveatoides Blanchard 1961	A. distincta Greene 1934	A. fraterculus (Wied. 1830)	A. hastata Stone 1942	A. leptozona Hendel 1914	A. macrura Hendel 1914	A. obliqua (Macquart 1835)	A. nr. pickeli Lima 1934	A. rheediae Stone 1942	A. serpentina (Wied. 1830)	A. sororcula Zucchi 1979	A. striata Schiner 1868	A. turpiniae Stone 1942	A. undosa Stone 1942	A. zenilidae Zucchi 1979	A. zeryni Lima 1934	C. capitata ♂	C. capitata ♀	Infestation (Mean No. of L3 / fruits)	Infestation (Mean No. of L3 / mass in g)	% VL3
Moraceae																												
<i>Ficus insipida</i>																												
Willdenow	s-n	3,111	13,400	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
<i>Sorocea sprucei</i>																												
<i>saxicola</i> (Baillon)																												
Macbride	s/p-n	497	1,787.6	72	69	8	32	—	—	—	—	—	—	—	—	—	—	—	29	—	—	—	—	—	—	0.14	0.04	95.83
Violales																												
Passifloraceae																												
<i>Passiflora</i>																												
<i>coccinea</i> Aublet	c/p-n	173	4,096.3	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
<i>Passiflora edulis</i>	c-n	102	10,149	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
"Pindaiva vermelha"	p-n	1,752	4,580	3	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—	—	—	—	—	—	—	0	0	66.67
Totals		53,538	669,441	22,505	13,443	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Abundance by fruit flies species						760	3,425	503	3	6	25	64	1	207	4	15	97	1,301	167	16	436	553	7	2,944	2,911	—	—	
Frequency by fruit flies species							73	4	1	4	2	6	1	10	2	3	6	26	6	5	4	15	4	17	16	—	—	
N° of plant genera infested							13	1	1	1	1	1	1	2	1	2	1	6	1	2	1	3	3	3	5	—	—	
N° of plant family infested							17	1	1	1	1	1	1	2	1	2	1	8	1	2	1	4	3	8	7	—	—	
N° of hosts colonized	9	30	1	1	2	1	1	1	7	1	2	2	10	3	3	1	6	3	10	8	—	—	—	—	—	—	—	

TABLE 3. TAXA OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF LANCE FLIES (DIPTERA: LONCHAEIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL; ORIGIN OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES					FRUGIVOROUS LONCHAEIDAE										INDICES / RATES			
	Environments	N° of fruits	Biomass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Lonchaea sp. (Fällen)	Neosilba spp. ♀	Neosilba bifida Strikis & Prado	N. certa (Walker)	N. glaberrima (Wied.)	N. inesperta Strikis a & Prado	N. pendula (Bezzi)	N. pradoi Strikis & Lereña	N. zadolicha McAlpine & Steyskal	Neosilba morphotype MSP1	Mean No. of L3/fruits	Mean No. of L3/in g	%VL3
Arecles																			
Arecaceae																			
Syagrus flexuosa (Martius) Beccari	p-n	620	4,200	9	0	—	0	—	—	—	—	—	—	—	—	—	0.01	0.00	0.00
Alloopterla leucocalyx (Drude) Kuntze	c-n	264	2,350	5	2	—	—	—	—	—	—	—	—	—	—	2	0.02	0.00	40.00
Celatrales																			
Hippocrateaceae																			
Cheilodinium cognatum (Miers) A.C. Smith	c-n	161	1,471	0	0	—	—	—	—	—	—	—	—	—	—	—	0.00	0.00	0.00
Salacia elliptica (Martius) G. Don	s-n	374	16,793	7	4	—	—	2	—	—	—	—	—	—	2	—	0.02	0.00	57.14
Ebenales																			
Ebenaceae																			
Diospyros hispida A. de Candolle	s/c-n	593	18,565	80	37	—	—	17	—	—	—	—	—	—	20	—	0.13	0.00	46.25
Ericales																			
Sapotaceae																			
Chrysophyllum gonocarpum (Martius & Eichler) Engler	s-n	521	22,822	0	0	—	—	—	—	—	—	—	—	—	—	—	0.00	0.00	0.00
Chrysophyllum soboliferum Rizzini	c-n	1,578	14,650	8	1	—	—	—	—	—	—	—	1	—	—	—	0.01	0.00	12.50
Pouteria glomerata (Miquel) Radlkofer	c-n	752	17,552	55	7	—	—	2	—	1	—	—	—	—	4	—	0.07	0.00	12.73
Pouteria ramiflora (Martius) Radlkofer	p-n	1,111	21,954	816	340	—	—	163	—	—	1	1	8	—	167	—	0.73	0.04	41.67
Pouteria torta (Martius) Radlkofer	c/s-n	1,600	34,594	589	366	—	—	195	—	3	19	2	26	—	121	—	0.37	0.02	62.14
Fabales																			
Fabaceae																			
Andira cuyabensis Bentham	s/c-n	66	1,915	12	12	—	—	5	—	—	—	—	6	—	1	—	0.18	0.01	100.00

*Host plant with 2 Figitidae, *Lopheucoila anastrephae*, parasitizing larvae of *Neosilba* spp.

TABLE 3. (CONTINUED) TAXA OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF LANCE FLIES (DIPTERA: LONCHAEIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL; ORIGIN OF PLANTS: N = NATIVE, E = EXOTIC.

SAMPLES				FRUGIVOROUS LONCHAEIDAE										INDICES / RATES						
PLANT TAXA	Environments	N° of fruits	Biomass of fruits (g)	Larvae (L3)	Emergед Adults	Associated Parasitoids	Lonchaea sp. (Fallen)	Neosilba spp. ♀ ♀	Neosilba bifida Strikis & Prado	N. certa (Walker)	N. glaberrima (Wied.)	N. inesperata Strikis a & Prado	N. pendula (Bezzi)	N. pradoi Strikis & Lereña	N. zadolicha McAlpine & Steyskal	Neosilba morphotype MSP1	Mean No. of L3/fruits	Mean No. of L3/in g	%VL3	
Flacourtiaceae																				
Banara arguta Briquet	s-n	880	1,100	4	2	—	—	1	—	—	—	—	1	—	—	—	—	0.00	0.00	50.00
Casearia sylvestris Swartz	c-n	2,009	1,970	9	1	—	—	1	—	—	—	—	—	—	—	—	—	0.00	0.00	11.11
Mimosaceae																				
*Inga laurina (Swartz) Willdenow	c/p/s-n	1,042	15,192	127	80	2	—	38	—	—	—	21	11	1	7	—	0.12	0.01	62.99	
Gentianales																				
Apocynaceae																				
Hancornia speciosa Gomez	c-n	395	8,770	5	2	—	—	—	—	—	—	—	—	—	2	—	0.01	0.00	40.00	
Loganiaceae																				
Strychnos pseudoquina Saint-Hilarie	c/s-n	1,335	17,160	27	14	—	—	6	—	—	—	2	—	—	6	—	0.02	0.00	51.85	
Rubiaceae																				
Alibertia edulis (A. Richard) A. Richard	c/p-n	304	19,307	32	30	—	—	19	—	—	5	—	—	—	6	—	0.11	0.00	93.75	
Genipa americana L.	c/p-n	741	82,773	13	8	—	—	4	—	—	1	—	—	—	3	—	0.02	0.00	61.54	
Tocoyena formosa (Chamisso & Schechtendal) K. Schumann	c/p-n	162	4,670	9	6	—	—	2	—	—	—	—	—	—	4	—	0.06	0.00	66.67	
Geraniales																				
Euphorbiaceae																				
Manihot sp.	c-n	396	6,350	0	0	—	—	—	—	—	—	—	—	—	—	—	0.00	0.00	0.00	
Guttiferales																				
Clusiaceae																				
Rheedia brasiliensis Planchon & Triana	p-n	460	1,312	0	0	—	—	—	—	—	—	—	—	—	—	—	0.00	0.00	0.00	

*Host plant with 2 Figitidae, *Lopheucoila anastrephae*, parasitizing larvae of *Neosilba* spp.

TABLE 3. (CONTINUED) TAXA OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF LANCE FLIES (DIPTERA: LONCHAEIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). LEGEND: C = CERRADO, S = PANTANAL; ORIGIN OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES					FRUGIVOROUS LONCHAEIDAE										INDICES / RATES			
	Environments	N° of fruits	Biomass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	<i>Lonchaea</i> sp. (Fallén)	<i>Neosilba</i> spp. ♀ ♀	<i>Neosilba bifida</i> Strikis & Prado	<i>N. certa</i> (Walker)	<i>N. glaberrima</i> (Wied.)	<i>N. inesperta</i> Strikis a & Prado	<i>N. pendula</i> (Bezzi)	<i>N. pradoi</i> Strikis & Lereña	<i>N. zadolicha</i> McAlpine & Steyskal	<i>Neosilba</i> morphotype MSP1	Mean No. of L3/fruits	Mean No. of L3/in g	%VL3
Magnoliales																			
Annonaceae																			
<i>Annona crassiflora</i> Martius	s-n	29	24,413	144	93	—	—	48	—	—	1	—	—	—	44	—	4.97	0.01	64.58
<i>Duguetia furfuracea</i> (Saint-Hilaire)	c-n	184	9,049	1983	1100	—	—	584	—	1	41	1	—	—	473	—	10.78	0.22	55.47
Benthams																			
Lauraceae																			
<i>Persea americana</i> Miller	c-e Mex.	31	21,100	60	49	—	—	27	—	—	—	—	—	—	22	—	1.94	0.00	81.67
Myrtales																			
Combretaceae																			
<i>Buchenavia</i> sp.	c-n	300	1,722	4	1	—	—	—	—	—	—	—	—	—	1	—	0.01	0.00	25.00
<i>Terminalia catappa</i> L.	c/p-eMal.	987	23,121	386	181	—	—	93	—	27	1	11	27	—	22	—	0.39	0.02	46.89
Melastomataceae																			
<i>Mouriri elliptica</i> Martius	c-n	378	5,697	20	5	—	—	4	—	—	—	—	—	—	1	—	0.05	0.00	25.00
Myrtaceae																			
<i>Syzygium cumini</i> (L.) Skeels	c-n	6,651	16,607	29	14	—	—	11	—	—	—	—	3	—	—	—	0.00	0.00	48.28
<i>Syzygium jambos</i> (L.) Alston	c/p-n	1,030	4,034	106	72	—	—	41	—	1	8	—	—	—	22	—	0.10	0.03	67.92
<i>Eugenia pitanga</i> (O.Berg) Kiaerskou	p-n	105	72	1	0	—	—	0	—	—	—	—	—	—	—	—	0.01	0.01	0.00
<i>Psidium cattleianum</i> Sabine	c-n	524	12,745	24	13	—	—	4	—	—	—	2	6	—	1	—	0.05	0.00	54.17
<i>Psidium guajava</i> L.	c/s-n	399	17,673	61	49	—	—	22	—	—	—	1	17	—	9	—	0.15	0.00	80.33
<i>Psidium kennedyanum</i> Morong	c/s-n	2,694	11,822	4	3	—	—	2	—	—	—	—	—	—	1	—	0.00	0.00	75.00
Rosales																			
Chrysobalaceae																			
<i>Licania tomentosa</i> (Benthams) Fritsch	c-n	916	36,900	6	10	—	—	8	—	—	—	—	—	—	2	—	0.01	0.00	96.30

*Host plant with 2 Figitidae, *Lopheucoila anastrephae*, parasitizing larvae of *Neosilba* spp.

TABLE 3. (CONTINUED) TAXA OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF LANCE FLIES (DIPTERA: LONCHAEIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). LEGEND: C = CERRADO, S = SIERRA, P = PANTANAL; ORIGIN OF PLANTS: N = NATIVE, E = EXOTIC.

SAMPLES										FRUGIVOROUS LONCHAEIDAE							INDICES / RATES														
PLANT TAXA																															
Environments										N° of fruits	Biomass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	Lonchaea sp. (Fallen)	Neosilba spp. ♀ ♀	Neosilba bifida Strikis & Prado	N. certa (Walker)	N. glaberrima (Wied.)	N. inesperta Strikis a & Prado	N. pendula (Bezzi)	N. pradoi Strikis & Lereña	N. zadolicha McAlpine & Steyskal	Neosilba morphotype MSP1	Mean No. of L3/fruits	Mean No. of L3/in g	%VL3				
Rutales																															
Malpighiaceae																															
Byrsonima orbignyana A. Jussieu										c/p-n	5,192	8,887	8	3	—	—	2	—	—	—	—	—	—	—	1	—	—	0.00	0.00	37.50	
Rutaceae																															
Citrus jambhiri Lush										c/p/s-n	79	11,113	20	14	—	—	5	—	—	—	—	—	—	—	4	—	—	0.25	0.00	70.00	
Citrus sinensis (L.) Osbeck										c/p-n	33	4,411	45	36	—	—	20	1	1	—	—	—	—	—	11	—	—	1.36	0.01	80.00	
Santalales																															
Olacaceae																															
Schoepfia sp.										s-n	1,601	3,715	57	42	—	—	29	—	—	—	—	—	—	—	8	—	—	0.04	0.02	73.68	
Ximenia americana L.										p-n	646	14,761	91	27	—	1	11	—	—	—	—	—	—	—	—	—	0.14	0.01	29.67		
Sapindales																															
Anacardiaceae																															
Anacardium humile Staint-Hilaire										c-n	715	21,336	35	10	—	—	4	—	—	—	—	—	—	—	1	—	—	0.05	0.00	28.57	
Anacardium othonianum Rizzini										c-n	157	9,000	6	2	—	—	—	—	—	—	—	—	—	—	—	—	—	0.04	0.00	33.33	
Mangifera indica L.										c-n	111	23,060	42	33	—	—	16	—	—	—	—	—	—	—	17	—	—	0.38	0.00	78.57	
Spondias dulcis Parkinson										c-e (Ind)	246	21,159	9	1	—	—	1	—	—	—	—	—	—	—	—	—	0.04	0.00	11.11		
Spondias purpurea L.										p-n	468	6,881	17	8	—	—	4	—	—	—	—	—	—	—	2	—	—	0.04	0.00	47.06	
Spondias lutea L.										c/s-n	120	1,024	0	0	—	—	—	—	—	—	—	—	—	—	—	—	—	0.00	0.00	0.00	
Tubiflorae																															
Convolvulaceae																															
Operculina alata (Hamilton) Urban										c-n	1,073	3,662	23	21	—	—	15	—	—	3	—	—	—	—	—	—	—	—	0.02	0.01	91.30
Solanaceae																															
Physalis angulata L.										c-n	7,254	5,303	893	889	—	—	484	—	—	16	—	—	—	—	—	—	—	—	0.12	0.17	99.55

*Host plant with 2 Figitidae, *Lopheucoila anastrephae*, parasitizing larvae of *Neosilba* spp.

TABLE 3. (CONTINUED) TAXA OF PLANTS (ANGIOSPERMAE), AND PATTERNS OF RELATIONSHIP WITH SPECIES OF LANCE FLIES (DIPTERA: LONCHAEIDAE) IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004). LEGEND: C = SIERRA, S = PANTANAL; ORIGIN OF PLANTS: N = NATIVE, E = EXOTIC.

PLANT TAXA	SAMPLES					FRUGIVOROUS LONCHAEIDAE										INDICES / RATES			
	Environments	N° of fruits	Biomass of fruits (g)	Larvae (L3)	Emerged Adults	Associated Parasitoids	<i>Lonchaea</i> sp. (Fallen)	<i>Neosilba</i> sp. ♀	<i>Neosilba bifida</i> Strikis & Prado	<i>N. certa</i> (Walker)	<i>N. glaberrima</i> (Wied.)	<i>N. inesperta</i> Strikis a & Prado	<i>N. pendula</i> (Bezzi)	<i>N. pradoi</i> Strikis & Lereña	<i>N. zadolichia</i> McAlpine & Steyskal	<i>Neosilba</i> morphotype MSP1	Mean No. of L3/fruits	Mean No. of L3/in g	%VL3
<i>Solanum sisymbriifolium</i> Lamark	c-n	616	693.7	38	22	—	—	15	—	—	—	7	—	—	—	—	0.06	0.05	57.89
Urticales																			
Moraceae																			
<i>Ficus insipida</i> Willdenow	s-n	3,111	13,400	127	91	—	—	56	—	5	18	—	1	—	11	—	0.04	0.01	71.65
<i>Sorocea sprucei saxicola</i> (Baillon)	s/p-n	497	1,787.6	9	2	—	—	1	1	—	—	—	—	—	—	—	0.02	0.01	22.22
Macbride																			
Violales																			
Passifloraceae																			
<i>Passiflora coccinea</i> Aubler	c/p-n	173	4,096.3	80	63	—	—	30	—	—	—	—	—	—	33	—	0.46	0.02	78.75
<i>Passiflora edulis</i> Sims	c-n	102	10,149	65	1	—	—	—	—	—	—	—	—	—	1	—	0.64	0.01	1.54
Family not identified																			
"Pindaiva vermelha"	p-n	1,752	4,580	9	5	—	—	—	—	—	—	—	5	—	—	—	0.01	0.00	55.56
Totals	—	53,538	669,443.6	6,209	3,770	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Abundance by Species	—	—	—	—	—	2	1	1,992	2	59	103	443	127	1	1,038	4	—	—	—
Frequency by Species	85	—	—	—	—	1	1	85	2	12	16	20	30	1	62	2	—	—	—
N° of Infested Genera of Plants	—	—	—	—	—	—	1	13	1	6	6	8	7	1	13	2	—	—	—
N° of Infested Family of Plants	24	—	—	—	—	—	1	23	1	8	6	10	9	1	21	2	—	—	—

*Host plant with 2 Figitidae, *Lopheucoila anastrephae*, parasitizing larvae of *Neosilba* spp.

and the recently described *Neosilba inesperata* Strikis & Prado (2009), which colonized 36, 18, and 14 hosts, respectively (Tables 3).

The infestation rate for the total of larvae (L3) in the 92 species of plants was 78.38% by tephritids and 21.62% by lonchaeids (Table 4).

Larval Viability and Duration of Pupal Period. The relative viability of the larvae from both families of Tephritoidea was 59.73% for the tephritids, and 60.72% for the lonchaeids (Table 4). The pupal period (on average) for *Anastrepha* species, *C. capitata* (including male and female), and *Neosilba* species were 15.85 d, 10.62 d, and 12.21 d, respectively (Table 5). Among *Anastrepha* species, *A. hastata* Stone had the longest pupal period (21.92 d), and *A. zenildae* (mean 13.61 d) the shortest among all species of *Anastrepha*. The females of *C. capitata* emerged after 11.03 d. Among the lonchaeids, *N. inesperata* had the longest pupal period (13.43 d), and *Neosilba glaberrima* (Wiedemann) the shortest, emerging at 11.48 d (Table 6).

DISCUSSION

From the 53 species of fruits infested by frugivorous larvae of Tephritoidea, 25 were attacked by Tephritidae plus Lonchaeidae, 23 only by *Neosilba* species (Lonchaeidae), and 5 only by Tephritidae species. The plant species infested by the highest diversity of Tephritoidea species were *Psidium guajava* (8 spp., 5 of Tephritidae and 3 of Lonchaeidae), *Terminalia catappa* (8 spp., 3 of Tephritidae and 5 of Lonchaeidae), and *Pouteria ramiflora* (Martius) Radlkofer (7 spp., 3 of Tephritidae and 4 of Lonchaeidae) (Table 2).

The species *Ceratitis capitata*, *A. sororcula*, *A. zenildae*, and *Anastrepha alveatoides* Blanchard were the most numerically abundant. The species with the widest host ranges were *A. sororcula*, *C. capitata*, and *A. zenildae*. *Ceratitis capitata* and *A. sororcula* both infested 10 species of host fruits. *Ceratitis capitata* was exclusive in *Chryso-*

phyllum gonocarpum and *Hancornia speciosa*. *Anastrepha sororcula* was the only species reared from fruits of *Casearia sylvestris* Swartz, *Physalis angulata*, and *Byrsonima orbignyana* A. Jussieu, and *A. obliqua* infested 7 species of host fruits (Table 2).

The highest general infestation rates by larvae occurred in *T. catappa* (0.41 L3/g of fruit), *A. cuyabensis* (0.34 L3/g of fruit) and *P. kennedyanum* (0.31 L3/g of fruit). The viability for all the larvae (Tephritidae plus Lonchaeidae) was 59.95%. The host fruits with highest viability rates were *Physalis angulata* (99.55%), *Licania tomentosa* (96.30%), and *Andira cuyabensis* (94.11%) (Table 2).

The host fruits with the highest rates of infestation (L3/g) were *Terminalia catappa* (Combretaceae, Afrotropical), *A. cuyabensis* (Fabaceae, Neotropical), and *Psidium kennedyanum* (Myrtaceae, Neotropical) (Table 2). Uramoto et al. (2008) also found high levels of infestation by Tephritidae in native Myrtaceae from the Atlantic Forest in the state of Espírito Santo, Brazil. Fruits of Neotropical species of Myrtaceae are known to keep a high diversity of Tephritoidea in natural environments (Uchôa et al. 2002; Uramoto et al. 2008), probably, because their fruits are nutritionally adequate, and some species (e.g., *Psidium* spp.), produce fruit more than a season during the year.

Tephritidae. The host fruits with highest Tephritidae diversity were *Psidium guajava*, and *P. kennedyanum* (Myrtaceae, Neotropical), with 5 spp. each, and *L. tomentosa* (Chrysobalanaceae, Neotropical), infested by 3 species (Table 2).

The species of Tephritidae were more abundant than those of Lonchaeidae for both number of larvae and adults. However, the Lonchaeidae larvae were more frequently collected. While Tephritidae were obtained on 73 fruit sampling expeditions (Table 2), Lonchaeidae were collected on 85 of 184 sampling trips (Table 3).

TABLE 4. INFESTATION RATES BY LARVAE (L3) OF FRUGIVOROUS TEPHRITOIDEA (DIPTERA), VIABILITY (%), AND RELATIVE ABUNDANCE (%) IN HOST FRUITS FROM SOUTH PANTANAL REGION, BRAZIL (APR1998 TO AUG 2000, OCT 2003, AND OCT 2004)

Families of Frugivorous Tephritoidea	3rd Instars				Emergence	
	No. of L3	Relative Abundance of Infestation (%)	Not emerged adults from L3 (%)	Viability of L3 (%)	No. of adults	Relative Abundance of adults (%)
Lonchaeidae	6,209	21.62	2,439 (39.28)	60.72	3,770	21.90
Tephritidae	22,505	78.38	9,062 (40.26)	59.73	13,443	78.10
Total	28,714	100	11,501 (40.05)	59.95	17,213	100

Larvae of Lonchaeidae Collected × Number of Adults of Lonchaeidae Emerged: ($\chi^2_{(1)} = 15.29$; $P < 0.01$).
Larvae of Tephritidae Collected × Number of Adults of Tephritidae Emerged: ($\chi^2_{(1)} = 10.00$; $P < 0.01$).
Total of Larvae of Collected Tephritoidea × Number Total of Adults Tephritoidea Emerged: ($\chi^2_{(1)} = 37.41$; $P < 0.01$).

TABLE 5. PUPAL PERIODS FOR GENERA OF FRUGIVOROUS FLIES (DIPTERA: TEPHRITOIDEA) IN HOST FRUITS SAMPLED IN 3 ENVIRONMENTS IN SOUTH PANTANAL REGION, BRAZIL (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004).

Genera of Tephritoidea	Descriptive			Comparison of the Means	Mean Pupal Period of Genus
	Mean (days)	Standard Deviation	n		
<i>Anastrepha</i> ♀ ♀	16.02	3.04	106	a	15.85
<i>Anastrepha</i> ♂ ♂	15.69	2.92	65	a	
<i>Neosilba</i> ♂ ♂	12.23	2.30	146	b	12.21
<i>Neosilba</i> ♀ ♀	12.20	2.05	81	b	
<i>Ceratitis capitata</i> ♀ ♀	11.03	2.25	16	c	10.62
<i>Ceratitis capitata</i> ♂ ♂	10.21	2.06	19	c	

Kruskal-Wallis Test ($\chi^2_{(2)} = 188.26$; $P < 0.001$), Mann-Whitney's Test $\alpha = 1\%$; ($P < 0.01$). Means followed by the same letter are not significantly different at 1% of probability.

Lonchaeidae. The host fruits with highest diversity of *Neosilba* species were *Pouteria torta* (Martius) Radlkofer (5 spp.) and *T. catappa* (5 spp.). The most abundant species of Lonchaeidae were *Neosilba zadolicha* Steyskal & McAlpine, and *N. pendula* (Table 3). Although in a old paper, the species of *Neosilba* McAlpine, as *Silba* (Lonchaeidae), were considered as secondary fruit infesters that use the oviposition wounds made by tephritids to gain access to fruit tissue (Souza et al. 1983), results in other papers are in disagree with this information (Malavasi & Morgante 1980; Norrbom & McAlpine 1997; Uchôa & Zucchi 1999; Uchôa et al. 2002; Strikis & Lerena 2009; Uchôa & Nicácio 2010). In fact, in this study a large numbers of fruit/lonchaeid combinations that have no associated tephritids suggest that that generalization can be reconsidered. Of course, *Neosilba* species have a less sclerotized aculeus in comparison to that of *Anastrepha* and *Ceratitis* species (tephritids), but further studies are needed to clarify if the females of *Neosilba* species are able to use muscular power to insert their eggs inside the fruit pulp or if the newly-hatched larvae are able to bore the fruit peel to reach and feed inside the fruits.

The species of *Neosilba* were characterized as polyphagous (*sensu* Aluja & Mangan 2008). *Neosilba zadolicha*, *N. pendula*, and *N. inesperata* were the species with the highest number of host plants (36, 18, and 14 species, respectively). These results show the potential of the species of *Neosilba* as pests of fruits and vegetables in the Neotropical Region (Uchôa et al. 2002; Strikis & Prado 2005; Strikis & Prado 2009; Strikis & Lerena 2009).

The highest rates of infestation by Lonchaeidae (L3/g) occurred in *Duguetia furfuracea* (Annonaceae, Neotropical), *Physalis angulata* (Pantropical), and *Solanum sisymbriifolium* Lamarck (Solanaceae, Neotropical), respectively. *D. furfuracea* was the only host fruit in which a seed-feeding species, a wasp, *Bephratelloides* sp. (Eurytomidae, Hymenoptera), was obtained. Spe-

cies of *Bephratelloides* are common pests on *Annona* seeds in the Neotropics (Peña & Bennet 1995).

The general pattern for Lonchaeidae was that the females of all *Neosilba* species emerged earlier than the conspecific males and on average, the pupal period was 12.21 d, differing significantly from the pupal period of Tephritidae species (Table 4).

Tephritidae *versus* Lonchaeidae. In this study, *Neosilba* species (Lonchaeidae) were the only tephritoids infesting species of Annonaceae, Arecaceae, Lauraceae, Loganiaceae, Rubiaceae, Rutaceae, and Passifloraceae. On the other hand, the tephritids were exclusive only in Clusiaceae and Euphorbiaceae. However, this does not mean that the former plant taxa could not be infested by tephritids nor that Clusiaceae and Euphorbiaceae are never adequate host fruits for lonchaeids either. For example, different *Neosilba* species have been reared from fruits or apical and axillary buds of *Manihot esculenta* Crantz (Euphorbiaceae), *C. capitata* and *A. turpiniae* were reared from fruits of *Citrus sinensis* (Rutaceae) (Uchôa et al. 2002), and *A. fraterculus* was reared from fruits of *Rollinia laurifolia* Schlechtendal (Annonaceae) (Uramoto et al. 2008).

In previous studies, a total of 35 plant species were evaluated for frugivorous insects in areas of Cerrado near the Pantanal, and frugivorous Tephritoidea were found in fruits of 29 species from 16 families. In that study 16 species of *Anastrepha*, *C. capitata* (Tephritidae), and 3 species of Lonchaeidae were recovered from their host fruits (Uchôa et al. 2002). Herein, in addition to the species found by Uchôa et al. (2002), 9 different species of Tephritidae were associated with their host plants (*Anastrepha alveatoides*, *A. distincta* Greene, *A. hastata*, *A. leptozona* Hendel, *A. macrura* Hendel, *A. rheediae* Stone, *A. serpentina* (Wiedemann), *A. undosa*, and *A. zernyi* Lima), and new association was found between *C. capitata* and *Hancornia speciosa* (Apocynaceae).

TABLE 6. NUMBER OF HOST FRUIT SAMPLES (ANGIOSPERMAE) FROM THE SOUTH PANTANAL REGION AND NEIGHBORHOOD, BRAZIL, NUMBER AND RELATIVE PERCENT OF REARED FLIES BY GENERA AND SPECIES (TEPHRITOIDEA: DIPTERA), THEIR PUPATION PERIOD (MEAN IN DAYS), STANDARD DEVIATION, AND COMPARISON OF MEANS (APR 1998 TO AUG 2000, OCT 2003, AND OCT 2004).

Species of frugivorous Flies	N° of samples	N° of individuals	(%) Relative occurrence for Genera	(%) Relative occurrence for species	Mean period of pupation (days)	Standard Deviation	Comparison of the means
<i>Anastrepha hastata</i>	4	25	0.74	0.31	21.92	0.59	A
<i>A. zernyi</i>	8	7	0.21	0.09	20.90	4.43	A
<i>A. fraterculus</i>	16	6	0.18	0.07	20.38	1.97	A
<i>A. rheediae</i>	5	15	0.44	0.19	19.50	278	A b
<i>A. undosa</i>	4	436	12.83	5.40	17.40	2.73	b c
<i>A. striata</i>	19	167	4.92	2.07	16.83	2.52	b c
<i>A. turpiniae</i>	18	16	0.47	0.20	16.10	2.25	c d E
<i>A. leptozona</i>	6	64	1.88	0.79	16.07	1.89	c d E
<i>A. obliqua</i>	25	207	6.09	2.56	15.69	1.64	c d E f
<i>A. albeatoides</i>	2	503	14.81	6.22	15.36	0.35	c d E f
<i>A. sororcula</i>	43	1,301	38.3	16.10	14.84	2.64	c d E f
<i>A. serpentina</i>	12	97	2.86	1.20	13.95	0.93	d E f g
<i>A. zenildae</i>	22	553	16.28	6.84	13.61	1.44	E f g H
<i>Neosilba inesperata</i>	50	443	24.99	5.48	13.43	3.39	E f g H
<i>Neosilba MSP1</i>	3	4	0.23	0.05	13.39	1.25	E f g H
<i>Neosilba certa</i>	32	59	3.33	0.73	12.74	1.89	f g H
<i>Neosilba zadolicha</i>	130	1,037	58.49	12.83	12.06	2.21	g H
<i>Neosilba pendula</i>	58	127	7.16	1.57	11.85	1.59	g H
<i>Neosilba glaberrima</i>	45	103	5.81	1.27	11.48	1.42	g H
<i>Ceratitis capitata</i> ♀	37	2,911	100	36.02	11.03	2.25	H
Total	177	8,080	-	100	—	—	

ANOVA = 1%; Duncan's Test (P < 0.05); Means followed by the same letter are not significantly different.

Anastrepha was the most abundant Tephritoidea genus reared from naturally infested host fruits in this study. A similar pattern was found by Malavasi et al. (1980) in a survey including 55 fruit species sampled in several states of Brazil. Herein, *A. sororcula* and *A. zenildae* were the most frequent (percentage of individual of a fly species in relation to the total number of individuals of all fruit fly species) and abundant (relationship of the number of individuals of each species of fruit fly by each species of host fruits) *Anastrepha* species.

The species of *Neosilba* were more generalists. They attacked more hosts (48) than the Tephritidae (30). Similar results were obtained in a previous survey in the Cerrado of Mato Grosso do Sul, Brazil (Uchôa et al. 2002).

Larval viability was lower for tephritids (59.73%) than for lonchaeids (60.72%) (Table 4). In this study the larval viability of the tephritids was higher than that reported by Malavasi & Morgante (1980) (10%). This suggests that the methodology herein employed is adequate for obtaining frugivorous insects.

The higher viability of the pre-pupal larval of Lonchaeidae species is probably due to the lower parasitism rates in this taxon than in Tephritidae species (M. A. Uchôa et al., unpublished data). No adult emerged from 40.05% of the pre-pupal larval of Tephritoidea. Despite of the lower viability (59.73%) of *Anastrepha* spp. and *Ceratitis capitata* larvae, their relative abundance in fruit infestation was higher (78.1%), in comparison with Lonchaeidae species (21.9%, Table 4).

The genera of Tephritoidea showed a highly significant difference in their pupation period. The species of *Anastrepha* had the longest pupation period, followed by *Neosilba* spp., and *C. capitata*, respectively. The males of all Tephritidae species (16 *Anastrepha* spp. and *C. capitata*) emerged before their conspecifics females. Although mating systems of the Lonchaeidae must be more ancient in relation to that of tephritids, because the lonchaeids are among the most primitive Tephritoidea, the opposite pattern was found in the species of *Neosilba*. All females emerged earlier than the conspecifics males, although the difference was not statistically significant (Table 5).

The average pupation period for the 20 species of frugivorous flies evaluated in this study showed significant differences among 3 groups of species: (*Anastrepha hastata* Stone to *Anastrepha fraterculus* (Wiedemann)), (*Anastrepha rheediae* Stone to *Anastrepha striata* Schiner), and (*Neosilba certa* (Walker) to *Ceratitis capitata*), ranging from 21.92 d for *A. hastata* to 11.03 d for *C. capitata* (Table 6). In this analysis, *Anastrepha macrura* Hendel (22 d) (1 ♀), *Anastrepha* n. sp. aff. *pickeli* Lima (18.33 d) (4 ♀), *Anastrepha distincta* Greene (13 d) (3 ♀), *Lonchaea* sp. (7.0 d) (1 ♀), and

Neosilba pradoi Strikis & Lerena (16.80 d) (1 ♂), were not included, because each of them occurred in fewer than 3 fruit samples.

Facholi-Bendassolli & Uchôa (2006) studied the reproductive behavior of *Anastrepha sororcula* and found a marked protandry: males reached sexual maturity at an average of 11.75 d after emergence, and females became sexually mature, on average, 19 d after emergence. There are no reports on the sexual maturation of *Neosilba* species, but as evidenced in this study, it is probable that the females of *Neosilba* reached sexual maturity earlier than the conspecific males, a case of protogyny. However, a more specific study is necessary to answer this question satisfactorily. The asynchronous development between males and females of these insects may play an important evolutionary role. If males and females of the same progeny (offspring) reach sexual maturity at different times in nature, the chance of inbred mating decreases, which increases the genetic variability of the species (Futuyma 1992).

The average for the pupal period (interval between L3 exiting from fruits and emergence of adults) showed significant differences between the evaluated genera: *Anastrepha* (15.85 d), *Ceratitis* (10.62 d, mean between ♂♂ and ♀♀) (Tephritidae), and *Neosilba* (12.21 d, mean between ♂♂ and ♀♀) (Table 5)). However, when this period was compared between the species of the 3 genera, there is no standard pattern for each genus, because there are strong variations among the congeneric species (Table 6). We do not know if this difference were influenced by the differences in ambient conditions from the sites of fruits collections.

On average, the duration of the pupal period showed highly significant differences between *C. capitata* and the *Anastrepha* species. *Ceratitis capitata* emerged after 10.62 d and the *Anastrepha* species after 15.85 d. The Tephritidae with the highest mean for pupal period were *Anastrepha hastata* Stone (21.92 d) and *Anastrepha zernyi* Lima (20.90 d). *Ceratitis capitata* (♀♀) and *A. zenildae* had the shortest pupal periods of 11.03 and 13.61 d, respectively (Table 6). The pupal period for *C. capitata* obtained in this study (10.62 d) is similar to that found by Papadopoulos et al. (2002), 10 d under laboratory conditions (25°C 2 and 65% 5 RH (relative humidity)).

ACKNOWLEDGMENTS

We thank the botanists: Ubirazilda Maria Rezende from the Herbarium of the Universidade Federal de Mato Grosso do Sul (UFMS) at Campo Grande-MS, and José Rubens Pirani from the Herbarium of the Universidade de São Paulo (USP, São Paulo), for identification of plant species; Pedro Carlos Strikis (UNICAMP, Campinas-SP), for identification of Lonchaeidae; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), for the grant award to

J. N. Nicácio; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), for the postdoctoral fellowship (Process N°. 1030/09-4 BEX) to M. A. Uchôa, and Allen L. Norrbom and James L. Nation, for valuable suggestions in the manuscript, making it more comprehensive.

REFERENCES CITED

- ALUJA, M., AND MANGAN, R. L. 2008. Fruit fly (Diptera: Tephritidae) host status determination: critical conceptual, methodological, and regulatory considerations. *Annu. Rev. Entomol.* 53: 473-502.
- CALKINS, C. O., AND MALAVASI, A. 1995. Biology and control of fruit fly (*Anastrepha*) in tropical and temperate fruit. *Rev. Brasileira Frut.* 17: 36-45.
- CANAL, N. A. 2010. New species and records of *Anastrepha* Schiner (Diptera: Tephritidae) from Colombia. *Zootaxa* 2425:32-44.
- CARVALHO, F. M. N., DE MARCO JR., P., AND FERREIRA, L. G. 2009. The Cerrado into-pieces: habitat fragmentation as a function of landscape use in the savannas of Central Brazil. *Biol. Conserv.* 142: 1392-1403.
- DAWSON, B., AND TRAPP, R. G. 2003. *Bioestatística básica e Clínica*. 3 ed. Editora McGraw-Hill-Interamericana. São Paulo. 364 pp.
- FACHOLI-BENDASSOLLI, M. C., AND UCHÔA, M. A. 2006. Comportamento sexual de *Anastrepha sororcula* Zucchi (Diptera, Tephritidae) em laboratório. *Rev. Brasileira Entomol.* 50: 406-412.
- FOOTE, R. H. 1980. Fruit Fly Genera South of the United States (Diptera: Tephritidae). USDA Tech. Bull. 1600. Washington-DC, USA. 79 pp.
- FUTUYMA, D. J. 1992. *Biologia Evolutiva*. 2 ed. Sociedade Brasileira de Genética (SBG)-CNPq. Ribeirão Preto. 631 pp.
- GARCIA, F. R. M., SAVARIS, M., AND BOGUS, G. M. 2008. First host plant record for *Anastrepha elegans* (Diptera: Tephritidae). *Florida Entomol.* 91: 317-318.
- GALATI, E. A. B., NUNES, V. L. B., BOGGIANI, P. C., DORVAL, M. E. C., CRISTALDO, G., ROCHA, H. C., OSHIRO, E. T., AND DAMASCENO-JUNIOR, G. A. 2006. Phlebotomines (Diptera: Psychodidae) in forested areas of the Serra da Bodoquena, state of Mato Grosso do Sul, Brazil. *Mem. Inst. Oswaldo Cruz* 101: 175-193.
- JUNK, W. J., DA CUNHA, C. N., WANTZEN, K. M., PETERMANN, P., STRÜSSMANN, C., MARQUES, M. I., AND ADIS, J. 2006. Biodiversity and its conservation in the Pantanal of Mato Grosso, Brazil. *Aquat. Sci.* 68: 378-309.
- LIMA, A. da C. 1934. Moscas-das-frutas do gênero *Anastrepha* Schiner, 1868. *Mem. Inst. Oswaldo Cruz* 28: 487-575.
- MALAVASI, A., AND MORGANTE, J. S. 1980. *Biologia de "moscas-das-frutas" (Diptera, Tephritidae). II: Índices de infestação em diferentes hospedeiros e localidades*. *Rev. Brasileira Biol.* 40: 17-24.
- MALAVASI, A., MORGANTE, J. S., AND ZUCCHI, R. A. 1980. *Biologia de "moscas-das-frutas" (Diptera, Tephritidae). I: Lista de hospedeiros e ocorrência*. *Rev. Brasileira Biol.* 40: 09-16.
- MCALPINE, J. F., AND STEYSKAL, G. C. 1982. A revision of *Neosilba* McAlpine with a key to the world genera of Lonchaeidae (Diptera). *Canadian Entomol.* 114: 105-137.
- MAROCO, J. 2007. *Análise Estatística com Utilização do SPSS*. 3 ed. Edições Sílabo. Lisboa. 822p.
- MYERS, N., MITTERMEIER, R. A., MITTERMEIER, C. G., DA FONSECA, G. A. B., AND KENT, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858.
- NORRBOM, A. L., AND MCALPINE, J. F. 1997. A review of Neotropical species of *Dasiops* Rondani (Diptera: Lonchaeidae) attacking *Passiflora* (Passifloraceae). *Mem. Entomol. Soc. Washington* 18: 189-211.
- NORRBOM, A. L., AND KORYTKOWSKI, C. A. 2009. A revision of the *Anastrepha robusta* species group (Diptera: Tephritidae). *Zootaxa* 2182: 1-91.
- NORRBOM, A. L., AND KORYTKOWSKI, C. A. 2011. New species of and taxonomic notes on *Anastrepha* (Diptera: Tephritidae). *Zootaxa* 2740: 1-23.
- NORRBOM, A. L., AND UCHÔA, M. A. 2011. New species and records of *Anastrepha* (Diptera: Tephritidae) from Brazil. *Zootaxa* 2835: 61-67.
- PAPADOPOULOS, N. T., KATSOYANNOS, I. B., AND CAREY, R. J. 2002. Demographic parameters of the Mediterranean fruit fly (Diptera: Tephritidae) reared in apples. *Ann. Entomol. Soc. America* 95: 564-569.
- PEÑA, J. E., AND BENNET, F. D. 1995. Arthropods associated with *Annona* spp. in the Neotropics. *Florida Entomol.* 78: 329-349.
- PESTANA, M. H., AND GAGEIRO, J. N. 2008. *Análise de Dados para Ciências Sociais: A Complementaridade do SPSS*. 5 ed. Edições Sílabo. Lisboa. 692 pp.
- POTT, A., AND POTT, V. J. 1994. *Plantas do Pantanal*. Embrapa, Brasília-DF. 320 pp.
- RADAM BRASIL. 1982. Ministério das Minas e Energia, Departamento Nacional de Produção Mineral. Projeto RADAMBRASIL, V. 28, Folha SF 21, Campo Grande: Geologia, geomorfologia, pedologia, vegetação e uso potencial da terra. Rio de Janeiro. 416 pp.
- SOUZA, H. M. L., CYTRYNOWICZ, M., MORGANTE, J. S., AND PAVAN, O. H. O. 1983. Occurrence of *Anastrepha fraterculus* (Wied.), *Ceratitis capitata* (Wied.) (Diptera, Tephritidae) and *Silba* spp. (Diptera, Lonchaeidae) eggs in oviposition bores on three host fruits. *Rev. Brasileira Entomol.* 27: 191-195.
- STONE, A. 1942. The Fruit Flies of the Genus *Anastrepha*. USDA Misc. Publ. Washington-DC, USA. 112 pp.
- STRIKIS, P. C., AND PRADO, A. P. 2005. A new species of the genus *Neosilba* (Diptera: Lonchaeidae). *Zootaxa* 828: 1-4.
- STRIKIS, P. C., AND PRADO, A. P. 2009. Lonchaeidae associados a frutos de nêspira, *Eryobotria japonica* (Thunb.) Lindley (Rosaceae), com descrição de uma espécie nova de *Neosilba* (Diptera: Tephritoidea). *Arq. Inst. Biol.* 76: 49-54.
- STRIKIS, P. C., AND LERENA, M. L. M. 2009. A new species of *Neosilba* (Diptera: Lonchaeidae) from Brazil. *Iheringia, Sér. Zool.* 99: 273-275.
- UCHÔA, M. A., AND NICÁCIO, J. N. 2010. New records of Neotropical fruit flies (Tephritidae), lance flies (Lonchaeidae) (Diptera: Tephritoidea), and their host plants in the South Pantanal and adjacent areas, Brazil. *Ann. Entomol. Soc. America* 103:723-733.
- UCHÔA, M. A., AND ZUCCHI, R. A. 1999. Metodología de colecta de Tephritidae y Lonchaeidae frugívoros (Diptera: Tephritoidea) y sus parasitoides (Hymenoptera). *An. Soc. Entomol. Brasil* 28: 601-610.

- UCHÔA, M. A., OLIVEIRA, I., MOLINA, R. M. S., AND ZUCCHI, R. A. 2002. Species diversity of frugivorous flies (Diptera: Tephritoidea) from hosts in the cerrado of the state of Mato Grosso do Sul, Brazil. *Neotropical Entomol.* 31:515-524.
- URAMOTO, K., MARTINS, D. S., AND ZUCCHI, R. A. 2008. Fruit flies (Diptera, Tephritidae) and their associations with native host plants in a remnant area of the highly endangered Atlantic Rain Forest in the State of Espírito Santo, Brazil. *Bull. Entomol. Res.* 98:457-466.
- URAMOTO, K., AND ZUCCHI, R. A. 2010. New species of *Anastrepha* Schiner (Diptera, Tephritidae) from remnant area of the Atlantic Rain Forest and surroundings in the state of Espírito Santo, Brazil. *Zootaxa* 2535: 49-60.
- ZUCCHI, R. A. 2008. Fruit flies in Brazil—*Anastrepha* species their host plants and parasitoids. Available at: www.lea.esalq.usp.br/anastrepha/, updated on May 04, 2011. <Accessed on May 26, 2011>.