



Species of Anastrepha (Diptera: Tephritidae), Their Host Plants, and Parasitoids in Small Fruit Production Areas in the State of Amapá, Brazil

Authors: Lemos, Lailson do Nascimento, Deus, Ezequiel da Glória de, Nascimento, Danilo Baia do, Jesus-Barros, Cristiane Ramos de, Costa-Neto, Salustiano Vilar da, et al.

Source: Florida Entomologist, 100(2) : 403-410

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0201>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Species of *Anastrepha* (Diptera: Tephritidae), their host plants, and parasitoids in small fruit production areas in the state of Amapá, Brazil

Lailson do Nascimento Lemos¹, Ezequiel da Glória de Deus¹, Danilo Baia do Nascimento¹, Cristiane Ramos de Jesus-Barros², Salustiano Vilar da Costa-Neto³, and Ricardo Adaime^{1,2,*}

Abstract

Fruit flies and associated parasitoids were determined in native and introduced fruit species in the state of Amapá, Brazil. Fruits were collected every 30 d, from Jan to Dec 2012, on 3 small farms in the municipalities of Mazagão, Porto Grande, and Santana. We collected 412 samples (78 species of plants belonging to 32 families and consisting of 4,554 fruits weighing 323.4 kg) and obtained 5,252 *Anastrepha* (Diptera: Tephritidae) puparia from 107 infested samples (20 plant species from 13 botanical families). We report here the occurrence of 11 species of *Anastrepha* and 5 species of parasitoids (4 Braconidae and 1 Figitidae species; both families in the order Hymenoptera). New hosts for *Anastrepha distincta* Greene, *Anastrepha leptozona* Hendel, *Anastrepha obliqua* (Macquart), and *Anastrepha serpentina* (Wiedemann) were documented in Brazil.

Key Words: fruit fly; infestation; host fruit; Amazon

Resumo

Moscas-das-frutas e parasitoides associados foram reportados em espécies vegetais frutíferas nativas e introduzidas no estado do Amapá, Brasil. Frutos foram coletados a cada 30 dias, de janeiro a dezembro de 2012, em 3 pequenas propriedades agrícolas nos municípios de Mazagão, Porto Grande e Santana. Nós coletamos 412 amostras (78 espécies de plantas pertencentes a 32 famílias, consistindo de 4.554 frutos, pesando 323,4 kg) e obtivemos 5.252 pupários de *Anastrepha*, a partir de 107 amostras infestadas (20 espécies vegetais de 13 famílias botânicas). Reportamos a ocorrência de 11 espécies de *Anastrepha* e 5 de parasitoides (4 Braconidae e 1 Figitidae; ambos Hymenoptera). Novos hospedeiros para *Anastrepha distincta* Greene, *Anastrepha leptozona* Hendel, *Anastrepha obliqua* (Macquart) e *Anastrepha serpentina* (Wiedemann) foram documentados para o Brasil.

Palavras Chave: moscas-das-frutas; infestação; fruto hospedeiro; Amazônia

Fruit flies (Diptera: Tephritidae) are among the principal pests in agriculture (Malavasi 2009). In addition to the direct damage caused by the development of larvae inside the fruits, there are significant monetary losses caused by quarantine restrictions for certain species of fresh fruit destined for export (Carvalho et al. 2000). Among the Tephritidae, the genus *Anastrepha* Schiner, endemic to the Neotropical region, has greatest economic impact in the Americas. *Anastrepha* is quite diverse, with 235 species described, and is widely distributed in the American tropical and subtropical regions (Aluja 1994; Uramoto et al. 2008; Uchôa & Nicácio 2010). In Brazil, 120 species have already been reported (Zucchi 2008).

In the Brazilian Amazon, 76 *Anastrepha* species are known (Adaime et al. 2016). Of these, 5 are considered of economic importance: *Anastrepha fraterculus* (Wiedemann), *Anastrepha obliqua* (Macquart), *Anastrepha pseudoparallela* (Loew), *Anastrepha striata* Schiner, and *Anastrepha zenildae* Zucchi (Uramoto & Zucchi 2009; Dutra et al. 2013; Adaime et al. 2016). In Amapá, 37 species of *Anastrepha* are already

documented, making this state the second one in species richness in the Amazon region (Zucchi 2008).

Despite the economic importance of Tephritidae, the association of fruit fly species with host plants has not been established for most of the known species in the Brazilian Amazon (Zucchi et al. 2011a). The identification of the host fruits is essential for the establishment of an adequate and efficient management program for the pest species (Jesus-Barros et al. 2012). From fruit collections, it is possible to determine the rates of fruit infestation by tephritids and the parasitism rates of the tephritids by parasitoids (Silva et al. 2011a). These relationships can also provide information on the life history of fruit fly species because the host plant, together with temperature, humidity and natural enemies, can affect larval development, adult fecundity, and survival rate. In addition, the understanding of these relationships provides information on the structure of the fruit fly community in the Brazilian Amazon, because the host plant may have a direct effect on the distribution and abundance of *Anastrepha* species. It is worth not-

¹Universidade Federal do Amapá, Programa de Pós-Graduação em Biodiversidade Tropical, Macapá, Amapá, 68902-280, Brazil; E-mail: l.n.lemos@hotmail.com (L. N. L.), egd_bio@hotmail.com (E. G. D.), danilo.baia@hotmail.com (D. B. N.)

²Embrapa Amapá, Macapá, Amapá, 68903-419, Brazil; E-mail: cristiane.jesus@embrapa.br (C. R. J.-B.), ricardo.adaime@embrapa.br (R. A.)

³Instituto de Pesquisas Científicas e Tecnológicas do Amapá, Macapá, Amapá, 68900-000, Brazil; E-mail: salucostaneto@gmail.com (S. V. C.-N.)

*Corresponding author; E-mail: ricardo.adaime@embrapa.br (R. A.)

ing that tephritids show different degrees of specialization on species of plants native to the American continent, ranging from monophagy to polyphagy (Duyck et al. 2004; Aluja & Mangan 2008; Malavasi 2009; Hafsi et al. 2016).

Fruit production areas in the Brazilian Amazon, especially in the state of Amapá, are mostly characterized by the use of agroforestry systems and urban and rural backyards usually surrounded by native vegetation. This design can favor infestation by fruit flies because there are about 200 fruit species in the Brazilian Amazon, of which half are native and have the potential to host tephritid species. In addition, the availability of cultivated fruits, together with the high diversity of native plant species that fruit throughout the year, may allow the succession of hosts and the maintenance of fruit fly populations (Silva & Ronchi-Teles 2000).

Host availability is one of the main factors affecting the population dynamics of fruit flies. Jesus-Barros et al. (2012), in a study carried out in the state of Amapá, observed that the highest rates of infestation by *Anastrepha* were recorded in native fruits of the Amazon region. Lemos et al. (2015), in a study carried out in the southeast region of Brazil, found that the population dynamics of *Anastrepha* in guava orchards was influenced by the availability of alternative hosts in the areas adjacent to the orchard.

The action of natural enemies is another biotic factor, as well as host availability and phenology, which has a strong influence on the population regulation of tephritid species (Leal et al. 2009; Aluja et al. 2012). In the Neotropical region, the braconid parasitoids are among the principal mortality factors of *Anastrepha* (Carvalho et al. 2010). *Doryctobracon areolatus* (Szépligeti) (Hymenoptera: Braconidae) has been shown to be the most abundant and widely distributed parasitoid in Brazil (Jesus-Barros et al. 2012). Nonetheless, natural parasitism in the various fruit producing regions in Brazil is variable because it is influenced by the host plant species, by the host fly, and by the place and time of collection (Ovruski et al. 2000; Bittencourt et al. 2012).

The aim of this research was to identify species of *Anastrepha* and their host plants and associated parasitoids in fruit production areas of 3 municipalities in the state of Amapá in the Brazilian Amazon.

Materials and Methods

STUDY SITE

We conducted this study on 3 farms located in the municipalities of Mazagão, Porto Grande, and Santana, state of Amapá, Brazil. All 3 properties contained commercial orchards with adjoining areas of native forest.

The area in Mazagão (0.1°S, 51.25°W) exhibits native floodplain forest vegetation, which undergoes daily flooding due to tides. Common species include *Platymiscium duckei* Huber (Fabaceae), *Virola surinamensis* (Rol.) Warb. (Myristicaceae), *Calycophyllum spruceanum* (Benth.) Hook. f. ex K. Schum. (Rubiaceae), and *Carapa guianensis* Aubl. (Meliaceae). The commercial orchard area occupies an area of approximately 10 ha, with cultivated fruit trees including Tahiti lime (*Citrus aurantifolia* Swingle 'Tahiti'; Rutaceae), passionfruit (*Passiflora edulis* Sims; Passifloraceae), and soursop (*Annona muricata* L.; Annonaceae) grown to make fruit concentrate or to be sold fresh.

The area in Porto Grande (0.6°N, 1.45°W) exhibits dense forest vegetation with numerous tree species including *Protium* species (Bursaceae), *Caryocar villosum* (Aubl.) Pers. (Caryocaraceae), and *Dipteryx odorata* (Aubl.) Willd. (Fabaceae). The property covers approximately 100 ha, of which 30% is used for the cultivation of fruit trees, mainly hog plum (*Spondias mombin* L.; Anacardiaceae), araza (*Eugenia stipitata* McVaugh; Myrtaceae), and soursop (*A. muricata*).

The area in Santana (0.03°S, 51.21°W) exhibits secondary forest vegetation with *Eschweilera tenuifolia* Miers (Lecythidaceae), *Ficus pertusa* L. f. (Moraceae), and *Pourouma guianensis* Aubl. (Urticaceae) as predominant species. The commercial orchard is approximately 20 ha, and the main species are acerola (*Malpighia emarginata* DC; Malpighiaceae), guava (*Psidium guajava* L.; Myrtaceae), and passionfruit (*P. edulis*).

As classified by Köppen, the climate type at the study sites is Am, characterized as hot and humid with a short dry season in the spring. Rainfall is distributed between 2 well-defined periods: a rainy season (Jan–Jul), which receives about 80% of all annual rainfall in the area, and a dry season (Aug–Dec). Temperatures are high throughout the year, with an annual average of 25 to 27 °C (low of 22 °C and high of 34 °C) (IBGE 2010).

EXPERIMENTAL PROCEDURES

Samples were obtained from fruit crops and adjacent areas of native vegetation. Once a month from Jan to Dec 2012, fruits were collected from various plant species either directly from the plant or by retrieving recently fallen fruit. Samples were transported in stackable plastic crates to the Entomology Laboratory at Embrapa Amapá, where the fruits were weighed on a digital scale. For preparation of the samples, the fruits were processed and individualized as per the sample size criterion established by Silva et al. (2011a): a) 15 units, for small fruits; b) 10 units, for medium-sized fruits; and c) 3 units, for large or elongated fruits. In each sample, each individual fruit was considered as 1 sub-sample.

The collected fruits were placed either individually in tubular, transparent plastic vials (8 × 6 cm DH) with vented lids lined with organza or, in the case of large or elongated fruits, on rectangular plastic trays (33 × 18 × 6 cm LWH) also covered with organza held in place by rubber bands. Both types of containers included a thin layer of sterilized sand as a substrate for pupation.

Every 3 d, the sand was examined, and tephritid puparia were removed with spatulas. The puparia were transferred to transparent plastic vials (8 cm diameter) with vented lids covered with organza and containing a thin layer of moistened vermiculite. The adult insects that emerged were killed, transferred to microcentrifuge tubes containing 70% ethanol, and labeled for subsequent identification.

The insects obtained were identified according to Zucchi et al. (2011b) and Marinho et al. (2011). Wild plant species were identified with identification keys and by comparison with specimens available at the Herbário Amapaense, the herbarium at the Amapá State Institute for Scientific and Technological Research (Macapá, Amapá, Brazil).

The following infestation rates were calculated: 1) percentage of fruits infested = (number of fruits infested ÷ number of fruits collected) × 100; 2) number of puparia per fruit = number of puparia obtained ÷ number of infested fruits in the sample.

Results

During the sampling period, 412 fruit samples (4,554 fruits; 323.4 kg) representing 78 plant species (51 native and 27 introduced) were obtained from 32 botanical families (Table 1). Infestation was observed in 107 (26.0%) collected samples, representing 20 plant species in 13 botanical families (Tables 1 and 2). In total, 5,252 puparia were obtained, from which emerged specimens of 11 species of *Anastrepha* and 5 species of parasitoids (4 Braconidae and 1 Figitidae species; both Hymenoptera). The species of *Anastrepha* obtained were: *Anastrepha antunesi* Lima, *Anastrepha coronilli* Carrejo & González, *Anastrepha*

Table 1. Number of fruit samples and individual fruits collected, and number infested by *Anastrepha* species, from 3 municipalities in the state of Amapá, Brazil (Jan–Dec 2012).

Scientific name	Common name [Portuguese]	Family	Origin N / I	Mazagão				Porto Grande				Santana			
				CS	CF	IS	IF	CS	CF	IS	IF	CS	CF	IS	IF
<i>Alibertia edulis</i> (Rich.) A. Rich.	Puruí	Rubiaceae	N	0	0	0	0	2	30	0	0	0	0	0	0
<i>Anacardium occidentale</i> L.	Caju	Anacardiaceae	I	1	10	0	0	0	0	0	0	2	20	0	0
<i>Annona mucosa</i> Jacq.	Biribá	Annonaceae	N	0	0	0	0	3	9	0	0	0	0	0	0
<i>Annona muricata</i> L.	Graviola	Annonaceae	I	1	3	0	0	4	12	0	0	2	6	1	1
<i>Artocarpus altifolius</i> (Parkinson) Fosberg	Fruta-pão	Moraceae	I	0	0	0	0	4	12	0	0	0	0	0	0
<i>Artocarpus heterophyllus</i> Lam.	Jaca	Moraceae	I	0	0	0	0	2	6	0	0	3	9	0	0
<i>Astrocaryum aculeatum</i> G. Mey.	Tucumã	Arecaceae	N	0	0	0	0	0	0	0	0	1	15	0	0
<i>Astrocaryum murumuru</i> Mart.	Murumuru	Arecaceae	N	6	60	0	0	0	0	0	0	0	0	0	0
<i>Attalea maripa</i> (Aubl.) Mart.	Inajá	Arecaceae	N	0	0	0	0	1	15	0	0	0	0	0	0
<i>Attalea phalerata</i> Mart. ex Spreng.	Urucuri	Arecaceae	N	7	70	0	0	0	0	0	0	0	0	0	0
<i>Averrhoa carambola</i> L.	Carambola	Oxalidaceae	I	0	0	0	0	9	90	0	0	0	0	0	0
<i>Bactris gasipaes</i> Kunth	Pupunha	Arecaceae	I	0	0	0	0	2	30	0	0	0	0	0	0
<i>Bactris maraja</i> Mart.	Marajá	Arecaceae	N	1	15	0	0	0	0	0	0	0	0	0	0
<i>Bellucia egensis</i> (DC.) Penneys, Michelangeli, Judd, Almeida	Verde-peludo	Melastomataceae	N	0	0	0	0	3	45	0	0	4	60	0	0
<i>Bellucia grossularioides</i> (L.) Triana	Goiaba-de-anta	Melastomataceae	N	0	0	0	0	2	30	2	6	0	0	0	0
<i>Byrsonima crassifolia</i> (L.) Kunth	Muruci	Malpighiaceae	N	0	0	0	0	0	0	0	0	5	75	0	0
<i>Calophyllum brasiliensis</i> Cambess.	Jacareúba	Calophyllaceae	N	1	15	0	0	0	0	0	0	0	0	0	0
<i>Capsicum chinense</i> Jacq.	Pimentinha	Solanaceae	I	8	120	0	0	1	15	0	0	3	45	0	0
<i>Carica papaya</i> L.	Mamão	Caricaceae	I	1	3	0	0	5	15	0	0	6	90	0	0
<i>Caryocar glabrum</i> (Aubl.) Pers.	Piquiarana	Caryocaraceae	N	0	0	0	0	1	10	0	0	0	0	0	0
<i>Cayaponia ferruginea</i> Gomes-Klein	Fruta de cutia	Curcubitaceae	N	0	0	0	0	0	0	0	0	1	15	0	0
<i>Cheiloclinium cognatum</i> (Miers) A.C. Sm.	Bacupari	Celastraceae	N	2	30	1	1	0	0	0	0	3	45	1	1
<i>Cissus amapaensis</i> Lombardi	Uva-do-mato	Vitaceae	N	0	0	0	0	0	0	0	0	2	30	0	0
<i>Citrus aurantifolia</i> Swingle 'Tahiti'	Limão taiti	Rutaceae	I	9	90	0	0	0	0	0	0	3	30	0	0
<i>Citrus aurantium</i> L.	Laranja-da-terra	Rutaceae	I	0	0	0	0	10	100	0	0	0	0	0	0
<i>Citrus latifolia</i> Tanaka	Limão galego	Rutaceae	I	0	0	0	0	0	0	0	0	2	20	0	0
<i>Citrus limonia</i> Osbeck	Limão-cravo	Rutaceae	I	8	80	0	0	8	80	0	0	11	110	0	0
<i>Citrus reticulata</i> Blanco	Tangerina	Rutaceae	I	2	20	0	0	4	40	0	0	5	50	0	0
<i>Citrus sinensis</i> (L.) Osbeck	Laranja	Rutaceae	I	4	40	0	0	0	0	0	0	1	10	0	0
<i>Coffea arabica</i> L.	Café	Rubiaceae	I	0	0	0	0	0	0	0	0	3	45	0	0
<i>Combretum laxum</i> Jacq.	Pombeiral	Combretaceae	N	0	0	0	0	2	30	0	0	0	0	0	0
<i>Conceveiba guianensis</i> Aubl.	Arraieira	Euphorbiaceae	N	0	0	0	0	0	0	0	0	1	15	0	0
<i>Cucumis anguria</i> L.	Maxixe	Curcubitaceae	N	5	75	0	0	1	15	0	0	1	15	0	0
<i>Cucumis sativus</i> L.	Pepino	Curcubitaceae	I	5	15	0	0	0	0	0	0	1	3	0	0
<i>Eschweillera tenuifolia</i> (O. Berg) Miers [flower bud]	Matá-matá	Lecythidaceae	N	0	0	0	0	0	0	0	0	4	60	0	0
<i>Eugenia stipitata</i> McVaugh	Araçá-boi	Myrtaceae	N	0	0	0	0	11	110	11	102	0	0	0	0
<i>Ficus pertusa</i> L. f.	Apuí	Moraceae	N	2	30	0	0	2	30	0	0	1	15	0	0
<i>Geissospermum sericeum</i> Miers	Quinarana	Apocynaceae	N	0	0	0	0	2	20	0	0	0	0	0	0
<i>Guarea guidonia</i> (L.) Sleumer	Jatáuba	Meliaceae	N	0	0	0	0	1	15	0	0	0	0	0	0
<i>Gustavia augusta</i> L.	Jenipaparana	Lecythidaceae	N	0	0	0	0	0	0	0	0	1	10	0	0
<i>Inga edulis</i> Mart.	Ingá-cipó	Fabaceae	N	2	6	2	6	0	0	0	0	5	15	3	15
<i>Licania laxiflora</i> Fritsch	Anoerá	Chrysobalanaceae	N	5	45	0	0	0	0	0	0	1	15	1	1

N = Native; I = Introduced; CS = collected samples; CF = collected fruits; IS = infested samples; IF = infested fruits.

Table 1. (Continued) Number of fruit samples and individual fruits collected, and number infested by *Anastrepha* species, from 3 municipalities in the state of Amapá, Brazil (Jan–Dec 2012).

Scientific name	Common name [Portuguese]	Family	Origin N / I	Mazagão					Porto Grande					Santana				
				CS	CF	IS	IF	CS	CF	IS	IF	CS	CF	IS	IF	CS	CF	IS
<i>Malpighia emarginata</i> DC	Acerola	Malpighiaceae	I	1	15	1	4	4	60	4	15	9	135	9	80			
<i>Mammea americana</i> L.	Abricó	Clusiaceae	I	0	0	0	0	3	30	0	0	0	0	0	0			
<i>Mangifera indica</i> L.	Manga comum	Anacardiaceae	I	2	20	0	0	1	10	0	0	2	20	0	0			
<i>Mangifera indica</i> L. 'Tommy Atkins'	Manga cavalo	Anacardiaceae	I	0	0	0	0	7	21	0	0	0	0	0	0			
<i>Manicaria saccifera</i> Gaertn	Bussu	Arecaceae	N	1	15	0	0	0	0	0	0	0	0	0	0			
<i>Manihot esculenta</i> Crantz	Mandioca	Euphorbiaceae	N	0	0	0	0	1	15	1	1	1	15	0	0			
<i>Manilkara elata</i> (Alemão ex Miq.) Monach	Maçaranduba	Sapotaceae	N	2	30	0	0	0	0	0	0	0	0	0	0			
<i>Manilkara zapota</i> (L.) P. Royen	Sapotilha	Sapotaceae	I	0	0	0	0	12	120	6	38	0	0	0	0			
<i>Metrodorea flovida</i> K. Krause	Laranjinha	Rutaceae	N	4	60	0	0	0	0	0	0	0	0	0	0			
<i>Oenocarpus bacaba</i> Mart.	Bacaba	Arecaceae	N	1	15	0	0	1	15	0	0	6	90	1	1			
<i>Parinari excelsa</i> Sabine	Isqueira	Chrysobalanaceae	N	1	15	0	0	0	0	0	0	0	0	0	0			
<i>Passiflora edulis</i> Sims [flower bud]	Maracujá	Passifloraceae	N	5	75	0	0	0	0	0	0	3	45	0	0			
<i>Passiflora edulis</i> Sims	Maracujá	Passifloraceae	N	8	80	0	0	0	0	0	0	2	20	0	0			
<i>Persea americana</i> Mill.	Abacate	Lauraceae	I	0	0	0	0	4	12	0	0	4	12	1	1			
<i>Physalis angulata</i> L.	Camapu	Solanaceae	I	1	15	0	0	0	0	0	0	0	0	0	0			
<i>Plinia cauliflora</i> (Mart.) Kausel	Jaboticaba	Myrtaceae	I	0	0	0	0	1	15	1	6	0	0	0	0			
<i>Pourouma guianensis</i> Aubl.	Mapatirana	Urticaceae	N	0	0	0	0	1	15	1	6	0	0	0	0			
<i>Pouteria caimito</i> (Ruiz & Pav.) Radlk.	Abiu	Sapotaceae	N	1	10	0	0	0	0	0	0	1	10	1	2			
<i>Pouteria gardneri</i> (Mart. & Miq.) Baehni	Aguaí-una	Sapotaceae	N	0	0	0	0	1	15	1	1	0	0	0	0			
<i>Pouteria macrophylla</i> (Lam.) Eyma	Cutite	Sapotaceae	N	0	0	0	0	4	40	0	0	0	0	0	0			
<i>Protium heptaphyllum</i> (Aubl.) Marchand	Breu vermelho	Burseraceae	N	1	15	0	0	0	0	0	0	0	0	0	0			
<i>Protium</i> sp.	Breu branco	Burseraceae	N	1	15	0	0	1	15	0	0	0	0	0	0			
<i>Psidium guajava</i> L.	Goíaba	Myrtaceae	I	4	40	4	33	0	0	0	0	12	120	12	118			
<i>Quararibea guianensis</i> Aubl.	Inajarana	Malvaceae	N	3	45	1	2	0	0	0	0	0	0	0	0			
<i>Solanum aethiopicum</i> L.	Jiló	Solanaceae	I	3	30	0	0	0	0	0	0	0	0	0	0			
<i>Solanum paniculatum</i> L.	Jurubea	Solanaceae	N	0	0	0	0	1	15	0	0	4	60	0	0			
<i>Spondias dulcis</i> Forst.	Cajarana	Anacardiaceae	I	0	0	0	0	11	110	11	4	0	0	0	0			
<i>Spondias mombin</i> L.	Taperebá	Anacardiaceae	N	3	45	3	24	6	90	6	77	5	75	5	43			
<i>Spondias purpurea</i> L.	Siriguela	Anacardiaceae	I	7	105	4	48	9	135	7	99	0	0	0	0			
<i>Strychnos asperula</i> Sprague & Sandwith	Gogó-de-guariba	Loganiaceae	N	0	0	0	0	0	0	0	0	1	15	0	0			
<i>Symphonia globulifera</i> L. f.	Anani	Clusiaceae	N	3	45	0	0	0	0	0	0	0	0	0	0			
<i>Syzygium cumini</i> (L.) Skeels	Jambolão	Myrtaceae	I	1	15	0	0	0	0	0	0	1	15	0	0			
<i>Syzygium jambos</i> (L.) Alston	Jambo vermelho	Myrtaceae	I	3	30	0	0	2	20	0	0	2	20	0	0			
<i>Tapirira obtusa</i> (Benth.) J. D. Mitch.	Fruto de pombo	Anacardiaceae	N	0	0	0	0	0	0	0	0	3	45	0	0			
<i>Theobroma cacao</i> L.	Cacau	Malvaceae	I	1	3	0	0	4	12	0	0	0	0	0	0			
<i>Trattinnickia rhoifolia</i> Willd.	Amesclão	Burseraceae	N	0	0	0	0	0	0	0	0	1	15	0	0			
<i>Trichilia rubra</i> C. DC.	Cachuá	Meliaceae	N	1	15	0	0	0	0	0	0	0	0	0	0			
<i>Trymatococcus amazonicus</i> Poepp. & Endl.	Inharé	Moraceae	N	1	15	0	0	0	0	0	0	0	0	0	0			
No. species = 78				129	1,495	17	126	154	1,524	51	356	129	1,535	39	296			
No. families = 32																		

N = Native; I = Introduced; CS = collected samples; CF = collected fruits; IS = infested samples; IF = infested fruits.

Table 2. Species of *Anastrepha*, host plants, and associated parasitoids in 3 municipalities in the state of Amapá, Brazil (Jan–Dec 2012).

FAMILY Species ^a	Infestation ^b		Anastrepha species ^c and parasitoid species ^d			
	Puparia (n)	% FI	P/F	Mazagão ^e	Porto Grande ^e	Santana ^e
ANACARDIACEAE						
<i>Spondias dulcis</i> ●	45	4.0	11.3	—	Ao(7 ♀), 5 ♂	—
<i>Spondias mombin</i> ●	918	60.0	6.4	Ao(9 ♀), Aa(2 ♀), 11 ♂ + Ob(12), Da(7)	Ao(99 ♀), Aa(20 ♀), Af(1 ♀), Ast(1 ♀), 96 ♂ + Ob(60), Da(16), Asa(7), Agp(2), Ua(1)	Ao(13 ♀), Aa(6 ♀), 18 ♂ + Ob(22), Da(18)
<i>Spondias purpurea</i> ●	243	61.3	1.7	Ao(4 ♀), 2 ♂	Ao(69 ♀), Af(4 ♀), 80 ♂ + Da(5), Ob(4)	—
ANNONACEAE						
<i>Annona muricata</i> ●	4	4.8	4	—	—	Ad(1)
ARECACEAE						
<i>Oenocarpus bacaba</i> ▲	1	0.8	1	—	—	0
CELASTRACEAE						
<i>Cheiloclinium cognatum</i> ▲	16	2.7	8	1 ♂	—	Ah(1 ♀), 2 ♂
CHRYSOBALANACEAE						
<i>Licania laxiflora</i> ▲	2	1.7	2	—	—	Ob(2)
EUPHORBIACEAE						
<i>Manihot esculenta</i> ●	1	3.3	1	—	Ap(1)	—
FABACEAE						
<i>Inga edulis</i> ●	643	100	30.6	Ad(184 ♀), 76 ♂	—	Ad(115 ♀), Af(2 ♀), 38 ♂
LAURACEAE						
<i>Persea americana</i> ●	1	4.2	1	—	—	0
MALPIGHIACEAE						
<i>Malpighia emarginata</i> ●	60	47.1	0.6	—	Ao(3 ♀), 1 ♂	Ao(5 ♀), 3 ♂
MALVACEAE						
<i>Quararibea guianensis</i> ▲	11	4.4	5.5	Ao(5 ♀)	—	—
MELASTOMATACEAE						
<i>Bellucia grossularioides</i> ▲	13	20.0	2.2	—	Ac(1 ♀), 2 ♂	—
MYRTACEAE						
<i>Eugenia stipitata</i> ●	885	92.7	8.7	—	Ao(155 ♀), Af(10 ♀), 168 ♂	—
<i>Plinia cauliflora</i> ●	15	40.0	2.5	—	Ao(6 ♀), 8 ♂	—
<i>Psidium guajava</i> ●	2,242	94.4	14.8	Ast(45 ♀), Af(3 ♀), 51 ♂	—	Ast(454 ♀), Af(18 ♀), Ad(1 ♀), Ao(1 ♀), Azn(1 ♀), 595 ♂ + Da(11), Ob(1)
SAPOTACEAE						
<i>Manilkara zapota</i> ●	86	31.7	2.3	—	Asp(4 ♀), Alp(1 ♀), 8 ♀	—
<i>Pouteria caimito</i> ●	18	50.0	1.8	—	—	Ast(8 ♀), 1 ♂
<i>Pouteria Gardneri</i> ▲	3	6.7	3	—	Asp(1 ♀) + Da(1)	—

^aSymbols following species names indicate ▲ = native host; ● = introduced host.

^bInfestation calculated as % FI = percentage of infested fruits; P/F = puparia per fruit.

^cFruit flies = Aa: *Anastrepha antunesi*, Ac: *Anastrepha coronilli*, Ad: *Anastrepha distincta*, Af: *Anastrepha fraterculus*, Ah: *Anastrepha leptozona*, Ao: *Anastrepha obliqua*, Ap: *Anastrepha pickeli*, Asp: *Anastrepha serpentina*, Ast: *Anastrepha striata*, Azn: *Anastrepha zenillidae*. Male flies (♂) were not identified to species.

^dParasitoids = Agp: *Aganaspis pelleranoi*, Asa: *Asobara anastrephae*, Da: *Doryctobracon areolatus*, Ob: *Opius bellus*, Ua: *Uteretes anastrephae*.

^eA dash (—) indicates that no species was found.

distincta Greene, *A. fraterculus*, *Anastrepha hastata* Stone, *Anastrepha leptozona* Hendel, *A. obliqua*, *Anastrepha pickeli* Lima, *Anastrepha serpentina* (Wiedemann), *A. striata*, and *A. zenildae* (Table 2).

The plant families Anacardiaceae, Myrtaceae, and Sapotaceae contained the greatest number of hosts of *Anastrepha* (3 species in each plant family). The plant families Annonaceae, Celastraceae, Euphorbiaceae, Fabaceae, Malpighiaceae, Malvaceae, and Melastomataceae each contained 1 host species (Table 2).

The highest percentages of infested fruits were recorded in *Inga edulis* Mart. (Fabaceae) (100%), *P. guajava* (94.4%), and *E. stipitata* (92.7%). In *I. edulis*, a high infestation rate was also obtained, as high as 30.6 puparia per fruit (Table 2). No adults emerged from the puparia of Tephritidae that were obtained from fruits of *Oenocarpus bacaba* Mart. (Arecaceae) and *Persea americana* Mill. (Lauraceae) (Table 2), whereas only the parasitoid *Opius bellus* Gahan (Hymenoptera: Braconidae) emerged from the puparia obtained from *Licania laxiflora* Fritsch (Chrysobalanaceae).

The most abundant species of *Anastrepha* were *A. striata* (508 ♀), *A. obliqua* (376 ♀), and *A. distincta* (301 ♀). The great majority of specimens of *A. striata* (98.2%) were from the infestation in *P. guajava*, but specimens were also obtained from *Pouteria caimito* (Ruiz & Pav.) Radlk. (Sapotaceae) and *S. mombin* (Table 2). *Anastrepha obliqua* was recorded in 7 hosts, predominately in *E. stipitata* (155 ♀) and *S. mombin* (121 ♀). *Anastrepha distincta* had the greatest incidence in *I. edulis* (184 ♀). *Anastrepha fraterculus* occurred in 2 species of Anacardiaceae, 2 of Myrtaceae and 1 of Fabaceae, being predominant in *P. guajava* (21 ♀). *Anastrepha serpentina* occurred in 2 species of Sapotaceae. *Anastrepha antunesi* occurred only in *S. mombin*. The species *A. coronilli*, *A. hastata*, *A. leptozona*, *A. pickeli*, and *A. zenildae* occurred in only 1 host each.

The species *A. antunesi*, *A. fraterculus*, *A. obliqua*, and *A. striata* were recorded in the 3 municipalities studied. *Anastrepha distincta* and *A. hastata* in 2 municipalities, and the others in only 1 (Table 3).

The species of parasitoids obtained were: *Asobara anastrephae* (Muesebeck) (Hymenoptera: Braconidae), *D. areolatus*, *O. bellus*, *Utetes anastrephae* (Viereck) (Hymenoptera: Braconidae), and *Aganaspis pelleranoi* (Brèthes) (Hymenoptera: Figitidae) (Table 2). *Opius bellus* was the most abundant species; we recovered 101 individuals from *Anastrepha* infesting 4 host plant species. *Doryctobracon areolatus* also was obtained from 4 plant species, but in lower abundance (67 specimens) than *O. bellus*.

Discussion

In this study, it was possible to characterize the host utilization by *Anastrepha* in areas with high diversity of fruit plants by means of broad sampling (78 plant species) on 3 fruit producing properties in the state of Amapá. Although the diversity of fruit plants available in the area was high, not all species were used as hosts (Table 1). According to Querino et al. (2010) and Hernández-Ortiz (1993), some species of fruit flies that occur in the Amazon have a high level of specificity in relation to hosts, with a strong preference for certain species or botanical families.

The percentages of infested fruits obtained from *I. edulis*, *P. guajava*, and *E. stipitata* are striking because they are high. However, it is not possible to compare them with most other studies carried out in Amapá because earlier studies were performed with pooled fruit

Table 3. Species of *Anastrepha* and their hosts in 3 municipalities in the state of Amapá, Brazil (Jan–Dec 2012).

Specie	Host		Municipality		
	Family	Species	Mazagão	Porto Grande	Santana
<i>Anastrepha antunesi</i>	Anacardiaceae	<i>Spondias mombin</i>	x	x	x
<i>Anastrepha coronilli</i>	Melastomataceae	<i>Bellucia grossularioides</i>		x	
<i>Anastrepha distincta</i>	Annonaceae	<i>Annona muricata</i> ^a			x
	Fabaceae	<i>Inga edulis</i>	x		x
	Myrtaceae	<i>Psidium guajava</i>			x
<i>Anastrepha fraterculus</i>	Anacardiaceae	<i>Spondias mombin</i>		x	
	Anacardiaceae	<i>Spondias purpurea</i>		x	
	Fabaceae	<i>Inga edulis</i>			x
	Myrtaceae	<i>Eugenia stipitata</i>		x	
	Myrtaceae	<i>Psidium guajava</i>	x		x
<i>Anastrepha hastata</i>	Celastraceae	<i>Cheiloclinium cognatum</i>	x		x
<i>Anastrepha leptozona</i>	Sapotaceae	<i>Manilkara zapota</i> ^a		x	
<i>Anastrepha obliqua</i>	Anacardiaceae	<i>Spondias dulcis</i>		x	
	Anacardiaceae	<i>Spondias mombin</i>	x	x	x
	Anacardiaceae	<i>Spondias purpurea</i>	x	x	
	Malpighiaceae	<i>Malpighia emarginata</i>		x	x
	Malvaceae	<i>Quararibea guianensis</i> ^a	x		
	Myrtaceae	<i>Eugenia stipitata</i>		x	
	Myrtaceae	<i>Plinia cauliflora</i>		x	
	Myrtaceae	<i>Psidium guajava</i>			x
<i>Anastrepha pickeli</i>	Euphorbiaceae	<i>Manihot esculenta</i>		x	
<i>Anastrepha serpentina</i>	Sapotaceae	<i>Manilkara zapota</i>		x	
	Sapotaceae	<i>Pouteria gardneri</i> ^a		x	
<i>Anastrepha striata</i>	Anacardiaceae	<i>Spondias mombin</i>		x	
	Myrtaceae	<i>Psidium guajava</i>	x		x
	Sapotaceae	<i>Pouteria caimito</i>			x
<i>Anastrepha zenildae</i>	Myrtaceae	<i>Psidium guajava</i>			x

^aUnpublished association for Brazil (*Anastrepha* and host).

samples, which prevents the estimation of the percentage of infested fruits. On the other hand, in relation to the number of puparia per fruit, the 3 species showed values similar to those already reported in Amapá (Silva et al. 2011b).

All recorded *Anastrepha* species had already been reported from the state of Amapá (Deus & Adaime 2013). In the present study, all 16 *P. guajava* samples collected were infested by *A. striata* (94.4% of the fruits). This close association of *A. striata* with *P. guajava* has been well documented in Amapá (Silva et al. 2011b) and elsewhere in the Brazilian Amazon (Silva et al. 2011c). It is a species widely distributed in Amapá, where 25 hosts (16 families) have already been registered (Silva et al. 2011b).

Anastrepha obliqua was obtained from 7 hosts, especially *E. stipitata* (41.1% of specimens) and *S. mombin* (32.6% of specimens). This species was also recorded from 4 other hosts, among them *Quararibea guianensis* Aubl. (Malvaceae). Infestation of *Q. guianensis* by an *Anastrepha* species has not previously been recorded.

Anastrepha distincta was the predominate fly infesting *I. edulis* (99.3% of the specimens), its main host in Amapá (Silva et al. 2011b). The species was also reared from *A. muricata*, which is a new association for Brazil.

Anastrepha fraterculus was recorded in 5 hosts, with 55.3% of its specimens being from *P. guajava*. The species is widely distributed in Amapá, although the population density is low (Deus & Adaime 2013).

Anastrepha antunesi was obtained only from *S. mombin*, a species with which it is frequently associated (Deus & Adaime 2013). In general, *A. antunesi* and *A. obliqua* are obtained from *S. mombin* fruits. Nascimento et al. (2015) collected 600 fruits of this plant and evaluated them individually. The authors recorded infestation by fruit flies in 151 fruits (25.2%) and obtained 298 puparia (mean of 1.97 puparia per fruit, with a maximum of 8 puparia per fruit). The simultaneous occurrence of *A. obliqua* and *A. antunesi* was recorded in only 2 of the 151 infested fruits (1.3%).

Anastrepha serpentina was obtained from fruits of *Manilkara zapota* (L.) P. Royen (Sapotaceae) and *Pouteria gardneri* (Mart. & Miq.) Baehni (Sapotaceae). The occurrence of *A. serpentina* in *P. gardneri* is not previously reported for Brazil. It should be noted that *A. serpentina* has already been reported in *Pouteria gardneriana* (A. DC.) Radlk. (Sapotaceae) in the state of Goiás (Zucchi 2008), but that plant species does not occur in the Amazon region.

The species *A. coronilli*, *A. hastata*, *A. pickeli*, and *A. zenildae* were represented by a single specimen each, in hosts already reported to support them (Deus & Adaime 2013). However, *A. leptozona* was recorded here for the first time in *M. zapota* in Brazil.

The action of parasitoids on *Anastrepha* species was also recorded in this study. Of the 179 specimens obtained, 56.4% were *O. bellus*, 38.0% *D. areolatus*, 3.9% *A. anastrephae*, 1.1% *A. pelleranoi*, and 0.6% *U. anastrephae* (Table 2). Together, *O. bellus* and *D. areolatus* accounted for 94.4% of the total specimens obtained. Due to their abundance, they are the species with the greatest potential to control the population of fruit flies under the conditions in Amapá (Deus & Adaime 2013). *Opius bellus* and *D. areolatus* occurred in the 3 municipalities studied. The other species occurred only in Porto Grande (Table 2). We obtained 81.0% of the parasitoid specimens reported in this study from *S. mombin*. This plant plays an important role as a natural reservoir of the fruit fly parasitoid populations in Amapá, which deserves further study (Sousa et al. 2016).

Acknowledgments

We thank our colleagues who helped with field work (Carlos Alberto Moraes, Manoel Jonas de Jesus Viana, and Marcelo Luiz de Oliveira) and laboratory work (Maria do Socorro Miranda de Sousa, Luana dos

Santos Pinheiro, Francisco Andrew Pacheco, Oziel Barroso Baía, Sarron Felipe, and Jonh Carlo Reis). We are grateful to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the Bolsa de Produtividade em Pesquisa (Productivity in Research Grant) awarded to R. Adaime. A. Leyva (USA) helped with English translation and editing of the manuscript.

References Cited

- Adaime R, Sousa MSM, Pereira JF. 2016. *Anastrepha* species and their hosts in the Brazilian Amazon, <http://anastrepha.cpafap.embrapa.br> (updated 3 Oct 2016; last accessed 29 Dec 2016).
- Aluja M. 1994. Bionomics and management of *Anastrepha*. Annual Review of Entomology 39: 155–178.
- Aluja M, Mangan RL. 2008. Fruit fly (Diptera: Tephritidae) host status determination: critical conceptual, methodological, and regulatory considerations. Annual Review of Entomology 53: 473–502.
- Aluja M, Ordano M, Guillén L, Rull J. 2012. Understanding long-term fruit fly (Diptera: Tephritidae) population dynamics: implications for areawide management. Journal of Economic Entomology 105: 823–836.
- Bittencourt MAL, Santos OO, Brito EA, Araújo EL, Marinho CF. 2012. Parasitóides (Braconidae) associados à *Anastrepha* (Tephritidae) em frutos hospedeiros do Litoral Sul da Bahia. Revista Ciência Agronômica 43: 811–815. [In Portuguese]
- Carvalho RS, Nascimento AS, Matrangolo WJR. 2000. Controle biológico, pp. 113–117 In Malavasi A, Zucchi RA [eds.], Moscas-das-frutas de Importância Econômica no Brasil. Conhecimento Básico e Aplicado. Holos, Ribeirão Preto, São Paulo, Brazil. [In Portuguese]
- Carvalho RS, Soares Filho WS, Ritzinger R. 2010. Uumbu-cajá como repositório natural de parasitoide nativo de moscas-das-frutas. Pesquisa Agropecuária Brasileira 45: 1222–1225. [In Portuguese]
- Deus EG, Adaime R. 2013. Dez anos de pesquisas sobre moscas-das-frutas (Diptera: Tephritidae) no estado do Amapá: avanços obtidos e desafios futuros. Biota Amazônia 3: 157–168. [In Portuguese]
- Dutra VS, Ronchi-Teles B, Garcia MVB, Adaime R, Silva JG. 2013. Native hosts and parasitoids associated with *Anastrepha fractura* and other *Anastrepha* species (Diptera: Tephritidae) in the Brazilian Amazon. Florida Entomologist 96: 270–273.
- Duyck P-F, David P, Quilici S. 2004. A review of relationships between interspecific competition and invasions in fruit flies (Diptera: Tephritidae). Ecological Entomology 29: 511–520.
- Hafsi A, Facon B, Ravigné V, Chiroleu F, Quilici S, Chermiti B, Duyck P-F. 2016. Host plant range of a fruit fly community (Diptera: Tephritidae): Does fruit composition influence larval performance? BMC Ecology 16:40
- Hernández-Ortiz V. 1993. Taxonomy, distribution, and natural host plants of *Anastrepha* fruit flies in Mexico, pp. 31–34 In Aluja M, Liedo P [eds.], Fruit Flies: Biology and Management. International Symposium on fruit flies of economic importance, 1990, Antigua, Guatemala. Springer-Verlag, New York, New York.
- IBGE. 2010. Produto. Mapas de climas do Brasil. Updated in 2010, http://www.ibge.gov.br/home/geociencias/default_prod.shtml#MAPAS (last accessed 11 Dec 2013). [In Portuguese]
- Jesus-Barros CR, Adaime R, Oliveira MN, Silva WR, Costa-Neto SV, Souza-Filho MF. 2012. *Anastrepha* (Diptera: Tephritidae) species, their hosts and parasitoids (Hymenoptera: Braconidae) in five municipalities of the state of Amapá, Brazil. Florida Entomologist 95: 694–705.
- Leal MR, Souza SAS, Aguiar-Menezes EL, Lima Filho M, Menezes EB. 2009. Diversidade de moscas-das-frutas, suas plantas hospedeiras e seus parasitóides nas regiões Norte e Noroeste do Estado do Rio de Janeiro, Brasil. Ciência Rural 39: 627–634. [In Portuguese]
- Lemos LUJ, Souza-Filho MF, Uramoto K, Lopes GN, Zucchi RA. 2015. Espécies de *Anastrepha* (Diptera: Tephritidae) em pomares de goiaba: diversidade, flutuação populacional e fenologia do hospedeiro. Arquivos do Instituto Biológico 82: 1–5. [In Portuguese]
- Malavasi A. 2009. Biologia, ciclo de vida, relação com o hospedeiro, espécies importantes e biogeografia de tefritídeos, pp. 1–15 In Malavasi A, Virginio J [eds.], Biologia, Monitoramento e Controle de Moscas-das-frutas: V Curso Internacional de Capacitação em Moscas-das-frutas, 21 a 29 de outubro de 2009. Biofábrica Moscamed Brasil, Juazeiro, Bahia, Brazil. [In Portuguese]
- Marinho CF, Silva RA, Zucchi RA. 2011. Chave de identificação de Braconidae (Alysiinae e Opiinae) parasitoides de larvas frugívoras na região Amazônica, pp. 91–101 In Silva RA, Lemos WP, Zucchi RA [eds.], Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais. Embrapa Amapá, Macapá, Amapá, Brazil. [In Portuguese]

- Nascimento DB, Adaime R, Cunha AC, Silva JG. 2015. Influência dos parâmetros biométricos de frutos de *Spondias mombin* L. sobre os índices de infestação por *Anastrepha* spp. (Diptera: Tephritidae) e parasitismo. *Biota Amazônia* 5: 83–87. [In Portuguese]
- Ovruski S, Aluja M, Sivinski J, Wharton R. 2000. Hymenopteran parasitoids on fruit-infesting Tephritidae (Diptera) in Latin America and the southern United States: diversity, distribution, taxonomic status and their use in fruit fly biological control. *Integrated Pest Management Reviews* 5: 81–107.
- Querino RB, Costa SGM, Ronchi-Teles B, Strikis P, Zucchi RA. 2010. Interação de larvas frugívoras (Diptera, Tephritidae e Lonchaeidae) e seus hospedeiros na Reserva Florestal Adolpho Ducke, Manaus, Amazonas, Brasil. *Teresina: Embrapa Meio-Norte (Boletim de Pesquisa e Desenvolvimento, 88)*. [In Portuguese]
- Silva NM, Ronchi-Teles B. 2000. Amapá, Amazonas, Pará, Rondônia e Roraima, pp. 203–209 *In* Malavasi A, Zucchi RA [eds.], *Moscas-das-frutas de Importância Econômica no Brasil. Conhecimento Básico e Aplicado*. Holos, Ribeirão Preto, São Paulo, Brazil. [In Portuguese]
- Silva RA, Deus EG, Raga A, Pereira JDB, Souza-Filho MF, Costa Neto SV. 2011a. Monitoramento de moscas-das-frutas na Amazônia: amostragem de frutos e uso de armadilhas, pp. 33–47 *In* Silva RA, Lemos WP, Zucchi RA [eds.], *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá, Amapá, Brazil. [In Portuguese]
- Silva RA, Deus EG, Pereira JDB, Jesus CR, Souza-Filho MF, Zucchi RA. 2011b. Conhecimento sobre moscas-das-frutas no Estado do Amapá, pp. 223–236 *In* Silva RA, Lemos WP, Zucchi RA [eds.], *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá, Amapá, Brazil. [In Portuguese]
- Silva RA, Lemos WP, Zucchi RA [eds.]. 2011c. *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá, Amapá, Brazil. [In Portuguese]
- Sousa MSM, Jesus-Barros CR, Yokomizo GK, Lima AL, Adaime R. 2016. Ocorrência de moscas-das-frutas e parasitoides em *Spondias mombin* L. em três municípios do estado do Amapá, Brasil. *Biota Amazônia* 6: 50–55. [In Portuguese]
- Uchôa MA, Nicácio JN. 2010. New records of Neotropical fruit flies (Tephritidae), lance flies (Lonchaeidae) (Diptera: Tephritoidea), and their host plants in the south Pantanal and adjacent areas, Brazil. *Annals of the Entomological Society of America* 103: 723–733.
- Uramoto K, Zucchi RA. 2009. Taxonomia de espécies de *Anastrepha* (Diptera, Tephritidae), pp. 7–11 *In* Malavasi A, Virgínio J [eds.], *Biologia, Monitoramento e Controle de Moscas-das-frutas: V Curso Internacional de Capacitação em Moscas-das-frutas*, 21 a 29 de outubro de 2009. Biofábrica Moscamed Brasil, Juazeiro, Bahia, Brazil. [In Portuguese]
- Uramoto K, Martins DS, Zucchi RA. 2008. Fruit flies (Diptera, Tephritidae) and their associations with native host plants in a remnant area of the highly endangered Atlantic rain forest in the state of Espírito Santo, Brazil. *Bulletin of Entomological Research* 98: 457–466.
- Zucchi RA. 2008. Fruit flies in Brazil: *Anastrepha* species their host plants and parasitoids, www.lea.esalq.usp.br/anastrepha/ (updated 10 Nov 2015; last accessed 20 May 2016).
- Zucchi RA, Silva RA, Deus EG. 2011a. Espécies de *Anastrepha* e seus hospedeiros na Amazônia brasileira, pp. 53–70 *In* Silva RA, Lemos WP, Zucchi RA [eds.], *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá, Amapá, Brazil. [In Portuguese]
- Zucchi RA, Uramoto K, Souza-Filho MF. 2011b. Chave ilustrada para as espécies de *Anastrepha* da região Amazônica, pp. 71–90 *In* Silva RA, Lemos WP, Zucchi RA [eds.], *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá, Amapá, Brazil. [In Portuguese]