



## **Assessing Genetic Diversity of Three Species of Potato Tuber Moths (Gelechiidae, Lepidoptera) in the Ecuadorian Highlands**

Authors: Fu, Zhen, Castillo Carrillo, Carmen I., Rashed, Arash, Asaquibay, César, Aucancela, Rodrigo, et al.

Source: Florida Entomologist, 103(3) : 329-336

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.103.0304>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Assessing genetic diversity of three species of potato tuber moths (Gelechiidae, Lepidoptera) in the Ecuadorian highlands

Zhen Fu<sup>1,5,\*</sup>, Carmen I. Castillo Carrillo<sup>2</sup>, Arash Rashed<sup>3</sup>, César Asaquibay<sup>2</sup>, Rodrigo Aucancela<sup>2</sup>, José Camacho<sup>2</sup>, Victoria López<sup>2</sup>, Verónica Quimbiamba<sup>2</sup>, Fausto Yumisaca<sup>2</sup>, Nancy Panchi<sup>4</sup>, and Claudio Velasco<sup>4</sup>

## Abstract

Three species of potato tuber moths, *Tecia solanivora* Povolny, *Symmetrischema tangolias* Gyen, and *Phthorimaea operculella* Zeller (all Lepidoptera: Gelechiidae), attack potato plants in the highlands of Ecuador and cause great economic losses. To understand their ecology and to develop precise integrated pest management strategies, a description of the molecular identification of each species and genetic diversity within populations is needed. In this study, we assessed the genetic diversity of a total of 112 moth samples of these 3 distinct species. Samples were collected from the 4 central provinces (Bolívar, Chimborazo, Cotopaxi, and Tungurahua) in the Ecuadorian highlands, where potato is a major crop for food security. Using polymerase chain reaction (PCR) and Sanger sequencing, we compared mitochondrial gene sequences among conspecific moth samples. Each of the 3 species exhibited different patterns regarding genetic diversity; more than 1 haplotype was present in *P. operculella* and *S. tangolias*, whereas all *T. solanivora* samples were found to be genetically identical. This initial effort of molecular characterization of the potato tuber moths will facilitate identifying incursion and potential migration route of Gelechiidae species as well as prevention of the pest outbreaks.

Key Words: *Tecia solanivora*; *Symmetrischema tangolias*; *Phthorimaea operculella*; haplotype; potato tuber moth

## Resumen

Tres especies de polillas de la papa, *Tecia solanivora* Povolny, *Symmetrischema tangolias* Gyen, y *Phthorimaea operculella* Zeller (todos Lepidoptera: Gelechiidae), atacan el cultivo de la papa en la sierra ecuatoriana causando grandes pérdidas económicas. Para comprender su ecología y desarrollar estrategias precisas de manejo integrado de plagas, se necesita una descripción de la identificación molecular de cada especie y la diversidad genética de poblaciones intraespecíficas. En este estudio, realizamos una caracterización molecular de un total de 112 muestras de polillas de estas 3 especies. Se recolectaron muestras de las 4 provincias centrales de los Andes ecuatorianos (Bolívar, Chimborazo, Cotopaxi, y Tungurahua) donde la papa es un cultivo importante para la seguridad alimentaria. Utilizando la reacción en cadena de la polimerasa y secuenciación Sanger, comparamos las secuencias de los genes mitocondriales entre las muestras de las 3 especies de polillas. Cada una de las 3 especies exhibió diferentes patrones con respecto a la diversidad genética; más de 1 haplotipo estuvo presente en *P. operculella* y *S. tangolias*, mientras que todas las muestras de *T. solanivora* fueron genéticamente idénticas. Este esfuerzo inicial de caracterización molecular de las polillas del tubérculo de la papa facilitará la identificación de la incursión y la posible ruta de migración de las especies de Gelechiidae y prevención de despuntes poblacionales de las plagas.

Key Words: *Tecia solanivora*; *Symmetrischema tangolias*; *Phthorimaea operculella*; haplotipo; polilla de la papa

Potato (*Solanum tuberosum* L.; Solanaceae) is an economically important staple food in South American countries, including Ecuador. Numerous insect pests attack potato plants and tubers during the growing season and post-harvest (i.e., storage). Three species of potato tuber moths (Lepidoptera: Gelechiidae), the Guatemalan potato tuber moth, *Tecia solanivora* Povolny, the Andean potato tuber moth,

*Symmetrischema tangolias* Gyen, and the common potato tuber moth, *Phthorimaea operculella* Zeller, are present in Ecuador and cause significant economic losses (Gallegos et al. 2004; Rondon & Gao 2018).

The Guatemalan potato tuber moth, *T. solanivora*, is known to have originated from Central America and is a highly diverse species in that region (Torres-Leguizamón et al. 2011). There are 60

<sup>1</sup>Department of Entomology, 166 FSHN, Washington State University, Pullman, Washington, 99164, USA; E-mail: zhen.fu@tamu.edu (Z. F.)

<sup>2</sup>Instituto Nacional de Investigaciones Agropecuarias, Estación Experimental Santa Catalina, Panamericana Sur km 1, Quito, 171107, Ecuador; E-mail: carmen.castillo@iniap.gob.ec (C. I. C. C.), cesar.asaquibay@iniap.gob.ec (C. A.), rodrigo.aucancela@iniap.gob.ec (R. A.), jose.camacho@iniap.gob.ec (J. C.), victoria.lopez@iniap.gob.ec (V. L.), vquimbiambaregion5@gmail.com (V. Q.), fausto.yumisaca@iniap.gob.ec (F. Y.)

<sup>3</sup>Department of Entomology, Plant Pathology and Nematology, 875 Perimeter Drive, MS 2329, University of Idaho, Moscow, Idaho 83844, USA; arashed@uidaho.edu (A. R.)

<sup>4</sup>Centro Internacional de la Papa (CIP), Quito, Ecuador E-mail: n.panchi@cgiar.org (N. P.), c.valasco@cgiar.org (C. V.)

<sup>5</sup>Current address: Department of Entomology, 120 Entomology Research Lab, Texas A&M University, College Station, Texas 77843, USA

\*Corresponding author; email: zhen.fu@tamu.edu

Supplementary material for this article in Florida Entomologist 103(3) (September 2020) is online at <http://purl.fcla.edu/fcla/entomologist/browse>


documented haplotypes of *T. solanivora* based on the cytochrome b (cytb) gene in Guatemala, Costa Rica, and Venezuela. The distribution of this moth stretches from Mexico in the north to Ecuador on its south boundary, covering a handful of Central and South American countries such as Panama, Costa Rica, Nicaragua, El Salvador, Honduras, Guatemala, Venezuela, and Colombia (Salasar & Escalante 1984; Rincón & López-Ávila 2004; Trujillo et al. 2004; Cruz et al. 2011). Across the Atlantic, *T. solanivora* was reported from the Canary Islands, Spain, in 1999, followed by another report of the pest from Spain's mainland in 2005 (EPPO 2015). In Ecuador, *T. solanivora* was spotted for the first time in 1996 in the northern province of Carchi (Gallegos 1997). It was thought that the range of this species in the Ecuadorian Andes is confined to regions that are below 3,300 masl (Gallegos et al. 2004). However, the latest survey debunked such notation because *T. solanivora* was found in potato fields in Chimborazo where elevation reaches 3,645 masl (Castillo Carrillo et al. 2018). *Tecia solanivora* is present in Imbabura, Pichincha, Cotopaxi, Tungurahua, and Bolívar provinces. In certain areas of the province of Bolívar, *T. solanivora* is the most common potato tuber moth species (Castillo Carrillo et al. 2018). In Azuay, the presence of *T. solanivora* has not been reported. There have been informal reports of *T. solanivora* being present in Cañar province. The genetic diversity outside of its native range is low, as only 1 haplotype, H6, has been found in samples collected from Colombia, Ecuador, and the Canary Islands (Torres-Lequizamón et al. 2011). The larva of *T. solanivora* feeds exclusively on potato tubers during its development in the field and during storage. It creates galleries in the tubers as it feeds, which render tubers unmarketable and results in considerable economic losses to farmers.

A recent survey covering 4 Ecuadorian provinces determined that the Andean moth, *S. tangolias*, is more common than *T. solanivora* in Chimborazo, Cotopaxi, and Tungurahua (Castillo Carrillo et al. 2018). *Symmetrischema tangolias* may be present in large numbers; in Chagrapamba, canton Pillaro, Tungurahua, over 100 male *S. tangolias* were collected in a single pheromone trap within 24 h (Castillo Carrillo et al. 2018). *Symmetrischema tangolias* is thought to be native to the Peruvian or Bolivian Andes and is broadly distributed in the

Andean zones of Colombia, Ecuador, Peru, and Bolivia. Beyond South America, *S. tangolias* is present in Australia, Tasmania, New Zealand, and Indonesia (Terauds et al. 1984; Martin 1999; Kroschel & Schaub 2013). The larva mines the potato plant stems, including tubers, both in the field and post-harvest. In addition to potato plants, *S. tangolias* can attack other solanaceous hosts, such as tomato (*Solanum lycopersicum* L.), pepino (*S. muricatum* Ait.), *S. aviculare* G. Forster, and *S. laciniatum* Ait. (all Solanaceae) (Kroschel & Schaub 2013). The first sighting of *S. tangolias* in Ecuador dates back to the early 1990s in the province of Carchi (P. Gallegos, personal communication).

Different from *T. solanivora* and *S. tangolias*, *P. operculella* is a cosmopolitan pest that causes considerable damage worldwide (Rondon & Gao 2018). *Phthorimaea operculella* can be found in over 90 countries across the Americas, Asia, Africa, and Australia (Kroschel et al. 2013; Rondon & Gao 2018). It feeds on solanaceous crops such as potatoes and tomatoes. Females oviposit on potato leaves and stems (including tubers). Their larvae mine leaves and tubers, resulting in economic damage. *Phthorimaea operculella* was reported as a significant pest of potato tubers in the early 1900s in South America (Graft 1917). In Ecuador, it was observed for the first time between 1984 and 1985 in the southern province of Cañar (P. Gallegos, personal communication). Generally, *P. operculella* is found less frequently than the other 2 moth species, *T. solanivora* and *S. tangolias*, in the potato producing regions in Ecuador, except for certain areas in Tungurahua (Castillo Carrillo et al. 2018).

Despite the economic importance and coexistence of all 3 species in the same region (Dangles et al. 2009; Castillo Carrillo et al. 2018), information regarding the identification and genetic diversity is largely missing except for *T. solanivora* (Torres-Lequizamón et al. 2011). This lack of accurate identification and knowledge of their biology and distribution contributes in part to inefficient management of these species. To establish a clear genetic description of these 3 insect species, we characterized intraspecific genetic diversity of the potato tuber moth samples collected from 4 Ecuadorian provinces using previously published primers that target the cytochrome oxidase I (COI) gene for *P. operculella* and *S. tangolias* and the cytb gene for *T. solanivora*.



**Fig. 1.** Use of pheromone traps to collect potato tuber moths in Cacha, Chimborazo, and Tishinguirí, Bolívar: (A) potato crops are grown in mountainous regions of Ecuador where smallholder terrace farming is predominant; (B) low-cost pheromone traps were made of plastic bottles; a species-specific lure is fixed to the bottle cap using a string, and thumb-sized holes were carved to allow moths to fly into the bottle that was filled with soapy water; (C) 3 different traps were placed (pointed by arrows) in each field at least 100 m apart to catch different moths.

**Table 1.** Primers used in this study.

Targeting species	Primer name	Sequences (5' to 3')	Product size (base pairs)	Reference
<i>Symmetrischema tangolias</i>	Sym-tan-S369	GCTGAATTAGGTAACCCAGGC	431	Sint et al. 2016
	Lep-gen-A309	GGTATTTGGTCAAATGAAAGTCC		Sint et al. 2016
<i>Phthorimaea operculella</i>	Pht-ope-S349	AGGAATAGTTGGAACCTCTCTAGT	469	Sint et al. 2016
	Lep-gen-A309	GGTATTTGGTCAAATGAAAGTCC		Sint et al. 2016
<i>Tecia solanivora</i>	TsF1-re	GCTAACATTGAGTTAGCTTTTAT	963	This study
	TsR-re	AAAAATTAGAGTATCTCAAAAT		This study

## Materials and Methods


### COLLECTION OF MOTHS

Live moths were collected using pheromone traps (ChemTica Internacional, Heredia, Costa Rica). These traps were specific to each species, and they were set up in potato fields in provinces of Bolívar, Chimborazo, Cotopaxi, and Tungurahua (Fig. 1, Table S1). Moths that were attracted to each species-specific pheromone trap were verified to be the correct species based on wing patterns (Sporleder et al. 2016a, b; Rondon & Gao 2018). Individual samples were placed in 1.5 mL micro-


centrifuge tubes (Hauppauge, New York, USA) and stored in 95% ethanol at 17 to 18 °C upon arrival at the laboratory. A total of 112 moths, including 47 *T. solanivora*, 37 *S. tangolias*, and 28 *P. operculella* samples that represented moth populations from 4 provinces were processed for DNA extractions and characterized with polymerase chain reaction (PCR) and Sanger sequencing.

### DNA EXTRACTION

Wings of a single moth were separated from the body, and the whole body was placed in a 1.5 mL microcentrifuge tube. The moth body was homogenized using a Mini-BeadBeater homogenizer (GlenMills, Clifton,



**Fig. 2.** Alignment of 5 COI haplotype sequences of *Symmetrischema tangolias* samples. KX443104.1 is a reference sequence from GenBank. Sequences of these 5 haplotypes were deposited in GenBank with accession numbers MN223391 to MN223395.



**Fig. 3.** Distribution of 5 haplotypes and sequence identity between each pair of *Symmetrischema tangolias* haplotypes. St-H1 was present in all 4 provinces; St-H2 was present in 3 provinces; the other 3 haplotypes were present only in 1 province for each haplotype. Arrows point to province(s) instead of a specific sampling location.

New Jersey, USA) for 2 min in 180 µL of ATL buffer (Qiagen DNeasy Blood & Tissue Kit, Qiagen, Germantown, Maryland, USA), with 5 glass beads (3 mm diam) added. The tissue homogenate was incubated overnight at 56 °C. The DNA extraction process was continued using a Qiagen DNeasy Blood & Tissue Kit. The DNA concentration was quantified using a NanoDrop 2000 (ThermoFisher Scientific, Frederick, Maryland, USA).

PCR AMPLIFICATION AND PURIFICATION

The total volume of all PCR reactions was 25 µL, which included 2.5 µL 10× PCR buffer, 1.5 µL 25mM MgCl<sub>2</sub>, 1.25 µL dNTPs, 0.5 µL of each 10 µM forward and reverse primer (Table 1), 0.25 µL Takara Ex Taq (Takara, Mountain View, California, USA), 1 µL extracted moth DNA, and 17.5 µL of molecular grade water. PCR buffer, MgCl<sub>2</sub>, and dNTPs were supplied with Takara Ex Taq; all concentrations listed above were the concentration prior to adding into the mix. The program for PCR was 94 °C for 2 min, then 35 cycles of 94 °C for 15 s, 55 °C for 60 s, and 72 °C for 60 s, followed by a final extension at 72 °C for 5 min, in a S1000 Thermal Cycler (Bio-Rad, Hercules, California, USA). The PCR products were visualized on 1.5% agarose gels.

For *T. solanivora*, we initially used primers TsF1 and TsR, according to Torres-Leguizamón et al. (2011). However, the gel electrophoresis indicated that there were multiple amplicons present. As a result, the sequences appeared to be poor quality, and the background noise was high. Thus, we redesigned primers TsF1-re and TsR-re (Table 1), which were used to amplify a 963 base pair section of the *T. solanivora* cytb gene. The PCR condition for redesigned primers was the same as described above, except the annealing temperature was raised from 55 °C to 59 °C.

For samples that formed a single clear band, 5 µL PCR product was purified using ExoSAP-IT (ThermoFisher Scientific, Frederick, Maryland, USA) according to the manufacturer protocol. Since ExoSAP-IT purified DNA was too concentrated for Sanger sequencing, we diluted them 10 times with molecular grade water. Two µL of the diluted DNA and 0.8 µL of either forward or reverse primers (10 µM) were added to a 0.2 mL microcentrifuge tube with 12.2 µL of molecular grade water to reach a total volume of 15 µL. Sanger sequencing with forward and reverse primer were performed separately for each DNA sample.

DATA ANALYSES AND ARCHIVING

Raw sequences were processed using the program BioEdit (vers. 7.0.5; Hall 1999). Consensus sequences were generated from the alignment of reverse and forward reads. Sequences from the same moth species were aligned using the program MUSCLE (Edgar 2004). Five sequences representing 5 unique haplotype of *S. tangolias* were deposited in GenBank with accession numbers MN223391 to MN223395. Three sequences representing 3 unique haplotypes of *P. operculella* were deposited in GenBank with accession numbers MN205567 to MN205569. The sequence of the only haplotype discovered within *T. solanivora* samples was archived in GenBank with accession number MN205566.

Results

We successfully amplified 42 of 47 *T. solanivora* samples with the newly designed primers TsF1-re and TsR-re. A single and clear band was seen on the gel for each successful amplification. All of the sequences

Po-H2	1	GAACCTCTCTAGTCTTTTAATTCGAGCAGAATTAGGAAACCCTGGATCTTTAATTGGGGATGATCAAATTTATAATACT
Po-H3	1	GAACCTCTCTAGTCTTTTAATTCGAGCAGAATTAGGAAACCCTGGATCTTTAATTGGGGATGATCAAATTTATAATACT
Po-H1	1	GAACCTCTCTAGTCTTTTAATTCGAGCAGAATTAGGAAACCCTGGATCTTTAATTGGGGATGATCAAATTTATAATACT
MF121882	1	GAACCTCTCTAGTCTTTTAATTCGAGCAGAATTAGGAAACCCTGGATCTTTAATTGGGGATGATCAAATTTATAATACT
Po-H2	81	ATTGTTACAGCTCAGCTTTTATTATAATTTTTTTATGGTTATACCTATTATAATTGGAGGATTGGTAATTGATTAGT
Po-H3	81	ATTGTTACAGCTCAGCTTTTATTATAATTTTTTTATGGTTATACCTATTATAATTGGAGGATTGGTAATTGATTAGT
Po-H1	81	ATTGTTACAGCTCAGCTTTTATTATAATTTTTTTATGGTTATACCTATTATAATTGGAGGATTGGTAATTGATTAGT
MF121882	81	ATTGTTACAGCTCAGCTTTTATTATAATTTTTTTATGGTTATACCTATTATAATTGGAGGATTGGTAATTGATTAGT
Po-H2	161	ACCATTAATATTAGGGGCTCCAGATATAGCTTTCCCCCGAATAAATAATATAAGTTTTTGATTATTACCACCCTCTCTTA
Po-H3	161	ACCATTAATATTAGGGGCTCCAGATATAGCTTTCCCCCGAATAAATAATATAAGTTTTTGATTATTACCACCCTCTCTTA
Po-H1	161	ACCATTAATATTAGGGGCTCCAGATATAGCTTTCCCCCGAATAAATAATATAAGTTTTTGATTATTACCACCCTCTCTTA
MF121882	161	ACCATTAATATTAGGGGCTCCAGATATAGCTTTCCCCCGAATAAATAATATAAGTTTTTGATTATTACCACCCTCTCTTA
Po-H2	241	CATTATTAATTTCAAGAAGAATTGTAGAAAATGGAGCTGGTACTGGTTGAAGTGTCTACCCTCCTTTATCTTCTAATATT
Po-H3	241	CATTATTAATTTCAAGAAGAATTGTAGAAAATGGAGCTGGTACTGGTTGAAGTGTCTACCCTCCTTTATCTTCTAATATT
Po-H1	241	CATTATTAATTTCAAGAAGAATTGTAGAAAATGGAGCTGGTACTGGTTGAAGTGTCTACCCTCCTTTATCTTCTAATATT
MF121882	241	CATTATTAATTTCAAGAAGAATTGTAGAAAATGGAGCTGGTACTGGTTGAAGTGTCTACCCTCCTTTATCTTCTAATATT
Po-H2	321	GCTCATGGGGTAGCTCTGTAGATTTAGCTATTTTTTCTCCTTCATCTAGCTGGTATTTTCATCAATTTTAGGAGCTATTAA
Po-H3	321	GCTCATGGGGTAGCTCTGTAGATTTAGCTATTTTTTCTCCTTCATCTAGCTGGTATTTTCATCAATTTTAGGAGCTATTAA
Po-H1	321	GCTCATGGGGTAGCTCTGTAGATTTAGCTATTTTTTCTCCTTCATCTAGCTGGTATTTTCATCAATTTTAGGAGCTATTAA
MF121882	321	GCTCATGGGGTAGCTCTGTAGATTTAGCTATTTTTTCTCCTTCATCTAGCTGGTATTTTCATCAATTTTAGGAGCTATTAA
Po-H2	401	TTTTATTACTACTATTATTAATATACGAATTAATGGACTTTCATT
Po-H3	401	TTTTATTACTACTATTATTAATATACGAATTAATGGACTTTCATT
Po-H1	401	TTTTATTACTACTATTATTAATATACGAATTAATGGACTTTCATT
MF121882	401	TTTTATTACTACTATTATTAATATACGAATTAATGGACTTTCATT

**Fig. 4.** Alignment of 3 unique COI haplotype sequences of *Phthorimaea operculella*. MF121882 is a reference sequence from GenBank. Sequences of these 3 haplotypes were deposited in GenBank with accession numbers MN205567 to MN205569.

were 100% identical to each other, with 100% similarity to several sequences that previously had been deposited in GenBank, including a mitochondrial genome of *T. solanivora* from Colombia (KT326187.1) and several previously reported *T. solanivora* samples from Ecuador (EF212202.1 and DQ780487.1). Lastly, the cytb sequence of *T. solanivora* in our study was 100% identical to H6, the only haplotype discovered from Ecuador, according to Torres-Leguizamón et al. (2011).


Out of 37 *S. tangolias* samples, 35 had a distinct band of the expected size of 431 base pairs. Five haplotypes were identified within these 35 samples. Sequence St-H1 was the dominant haplotype, representing over 75% of the collected samples in the 4 provinces (Figs. 2, 3). Sequence St-H2 was present in 4 samples from 3 provinces (Tungurahua, Bolívar, and Cotopaxi). Sequence St-H2 was only one nucleotide different from sequence St-H1 (Fig. 2). Sequence St-H3 was the most distant haplotype in relation to the other 4 haplotypes. The identity between St-H3 and any haplotype was less than 96% (Fig. 3). Only 2 samples collected from Chimborazo were characterized as St-H3 haplotype. One sample from Tungurahua and 1 sample from Cotopaxi were characterized as St-H4 and St-H5, respectively. These 2 haplotypes were 99% identical with 3 bases different from each other (Fig. 2).

We processed 28 samples of *P. operculella*, of which 25 showed a distinct band at the expected 469 base pairs. Three haplotypes were present within 25 samples (Figs. 4, 5). The first haplotype, Po-H1, was the most dominant haplotype; 19 out of 25 *P. operculella* samples spanning 4 provinces were characterized as Po-H1. The sequence of Po-H1 was 100% identical to that of MF121882, which was deposited in

GenBank, with only 1 nucleotide different from the other 2 haplotypes, Po-H2 and Po-H3 (Fig. 4). Six samples were characterized as Po-H2 and Po-H3, the 2 haplotypes that are differentiated by only 2 nucleotides (Fig. 4). The 2 samples which were characterized as Po-H2 were present only in Tungurahua. Four samples that belong to Po-H3 were present in Cotopaxi and Chimborazo.

## Discussion

Pest management for potato production in the Andean region of Ecuador is important because potato is a major staple food and is of significant economic value. Our study made the initial effort to document and characterize 2 species of potato tuber moths, *S. tangolias* and *P. operculella*, in Ecuador using PCR and sequencing based on the COI gene. Additionally, we compared our results of *T. solanivora* cytb sequences to a previous study which scrutinized haplotypes of *T. solanivora* in Central and South American countries. Five haplotypes were discovered from the 35 sequenced Andean potato tuber moth *S. tangolias*. The presence of 3 haplotypes was detected in the collected *P. operculella* moth samples. All of the collected Guatemalan potato tuber moths, *T. solanivora*, were genetically identical for the cytb gene sequenced. Mitochondrial gene sequences presented here and archived in GenBank will facilitate further development of molecular markers for diagnostic purposes as well as for scrutinizing phenotypic variation among haplotypes within a species.



**Fig. 5.** Distribution of 3 haplotypes and sequence identity between each pair of *Phthorimaea operculella* haplotypes. Po-H1 was present in all 4 provinces, Po-H3 was present only in Cotopaxi and Chimborazo; Po-H2 was limited to Tungurahua.

Interestingly, one of the *S. tangolias* haplotypes, St-H3, displayed a much greater divergence compared to the other intraspecific haplotypes (> 3%). Both samples that were characterized as St-H3 were from the same province, Chimborazo. When we queried the sequences of these 2 samples in GenBank, the closest homologs (96.5% identity) were 2 *S. tangolias* samples collected from northern Chile, KY951811.1 and KY951905.1. The sequence divergence of 3% also is much higher than what was reported in 3 Lepidopteran families. Hajibabaei et al. (2006) examined several specimens of Hesperidae, Sphingidae, and Saturniidae in a tropical region, and the reported average within-species divergences of COI sequences were 0.17%, 0.43%, and 0.46%, respectively. Contrarily, Cognato (2006) detected a greater range of intraspecific divergence, i.e., 0.4% to 4.8% from Prodoxidae (yucca moth), with an average of 2.2% based on the COI gene. Cognato (2006) also showed overlap between intraspecific and interspecific sequence divergence within Prodoxidae. The present survey also brought our attention to the issues associated with species boundaries and cryptic species despite examining the same region of the COI gene, i.e., the section of COI gene amplified in Hajibabaei et al. (2006) overlapped with our study. Although delimiting species boundaries of *S. tangolias* samples is beyond the scope of this work, we still need to interpret the results with caution. We suggest future work could focus on broader sampling efforts followed by detailed morphological characterization and molecular identification.

Genetic variation and diversity reflect the evolutionary history and ecological adaptation of insect pests, which are vital to implementing effective pest management programs. Many lepidopteran pests are highly mobile and have a great propensity to invade new agricultural regions. Using COI genetic marker, Naik et al. (2020) found 2 common haplotypes of cotton pink ballworms, *Pectinophora gossypiella* Saunders (Lepidoptera: Gelechiidae), across cotton-growing regions of India. The star-shaped haplotype network, with the most common haplotype at the center surrounded by rare haplotypes with a few mutational steps, suggested a recent population expansion of this Gelechiidae moth. In a similar study based on genetic sequences of 2 mitochondrial genes (COII and Nad4), Liu et al. (2010) discovered similar star-shaped phylogeny of *P. gossypiella* across multiple provinces in China. Overall, low genetic diversity of *P. gossypiella* population was attributed to invasion bottlenecks (Liu et al. 2010). The haplotype prevalence in both *S. tangolias* and *P. operculella* in our study were similar to what was described in Liu et al. (2010) and Naik et al. (2020), and that was the presence of 1 dominant haplotype along with multiple less common haplotypes. It is possible that these 2 species underwent population expansion, rapid growth from a small-sized founder population (Graputo et al. 2005; Saw et al. 2006). In order to elucidate the evolutionary history of *S. tangolias* and *P. operculella*, future work is needed to evaluate the population genetics based on both historical samples and samples from broader geographic regions, including those from their native ranges. Genetic variations within species also have been linked

to some behavioral differences. Examples are the observed differences in host preference by 2 fall armyworm strains (Pashley 1986) and the presence of allochronic populations of processionary moth (Santos et al. 2013; Leblois et al. 2018). There were only a handful of studies that examined genetic diversity and variation of Gelechiidae (Liu et al 2010; Medina et al. 2010; Torres-Leguizamón et al. 2011; Naik et al 2020). It is in our future interest to test whether haplotypes of each potato tuber moth species show differences in host preference, behavior, and life-history traits.

Multiple potato tuber moth species may co-exist in potato fields, as demonstrated in our survey. Due to large-scale cultivation of plants in agroecosystems, the presence of multiple insect pest species attacking a single crop becomes a likely scenario (Daamen & Stol 1994; Davidson et al. 2007; Dangles et al. 2008). This could bring bad news for the farmers because the presence of multiple pests, with their species-specific ecology/biology, may exacerbate the effectiveness of implemented management practices. It has been shown recently that the presence of *T. solanivora*, *S. tangolias*, and *P. operculella* together can lead to greater damage to potato crops than when each pest is present alone (Dangles et al. 2009). The underlying mechanisms of this additive effect of multiples pest species remain elusive. Complete resource use and interaction of insect pests may contribute to losses in field crops. Further research is warranted to determine the extent of the additive effect of crop loss due to the coexistence of 3 potato tuber moths in Ecuador.

## Acknowledgments

This work was funded by the International Fund for Agricultural Development (FIDA) through the project “Strengthening innovation to improve income, food security and resilience of potato producers in Bolivia, Ecuador and Peru” of the Centro Internacional de la Papa (CIP), and was executed in collaboration with the Instituto Nacional de Investigaciones Agropecuarias (INIAP). Collections were made under the general agreement for access to resources between the National Institute of Agronomic Research and the Ministry of Environment of Ecuador with reference number MAE-DNB-CM-2015-0024. We would like to dedicate this work to author Rodrigo Aucancela, who lost his life to COVID-19 after submission of this manuscript. Rodrigo (shown in Fig. 1) was a research technician in Instituto Nacional de Investigaciones Agropecuarias, in Chimborazo Province, Ecuador. Rodrigo will continue to inspire us to conduct applied research to reduce the impact of pests and diseases for rural farmers.

## References Cited

- Castillo Carrillo C, Rebaudo F, Asaquiabay C, Aucancela R, Camacho J, López V, Quimbamba V, Yumisaca F, Panchi N, Velasco C. 2018. Informe final de la actividad “Distribución y caracterización molecular de tres especies de polillas de la papa en Ecuador” del proyecto “Fortalecimiento de la innovación para mejorar los ingresos, la seguridad alimentaria y la resiliencia de productores de papa en Bolivia, Ecuador y Perú.” Instituto Nacional de Investigaciones Agropecuarias- Centro Internacional de la Papa (INIAP-CIP). Quito, Ecuador.
- Cognato AI. 2006. Standard percent DNA sequence difference for insects does not predict species boundaries. *Journal of Economic Entomology* 99: 1037–1045.
- Cruz E, Castillo A, Malo E. 2011. First report of *Tecia solanivora* (Lepidoptera: Gelechiidae) attacking the potato *Solanum tuberosum* in Mexico. *Florida Entomologist* 94: 1055–1056.
- Daamen RA, Stol W. 1994. Surveys of cereal diseases and pests in the Netherlands. 6. Occurrence of insect pests in winter wheat. *Netherlands Journal of Plant Pathology* 99: 51–56.
- Dangles O, Carpio C, Barragan AR, Zeddam JL, Silvain JF. 2008. Temperature as a key driver of ecological sorting among invasive pest species in the tropical Andes. *Ecological Applications* 18: 1795–1809.
- Dangles O, Mesías V, Crespo Perez V, Silvain JF. 2009. Crop damage increases with pest species diversity: evidence from potato tuber moths in the tropical Andes. *Journal of Applied Ecology* 46: 1115–1121.
- Davidson SA, Peairs FB, Khosla R. 2007. Insect pest densities across site-specific management zones of irrigated corn in Northeastern Colorado. *Journal of Economic Entomology* 100: 781–789.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797.
- EPPO – EPPO Global Database. 2015. Reporting Service no. 11. <https://gd.eppo.int/reporting/article-5149> (last accessed 2 Jun 2020).
- Gallegos P. 1997. Informe anual DNPV y PNRT. EESC, INIAP. Quito, Ecuador.
- Gallegos P, Suquillo J, Chamorro F, Asaquiabay C. 2004. Alternativas de manejo de *Tecia solanivora* (Povolny) (Lepidoptera: Gelechiidae) en Ecuador. *Memorias del II Taller Internacional de Polilla Guatemalteca*. Pontificia Universidad Católica del Ecuador. Quito, Ecuador.
- Graft JE. 1917. The potato tuber moth. USDA Technical Bulletin 427. Washington, DC, USA.
- Grapputo A, Boman S, Lindstroem L, Lyytinen A, Mappes J. 2005. The voyage of an invasive species across continents: genetic diversity of North American and European Colorado potato beetle populations. *Molecular Ecology* 14: 4207–4219.
- Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN. 2006. DNA barcodes distinguish species of tropical Lepidoptera. *Proceedings of the National Academy of Sciences of the United States of America* 103: 968–971.
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Kroschel J, Schaub B. 2013. Biology and ecology of potato tuber moths as major pests of potato, pp. 165–192. In Giordanengo P, Vincent C, Alyokhin I (eds.), *Pests of Potato: Global Perspectives on Biology and Management*. Academic Press, London, United Kingdom.
- Kroschel J, Sporleder M, Tonnang HEZ, Juarez H, Carhuapoma P, Gonzales JC, Simon R. 2013. Predicting climate-change-caused changes in global temperature on potato tuber moth *Phthorimaea operculella* (Zeller) distribution and abundance using phenology modeling and GIS mapping. *Agricultural and Forest Meteorology* 70: 228–241.
- Leblois R, Gautier M, Rohfritsch A, Foucaud J, Burban C, Galan M, Loiseau A, Sauné L, Branco M, Gharbi K, Vitalis R, Kerdelhué C. 2018. Deciphering the demographic history of allochronic differentiation in the pine processionary moth *Thaumetopoea pityocampa*. *Molecular Ecology* 27: 264–278.
- Liu Y, Hou M, Wu K. 2010. Genetic variation of mitochondrial DNA in Chinese populations of *Pectinophora gossypiella* (Lepidoptera: Gelechiidae). *Environmental Entomology* 39: 1344–1351.
- Martin NA. 1999. Arthropods and molluscs associated with poroporo (*Solanum aviculare* and *S. laciniatum*): an annotated species list. *Journal of the Royal Society of New Zealand* 29: 65–76.
- Medina RF, Rondon SI, Reyna SM, Dickey AM. 2010. Population structure of *Phthorimaea operculella* (Lepidoptera: Gelechiidae) in the United States. *Environmental Entomology* 39: 1037–1042.
- Naik VC, Pusadkar PP, Waghmare ST, Raghavendra KP, Kranthi S, Kumbhare S, Nagare VS, Kumar R, Prabhulinga T, Gokte-Narkhedkar N, Waghmare VN. 2020. Evidence for population expansion of cotton pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) in India. *Scientific Reports* 10: 4740. <https://doi.org/10.1038/s41598-020-61389-1>
- Pashley DP. 1986. Host-associated genetic differentiation in fall armyworm (Lepidoptera: Noctuidae): a sibling species complex? *Annals of the Entomological Society of America* 79: 898–904.
- Rincón D, López-Ávila A. 2004. Dimorfismo sexual en pupas de *Tecia solanivora* (Povolny) (Lepidoptera: Gelechiidae). *Revista Corpoica* 5: 41–42.
- Rondon SI, Gao Y. 2018. The journey of the potato tuberworm around the world, pp. 1–37. In Khan Perveen F [ed.], *Moths-Pests of Potato, Maize and Sugar Beet*. IntechOpen. doi: 10.5772/intechopen.81934
- Salasar J, Escalante W. 1984. La polilla guatemalteca de la papa, *Scrobipalopsis solanivora*, nueva plaga del cultivo de la papa en Venezuela. *Sociedad Venezolana de Ingenieros Agrónomos* 9: 24–28.
- Santos HM, Paiva MR, Rocha S, Kerdelhué C, Branco M. 2013. Phenotypic divergence in reproductive traits of a moth population experiencing a phenological shift. *Ecology and Evolution* 3: 5098–5108.
- Saw J, Endersby NM, McKechnie SW. 2006. Low mtDNA diversity among widespread Australian diamondback moth *Plutella xylostella* (L.) suggests isolation and a founder effect. *Insect Science* 13: 365–373.
- Sint D, Sporleder M, Wallinger C, Zegarra O, Oehm J, Dangi N, Traugott M. 2016. A two-dimensional pooling approach towards efficient detection of parasitoid and pathogen DNA at low infestation rates. *Methods in Ecology and Evolution* 12: 1548–1557.

- Sporleder M, Carhuapoma P, Kroschel J. 2016a. Guatemalan potato tuber moth, *Tecia solanivora* (Povolny 1973), pp. 24–38 *In* Kroschel J, Mujica N, Carhuapoma P, Sporleder M [eds.], Pest Distribution and Risk Atlas for Africa. Potential Global and Regional Distribution and Abundance of Agricultural and Horticultural Pests and Associated Biocontrol Agents under Current and Future Climates. International Potato Center (CIP), Lima, Peru.
- Sporleder M, Carhuapoma P, Kroschel J. 2016b. Andean potato tuber moth, *Symmetrischema tangolias* (Gyen 1913), pp. 39–53 *In* Kroschel J, Mujica N, Carhuapoma P, Sporleder M [eds.], Pest Distribution and Risk Atlas for Africa. Potential Global and Regional Distribution and Abundance of Agricultural and Horticultural Pests and Associated Biocontrol Agents under Current and Future Climates. International Potato Center (CIP), Lima, Peru.
- Terauds A, Rapley PEL, Williams MA, Ireson JE, Miller LA, Brieze-Stegeman R, McQuillan PB. 1984. Insect pest occurrences in Tasmania 1982/83. <https://www.plantwise.org/KnowledgeBank/pestalert?pan=19850521177> (last accessed 7 Jun 2020).
- Torres-Leguizamón M, Dupas S, Dardon D, Gómez Y, Nino L, Carnero A, Lery X. 2011. Inferring native range and invasion scenarios with mitochondrial DNA: the case of *T. solanivora* successive north-south step-wise introductions across Central and South America. *Biological Invasions* 13: 1505–1519.
- Trujillo E, Ríos D, Cabrera R. 2004. Distribución de *Tecia solanivora* (Povolny) (Lepidoptera: Gelechiidae) en Tenerife, Islas Canarias, España. *Memorias del II Taller Internacional de Polilla Guatemalteca*. Pontificia Universidad Católica del Ecuador. Quito, Ecuador.