

Diet with Sucrose Ameliorates Solenopsis Invicta Virus 3 (Solinviviridae: Invictavirus) Infection in Solenopsis invicta (Hymenoptera: Formicidae) Worker Ants

Author: Valles, Steven M.

Source: Florida Entomologist, 104(3) : 239-242

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.104.0314>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Diet with sucrose ameliorates *Solenopsis invicta* virus 3 (Solinviviridae: Invictavirus) infection in *Solenopsis invicta* (Hymenoptera: Formicidae) worker ants

Steven M. Valles^{1,*}

Solenopsis invicta Buren (Hymenoptera: Formicidae) is an invasive ant pest that was introduced into the US near Mobile, Alabama, USA, in the early 1900s (Tschinkel 2006). Control efforts and damage repair from *S. invicta* were estimated to exceed \$5 billion annually in 2003 (Pereira 2003), which would exceed \$8.5 billion today after adjusting for inflation. Although insecticides are highly effective at controlling this ant pest, they must be used on a regular interval to maintain control. If insecticide use is stopped, fire ant populations will invariably re-infest the area. The invasive success of *S. invicta* has been attributed to human-assisted spread, pre-adaptation for colonization of new areas (Ascunce et al. 2011), and a lack of natural enemies during founding (Porter et al. 1997).

Solenopsis invicta virus 3 (Solinviviridae: Invictavirus) has been shown to be an effective natural control agent for *S. invicta* in the laboratory (Valles et al. 2013). The virus exhibits desirable characteristics as a control agent, including species specificity, ready transmission, pathogenicity, and virulence (Porter et al. 2013; Valles et al. 2013, 2014; Valles & Oi 2014; Porter et al. 2015). The *Solenopsis invicta* virus 3 infection is stage-dependent; virus replication occurs only in adult ant stages. Infected adult workers stop foraging for food (Chen et al. 2012), which results in a cascade of events including starvation, a severe reduction in queen fecundity (Valles et al. 2013), and ultimate colony collapse (Valles et al. 2014). During laboratory tests involving *Solenopsis invicta* virus 3, it was noticed that the impact of *Solenopsis invicta* virus 3 on worker mortality appeared to be reduced when a sugar solution was available to the infected ant colonies as part of their diet. Therefore, the objective of this work was to test the hypothesis that the availability of a sucrose solution as part of the diet could influence the ability of *S. invicta* workers to better tolerate *Solenopsis invicta* virus 3 infection.

Two polygyne *S. invicta* colonies were collected from the field in Gainesville, Florida, USA, removed from the nest soil by flooding, placed into rearing trays, and provided an ad libitum diet of crickets (*Acheta domesticus* [L.]; Orthoptera: Gryllidae), 10% sucrose, and water. A sample of worker ants ($n = 15$) from each colony was examined by microscopy (Knell & Allen 1977) for the presence of the microsporidian, *Kneallhazia solenopsae* (Knell, Allen, & Hazard) (Microsporida: Thelohaniidae) (Williams et al. 1999). Reverse Transcription-Polymerase Chain Reaction (RT-PCR) also was conducted on the sample to detect *Solenopsis invicta* virus 1, *Solenopsis invicta* virus 2, and *Solenopsis*

invicta virus 3 (Valles et al. 2009). Both colonies were found to be free of infection by these viruses and the microsporidian.

The ant colonies were permitted to acclimate to the laboratory for 2 wk. The *S. invicta* colonies were divided equally, and the queens removed. One sub-colony from each parent colony was infected with *Solenopsis invicta* virus 3 and the other sub-colonies were retained as control groups. A homogenate prepared from *Solenopsis invicta* virus 3-infected *S. invicta* colonies was used as the source of *Solenopsis invicta* virus 3 inoculum, which can successfully transmit the virus to uninfected ant colonies in the laboratory (Valles & Hashimoto 2009). *Solenopsis invicta* virus 3-infected worker ants (about 20 g) were blended with 40 mL of 10% (w/v) sucrose prepared with deionized water for 1 min at high speed. The homogenate was filtered through 3 layers of cheesecloth and then filtered by vacuum in a Buchner funnel (ThermoFisher Scientific Company, Waltham, Massachusetts, USA) through a number 1 Whatman paper (SigmaAldrich, St. Louis, Missouri, USA). The homogenate (10 mL) was mixed with 10 freeze-killed adult house crickets (*A. domesticus*) that were pulverized with a mortar and pestle to create a crude paste. Each of the 2 sub-colonies received 5 mL of homogenate paste for 24 h, after which the remaining homogenate paste was removed. A control group homogenate paste was prepared identically using *Solenopsis invicta* virus 3-uninfected worker ants, and the control sub-colonies were exposed in identical fashion. After 5 d, 10 individual worker ants were tested by quantitative polymerase chain reaction (qPCR) for the presence of *Solenopsis invicta* virus 3 to verify successful infection transmission (Valles & Hashimoto 2009). The sub-colonies were shown to have a weak, but established, *Solenopsis invicta* virus 3 infection. All 10 individual workers (100%) sampled from each sub-colony were shown to be infected. The mean (\pm standard deviation) *Solenopsis invicta* virus 3 titer in each worker of colony 1 and colony 2 was $8.95 \times 10^2 \pm 5.31 \times 10^2$ and $3.72 \times 10^3 \pm 8.75 \times 10^3$ *Solenopsis invicta* virus 3 genome equivalents per ng RNA, respectively.

Fragment colonies were prepared from each of the *Solenopsis invicta* virus 3-infected and -uninfected sub-colonies as experimental units. Each fragment was comprised of 1 mL of brood and 2 mL of workers. All colony fragments received a diet of a cricket every other d and water in a cotton-stoppered test tube. The treatment groups were distinguished by the addition, or not, of a 10% (w/v) sugar solution as

¹Center for Medical, Agricultural and Veterinary Entomology, USDA-ARS, Gainesville, Florida, 32608, USA; E-mail: steven.valles@usda.gov (S. M. V.)

*Corresponding author; E-mail: steven.valles@usda.gov

part of the diet, and included *Solenopsis invicta* virus 3-infected fragment colonies provided a 10% sugar solution ($n = 6$; 3 replicates per sub-colony), *Solenopsis invicta* virus 3-infected fragment colony without a 10% sugar solution ($n = 6$; 3 replicates per sub-colony), *Solenopsis invicta* virus 3-uninfected fragment colonies provided a 10% sugar solution ($n = 3$; 3 replicates per sub-colony), *Solenopsis invicta* virus 3-uninfected fragment colonies without a 10% sugar solution ($n = 3$; 3 replicates per sub-colony). One of the control groups became infected with *Solenopsis invicta* virus 3 during the course of the study and was not used. Worker mortality was recorded 1, 3, 5, 10, 12, 14, 17, 19, and 21 d after the introduction of specific diets, and was expressed as cumulative mortality over the duration of the study. *Solenopsis invicta* virus 3 was quantified in the ants that died. Total RNA was extracted from the pooled dead worker ants (pooled by d and replicate) and used as a template for reverse transcription and subsequent qPCR (Valles & Hashimoto 2009). *Solenopsis invicta* virus-3 also was quantified in live and dead workers from each colony fragment on d 24.

Cumulative mortality among groups (*Solenopsis invicta* virus-infected with sugar supplement; *Solenopsis invicta* virus-infected without sugar supplement; *Solenopsis invicta* virus-uninfected with sugar supplement; *Solenopsis invicta* virus-uninfected without sugar supplementation) on d 21 was compared by Analysis of Variance followed by

Scheffe's mean separation test (SAS 2009). Mortality by d (1, 3, 5, 10, 12, 14, 17, 19, and 21) for each treatment (*Solenopsis invicta* virus-infected with sugar supplement; *Solenopsis invicta* virus-infected without sugar supplement; *Solenopsis invicta* virus-uninfected with sugar supplement; *Solenopsis invicta* virus-uninfected without sugar supplement) was compared by Analysis of Variance followed by Scheffe's mean separation test (SAS 2009). Student's *t*-test (SAS 2009) was used to compare the quantity of *Solenopsis invicta* virus 3 in *Solenopsis invicta* virus-infected with sugar supplement and *Solenopsis invicta* virus-infected without sugar supplement groups for each d (*Solenopsis invicta* virus 3 was not present in the control groups). Student's *t*-test also was used to compare the quantity of *Solenopsis invicta* virus 3 detected in dead workers (*Solenopsis invicta* virus-infected with sugar supplement versus *Solenopsis invicta* virus-infected without sugar supplement) and live workers (*Solenopsis invicta* virus-infected with sugar supplement versus *Solenopsis invicta* virus-infected without sugar supplement) collected on d 24.

Cumulative mortality increased significantly over the 3-wk monitoring period in the *Solenopsis invicta* virus 3-infected colonies with ($F = 16.7$; $df = 8,45$; $P < 0.0001$) and without ($F = 11.6$; $df = 8,45$; $P < 0.0001$) sugar supplementation and the *Solenopsis invicta* virus 3-uninfected colonies without sugar supplementation ($F = 11.6$; $df = 8,18$; $P < 0.0001$) (Fig. 1). There was no significant difference ($F = 2.24$; $df =$




Fig. 1. Cumulative *Solenopsis invicta* worker ant mortality among *Solenopsis invicta* virus 3-infected and -uninfected colonies provided a diet of crickets (*Acheta domesticus*) and either supplemented (open symbols) or not supplemented (solid symbols) with a 10% sucrose solution. Analysis of Variance by treatment was conducted for d 21 values and found to be significant ($F = 10.0$; $df = 3,14$; $P < 0.0009$). Scheffe's multiple comparison procedure was used to separate the means. Symbols with the same letter are not statistically different.

8,18; $P < 0.074$) in cumulative mortality between the *Solenopsis invicta* virus 3-uninfected colonies with and without sugar supplementation. Mortality on d 21 reached a mean of 111.8 ± 22.0 worker ants in *Solenopsis invicta* virus 3-treated colonies without sugar supplementation, which was significantly ($F = 10.0$; $df = 3,14$; $P < 0.0009$) greater than the *Solenopsis invicta* virus 3-infected colonies with sugar supplementation (33.5 ± 5.8), *Solenopsis invicta* virus 3-uninfected colonies without sugar supplementation (17.3 ± 10.7), and *Solenopsis invicta* virus 3-uninfected colonies with sugar supplementation (6.7 ± 2.5).

The quantity of *Solenopsis invicta* virus 3 in the dead worker ants generally was higher (by about 1 order of magnitude) in colonies without sugar supplementation compared with colonies with sugar supplementation (Fig. 2). However, significant differences were noted only on d 12 ($t = 2.52$; $df = 9$; $P < 0.033$) and 19 ($t = 2.44$; $df = 9$; $P < 0.037$). *Solenopsis invicta* virus 3 was not detected in either of the control (virus-uninfected) colonies (Fig. 2). A final comparison of *Solenopsis invicta* virus 3 quantity was made on d 24 among live and dead workers in the *Solenopsis invicta* virus 3-infected colonies with and without sugar supplementation. Among dead workers collected on that d, the mean ($2.21 \times 10^6 \pm 6.38 \times 10^6$ *Solenopsis invicta* virus 3 genome equivalents per ng RNA) amount of *Solenopsis invicta* virus 3 was greater significantly ($t = -3.14$; $df = 9$; $P = 0.012$) in colonies without sugar supple-

mentation compared with colonies with sugar supplementation ($8.55 \times 10^5 \pm 6.06 \times 10^5$ *Solenopsis invicta* virus 3 genome equivalents per ng RNA). Among live worker ants in the *Solenopsis invicta* virus 3-infected colonies, there was no significant difference in the quantity of *Solenopsis invicta* virus 3 detected in colonies without sugar supplementation ($3.17 \times 10^6 \pm 2.14 \times 10^6$ *Solenopsis invicta* virus 3 genome equivalents per ng RNA) or with sugar supplementation ($2.53 \times 10^5 \pm 1.47 \times 10^5$ *Solenopsis invicta* virus 3 genome equivalents per ng RNA). However, the trend was higher in the colonies without sugar supplementation.

Adequate nutrition is essential for normal insect development and propagation (Dadd 1973), but it also influences other aspects of survival, including resistance to pathogen infection (Muturi et al. 2011; Di Pasquale et al. 2013). *Solenopsis invicta* virus 3-treated colonies without a sugar supplement to their diet exhibited significantly higher mortality rates and generally higher *Solenopsis invicta* virus 3 titers than their counterparts that were provided a sugar supplement. During virus development, especially in the chronic to acute phase of pathogenesis, glucose demands may increase significantly to support the energetic and anabolic demands of virus replication (Wang et al. 2019). These increased metabolic demands often are detrimental to the insect host. Honey bees infected with various RNA viruses (similar to *Solenopsis invicta* virus 3) exhibit higher mortality with poor qual-




Fig. 2. *Solenopsis invicta* virus 3 genome equivalents per ng RNA from dead *Solenopsis invicta* worker ants among *Solenopsis invicta* virus 3-infected and -uninfected colonies provided a diet of crickets (*Acheta domesticus*) and either supplemented (open symbols) or not supplemented (solid symbols) with a 10% sucrose solution. *Solenopsis invicta* virus 3 was not detected in the *Solenopsis invicta* virus 3-uninfected group. Student's *t*-test was conducted to compare the virus quantity in colonies with and without the sucrose supplement by d. *Solenopsis invicta* virus 3 genome equivalents per ng RNA was greater significantly in colonies without sugar supplementation on d 12 ($t = 2.5$; $df = 9$; $P < 0.033$) and 19 ($t = 2.4$; $df = 9$; $P < 0.037$).

ity diets (Schmidt et al. 1987; Dolezal et al. 2019). Poorly nourished bees mount a weaker immune response (Alaux et al. 2010) and exhibit higher virus titers (DeGrandi-Hoffman et al. 2010). Sucrose supplementation during virus infection may help to alleviate the impact of greater biosynthetic material demands from the host by a virus in the active replication phase, as observed with *Solenopsis invicta* virus 3-infected *S. invicta*. This knowledge may prove important when developing *Solenopsis invicta* virus 3 as a natural control agent against *S. invicta*. It is known that the prevalence of *Solenopsis invicta* virus 3 among *S. invicta* is greater during the winter months (Valles et al. 2010). Scarcity of carbohydrate sources (for example, honeydew [Helms & Vinson 2002]) during the winter may make *S. invicta* less able to resist *Solenopsis invicta* virus 3 infection.

Summary

Mortality and virus titer were monitored in *Solenopsis invicta* colony fragments to examine the impact of diet sucrose supplementation. Mortality on d 21 reached a mean of 111.8 ± 22.0 worker ants in *Solenopsis invicta* virus 3-treated colonies without sugar supplementation, which was significantly greater ($F = 10.0$; $df = 3,14$; $P < 0.0009$) than the *Solenopsis invicta* virus 3-infected colonies with sugar supplementation (33.5 ± 5.8), *Solenopsis invicta* virus 3-uninfected colonies without sugar supplementation (17.3 ± 10.7), and *Solenopsis invicta* virus 3-uninfected colonies with sugar supplementation (6.7 ± 2.5).

Key Words: nutrition; RNA virus; fire ant; virus replication; virus pathogenesis

Sumario

Se monitorearon la mortalidad y la toxicidad del virus en fragmentos de colonias de *Solenopsis invicta* para examinar el impacto de la suplementación con sacarosa en la dieta. La mortalidad en el día 21 alcanzó un promedio de $111,8 \pm 22,0$ en hormigas obreras en colonias tratadas con *Solenopsis invicta* virus 3 sin suplementos de azúcar, que fue significativamente mayor ($F = 10,0$; $gl = 3,14$; $P < 0,0009$) que el *Solenopsis invicta* virus 3-colonias infectadas con suplementación con azúcar ($33,5 \pm 5,8$), colonias no infectadas con el virus *Solenopsis invicta* 3-colonias no infectadas sin suplementación con azúcar ($17,3 \pm 10,7$), y colonias no infectadas con el virus *Solenopsis invicta* 3-con suplemento con azúcar ($6,7 \pm 2,5$).

Palabras Clave: nutrición; Virus de ARN; hormiga de fuego; replicación de virus; patogenia del virus

References Cited

- Alaux C, Ducloz F, Crauser D, Le Conte Y. 2010. Diet effects on honeybee immunocompetence. *Biology Letters* 6: 562–565.
- Ascunce MS, Yang CC, Oakey J, Calcaterra L, Wu WJ, Shih CJ, Goudet J, Ross KG, Shoemaker D. 2011. Global invasion history of the fire ant *Solenopsis invicta*. *Science* 331: 1066–1068.
- Chen Y, Becnel JJ, Valles SM. 2012. RNA viruses infecting pest insects, pp. 133–170. In Vega F, Kaya HK [eds.], *Insect Pathology*, Second edition. Elsevier, Amsterdam, Netherlands.
- Dadd RH. 1973. Insect nutrition: current developments and metabolic implications. *Annual Review of Entomology* 18: 381–420.
- DeGrandi-Hoffman G, Chen Y, Huang E, Huang MH. 2010. The effect of diet on protein concentration, hypopharyngeal gland development and virus load in worker honey bees (*Apis mellifera* L.). *Journal of Insect Physiology* 56: 1184–1191.
- Di Pasquale G, Salignon M, Le Conte Y, Belzunces LP, Decourtaye A, Kretzschmar A, Suchail S, Brunet JL, Alaux C. 2013. Influence of pollen nutrition on honey bee health: do pollen quality and diversity matter? *PLoS ONE* 8: e72016. doi: 10.1371/journal.pone.0072016
- Dolezal AG, Carrillo-Tripp J, Judd TM, Allen Miller W, Bonning BC, Toth AL. 2019. Interacting stressors matter: diet quality and virus infection in honeybee health. *Royal Society Open Science* 6: 181803. doi: 10.1098/rsos.181803
- Helms KR, Vinson SB. 2002. Widespread association of the invasive ant *Solenopsis invicta* with an invasive mealybug. *Ecology* 83: 2425–2438.
- Knell JD, Allen GE. 1977. Light and electron microscope study of *Thelohania solenopsae* n. sp. (Microsporidia: Protozoa) in the red imported fire ant, *Solenopsis invicta*. *Journal of Invertebrate Pathology* 29: 192–200.
- Muturi EJ, Kim CH, Alto BW, Berenbaum MR, Schuler MA. 2011. Larval environmental stress alters *Aedes aegypti* competence for Sindbis virus. *Tropical Medicine International Health* 16: 955–964.
- Pereira RM. 2003. Areawide suppression of fire ant populations in pastures: project update. *Journal of Agricultural and Urban Entomology* 20: 123–130.
- Porter SD, Valles SM, Oi DH. 2013. Host specificity and colony impacts of *Solenopsis invicta* virus 3. *Journal of Invertebrate Pathology* 114: 1–6.
- Porter SD, Williams DF, Patterson RS, Fowler HG. 1997. Intercontinental differences in the abundance of *Solenopsis* fire ants (Hymenoptera: Formicidae): escape from natural enemies? *Environmental Entomology* 26: 373–384.
- Porter SD, Valles SM, Wild AL, Dieckmann R, Plowes NJR. 2015. *Solenopsis invicta* virus 3: further host-specificity tests with native *Solenopsis* ants (Hymenoptera: Formicidae). *Florida Entomologist* 98: 122–125.
- SAS. 2009. Base SAS 9.2 Procedures Guide. SAS Institute, Cary, North Carolina, USA.
- Schmidt JO, Thoenes SC, Levin MD. 1987. Survival of honey-bees, *Apis mellifera* (Hymenoptera, Apidae), fed various pollen sources. *Annals of the Entomological Society of America* 80: 176–183.
- Tschinkel WR. 2006. The fire ants. The Belknap Press of Harvard University Press, Cambridge, Massachusetts, USA.
- Valles SM, Hashimoto Y. 2009. Isolation and characterization of *Solenopsis invicta* virus 3, a new positive-strand RNA virus infecting the red imported fire ant, *Solenopsis invicta*. *Virology* 388: 354–361.
- Valles SM, Oi DH. 2014. Successful transmission of *Solenopsis invicta* virus 3 to field colonies of *Solenopsis invicta* (Hymenoptera: Formicidae). *Florida Entomologist* 97: 1244–1246.
- Valles SM, Oi DH, Porter SD. 2010. Seasonal variation and the co-occurrence of four pathogens and a group of parasites among monogyne and polygyne fire ant colonies. *Biological Control* 54: 342–348.
- Valles SM, Porter SD, Firth AE. 2014. *Solenopsis invicta* virus 3: pathogenesis and stage specificity in red imported fire ants. *Virology* 461: 66–71.
- Valles SM, Porter SD, Choi MY, Oi DH. 2013. Successful transmission of *Solenopsis invicta* virus 3 to *Solenopsis invicta* fire ant colonies in oil, sugar, and cricket bait formulations. *Journal of Invertebrate Pathology* 113: 198–204.
- Valles SM, Varone L, Ramirez L, Briano J. 2009. Multiplex detection of *Solenopsis invicta* viruses -1, -2, and -3. *Journal of Virological Methods* 162: 276–279.
- Wang LL, Swevers L, Rombouts C, Meeus I, Van Meulebroek L, Vanhaecke L, Smaghe G. 2019. A metabolomics approach to unravel Cricket Paralysis Virus infection in silkworm Bm5 cells. *Viruses* 11: 861. doi: 10.3390/v11090861
- Williams DF, Oi DH, Knue GJ. 1999. Infection of red imported fire ant (Hymenoptera: Formicidae) colonies with the entomopathogen *Thelohania solenopsae* (Microsporidia: Thelohaniidae). *Journal of Economic Entomology* 92: 830–836.