

Reversing the Decline of Secondary Spruce Forests in Slovakia's Western Carpathians

Authors: Antoni, Jozef, Šomšák, Ladislav, and Jansk'y, Libor

Source: Mountain Research and Development, 20(2): 130-131

Published By: International Mountain Society

URL: https://doi.org/10.1659/0276-4741(2000)020[0130:RTDOSS]2.0.CO;2

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Jozef Antoni Ladislav Šomšák Libor Janský

130

Reversing the Decline of Secondary Spruce Forests in Slovakia's Western Carpathians

The Western Carpathian forests are of extraordinary importance, not only as production forests but also in terms of general and local public interest. Apart from being the source of economically vital products for local people, they play a very important role for neighboring countries in protecting soil and water resources in which these countries also have an interest. The total forest acreage in Slovakia has increased by more than 10% during the last 40 years, and total growing stock represents 190 m³ per hectare. But the condition of forests in the last decades has markedly deteriorated (Figures 1 and 2). Slovakia is currently exploring ways and means of restoring its forests to a healthy condition.



FIGURE 1 The beginning of the collective spruce decline in 1988, Nálepkovo district, Western Carpathians, Slovak Republic. (Photo by Ladislav Šomšák)



FIGURE 2 Disastrous gap as a result of collective decline of spruce stands in 1993, Nálepkovo district, Western Carpathians, Slovak Republic. (Photo by Ladislav Šomšák)

A long tradition of research and training

The Slovak Republic is a country with an established forestry tradition in the Western Carpathians. In 1762, on the basis of a decision made by the Empress Maria Theresa, a Mining and Forestry Academy was founded in Banská Šiavnica. In 1824, the Forestry Institute was added. It was the world's first institution of higher learning of its type; foresters from almost every European country studied there. By the end of the 19th century, an independent Research Forestry Institute had also been founded in Banská Štiavnica—one of the first in Europe.

Carpathian forests in the Slovak Republic cover approximately 2 million hectares-this represents 41% of the country's total surface. About 40% of this forest cover is state forests; the remaining 60% consists mainly of forests that belong to cooperatives, individuals, municipal communities, or churches. By comparison with surrounding countries, a special feature of the forests in Slovakia is the variety of different natural conditions and types of forests in a relatively small area, from lowland to mountain formations. Furthermore, a wide range of original tree species and mountain communities has been preserved. Mixed stands of spruce, fir, and beech are the typical forest community of mountain areas in Slovakia.

European-wide deterioration

The massive and disastrous decline of the mountain forests of Central Europe is not a new phenomenon. In the past, it occurred particularly in forests whose composition had changed. Wind and snowstorm disasters damaged the forests to such an extent that it was impossible to process the fallen wood in time; damaged areas developed into centers for the dispersion of bark beetles to surrounding stands, causing harm to the spruce monocultures planted for productive purposes. Although such forests may seem to be healthy and stable, when they are 80–100 years old, epidemics of wood beetle (mainly *Ips typographus*) can appear rapidly, damaging hundreds of hectares.

In recent years, the most severe deterioration has been observed in the mountains of Germany, the Czech Republic, Poland, and Slovakia. The main factors leading to the declining condition of tree species have been drought and subsequent insect attacks, exacerbated by air pollution from distant sources as well as local industries. In Slovakia, coniferous forests dominated by Norway spruce (Picea abies) were hit first. In the Western Carpathians, the condition of secondary spruce forests has declined over the past 20 years, but wind and snowstorms have not been the primary cause of bark beetle dispersion. In 1996, the infestation of spruce forests in Slovakia decreased by 29% (570,000 m³) compared with the previous year but is still at the second highest level since 1972, when the physiological decline of these forests began. Rare, randomly found cases of forest damage occurred on a large scale starting around 1985 and were identified as the first and fourth stages of forest damage severity. In 1989, the general forest tree injury for all stages of severity accounted for 85% of all damage. Present monitoring reveals a relative stagnation in the condition of forests; however, further deterioration is expected for the next 2–5 years.

Nálepkovo forest: A very slow process of recovery

The municipal spruce forest of Nálepkovo, situated in central-eastern Slovakia at wood-based panels.

expansion of bark beetle populations, har-

vests of wood were four to five times high-

processing industries processed the wood

Reclamation experiments began in 1995–1996, when sycamore maple and lin-

den trees able to survive under such con-

ditions and to act as hyperaccumulators of

pollutants from soil were planted in small

groups. Subsequently, a new tree composi-

tion for the damaged areas was proposed,

with 20% of each of the following species:

common beech (Fagus silvatica), silver fir

(Abies alba), Norway spruce (Picea abies),

sycamore maple (*Acer pseudoplatanus*), and linden (*Tilia cordata*). These were planted

in small groups at two monitoring plots at

altitudes of 600 and 900 m. Due to the

high acidity of the soil, half of each plot was limed (4 tonnes/ha). Scientists from

the Department of Soil Science, Faculty of

Natural Science, Comenius University in

Bratislava, will carry out further monitor-

ing and analytical activities to continue to

evaluate the causes of collective decline of

ues across the region. Reclamation experi-

ments—including liming, as the first 'gentle' action to stop the process—have not

influenced environmental conditions at

the site, in the local area, or downstream.

Fish breeding, game keeping, and tourism

have all been affected and, with these, the

in identifying and addressing the causes of

local communities and in further improve-

ment of the environmental conditions of

livelihoods of local people. Any progress

forest decline will be of great value to

The decline of spruce forests contin-

Norway spruce in the area.

into industrial roundwood, lumber, and

er than allowed. Local forest and wood

FIGURES 3 AND 4

Consequences of the group decline of spruce forests during 1986–1996, Nálepkovo district, Western Carpathians, Slovak Republic. (Photos by Ladislav Šomšák)



131



FURTHER READING

Butt N, Price MF, editors. 2000. Mountain People, Forests and Trees: Strategies for Balancing Local Management and Outside Interests. Report of the Mountain Forum electronic conference, April–May 1999. Franklin, WV: The Mountain Institute. European Commission. 1996. Forest Condition in Europe. Results of the 1995 Survey. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. Brussels: UN Economic Commission for Europe, European Commission.

FAO. 1997. Agricultural Reforms in the Transition Economies of Central and Eastern Europe. Rome: FAO.

Novotný J, editor. 1997. Forestry of the Slovak Republic. Bratislava: Poly-Kontakt. Novotný J, et al. 1997. The most significant injurious factors in the forests of Slovakia—the present situation. Slovak Forest 53:14–17.

Šomšák L, editor. 1997. Decline of secondary spruce forest of the Nálepkovo village. Proceedings of Conference on 16 December 1997. Bratislava: Department of Soil Science, Faculty of Science, Comenius University.

Valtýni J. 1986. Importance of Forest for Water Management and Water Protection. Volume 38, *Forest Studies*. Bratislava: Príroda.

an altitude of 450-1200 m and 3200 ha in area (60% with slopes of $20-45^{\circ}$), has been damaged significantly. The composition of today's forest is different from the original forest, dominated by common beech (Fagus silvatica), hornbeam (Carpinus betulus), silver fir (Abies alba), linden (Tilia cordata), and sycamore maple (Acer pseudoplatanus). Three to four centuries ago, this forest was converted to a secondary spruce forest in order to produce more coniferous timber for the region's expanding mining industry. The yearly allowable harvest is now about 5000 m³, 70% for logs and 30% for fuelwood. Over the centuries, the coniferous monocultures have increased the soil acidity significantly, from pH 5-6 to pH 3.8-4. This increased acidification seems to be one cause of the decline in the region's spruce forests.

In 1990, the Forest State Authority declared the Nálepkovo forest to be "extremely endangered" (Figures 3 and 4) and began research to evaluate the causes of this condition. The following causes of forest decline were explored: drought, soil acidification, toxic pollutants (mostly heavy metals and arsenic elements from local or remote sources), the degree of organic matter accumulation, disorders in microbiological processes, and secondary succession changes.

The participation of local people and communities in the research has been significant and very helpful in routine monitoring and in collecting various environmental data, including climatic data. From the beginning, the Nálepkovo municipal authorities covered all research expenses.

In 1995–1997, because of the need to extract dead trees in order to limit the

Libor Janský

the entire region.

The United Nations University, 53-70, Jingumae 5-chome, Shibuya-ku, Tokyo 150-8925, Japan. Jansky@hq.unu.edu Libor Janský is Senior Academic Program Officer in the field of environment and sustainable development at the

United Nations University in Tokyo, Japan. A forester by training, he received his PhD from the Faculty of Forestry, University of Zvolen, Slovakia. Prior to joining UNU in 1999, he was involved in research and teaching at the Faculty of Natural Science, Comenius University in Bratislava, Slovakia, focusing on aspects of soil and water conservation and land reclamation issues in agriculture and forestry. He has held several visiting research positions in Japan and the UK.

AUTHORS

Jozef Antoni

Municipal Forests, Nálepkovo, Slovak Republic. Jozef Antoni is Director of the Municipal Forests in Nálepkovo, Slovak Republic.

Ladislav Šomšák

Faculty of Sciences, Comenius University, 842 15 Bratislava, Slovak Republic.

Ladislav Šomšák is Professor of Geobotany at the Faculty of Natural Science, Comenius University in Bratislava, Slovak Republic.

Downloaded From: https://complete.bioone.org/journals/Mountain-Research-and-Development on 01 Jun 2025 Terms of Use: https://complete.bioone.org/terms-of-use