

Isolation and Characterization of Multiple-Lectins from Serum of the Desert Locust *Schistocerca gregaria* (Orthoptera: Acrididae)

Authors: Dorrah, Moataza A., Ayaad, Tahany H., Mohamed, Amr A., and Bassal, Taha T.M.

Source: Journal of Orthoptera Research, 18(1) : 103-112

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/034.018.0113>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Isolation and characterization of multiple-lectins from serum of the desert locust *Schistocerca gregaria* (Orthoptera: Acrididae)

Accepted March 31, 2009

MOATAZA A. DORRAH, TAHANY H. AYAAD, AMR A. MOHAMED AND TAHA T.M. BASSAL*

Entomology Department, Faculty of Science, Cairo University

*Author for correspondence. Email: tahabasal@gmail.com

Abstract

Three lectins, designated as Sg₁, Sg₂ and Sg₃, were identified in the serum of the desert locust *Schistocerca gregaria*. With the use of (NH₄)₂SO₄ fractionation, ion-exchange chromatography on DEAE-cellulose and affinity chromatography on CNBr-activated Sepharose 4B, three pools, each containing one of the putative lectins, were obtained. IEF, native PAGE and SDS/PAGE revealed that the three pools contained Sg₁, Sg₂ and Sg₃, respectively, with pI = 6.39, 8.11 and 6.8; native PAGE R_m = 0.44, 0.32 and 0.32; SDS/PAGE subunits approximate molecular weights = 21.5, 44.5, and 43.9 kDa, respectively. Under reducing conditions, SDS/PAGE has separated each of Sg₂ and Sg₃ into two bands that indicate the presence of covalent interactions between their subunits, which are not present in Sg₁. The three lectins are more highly specific for rabbit RBCs than for those of other vertebrates. The HA of lectins are Ca²⁺-dependent, heat-labile, and are inhibited strongly by α-linked-D-galactosides, followed by L-(+)-rhamnose.

Keywords

multiple lectins, isolation, serum, hemagglutination, sugar inhibition, α-linked-D-galactosides, rhamnose, *S. gregaria*

Abbreviations

HA, hemagglutinating activity
IEF, isoelectric focusing
PAGE, polyacrylamide gel electrophoresis
pI, isoelectric point
RM, relative mobility
SDS, Sodium dodecyl sulfate

Introduction

Lectins can be defined as proteins that recognize specific carbohydrate structures and thereby agglutinate cells by binding to cell-surface sugars, glycoproteins, and other glycoconjugates (Lis & Sharon 1998). They are usually structurally complex molecules with one or more carbohydrate-recognition domains and therefore with possible multiple binding sites (Gillespie *et al.* 1997, Dodd & Drickamer 2001). Among these lectins, those that require Ca²⁺ for their activity are called C-type lectins (Drickamer 1988). Animal lectins have been found in various invertebrates as well as vertebrates (Barondes 1984), either in soluble or in membrane-bound form. These lectins can play a variety of physiological roles: in particular, they are crucial in the innate immune system where they bind, as referred to above, to the carbohydrates present on the surface of potential pathogens (Rudd *et al.* 2001).

The most extensively studied functions of hemolymph lectins have been their roles in insect immune defense systems (Ratcliffe & Rowley 1980, Vasta *et al.* 2007). Furthermore, induced or endogenous hemolymph lectins and/or hemocyte-associated lectins have been shown to be involved in phagocytosis (Kawasaki *et al.* 1993, Wheeler *et al.* 1993, Wilson *et al.* 1999), encapsulation (Komano & Natori 1985), nodule formation (Kyriakides *et al.* 1993), activation of the prophenoloxidase system (Chen *et al.* 1995) and hemolymph coagulation (Minnick *et al.* 1986).

Relatively few insect lectins have been purified and characterized. The majority of insect lectins characterized to date have been detected in the hemolymph and in most cases they are thought to be synthesized in the fat body (Kubo *et al.* 1984, Amanai *et al.* 1991) and/or the hemocytes (Amiranti 1976, Stiles *et al.* 1988, Amanai *et al.* 1991, Boucias & Pendland 1993). Lectins have been isolated and characterized from some insects, *e.g.*, dipterans (Komano *et al.* 1980, Stynen *et al.* 1985, Ingram & Molyneux 1990, McKenzie & Preston 1992, Haq *et al.* 1996, Chen & Billingsley 1999, Volf *et al.* 2002), lepidopterans (Pendland & Boucias 1986, Castro *et al.* 1987, Qu *et al.* 1987, Amanai *et al.* 1990, Gül & Ayvali 2002, Ourth *et al.* 2005, Chai *et al.* 2008), orthopterans (Stebbins & Hapner 1985, Drif & Brehelin 1994, Ayaad 2004), dictyopterans (Kubo & Natori 1987, Chen *et al.* 1993), phasmids (Richards *et al.* 1988), and hemipterans (Gomes *et al.* 1991). Their structure, function, and carbohydrate-binding properties still need further investigation to clarify their role in the innate immune system of insects.

The present work reports on isolation and on the biological, physicochemical and molecular characterization of multiple lectins from serum of the 5th-instar desert locust *S. gregaria*.

Materials and Methods

Insects, sample preparation and biological and physicochemical tests.—Insect rearing, preparation of hemolymph sera, preparation of erythrocyte suspension, assay of hemagglutinating activity (HA), effect of divalent cations, inhibition assays of sugars and glycoconjugates, and stability tests — were carried out for the whole-hemolymph sera lectins (Ayaad *et al.* 2009 this issue). Recounted briefly: solutions of the isolated lectins were used instead of the prepared sera. Rabbit erythrocytes, 2% suspension in TBS/Ca²⁺-Mg²⁺ (pH 7.0) were used for any of the HA assays. The effect on HA of the bivalent cations (20 mM) Ca²⁺, Mg²⁺, Zn²⁺, and Mn²⁺, and EDTA (5–10 mM) was tested at pH 7.0. The inhibitory effect of carbohydrates on HA was tested in the presence of free and conjugated sugars in TBS/Ca²⁺-Mg²⁺ at pH 7.0. Heating for 25 min at 60 and at 100°C, and storage at both 4

and at -20°C over a period of 1 week to 3 months, were tested for their effect on HA (Ayaad *et al.* 2009 this issue).

Determination of total protein concentration.—The total protein concentration was determined according to the method of Bradford (1976), using Coomassie Brilliant Blue (COBB). Bovine serum albumin (BSA) fraction V (Sigma-Aldrich), dissolved in 0.15 M NaCl, was used as a protein standard.

Isolation of lectins [(NH₄)₂SO₄ – salting out of serum proteins].—Insect serum (25 ml) was diluted with an equal volume of 0.02 M TBS pH 7.0 containing 2 mM CaCl₂ and 1 mM MgCl₂ (TBS/Ca²⁺-Mg²⁺). Proteins were precipitated by addition of (NH₄)₂SO₄ (Sigma-Aldrich) at a concentration of 70%, with gentle stirring at 4°C. The resulting precipitate was collected by centrifugation and dialyzed in Visking® dialysis tubing (mol. wt cut-off [MWCO]: 12000-14000 Da) (Serva Electrophoresis GmbH, Germany) against distilled water for 24 h at 4°C with three changes. The dialyzed solution was then centrifuged at 8000 rpm for 5 min at 4°C to remove insoluble materials.

Ion-exchange chromatography of lectins.—Twenty grams of DEAE-cellulose (Amersham Pharmacia Biotech) were washed in 0.5 M NaOH for 10 min, then rinsed in distilled water, followed by 0.8 M HCl, and finally 0.5 M NaOH, to remove contaminants. The washed matrix was thoroughly rinsed free of NaOH with distilled water and mixed with a sufficient quantity of starting buffer (0.01 M TBS containing 0.01 M CaCl₂ and 0.01 M MgCl₂, pH 7.0) to produce a thin suspension. The treated matrix was packed into a column (20 cm × 1.0 cm i.d.) and thoroughly washed with the starting buffer. A sample from the last step of (NH₄)₂SO₄ fractionation (approximately 20 ml of 2.85 mg protein/ml) was applied to the DEAE-cellulose column. The column was then washed with the starting buffer until the baseline of absorption at 280 nm was achieved, and then elution carried out with a linear gradient of 0-0.25 M NaCl in 200 ml of the starting buffer. The flow rate was adjusted to 10 ml/h. Fractions (tubes) constituting the separated peaks were tested separately for HA against rabbit RBCs; all fractions displaying HA were then pooled for use in affinity chromatography.

Affinity chromatography of lectins.—Lectins of the same peak were isolated from the other proteins contained therein, by use of CNBr-activated Sepharose 4B (Sigma-Aldrich) affinity chromatography as described by Komano *et al.* (1980) with slight modifications. CNBr-activated Sepharose 4B matrix was packed into a column (10X1 cm i.d.) and extensively washed with about 500 ml of TBS/Ca²⁺-Mg²⁺, pH 7.0 containing 0.02% NaN₃ (Sigma-Aldrich). Each separate pool of fractions, having HA from the DEAE-cellulose column, was separately applied (slowly) to the CNBr-Sepharose 4B column at 4°C. This latter column was washed successively with TBS, to which 0.5 M NaCl was added, until no further protein was detected in the eluate by monitoring the absorbance at 280 nm. Lectins (the bound material) were then eluted with 0.3 M raffinose solution in TBS/Ca²⁺-Mg²⁺ (Ayaad 2004, Ayaad *et al.* 2009). The collected fractions of each peak were pooled and each pool separately dialyzed extensively against TBS/Ca²⁺-Mg²⁺, then against de-ionized distilled water to remove raffinose; then HA of each pool was assayed against rabbit RBCs. Each pool of fractions of the resulting peaks was lyophilized and stored at -20°C until use.

IEF of lectins.—The isoelectric points of lectins, contained in pools collected from affinity chromatography columns, were estimated

on a pH 3.6 to 9.3 gradient polyacrylamide gel, according to the method of O'Farrell (1975), in slab gels (140 × 120 × 0.75 mm). Electrophoresis ran at 350 V for 17-18 h, yielding approximately 6000 V/h at 22°C. The pI values of the isolated lectins were determined by running a mixture of standard proteins of known isoelectric points on the same gel. We obtained a number of mixtures of proteins with different pI values (IEF MIX-Sigma, USA, pH 3.6-9.3) containing amyloglucosidase (pI 3.6), trypsin inhibitor (pI 4.6), β-lactoglobulin A (pI 5.1), carbonic anhydrase II (bovine, pI 5.9), carbonic anhydrase I (human, pI 6.6), myoglobin (pI 6.8, 7.2), lectin from *Lens culinaris* (pI 8.2, 8.6, 8.8), and trypsinogen (pI 9.3); identifications of the different pI values were made using an image densitometer G 700 (Bio-Rad, USA).

Native PAGE of lectins.—The identity of the destined lectins contained in the pools collected from affinity chromatography columns (raffinose-eluates) was confirmed by native PAGE. Electrophoresis was carried out on 10% polyacrylamide gel under nondenaturing conditions, according to the method of Schagger and von Jagow (1991), using a Tris-glycine running buffer without SDS.

After isolation of protein bands by the nondenaturing PAGE, the characteristic bands were detected on the gel by staining and matching procedures, then were sliced away from the remainder of the gel with a sharp razor blade. Each of these specific gel areas was cut into pieces, approximately 1mm² for each, and the pieces of each band soaked in 2 ml of distilled water in a separate tube overnight at room temperature. Each tube was centrifuged at 6000 rpm for 10 min; the eluate (supernatant) was transferred into a new tube. The eluted proteins were concentrated by speed vacuum for 30 min and then stored at -20°C until used.

SDS/PAGE of lectins (under nonreducing and reducing conditions).—SDS/PAGE of isolated lectins was carried out by the discontinuous buffer system of Laemmli (Laemmli 1970). Some samples were denatured with 2% SDS containing 5% β-mercaptoethanol by boiling for 3 min. Treated samples were centrifuged at 10000 g for 5 min before being loaded onto the gels. Electrophoresis was carried out at a constant voltage of 200 V for 90 min. The gels were calibrated with standard molecular weight proteins [New England Biolabs Ltd. (low range: 116, 91, 46.4, 34.3, 28.7, and 21 kDa) and/or Titan Biotech Ltd. (high and low range: 200, 116, 97, 66, 45, and 21 kDa)] and the quantifications of the different molecular weights were made using an image densitometer G 700 (Bio-Rad).

Results and Discussion

The isolated lectins

Isolation of lectins.—The methods and procedures used for isolation of the 5th-instar *S. gregaria* lectins are presented in Figure 1.

After (NH₄)₂SO₄ fractionation and DEAE-cellulose chromatography, the HA was found to be present in fractions of peak I (18 ml), peak II (24 ml), and peak III (18 ml), but not in peaks IV and V (Fig. 2). Therefore, fraction pools of each of peaks I, II, and III were indicated to contain lectins. These pools were separately, and slowly applied to the prepared CNBr-activated Sepharose 4B columns for affinity chromatography.

On washing each column successively with TBS to which 0.5 M NaCl was added until no further protein was detected, it was shown that most proteins were not adsorbed to the column, appeared in the through-flow fractions, and had no HA (Fig. 3a, b, c). When

Fig. 1. Summary of methods and procedures used to isolate serum lectins of 5th instar *S. gregaria*.




Table 1. Purification schedule of the 5th instar *S. gregaria* serum lectins.

Purification step	Volume (ml)	Protein (mg/ml)	Hemagglutinating activity (HA)	Specific activity (HA/mg protein)	Yield (%)
A- Crude lectins and salting out					
Whole serum (crude lectins)	25	2.28	512	224.5	100
70% saturated (NH ₄) ₂ SO ₄	150	0.62	256	412.9	50
B- DEAE cellulose chromatography					
Peak I	18	1.99	8	4.02	1.562
Peak II	24	1.73	16	9.24	3.125
Peak III	18	2.04	16	7.84	3.125
C- Affinity chromatography					
Pool 1	7.5	0.85	128	150	25
Pool 2	7.5	0.66	256	387.8	50
Pool 3	6	0.55	256	465.4	50

Table 2. Hemagglutinating activity (HA)^a of the 5th instar *S. gregaria* serum lectins contained in the collected pools from affinity chromatography, against a group of vertebrate RBCs.

	HA ^b		
	P ₁	P ₂	P ₃
Rabbit	128	256-512	256
Sheep	4	32	16
Human:			
A ⁺	4	4	0
B ⁻	0	16	0
O ⁻	4	4	0
Rat	0	4	0
Guinea pig	0	0	0
Horse	0	0	0

^a Measured in a sample pool of 20 insects for each RBC type.

^b Standard assay condition: using TBS/Ca-Mg. HA expressed as log₂ titer.

the bound material (lectins) was then eluted with 0.3 M raffinose solution in TBS/ Ca²⁺-Mg²⁺, small protein peaks containing HA were realized (Fig. 3a, b, c). Fractions constituting each peak were considered as different pools [pool 1 (P₁), pool 2 (P₂), and pool 3 (P₃)], each containing different lectins. After dialysis to remove raffinose, the three resulting pools exhibited prominent HA; therefore, they seemed to contain different lectins. The specific activity of lectins recovered from serum by use of these methods and procedures showed a 1.5 to 4-fold increase in the specific activity (Table 1), indicating prominent reproducibility.

HA of the isolated lectin pools.—The determined HA patterns of lectins, of the 5th instar *S. gregaria*, contained in the three collected pools from affinity chromatography, are presented against a range of vertebrate RBCs, (Table 2). The isolated lectin pools have the highest HA against rabbit RBCs, when compared to the other RBC types, in the order P₂ > P₃ > P₁. For the other RBC types, sometimes there was a very low level of HA, or even none. These results demonstrate a high specificity of the 5th instar *S. gregaria* isolated serum lectins to RBCs of rabbit, when compared to those of the other vertebrates (Table 2).

Homogeneity and molecular characteristics of lectins

Homogeneity and molecular characteristics of the isolated lectins contained in the pools P₁, P₂ and P₃ were monitored and confirmed by use of IEF, and both the native PAGE and SDS/PAGE (under nonreducing and reducing conditions).

Isoelectric focusing (IEF).—The obtained results (Fig. 4) reveal some characteristics of these lectins. P₂ and P₃ both focused as single bands at pH 8.11 and pH 6.84, respectively. However, P₁ showed a major band (~84%) focused at pH 6.39 and a minor band (16%) focused at pH 6.53 and realized in association to the major one. Therefore, each of the P₂ and P₃ seems to be formed of a single and separate lectin. The nature of the two-band components of P₁ were confirmed by native PAGE, accompanied with HA determination to each separated band (see below).

The pI values of P₂ protein appear to lie in the slightly alkaline




Fig. 2. Ion-exchange chromatography on DEAE-cellulose, of the 5th instar *S. gregaria* serum proteins, resulting from salting out by 70% (NH₄)₂SO₄. Elution with a linear gradient of 0 – 0.25 M NaCl. Fractions of peaks I, II, III have HA against rabbit RBCs; but those of peaks IV, V have none.




Fig. 3. Affinity chromatography of lectins, contained in peaks I, II, III (a, b, c, respectively) of proteins separated from serum of 5th instar *S. gregaria* by DEAE-cellulose column, on CNBr-activated Sepharose 4B. Lectins (bound components) were eluted with 0.3 M raffinose solution in TBS/ Ca²⁺-Mg²⁺ (indicated by arrows).




Fig. 4. IEF/PAGE of the 5th instar *S. gregaria* lectins contained in the three pools P_1 , P_2 , and P_3 (lanes 3, 2, and 1, respectively) collected from Sepharose 4B affinity chromatography columns, and also of the whole-serum proteins (lane 4). IEF standards (lane 5) (Sigma) with different pH ranges are shown to the right. The gel was stained with COBB R-250.

range, whereas those of both P_1 and P_3 appear to lie in the slightly acidic range. In other insects, the pI values range from 4.7 to 6.4, as in the orthopterans *M. sanguinipes* (Stebbins & Hapner 1985), and *L. migratoria* (Drif & Brehelin 1994), the dictyopteran *B. discoidalis* (Chen *et al.* 1993), the lepidopterans *M. sexta* (Minnick *et al.* 1986), and *H. cecropia* (Castro *et al.* 1987), the coleopteran *A. dichotoma* (Umetsu *et al.*, 1984), and the dipteran *G. fuscipes* (Ingram & Molyneux 1990).

Native PAGE.—The identity of the lectins contained in the three pools P_1 , P_2 and P_3 , collected from affinity chromatography columns (raffinose-elutents) (Fig. 3 a-c) was confirmed by native PAGE. The electrophoretograph obtained (Fig. 5) shows that P_2 and P_3 appear as single bands. This confirms the nature of both P_2 and P_3 that appeared on IEF gel. Therefore, each of these two bands was considered as a discrete lectin and they are designated as Sg_2 and Sg_3 , respectively. Both bands have R_m values of almost 0.32, (Fig. 5) that were revealed in the high molecular weight range.

On the other hand, P_1 was separated into two bands with different relative mobility values of 0.38 (P_{1a}) and 0.44 (P_{1b}), respectively. The first band P_{1a} (R_m = 0.38) did not display any HA, whereas the second band P_{1b} (R_m = 0.44) enclosed HA, and was considered as a third lectin in the serum of *S. gregaria*; it was designated as Sg_1 . The nonagglutinating band may be a nonlectin glycoprotein that binds to, and is eluted from, the ion exchange and the affinity columns, in association with the isolated lectin Sg_1 . Co-isolated proteins, during the isolation of multiple lectins, that are eluted from both ion exchange and affinity chromatography, present a case observed before in other insects such as *E. tiaratum* (Richards *et al.* 1988) and *B. discoidalis* (Wilson *et al.* 1999).

Appearance of native PAGE patterns of Sg_1 (P_{1b}), Sg_2 and Sg_3 as a single band with high molecular weight range, indicates that each band may be formed of aggregates (oligomers) of subunits. This assumed oligomeric native form of lectins of the 5th instar *S. gregaria* serum is reported before in other insects, *e.g.*, the orthopterans *M. sanguinipes* (Stebbins & Hapner 1985), *L. migratoria* (Drif & Brehelin 1994), and adult *S. gregaria* (Ayaad 2004), and in the dipteran *G. fuscipes* (Ingram & Molyneux 1990).

SDS/PAGE.—When the lectins Sg_1 (P_{1b}), Sg_2 and Sg_3 were sliced away from the native PAGE, eluted, then electrophoresed by SDS/PAGE, under both nonreducing (Fig. 6), and reducing (Fig. 7) conditions, using 10% separating gel and 5% stacking gel, additional characteristics were revealed. Under the nonreducing conditions, each of the Sg_1 , Sg_2 , and Sg_3 was separated by SDS/PAGE as a single band with approximate molecular weights of 21.5, 44.5, and 43.9 kDa respectively (Fig. 6). This may indicate that subunits of each lectin are of approximately the same molecular weight and are held together by noncovalent interactions. Noncovalent linking of protein subunits was reported in lectins of other insects, *e.g.*, the dipteran *S. peregrina* (Komano *et al.* 1980), lepidopterans *H. cecropia* and *S. exigua* (Castro *et al.* 1987, Boucias & Pendland 1993), and the phasmid *E. tiaratum* (Richards *et al.* 1988).

Under the reducing conditions, SDS/PAGE separation (Fig. 7) showed a single band only from Sg_1 (21.7 kDa), whereas each of Sg_2 and Sg_3 were separated into two bands. Those resulting from Sg_2 have approximate molecular weight of 32.6 and 31.7 kDa, and those of Sg_3 are of 32.9 and 31.5 kDa. The separation of two different bands with two different molecular weights from each of the Sg_2 and Sg_3 (Fig. 8) indicates that each of these latter is formed of two different types of subunits linked with disulphide bonds. This character was reported before for some insect lectins, for example, the orthopterans *T. commodus* (Hapner & Jermyn 1981), and *M. sanguinipes* (Stebbins & Hapner 1985), BDL1, and BDL2 of the dictyopteran *B. discoidalis* (Chen *et al.* 1993), and allo A-I and -II of the coleopteran *A. dichotoma* (Umetsu *et al.* 1984).

The low molecular-weight range of the single-banded Sg_1 separated by the nonreducing and reducing SDS/PAGE (21.5 to 21.7 kDa) is near to those of other insects, for instance, 30-kDa lectin of *P. americana* (Kubo & Natori, 1987), 20-kDa lectin of *S. peregrina* (Fujita *et al.* 1998).

Physicochemical properties of the isolated lectins

Stability.— The obtained data revealed that HA was completely abolished after 25 min exposure to 100°C, but reduced to 75 % only upon exposure to 60°C for the same time. On the other hand, it was observed that HA was stable when exposed to 25°C (room temperature) for the same period of time. These observations indicate that lectins of the 5th instar *S. gregaria* are heat-labile in nature. Heat instability is characteristic for lectins of some other insects, *e.g.*, the orthopterans *T. commodus*, (Hapner & Jermyn 1981), and *M. sanguinipes* (Stebbins & Hapner 1985), the phasmid *E. tiaratum* (Richards *et al.* 1988), and the dipteran *G. fuscipes* (Ingram & Molyneux 1990). However, in contrast to these cases, in the coleopteran *L. decemlineata* (Minnick *et al.* 1986), and the orthopteran *L. migratoria* (Drif & Brehelin 1994) the lectins were reported to be heat resistant when subjected to elevated temperatures of 70–100°C.

For the storage temperature and period, the HA of lectins of 5th instar *S. gregaria* were stable when stored at -20°C, with an extremely slow insignificant decline observed on prolonged storage (3 mo).




Fig. 5. Native PAGE of the 5th instar *S. gregaria* serum proteins (lane 1), and the lectins contained in the pools P₁, P₂, and P₃ (lanes 2, 3, and 4 respectively) collected from Sepharose 4B affinity chromatography columns. The gel was stained with COBB R-250.

Since the storage medium of the isolated lectins under investigation was without any additive-sugars, it seems that these lectins do not need the presence of sugars for their stability. This situation is analogous to that reported in *S. peregrina* (Komano *et al.* 1980). However, in contrast to this case, isolated lectins from other insects require addition of sugar to the storage medium to maintain their activity. For example, in *T. commodus* (Hapner & Jermyn 1981), the lectin required the addition of galactose. Also the lectins of *M. sanguinipes* (Stebbins & Hapner 1985), once isolated, were so unstable that disaggregation of the multimer occurred readily in the absence of D-galactose; and in *E. tiaratum* instability was observed when the lectin was not stored in lactose (Richards *et al.* 1988).

Divalent cation requirement for HA.— The present data show that the isolated lectins Sg₁, Sg₂, and Sg₃ require Ca²⁺ to express their HA. Some cations such as Mg²⁺ or Zn²⁺ can only partly replace Ca²⁺; however other divalent cations, such as Mn²⁺, have no effect (Table 3). As expected, addition of EDTA caused complete inhibition to HA. Also removal of Ca²⁺ by dialyzing the lectin solutions against either TBS alone or TBS with 10 mM EDTA, similarly resulted in a complete loss of HA of these lectins. EDTA presumably exerted its effect through chelation of divalent cations that structurally associated with the lectins. The HA of C-type lectins of *T. commodus*, *M. sanguinipes*, *S. exigua*, *E. tiaratum* and adult *S. gregaria* were also lost in the presence of EDTA (Pendland & Boucias 1986, Stebbins &

Hapner 1985, Ayaad 2004, Richards *et al.* 1988, Hapner & Jermyn 1981). In most insect lectins a complete loss of HA was observed in the presence of EDTA (Ratcliffe *et al.* 1985). Therefore, for the lectins isolated from serum of the 5th instar *S. gregaria*, the observed requirement of Ca²⁺ to enhance and maintain HA of the three lectins Sg₁, Sg₂, and Sg₃, prominently indicates that these lectins are Ca²⁺-dependent. This property is a general characteristic of a group of lectins denoted as C-type lectins. This type of lectin was originally named to reflect the special importance of Ca²⁺ in the mechanism of carbohydrate binding (Zelensky & Gready 2005). Each of the known C-type lectins has carbohydrate-recognition domain structures containing a conserved Ca²⁺-binding site; and may also contain a second site (Drickamer 1999).

Sugar specificity and inhibition to HA.—The obtained profile of sugar inhibition to HA of lectins isolated from serum of 5th instar *S. gregaria* against rabbit RBCs shows certain characteristics. The inhibition of sugars and of glycoconjugates to HA of Sg₁, Sg₂ and Sg₃ showed great resemblance to each other (Table 4). The HA of these lectins was preferentially inhibited (IC₅₀ = 6.25 mM) by raffinose, α-m-nitrophenyl-D-galactose, α-p-nitrophenyl-D-galactose, and α-methyl-D-galactose, and the monosaccharide L-(+)-rhamnose. None of these lectins were inhibited by D-(+)-mannose, trehalose and N-acetylglucosamine (200 mM), or laminarin (> 1 %). Low sensitivities (IC₅₀ = 50-100 mM) were observed in case of the free sugars D-galactose, D-glucose, sucrose, lactose, and also by N-acetyl-D-galactosamine. These results allow speculation that the carbohydrate-binding site of the carbohydrate recognition domain of these lectins prefers (within the range of the tested sugars) α-linked-D-galactosides over




Fig. 6. SDS-PAGE analysis (under nonreducing conditions) of lectins Sg₁, Sg₂, and Sg₃ (lanes 1, 2 and 3 respectively) isolated from serum of the 5th instar *S. gregaria*. The gel was stained with COBB R-250. Molecular weights (kDa) of protein standards (lane M) are shown on the left.

Table 3. Effects of divalent cations and EDTA (cation chelator) on hemagglutinating activity (HA) of the lectins Sg₁, Sg₂, and Sg₃ isolated from serum of 5th instar *S. gregaria*.

	HA		
	Sg ₁	Sg ₂	Sg ₃
Ca ²⁺ - free medium	0	0	4
20 mM Ca ²⁺ ^a	128	256-512	256
20 mM Mg ²⁺	64-128	128	128
20 mM Zn ²⁺	128	128	128
20 mM Mn ²⁺	4	4	4
10 mM EDTA	0	0	0
5 mM EDTA	4	4	4

^a Standard assay condition; i.e., reference value of HA.

both the β-form and also the free D-galactose.

The affinity of the 5th instar isolated *S. gregaria* serum lectins toward α-linked galactosides is a feature that has also been reported in other acridids, such as *L. migratoria* (Drif & Brehelin 1994) and adult *S. gregaria* (Ayaad 2004). On the other hand, affinity of lectins to β-linked-D-galactosides, such as lactose and lactulose, was recorded in the coleopteran *A. dichotoma* (Umetsu *et al.* 1984), its lectins being inhibited by this group of carbohydrates. The latter lectin specificity is also known in vertebrates, where the best ligands are β-galactosides (Barondes 1984). Some lectins purified from other insect species, especially from orthopterans (Stebbins & Hapner 1985, Lackie 1981, Jurenka *et al.* 1982, Hapner 1983, Drif & Brehelin 1989) show an affinity for a broad spectrum of carbohydrates. Another orthopteran *T. commodus* (Hapner & Jermyn 1981), and the dipteran *P. duboscqi* (Volf *et al.* 2002) possess lectins with amino sugar-binding affinity. In numerous other insect species, mainly lepidopterans, hemolymph lectins show affinity for galactose and lactose (Pendland & Boucias 1986) or to glucosides (Minnick *et al.* 1986, Qu *et al.* 1987). Lectins from another group

Table 4. Inhibition by sugars and glycoconjugates to hemagglutinating activity (HA), of lectins Sg₁, Sg₂, and Sg₃, isolated from serum of the 5th instar *S. gregaria*, against rabbit RBCs.

	Minimum concentration ^a (mM/or %) required for 50 % inhibition (IC ₅₀) to HA		
	Sg ₁	Sg ₂	Sg ₃
Monosaccharide			
D-(+)-galactose	100	100	100
D-(+)-glucose	>100	50	>100
L-(+)-rhamnose	25	≥ 6	25
Oligosaccharides			
Lactose	100	100	100
Raffinose	12	6	6
Sucrose	100	100	>100
N-acetylated sugars			
N-acetyl-D-galactosamine	100	50	>50
Others			
α-methyl-D-galactose	12	12	25
α-m-nitrophenyl-D-galactose	6	6	6
α-p-nitrophenyl-D-galactose	6	< 6	< 6

^a Data presented are from experiments repeated three times. Values > 200 mM [(D-(+)-mannose, trehalose, and N-acetyl-D-glucosamine] or >1 % (zymosan and laminarin) indicate that no inhibition of agglutination was observed. All inhibitions were of two wells unless otherwise indicated.

of insects have binding affinity to mannose, e.g., the dictyopteran *B. discoidalis* (Chen *et al.* 1993), and the lepidopteran *H. virescens* (Ourth *et al.* 2005), and to mannan, e.g., the dipteran *A. stephensi* (Chen & Billingsley 1999).

In this context, C-type lectins are known to have a common carbohydrate recognition domain (Weis *et al.* 1991, Drickamer 1992), or two domains (Yu & Kanost 2000), that may increase their binding affinity to carbohydrates. This domain contains a conserved Ca²⁺-binding site, and may contain a second one (Drickamer 1999). The binding mechanism involves interaction with carbohydrate and the conserved Ca²⁺, which may form hydrogen bonding with acid and amide side groups (Weis & Drickamer 1996).

The ligand specificity of the carbohydrate recognition domain is largely dependent on the position of hydroxyl groups on the free or conjugated sugar, particularly the 3-OH and 4-OH (Ng *et al.* 1996, Kolatkar & Weis 1996). In addition to these interactions further specificity may be achieved by hydrogen bonds and electrostatic interactions with the surface of the protein (Ng & Weis 1997, Kolatkar *et al.* 1998). In the present work, agglutination of rabbit RBCs by C-type lectins Sg₁, Sg₂, and Sg₃ of the serum of the 5th instar *S. gregaria* was inhibited most efficiently by the α-linked-D-galactosides followed by L-(+)-rhamnose. However, the free monosaccharide D-(+)-galactose and the β-linked-D-galactosides (as in lactose) were of very low efficiency. In the latter cases, the orientation of 3-OH and 4-OH are the same for the same sugar, D-(+)-galactose; and the same was observed for 2-OH and 4-OH in D-(+)-galactose and L-(+)-rhamnose. Therefore, not only the orientation of 2-OH, 3-OH and 4-OH, but also the configuration (α or β) and the nature of the substituents at C₁ or the functional group at C₆, seem to affect the binding specificity and affinity of the 5th instar *S. gregaria* lectins to carbohydrates.




Fig. 7. SDS-PAGE analysis (under reducing conditions) of lectins Sg₁, Sg₂, and Sg₃ (lanes 1, 2, and 3, respectively) isolated from serum of the 5th instar *S. gregaria*. The gel was stained with COBB R-250. Molecular weights (kDa) of protein standards (lane M) are shown on the left.

According to these parameters, carbohydrate-binding specificity is crucial in recognition of these cell-surface carbohydrates, including those of the invading pathogens (McGreal *et al.* 2004). Therefore, animal C-type lectins are among the important proteins in pathogen recognition and cellular interaction (Rudd *et al.* 2001, Weis *et al.* 1998) by binding to the carbohydrate component of the surface molecular patterns of these targets (Janeway 1989).

References

- Amanai K., Sakurai S., Ohtaki T. 1990. Purification and characterization of haemagglutinin in the hemolymph of the silkworm, *Bombyx mori*. *Comparative Biochemistry and Physiology* 97B: 471-476.
- Amanai K., Sakurai S., Ohtaki T. 1991. Site of hemolymph lectins production and its activation in vitro by 20-hydroxyecdysone. *Archives of Insect Biochemistry and Physiology* 17: 39-51.
- Amiranti G.A. 1976. Production of heteroagglutinins in haemocytes of *Leucophaea maderae* L. *Experientia* 32: 526-528.
- Ayaad T.H. 2004. Isolation, characterization, and N-terminal amino acid sequence of lectin from plasma of *Schistocerca gregaria*. *Efflatounia* 4: 9-22.
- Ayaad T.H., Dorrah M.A., Mohamed A.A., Bassal T.T.M. 2009. Specificity and developmental changes of hemagglutination activity of serum of the desert locust *Schistocerca gregaria* (Orthoptera: Acrididae). *Journal of Orthoptera Research* 18: 51-56.
- Barondes S.H. 1984. Soluble lectins: a new class of extracellular proteins. *Science* 223: 1259-1264.
- Boucias D.G., Pendland J.C. 1993. The galactose binding lectin from the beet armyworm, *Spodoptera exigua*: distribution and site of synthesis. *Insect Biochemistry and Molecular Biology* 23: 233-242.
- Bradford M.M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.
- Castro V.M., Boman H.G., Hammarström S. 1987. Isolation and characterization of isolectins with galactose/N-acetyl galactosamine specificity from hemolymph of the giant silkworm *Hyalophora cecropia*. *Insect Biochemistry* 17: 513-523.
- Chai L-Q., Tian Y-Y., Yang D-T., Wang J-X., Zhao X-F. 2008. Molecular cloning and characterization of a C-type lectin from the cotton bollworm, *Helicoverpa armigera*. *Developmental and Comparative Immunology* 32: 71-83.
- Chen C., Billingsley P.F. 1999. Detection and characterization of a mannan-binding lectin from mosquito, *Anopheles stephensi* (Liston). *European Journal of Biochemistry* 263: 360-366.
- Chen C., Durrant H.J., Newton R.P., Ratcliffe N.A. 1995. A study of novel lectins and their involvement in the activation of the prophenoloxidase system in *Blaberus discoidalis*. *Biochemical Journal* 310: 23-31.
- Chen C., Ratcliffe N.A., Rowley A.F. 1993. Detection, isolation and characterization of multiple lectins from the hemolymph of the cockroach, *Blaberus discoidalis*. *Biochemical Journal* 294: 181-190.
- Dodd R.B., Drickamer K. 2001. Lectin-like proteins in model organisms: implications for evolution of carbohydrate-binding activity. *Glycobiology* 11: 71R-79R.
- Drickamer K. 1988. Two distinct classes of carbohydrate-recognition domains in animal lectins. *Journal of Biological Chemistry* 263: 9557-9560.
- Drickamer K. 1992. Engineering galactose-binding activity into a C-type mannose-binding protein. *Nature* 360: 183-186.
- Drickamer K. 1999. C-type lectin-like domains. *Current Opinion in Structural Biology* 9: 585-590.
- Drif L., Brehelin M. 1989. Agglutinin mediated immune recognition in *Locusta migratoria* (Insecta). *Journal of Insect Physiology* 35: 729-736.
- Drif L., Brehelin M. 1994. Purification and characterization of an agglutinin from the hemolymph of *Locusta migratoria* (Orthoptera). *Insect Biochemistry and Molecular Biology* 24: 283-228.
- Fujita Y., Kurata S., Homma K., Natori S. 1998. A novel lectin from *Sarcophaga* its purification, characterization, and cDNA cloning. *Journal of Biological Chemistry* 273: 9667-9672.
- Gillespie J.P., Kanost M., Trenczek T. 1997. Biological mediators of insect immunity. *Annual Review of Entomology* 42: 611-643.
- Gomes Y. De M., Furtado A.F., Coelho L.B.B. 1991. Partial purification and some properties of a hemolymph lectin from *Panstrongylus megistus* (Hemiptera, Reduviidae). *Applied Biochemistry and Biotechnology* 31: 97-107.
- Gül N., Ayvalı C. 2002. Purification and determination of the molecular structure of hemolymph lectin of *Agrotis segetum* (Denis and Schiff.). *Turkish Journal of Biology* 26: 49-55.
- Hapner K.D. 1983. Hemagglutinin activity in the hemolymph of individual Acrididae (grasshoppers) specimens. *Journal of Insect Physiology* 29: 101-106.
- Hapner K.D., Jermyn M.A. 1981. Hemagglutinin activity in the hemolymph of *Teleogryllus comodus* (Walker). *Insect Biochemistry* 11: 287-295.
- Haq S., Kubo T., Kurata S., Kobayashi A., Natori S. 1996. Purification, characterization, and cDNA cloning of a galactose specific C-type lectin from *Drosophila melanogaster*. *Journal of Biological Chemistry* 271: 20213-20218.
- Ingram G.A., Molyneux D.H. 1990. Lectins (haemagglutinins) in the haemolymph of *Glossina fuscipes fuscipes*: isolation, partial characterization, selected physico-chemical properties and carbohydrate-binding specificities. *Insect Biochemistry* 20: 13-27.
- Janeway C.A., Jr 1989. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harbor Symposia on Quantitative Biology* 54: 1-13.
- Jurenka R., Manfredi K., Hapner K.D. 1982. Haemagglutinin activity in Acrididae (grasshopper) haemolymph. *Journal of Insect Physiology* 28: 177-181.
- Kawasaki K., Kubo T., Natori S. 1993. A novel role of *Periplaneta* lectin as an opsonin to recognize 2-keto-3-deoxy octonate residues of bacterial lipopolysaccharide. *Comparative Biochemistry and Physiology* 106B: 675-680.
- Kolattkar A.R., Weis W.I. 1996. Structural basis of galactose recognition by C-type animal lectins. *Journal of Biological Chemistry* 271: 6679-6685.
- Kolattkar A.R., Leung A.K., Isecke R., Brossmer R., Drickamer K., Weis W.I. 1998. Mechanism of N-Acetylgalactosamine binding to a C-type animal lectin carbohydrate-recognition domain. *Journal of Biological Chemistry* 273: 19502-19508.
- Komano H., Natori S. 1985. Participation of *Sarcophaga peregrina* humoral lectin in the lysis of sheep red blood cells injected into the abdominal cavity of larvae. *Developmental and Comparative Immunology* 9: 31-40.
- Komano H., Mizuno D., Natori S. 1980. Purification of lectin induced in the hemolymph of *Sarcophaga peregrina* larvae on injury. *Journal of Biological Chemistry* 255: 2919-2924.
- Kubo T., Natori S. 1987. Purification and some properties of a lectin from the hemolymph of *Periplaneta americana* (American cockroach). *European Journal of Biochemistry* 16: 75-82.
- Kubo T., Komano H., Okada O., Natori S. 1984. Identification of haemagglutinating protein and bactericidal activity of *Sarcophaga peregrina* haemolymph of adult on injury of body wall. *Developmental and Comparative Immunology* 8: 283-291.
- Kyriakides T.R., McKillip J.I., Spence K.D. 1993. In vivo distribution of immune protein scolexin in bacteria-injected *Manduca sexta* larvae. *Tissue and Cell* 25: 423-434.
- Lackie A.M. 1981. The specificity of the serum agglutinins of *Periplaneta americana* and *Schistocerca gregaria* and its relationship to the insect's immune response. *Journal of Insect Physiology* 27: 139-143.
- Laemmili U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
- Lis H., Sharon N. 1998. Lectins: carbohydrate-specific proteins that mediate cellular recognition. *Chemical Reviews* 98: 637-674.

- McGreal E.P., Martinez-Pomares L., Gordon S. 2004. Divergent roles for C-type lectins expressed by cells of the innate immune system. *Molecular Immunology* 41:1109-1121.
- McKenzie A.N.J., Preston T.M. 1992. Purification and characterization of a galactose-specific agglutinin from the haemolymph of the larval stages of the insect *Calliphora vomitoria*. *Developmental and Comparative Immunology* 16: 31-39.
- Minnick M.E., Rupp R.A., Spence K.D. 1986. A bacterial-induced lectin which triggers hemocyte coagulation in *Manduca sexta*. *Biochemical and Biophysical Research Communication* 137: 729-736.
- Ng K.K., Weis W.I. 1997. Structure of a selectin-like mutant of mannose-binding protein complexed with sialylated and sulfated Lewis(x) oligosaccharides. *Biochemistry* 36: 979-988.
- Ng K.K., Drickamer K., Weis W. 1996. Structural analysis of monosaccharide recognition by rat liver mannose-binding protein. *Journal of Biological Chemistry* 271: 663-674.
- O'Farrell P.H. 1975. High resolution two-dimensional electrophoresis of proteins. *Journal of Biological Chemistry* 250: 4007-4021.
- Ourth D.D., Narra M.B., Chung K.T. 2005. Isolation of mannose-binding C-type lectin from *Heliothis virescens* pupae. *Biochemical and Biophysical Research Communication* 335: 1085-1089.
- Pendland J.C., Boucias D.G. 1986. Characteristics of a galactose binding hemagglutinin (lectin) from hemolymph of *Spodoptera exigua* larvae. *Developmental and Comparative Immunology* 10: 477-487.
- Qu X-M., Zhang C-F., Komano H., Natori S. 1987. Purification of a lectin from the hemolymph of Chinese oak silkworm (*Antheraea pernyi*) pupae. *Journal of Biochemistry* 101: 545-551.
- Ratcliffe N.A., Rowley A.F. 1980. Insect erythrocyte agglutinins. In vitro opsonization experiments with *Clitumnus extradentatus* and *Periplaneta americana* haemocytes. *Immunology* 40: 483-492.
- Ratcliffe N.A., Rowley A.F., Fitzgerald S.W., Rhodes C.P. 1985. Invertebrate immunity: basic concepts and recent advances. *International Review of Cytology* 97: 183-350.
- Richards E.H., Ratcliffe N.A., Renwanz L. 1988. Isolation and characterization of a serum lectin from the giant stick insect *Extatosoma tiaratum*. *Insect Biochemistry* 18: 691-700.
- Rudd P.M., Elliott T., Cresswell P., Wilson I.A., Dwek R.A. 2001. Glycosylation and the immune system. *Science* 291: 2370-2376.
- Schägger H., von Jagow G. 1991. Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Analytical Biochemistry* 199: 223-231.
- Stebbins M.R., Hapner K.D. 1985. Preparation and properties of haemagglutinin from haemolymph of Acrididae (grasshoppers). *Insect Biochemistry* 15: 451-462.
- Stiles B., Bradley R.S., Stuart G.S., Hapner K.D. 1988. Site of synthesis of the haemolymph agglutinin of *Melanoplus differentialis* (Acrididae: Orthoptera). *Journal of Insect Physiology* 34: 1077-1085.
- Stynen D., Vansteenwegen K., De Loof A. 1985. Anti-galactose lectin in the haemolymph of *Sarcophaga bullata* and three other calliphorid flies. *Comparative Biochemistry and Physiology* 1381: 171-176.
- Umetsu K., Kosaka S., Suzuki T. 1984. Purification and characterization of a lectin from the beetle *Allomyrina dichotoma*. *Journal of Biochemistry (Tokyo)* 95: 239-246.
- Vasta G.R., Ahmed H., Tasumi S., Odom E.W., Saito K. 2007. Biological roles of lectins in innate immunity: molecular and structural basis for diversity in self/non-self recognition, pp. 389-406. In: Lambiris, J.D. (Ed.) *Current Topics in Innate Immunity*. Springer, New York.
- Volf P., Skarupova S., Man P. 2002. Characterization of the lectin from females of *Phlebotomus duboscqi* sand flies. *European Journal of Biochemistry* 269: 6294-6301.
- Weis W.I., Drickamer K., 1996. Structural basis of lectin-carbohydrate recognition. *Annual Review of Biochemistry* 65: 441-473.
- Weis W.I., Kahn R., Fourme R., Drickamer K., Hendrickson W.A. 1991. Structure of the calcium-dependent lectin domain from a rat mannose-binding protein determined by MAD phasing. *Science* 254: 1608-1615.
- Weis W.I., Taylor M.E., Drickamer K. 1998. The C-type lectin superfamily in the immune system. *Immunological Reviews* 163: 19-34.
- Wheeler M.B., Stuart G.S., Hapner K.D. 1993. Agglutinin mediated opsonization of fungal blastospores in *Melanoplus differentialis* (Insecta). *Journal of Insect Physiology* 39: 477-483.
- Wilson R., Chen C., Ratcliffe N.A. 1999. Innate immunity in insects: The role of multiple, endogenous serum lectins in the recognition of foreign invaders in the cockroach, *Blaberus discoidalis*. *Journal of Immunology* 157: 1590-1596.
- Yu X-Q., Kanost M.R. 2000. Immulectin-2, a lipopolysaccharide-specific lectin from an insect, *Manduca sexta*, is induced in response to gram-negative bacteria. *Journal of Biological Chemistry* 275: 37373-37381.
- Zelensky A.N., Gready J.E. 2005. The C-type lectin-like domain superfamily. *FEBS Journal* 272: 6179-6217.