

## Demographic Analysis of Imperiled Eastern Massasaugas (Sistrurus catenatus catenatus)

Authors: Jones, Peter C., King, Richard B., and Sutton, Scott

Source: Journal of Herpetology, 51(3): 383-387

Published By: Society for the Study of Amphibians and Reptiles

URL: https://doi.org/10.1670/15-058

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Demographic Analysis of Imperiled Eastern Massasaugas (Sistrurus catenatus catenatus)

PETER C. JONES,<sup>1,2</sup> RICHARD B. KING,<sup>1</sup> AND SCOTT SUTTON<sup>3</sup>

<sup>1</sup>Department of Biological Sciences, Northern Illinois University, DeKalb, Illinois, USA <sup>3</sup>Parks Canada Agency, Georgian Bay Islands National Park, Midland, Ontario, Canada

ABSTRACT.—As quantitative modeling of wildlife populations increases, the need for accurate and precise estimates of demographic rates for these populations also grows. Eastern Massasaugas (*Sistrurus catenatus*) are an imperiled rattlesnake species found mainly in the Great Lakes region of North America. We focused on an Eastern Massasauga population found on Beausoleil Island in Georgian Bay that was the subject of a 30-yr mark–recapture study for demographic analysis. We estimated multiple demographic values including annual adult survival, the temporal process variance of survival, population size, and population growth rate. Annual adult survival did not differ significantly between sexes (males 0.74; females 0.73). The process variance of annual adult survival for males was 0.006 and was inestimable for females. This is the first estimate of process variance for Eastern Massasaugas and one of the few such estimates for a snake species. The use of the process variance of survival in population prediction resulted in a large decrease in estimated extinction risk compared to use of the total variance of survival (3% and 21% extinction risk, respectively). Population size ranged from 35-77 (mean = 55), and realized population growth rate was equal to 1.02. Our analysis showed the Beausoleil Island population was stable up to the end of the study (2008). Demographic estimates can be used to guide management and conservation of this species.

Precise estimates of demographic rates (e.g., survival and fecundity) are critical for the application of conservation tools like population viability analysis (PVA; Coulson et al., 2001). PVAs created with poor parameter estimates often result in inaccurate predictions of extinction rates and future population size (Beissinger and Westphal, 1998; Coulson et al., 2001; Ellner et al., 2002). Stochastic population models also require estimates of the process variance (e.g., temporal and spatial variance) of demographic rates. For example, a population model incorporating environmental stochasticity requires an estimate of the temporal process variance of survival as well as other demographic rates. Total variance is made up of the process variance(s) and the sampling variance. Because of the inclusion of sampling variance, using total variance in a population model causes an inflation of the probability of population decline and extinction (Gould and Nichols, 1998). In contrast, process variance accounts for the real variation in a parameter and is better suited for use in a PVA than the total variance (Gould and Nichols, 1998). Although population models for Eastern Massasaugas (Sistrurus catenatus catenatus) exist, models incorporating environmental stochasticity used the total variance of survival rather than the process variance (Middleton and Chu, 2004; Miller et al., 2005; Bissel, 2006; Bailey et al., 2011).

Other demographic characteristics like population size and growth rate also are useful parameters to estimate. Population size estimates are useful for determining trends in population size over the years of the study and whether the population is at a critically small size. Population growth rates also can be compared with projected growth rates from population models as a method to check the accuracy of the population model or to determine future trends in population growth (Miller et al., 2011).

Demographic estimates are needed for the conservation and management of imperiled species such as Eastern Massasaugas (Crother et al., 2012). Eastern Massasaugas are small, thickbodied rattlesnakes found in the upper Midwest and Great Lakes regions of North America (Szymanski, 1998). Despite a relatively large range, populations are highly fragmented, and many populations appear to be in decline (Szymanski, 1998; Rouse and Wilson, 2002). Eastern Massasaugas are afforded some level of protection in every state or province in which the species resides (Szymanski, 1998). They are a candidate species for listing under the U.S. Endangered Species Act of 1973 (U.S. Fish and Wildlife Service, 2010) and are listed as threatened under Canada's Federal Species at Risk Act (Environment Canada, 2012). Because of this status, there is a demand for the application of tools like PVA and, therefore, estimates of demographic characteristics such as survival to inform management decisions.

Here, we use data from a long-term mark–recapture study of a northern population of Eastern Massasauga rattlesnakes to estimate a number of demographic parameters, including adult survival, sex differences in adult survival, temporal process variance of survival, population growth rate, and population size.

### MATERIALS AND METHODS

Study Site.—Data were collected from an Eastern Massasauga rattlesnake population on Beausoleil Island in southeastern Georgian Bay, Canada (Fig. 1). The island is ~1,100 ha and has a moderate climate with reduced temperature extremes, delayed warming in spring, and milder fall temperatures compared to areas further inland. The southern two-thirds of the island consist of drumlins, till, and sand cover, whereas the northern portion consists of till and sand patches, ponds, streams, wetlands, and two small lakes. Vegetation also differs from north to south with mixed and coniferous forest in the north and deciduous stands in the south. Beausoleil Island has a long history of human occupation that includes multiple Native American tribes, the French, and the English. Parts of the island were farmed in the past, and some logging occurred; however, the island was used for recreation since the 1920s, and it was established as a national

<sup>&</sup>lt;sup>2</sup>Corresponding Author. E-mail: pmcjones@gmail.edu DOI: 10.1670/15-058

Downloaded From: https://complete.bioone.org/journals/Journal-of-Herpetology on 06 Jul 2025 Terms of Use: https://complete.bioone.org/terms-of-use



FIG. 1. Range map of Eastern Massasaugas (*Sistrurus catenatus*). The Georgian Bay population location is denoted by the star.

park in 1929. Currently, there are roads, trails, staff residences, and campgrounds on the island.

Field Methods.—National park staff collected mark–recapture data from 1978–2008. The staff used three different marking methods over the years of the study: 1) from 1978–1985, plastic colored disks attached to rattle segments with monofilament; 2) from 1985–1992, subcaudal scale branding; and 3) from 1992–2008, passive integrated transponder (PIT) tags. Staff measured snakes for snout–vent length (SVL) and determined sex via probing the base of the tail at capture. Because these changing marking methods have different levels of efficacy, and to minimize the number of parameters in the models, we restricted the data analyzed to adult snakes in the 1992–2008 period. We defined adult snakes as animals with SVL > 350 mm based on a natural break in the size-frequency distribution observed in this population.

*Data Analysis.*—To estimate survival and recapture rates, we fit Cormack–Jolly–Seber (CJS) models to the data with the package "RMark" (Laake, 2013) for R 3.0.1 (R Core Team, 2014) and Program MARK 7.1 (White and Burnham, 1999). We fit the models using the log link function and second partial derivative method for variance estimation. We also estimated the process variance of the annual survival rate via variance components analysis of the most parsimonious model within MARK (Burnham et al., 1987; White et al., 2001).

The global model for this data set was the fully parameterized three-way interaction of sex, transience effect, and time for survival and the sex-by-time interaction for recapture rate. In terms of CJS models, transience is defined as individuals that are captured once but not recaptured throughout the rest of the study (Choquet et al., 2005). To account for these individuals, we included a transience effect (indicated by "tran" in the model parameters) in the candidate model set. The transience effect replaced typical time dependence in survival with survival over two different time periods: survival over the first year since marking and survival over subsequent years. We compared models using Akaike's Information Criterion corrected for small sample sizes (AIC<sub>c</sub>). Our candidate set included all models nested within the global model (Table 1). We tested the global model for goodness-of-fit including testing for the transience effect (TEST3.SR and TEST3.Sm), testing for a trap dependence effect (TEST2.CT and TEST2.CL), and overall goodness-of-fit

TABLE 1. Summary of Cormack-Jolly-Seber model results for Eastern Massasaugas (*Sistrurus catenatus catenatus*). Phi is the survival rate and p is the recapture rate. The "tran" effect represents the time since marking effect added to compensate for snake transience.

-					
Model	Number of parameters	AIC <sub>c</sub>	Delta AIC <sub>c</sub>	Model weight	Deviance
Phi(tran) p(sex)	4	847.51	0.00	0.79	418.69
$Phi(sex \times tran)p(sex)$	6	850.22	2.71	0.20	417.29
Phi(sex) p(sex)	4	865.64	18.14	0.00	436.82
Phi(tran) p(time)	18	866.47	18.96	0.00	407.96
$Phi(sex \times tran)$	20	870.04	22.53	0.00	407.12
p(time)	-0	0.0.01		0.00	107112
Phi(time) p(sex)	18	875.86	28.36	0.00	417.36
$Phi(sex \times tran) p(sex)$	36	879.92	32.42	0.00	379.91
× time)	00	0	02.12	0.00	0.,,,,1
Phi(tran) $p(sex \times$	34	879.99	32.48	0.00	384.78
time)				0.00	
Phi(sex) p(time)	18	881.63	34.13	0.00	423.13
$Phi(sex \times time) p(sex)$	34	890.13	42.62	0.00	394.93
$Phi(sex) p(sex \times time)$	34	893.43	45.92	0.00	398.23
Phi(time) p(time)	31	893.45	45.95	0.00	405.37
Phi(time) $p(sex \times$	47	908.46	60.96	0.00	381.00
time)				0.00	
$Phi(sex \times time)$	47	909.92	62.42	0.00	382.46
p(time)				0.00	
$Phi(sex \times tran \times$	66	916.10	68.59	0.00	336.98
time) p(sex)				0.00	
$Phi(sex \times time) p(sex$	62	930.39	82.88	0.00	362.63
× time)			02.00	0.00	
Phi(sex $\times$ tran $\times$	76	939.62	92.12	0.00	330.89
time) p(time)					
······					

(sum of all tests) with U-CARE (Choquet et al., 2009). We used the median c-hat test within MARK to estimate overdispersion in the global model.

We generated population estimates using the Jolly–Seber model via the program JOLLY (http://www.mbr-pwr.usgs. gov/software.html; Pollack et al., 1990) with 95% confidence intervals calculated using Manly's method (Manly, 1984). We also calculated average population size over the study years via the Horvitz–Thompson estimator (Horvitz and Thompson, 1952). We estimated the population growth rate with the Pradel survival and population growth rate model within MARK (Pradel, 1996). The same model fitting procedure used for the CJS models above was used for the Pradel models.

#### RESULTS

The mark-recapture data set consisted of 262 marked snakes (160 males and 102 females) captured 431 times from 1992-2008. The sum of all tests with pooled sexes indicated no sex effect ( $\chi^2$ = 45.43, df = 55, P = 0.82); thus, all further goodness-of-fit tests were performed on the pooled sex data set. The test for transience was significant ( $\chi^2 = 30.31$ , df = 15, *P* = 0.01); the test for trap-dependence was non-significant ( $\chi^2 = 7.23$ , df = 14, P = 0.92). Seventeen models were fit to the data (Table 1). The most parsimonious (model weight = 0.79) contained the parameters "tran" for survival rate and "sex" for recapture rate. The second most parsimonious model (model weight of 0.20) contained the parameters "sex" and "tran" in interaction for survival rate and "sex" for recapture rate. We used model averaging to calculate survival and recapture estimates (Table 2). The survival rate estimate for the first year since marking was lower than the survival rate estimate in subsequent years for both males and females, but there were only small differences in magnitude between males and females for either survival rate estimate



FIG. 2. Population size estimates (solid line), 95% confidence intervals (gray shading) and minimum number known alive (dotted line) from 1993–2007 for Eastern Massasaugas (*Sistrurus catenatus*) in Canada.

(Table 2). Recapture rates were consistent across years, but differed between the sexes (Table 2).

The adult survival model "Phi(sex × tran × time) p(time)" (sex, age, and time dependent survival and time dependent recapture rates) was used to calculate the process variance of the survival rate within MARK because time-dependence is necessary for the estimation process (White et al., 2001). The process variance for males was 0.006 and accounted for ~11% of the total variance (0.054). The process variance for females was unestimable, perhaps attributable to insufficient data.

TABLE 2. Model averaged survival and recapture rate estimates of the Cormack-Jolly-Seber models for Eastern Massasaugas (*Sistrurus catenatus* catenatus).

Parameter	Estimate	SE	95% CI	Process variance
Survival				
Males				
First year since marking	0.44	0.05	0.35 - 0.54	-
> 1 year since marking	0.74	0.03	0.67 - 0.80	0.006
Females				
First year since marking	0.47	0.06	0.35-0.59	-
> 1 year since marking	0.73	0.03	0.64 - 0.80	-
Recapture				
Males	0.53	0.05	0.44 - 0.62	
Females	0.43	0.06	0.32-0.55	
Females	0.43	0.06	0.32-0.55	

Population size estimates from Program JOLLY ranged from 35–77 (mean = 55) over the study years (Fig. 2). The Horvitz–Thompson estimate for average population size over the study years was 54  $\pm$  4.03 SE. Two Pradel survival and population growth models based on the most parsimonious CJS model (Phi[tran] p[sex]) were fit within MARK: 1) with time-dependent population growth rates and 2) with time-independent population growth rate was clearly the most parsimonious (Table 3). The growth rate estimate was 1.02 (95% CI = 0.99–1.04), indicating a stable to slightly increasing population.

TABLE 3. Summary of Pradel model results for Eastern Massasaugas (*Sistrurus catenatus catenatus*). Phi is the survival rate, p is the recapture rate, and  $\lambda$  is the population growth rate. The model structure abbreviations are as follows: "t" is time dependence and "." is time independence.

Model	AICc	Delta AICc	AICc Weights	Number of parameters	Deviance
Phi(tran) p(sex) λ(.)	2,342.662	0	1.00	5	470.97
Phi(tran) $p(sex)$ $\lambda(t)$	2,353.838	11.18	0.00	20	450.23

Downloaded From: https://complete.bioone.org/journals/Journal-of-Herpetology on 06 Jul 2025 Terms of Use: https://complete.bioone.org/terms-of-use

#### DISCUSSION

This study provides an unusually complete analysis of survival and population size of an Eastern Massasauga population near the northern limits of the species' range. Importantly, we demonstrate a significant effect of transience on adult survival. The models including the transience effect contained two separate survival rates that varied greatly from each other. The first survival rate (0.44 for males; 0.47 for females) is the apparent survival of snakes during their first year after marking; the second survival estimate (0.74 for males; 0.73 for females) is the survival during subsequent years (i.e., of resident snakes). Apparent survival is the product of the actual survival probability and probability of transience. The transience effect in the model effectively partitioned the effect of transient snakes in the apparent survival rate of the first year since marking. Because we are interested only in the probability of survival over the bulk of the study, the apparent survival rate of the first year since marking can be considered a nuisance parameter. We consider the second survival estimate to reflect more accurately the true survival rate. One biological interpretation of this transience effect is that two subclasses of snakes are present; a "resident" subclass that was sampled consistently throughout the study and a "transient" subclass consisting of individuals that were sampled only rarely (e.g., as they moved onto and off of Beausoleil Island or as they moved in and out of more consistently sampled parts of the island). Whether such transience characterizes Eastern Massasaugas or other snake species more generally warrants investigation. Although this study considers the Eastern Massasaugas residing on Beausoleil Island as a single population, there is movement to and from the island (K. Prior, pers. comm.). The population on Beausoleil Island is part of a larger, roughly defined metapopulation found on the eastern side of Georgian Bay (Szymanski, 1998; Parent and Weatherhead, 2000).

Previously, Middleton and Chu (2004) attempted to use this same data set to estimate a survival rate for a PVA; however, the spreadsheet containing the data was corrupted (possibly attributable to a sorting error) prior to their analysis (Middleton and Chu, 2004). They were able to extract some demographic data and perform a time series analysis on the data set, but they did not estimate survivorship (Middleton and Chu, 2004). Instead, they based their survivorship estimate (0.69) on information from multiple sources, including components of the data set analyzed in the current study (Middleton and Chu, 2004). Brennan and Tischendorf (2004) and Miller et al. (2005) based their model parameters on Middleton and Chu's work and used an adult survival rate of 0.69 as well. The annual survival rate estimated in the current study was somewhat higher than the rates used in those Eastern Massasauga PVAs, although the difference between 0.74 and 0.69 may not be significant. Other adult survival estimates for Eastern Massasauga populations range from 0.33-0.95 (table 1 in Jones et al., 2012) and encompass our estimates for both males and females. Adult survival estimates generated using known-fate models applied to radiotelemetry data show a geographic trend of increasing survival from the southwest to the northeast (Jones et al., 2012). Our average adult annual survival estimate was at the higher end of the range of survival estimates for other Eastern Massasauga populations in Ontario (0.62-0.79) and was higher than the average survival across populations of 0.67 (Jones et al., 2012).

Calculation of the process variance requires long-term data sets. Few other estimates of the process variance of survival have been published for snakes or other reptiles (Whiting et al., 2008; Miller et al., 2011; Stanford, 2012). Process variance of adult and subadult survival for four populations of *Nerodia sipedon insularum* ranged 0.0005–0.125 (Stanford, 2012). Process variance for adult survival in *Nerodia harteri paucimaculata* was 0.0073 over 9 yr (Whiting et al., 2008). Process variance of adult survival in five populations of *Thamnophis elegans* range 0.0003–0.00353 over 8–13 yr (Miller et al., 2011). We were unable to estimate process variance for females; thus, we focus on the male estimate here. The estimate of 0.006 for the male Georgian Bay population falls within the process variance range for *N. s. insularum*, indicating similar variation in survival over time but was much higher than estimates for *T. elegans* or *N. h. paucimaculata* populations, indicating much more variation in survival over time. The unrealistic estimate of process variance for females may be attributable to the smaller sample size of females.

Previous Eastern Massasauga PVAs used the total variance of population parameters (table 1 in Jones et al., 2012). Total variance is made up of the process (= temporal) variance and the sampling (= error) variance, and its use in a PVA will result in an inflated estimate of extinction risk (Gould and Nichols, 1998). As a demonstration of this inflation, we created two population models in VORTEX 10.0.7.9 (Lacy and Pollak, 2014) based on Eastern Massasauga demographic values (Parent and Weatherhead, 2000; Dreslik et al., 2011), including the adult survival and variance estimates from this study. Parameters were adjusted to give the models a deterministic population growth rate of 1.00. The models were identically parameterized except we used the total variance of adult survival as the measure of environmental stochasticity in the first and the process variance of adult male survival in the second (variance was set to 0 for all other survival parameters). Over 1,000 simulations, the probability of extinction over a 50-yr period was 20.8% for the model using total variance, but the probability of extinction was only 3.0% for the model using process variance.

The mean population size estimates of 55 and 54 are quite small, and accordingly, this population is at risk of extinction (Seigel and Sheil, 1999; Mauger and Wilson, 1999). Although our estimation of population size did not use a maximum likelihood methodology, the relative stability of our population size estimates over the years is supported by the population growth rate estimate from the Pradel model. Genetic data suggest Eastern Massasauga populations are frequently small (100 to a few thousand individuals) and relatively isolated (Chiucchi and Gibbs, 2010). In a small population such as this, a 1-2% increase in annual adult mortality can cause large decreases in population size and ultimately population extinction (Seigel and Sheil, 1998). Given the Eastern Massasaugas' protected status in Canada, however, the population may remain stable or even increase in size if protection continues, because this population resides on government land and our long-term population growth rate estimate is slightly greater than 1.00.

Reconstruction and analysis of this 30-yr Eastern Massasauga data set produced multiple demographic values, including the first estimate of temporal process variance in survival for this species and one of the few for any snake species. These demographic values are useful not only for assessing the health of this population of Eastern Massasaugas but also for comparison with known values of other populations and in future analyses such as population projection models and PVAs. As for the Beausoleil Island population itself, retrospective analysis showed the population was relatively stable for over a

#### LITERATURE CITED

decade. If the population remains undisturbed, this stability

may continue. Further monitoring and conservation-related

management will help ensure this Eastern Massasauga popu-

Acknowledgments.—We thank M. Desjardins, K. Prior, and A.

Promaine for their assistance in obtaining the dataset analyzed

here and reviewing the manuscript. We also thank the park staff

lation's persistence.

for assistance with the map.

- BAILEY, R. L., H. CAMPA, III, T. M. HARRISON, AND K. BISSELL. 2011. Survival of eastern massasauga rattlesnakes (*Sistrurus catenatus catenatus*) in Michigan. Herpetologica 67:167–173.
- BEISSINGER, S. R., AND M. I. WESTPHAL. 1998. On the use of demographic models of population viability in endangered species management. Journal of Wildlife Management 62:821–841.
- BISSELL, K. M. 2006. Modeling Habitat Ecology and Population Viability of the Eastern Massasauga Rattlesnake in southwestern lower Michigan. Unpubl. master's thesis, Michigan State University, USA.
- BRENNAN, J. M., AND L. TISCHENDORF. 2004. Eastern Massasauga Rattlesnake *Sistrurus catenatus catenatus* Population Viability in the Ojibway Prairie Complex. Environment Canada, Canada.
- BURNHAM, K. P., D. R. ANDERSON, G. C. WHITE, C. BROWNIE, AND K. H. POLLOCK. 1987. Design and analysis methods for fish survival experiments based on release–recapture. American Fisheries Society Monograph 5:1–437.
- CHIUCCHI, J. E., AND H. L. GIBBS. 2010. Similarity of contemporary and historical gene flow among highly fragmented populations of an endangered rattlesnake. Molecular Ecology 19:5345–5458.
- CHOQUET, R., A. M. REBOULET, J. D. LEBRETON, O. GIMENEZ, AND R. PRADEL. 2005. U-CARE 2.2 User's Manual. CEFE, France.
- CHOQUET, R., J. D. LEBRETON, O. GIMENEZ, A. M. REBOULET, AND R. PRADEL. 2009. U-CARE: utilities for performing goodness of fit tests and manipulating CApture–REcapture data. Ecography 32:1071–1074.
- COULSON, T., G. M. MACE, E. HUDSON, AND H. POSSINGHAM. 2001. The use and abuse of population viability analysis. Trends in Ecology and Evolution 16:219–221.
- CROTHER, B. I., J. BOUNDY, F. T. BURBRINK, J. A. CAMPBELL, AND R. A. PYRON. 2012. Squamata–snakes. Pp. 32–51 in B. I. Crother (ed.), Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico, with Comments Regarding Confidence in Our Understanding. SSAR Herpetological Circular 39. Society for the Study of Amphibians and Reptiles, USA.
- DRESLIK, M. J., S. J. WYLIE, M. A. DAVIS, D. B. WYLIE, AND C. A. PHILLIPS. 2011. Demography of the eastern massasauga (*Sistrurus c. catenatus*) at Carlyle Lake, Illinois. INHS Technical Report 2011(14):1–32.
- ELLNER, S., J. FIEBERG, D. LUDWIG, AND C. WILCOX. 2002. Precision of population viability analysis. Conservation Biology 16:258–261.
- ENVIRONMENT CANADA. 2012. Species at risk public registry: eastern massasauga profile [Internet]. Available from: http://www.sararegistry.gc.ca/species/speciesDetails\_e.cfm?sid=277. Archived by WebCite at http://www.webcitation.org/6jepRsyzk on 10 August 2016.
- GOULD, W. R., AND J. D. NICHOLS. 1998. Estimation of temporal variability of survival in animal populations. Ecology 79:2531–2538.
- HORVITZ, D. G., AND D. J. THOMPSON. 1952. A generalization of sampling without replacement from a finite universe. Journal of the American Statistical Association 47:663–685.
- JONES, P. C., R. B. KING, R. L. BAILEY, N. D. BIESER, K. BISSELL, H. CAMPA, III, T. CRABILL, M. D. CROSS, B. A. DEGREGORIO, M. J. DRESLIK. 2012. Rangewide analysis of eastern massasauga survivorship. Journal of Wildlife Management 76:1567–1586.
- LAAKE, J. L. 2013. RMark: an R interface for analysis of capture–recapture data with MARK. AFSC Processed Rep 2013-01:1–25.
- LACY, R. C., AND J. P. POLLAK. 2014. VORTEX: A Stochastic Simulation of the Extinction Process. Version 10.0. Chicago Zoological Society, USA.

- MANLY, B. F. J. 1984. Obtaining confidence limits on parameters of the Jolly–Seber model for capture–recapture data. Biometrics 40:749– 758.
- MAUGER, D., AND T. P. WILSON. 1999. Population characteristics and seasonal activity of *Sistrurus catenatus catenatus* in Will County, Illinois: implications for management and monitoring. Pp. 110–124 in B. Johnson and M. Wright (eds.) Second International Symposium and Workshop on the Conservation of the Eastern Massasauga Rattlesnake, *Sistrurus catenatus catenatus*: Population and Habitat Management Issues in Urban, Bog, Prairie and Forested Ecosystems. Toronto Zoo, Canada.
- MIDDLETON, J., AND J. Y. CHU. 2004. Population viability analysis (PVA) of the eastern massasauga rattlesnake, *Sistrurus catenatus catenatus*, in Georgian Bay Islands National Park and elsewhere in Canada. Eastern Massasauga Rattlesnake Species Recovery Team, Canada.
- MILLER, D. A., W. R. CLARK, S. J. ARNOLD, AND A. M. BRONIKOWSKI. 2011. Stochastic population dynamics in populations of western terrestrial garter snakes with divergent life histories. Ecology 92:1658–1671.
- MILLER, P., AND PARTICIPANTS OF THE THIRD INTERNATIONAL EASTERN MASSASAUGA SYMPOSIUM. 2005. Population Viability Assessment for the Eastern Massasauga Rattlesnake (*Sitrurus catenatus catenatus*) on the Bruce Peninsula, Ontario, Canada. Conservation Specialist Group, USA.
- PARENT, C., AND P. J. WEATHERHEAD. 2000. Behavioral and life history responses of eastern massasauga rattlesnakes (*Sistrurus catenatus*) catenatus) to human disturbance. Oecologia 125:170–178.
- POLLOCK, K. H., J. D. NICHOLS, C. BROWNIE, AND J. E. HINES. 1990. Statistical inference for capture–recapture experiments. Wildlife Monographs 107:1–97.
- PRADEL, R. 1996. Utilization of capture-mark-recapture for the study of recruitment and population growth rate. Biometrics 52:703–709.
- R CORE TEAM, 2014. R: A Language and Environment for Statistical Computing [Internet]. R Foundation for Statistical Computing, Vienna, Austria. Available from: http://www.R-project.org/.
- ROUSE, J. D., AND R. J. WILLSON. 2002. Update COSEWIC status report on the massasauga *Sistrurus catenatus* in Canada. Pp. in COSEWIC Assessment and Update Status Report on the Massasauga *Sistrurus catenatus* in Canada. Committee on the Status of Endangered Wildlife in Canada, Canada.
- SEIGEL, R. A., AND C. A. SHEIL. 1999. Population viability analysis: applications for the conservation of massasaugas. Pp. 17–22 in B. Johnson and M. Wright (eds.), Second International Symposium and Workshop on the Conservation of the Eastern Massasauga Rattlesnake, *Sistrurus catenatus catenatus*: Population and Habitat Management Issues in Urban, Bog, Prairie, and Forested Ecosystems. Toronto Zoo, Canada.
- STANFORD, K. 2012. Spatial and Temporal Variation in Demographic Parameters of the Lake Erie Watersnake (*Nerodia sipedon insularum*). Unpubl. Ph.D. diss., Northern Illinois University, USA.
- SZYMANSKI, J. A. 1998. Status Assessment for the Eastern Massasauga (Sistrurus c. catenatus). U.S. Fish and Wildlife Service, USA.
- U.S. FISH AND WILDLIFE SERVICE. 2010. Endangered and threatened wildlife and plants; review of native species that are candidates for listing as endangered or threatened; annual notice of findings on resubmitted petitions; annual description of progress on listing actions. Federal Register 75:69222–69294.
- WHITE, G. C., AND K. P. BURNHAM. 1999. Program MARK: survival estimation from populations of marked animals. Bird Study 46 (Suppl.):120–138.
- WHITE, G. C., K. P. BURNHAM, AND D. R. ANDERSON. 2001. Advanced features of Program Mark. Pp. 368–377 in R. Field, R. J. Warren, H. Okarma, and P. R. Sievert (eds.), Wildlife, Land, and People: Priorities for the 21st Century. Proceedings of the Second International Wildlife Management Congress. Wildlife Society, USA.
- WHITING, M. J., J. R. DIXON, B. D. GREENE, J. M. MUELLER, O. W. THORNTON JR., J. S. HATFIELD, J. D. NICHOLS, AND J. E. HINES. 2008. Population dynamics of the concho water snake in rivers and reservoirs. Copeia 2008:438–445.

Accepted: 24 February 2017.

Published online: 28 July 2017.