

Habitat Preferences of Butterflies in the Bumbuna Forest, Northern Sierra Leone

Authors: Sundufu, Abu James, and Dumbuya, Rashida

Source: Journal of Insect Science, 8(64) : 1-17

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.008.6401>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Habitat preferences of butterflies in the Bumbuna forest, Northern Sierra Leone

Abu James Sundufu^a and Rashida Dumbuya^b

Department of Biological Sciences, School of Environmental Sciences, Njala University, Sierra Leone

Abstract

The habitat preferences of the butterfly fauna were studied in the Bumbuna Forest Reserve in northern Sierra Leone. The intact forest reserve and a secondary forest regrowth, disturbed as a result of slash-and-burn agriculture, were compared to savanna habitats. Of the 290 specimens collected, 195 butterfly species were included, of which significant proportion were Nymphalidae. Of the 147 forest species, 111 (75.5%) showed preferences for the forest habitats, while 70 (47.6%) and 34 (23.1%) preferred disturbed and savannah habitats, respectively. Numerically, a comparable proportion of savannah species were recorded in the 18 disturbed (73.9%) and 16 savannah habitats (63.2%). Accumulated species richness and diversity indices were lower in the disturbed habitats compared to the forest reserve, but lowest in the savanna habitats. However, a large proportion of forest species, especially those with either a more restricted geographic range or species for which no information on geographic distribution was available, were exclusively captured in the forest patches. The survey indicated the presence of a rich butterfly fauna, which should be systematically collected for further research and study in order to build a good taxonomic database for Sierra Leone.

Keywords: species richness, species diversity

Correspondence: ^ajasundufu@yahoo.com, ^brashidat_2006@yahoo.com

Received: 8 July 2007 | **Accepted:** 7 September 2007 | **Published:** 24 October 2008

Copyright: This is an open access paper. We use the Creative Commons Attribution 2.5 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Volume 8, Number 64

Cite this paper as:

Sundufu A, Dumbuya R. 2008. Habitat preferences of butterflies in the Bumbuna forest, Northern Sierra Leone. 17pp. *Journal of Insect Science* 8:64, available online: insectscience.org/8.64

Introduction

Tropical forest ecosystems are under enormous pressure all over the world. Many forest areas in the tropics may only persist as production areas (Brown 1997, Hunter 1999), and pressure on unprotected forests is very likely to escalate (Terborgh 1999, Lewis 2000). Despite their generally recognized importance for global diversity (Sutton and Collins 1991; World Conservation Monitoring Centre 1992), no more than 4% of tropical forests are situated within the boundaries of reserves or national parks (Whitmore and Sayer 1992). Even the best protected areas might not be adequate to maintain the original ecosystems because of their small size and difficult political and social circumstances (Terborgh 1999). Although the magnitude of biodiversity present on Earth is largely unknown (Dobson 1995) and its estimates remain highly controversial (May 1990; Stork 1988), it is generally accepted that much, if not most, of the global diversity in terms of numbers of species is represented by arthropods inhabiting tropical rainforests (Wilson 1988). Still, few data are available about the effects of forest disturbance upon these species-rich insect faunas (Klein 1989; Hollo-way et al. 1992; Eggleton et al. 1995;). Butterflies, however, are comparatively well studied. Butterfly species composition in disturbed and undisturbed forests has been investigated for example in Southeast Asia (Spitzer et al. 1993; Hill et al. 1995; Beck and Schulze 2000), Madagascar (Kremen 1992), and the Neotropics (Lovejoy et al. 1986; Brown 1991; Sparrow et al. 1994; DeVries et al. 1997; Wood and Gillman 1998). Several studies showed that low disturbance levels have a positive effect on diversity and abundance of rainforest butterflies (Lovejoy et al. 1986; Brown 1991; Sparrow et al. 1994; Wood and Gillman 1998). These results are in accordance with the intermediate disturbance theory (Connell 1978) and have parallels in temperate forest habitats, where forest management providing a large range of shade levels has been found to increase the number of habitats suitable to different butterfly species (Warren 1985). In contrast, other studies indicate adverse effects of disturbance on tropical butterfly communities (Thomas 1991; Spitzer et al. 1993, 1997; Kremen 1994; Hill et al. 1995, 2001; Brown 1997; Hill and Hamer 1998; Hamer and Hill 2000, Lewis 2000; Fermon et al. 2000, 2001), indicating an increase in diversity and/or abundance of widespread, common butterfly species and a decline in restricted range species after disturbance. Although deforestation rates are highest in several West African regions, little is known about the effects of forest disturbance on afrotropical butterflies (Larsen 1995a). In Madagascar, disturbed forest habitats and edges were equally found to be richer in species than undisturbed areas (Kremen 1992, 1994). Also in southern Nigeria, Larsen et al. (1979) found a surprisingly rich butterfly fauna in mixed secondary growth within the rainforest zone.

In south-central Benin, overall butterfly species richness was higher in clearings than closed forest, however, a high proportion of forest understorey species with a restricted geographic range were exclusively captured in closed forest patches (Fermon et al. 2001). Whereas there is still much work to do to describe the biodiversity of unmodified forest systems, questions concerning ecologically sound management plans cannot be answered without proper assessment in managed forest areas. Considering the high deforestation rates and the fact that a combination of ecology and economy is often the only strategy to protect the rich rainforest biodiversity in many developing countries (Brown 1997), these assessment studies will become increasingly important. The present study mainly documents habitat specificity and diversity of butterflies in the disturbed Bumbuna Forest Reserve in northern Sierra Leone. The study took place within the framework of an Environmental Impact Assessment Survey (Bumbuna Hydroelectric Project) and the data reported here will be included in this survey (TB Larsen in preparation).

Materials and Methods

Study area

Sierra Leone is located on the Atlantic Coast of West Africa, and lies at the western end of the Upper Guinea Forest Block. It is one of the more severely deforested countries in the region (Barrie 2002). Bumbuna is located in the Northern Province along the valley of river Seli. It is a Headquarter town in the Kalansogoia Chiefdom with a total of 65 villages. The total population is 1,700 in about 400 households, the majority of who are farmers. The climate in the study area reflects the general climatic pattern in Sierra Leone, which can be classified as a tropical savannah climate with a distinct tropical wet and dry season. The wet season starts in May and ends in October. Thunderstorms, accompanied by heavy rains, characterize both the start and end of this period. The dry season (November-April) is usually interspersed with the harmattan, a dry dust-laden wind blowing from the Sahara, which occurs between late December and early February, bringing low humidity and relatively cool night temperatures. Highest temperatures are in March with 35°C. Rainfall in the area indicates an average annual precipitation of 2635 mm with maximum in August of 600 mm. The vegetation of the study area is a forest-savannah mosaic type consisting of patches of closed forest communities and serial stages interspersed within savannah woodlands. Closed moist forest regrowth and thicket (secondary forests) and savannah woodland are the three major plant communities occurring predominantly in the area. Three other plant communities present to a more limited extent were (i) fringing forests along rivers and streams (gallery forests), (ii) inland valley swamps (cultivated and natural) and (iii) upland grassland and/or sedges on granite outcrops.

Study site

Many butterflies are localized or restricted to specific habitat types. For this survey, sample sites included: two types of forests (Rashida forest, Radio Hill), three types of savannahs (Kasokira road, savannah to Makeni, savannah-Binkolo to Kafogo), and three types of disturbed habitats (Road leading to Kasassi, Kabari village, Kafogo forest) (Figure 1). The forest habitats are not virgin forests but rather high forests with vegetation cover mostly canopy and sub-canopy. Although, it had not been disturbed for at least 25 years, it is presently under going felling. The 4 sq km Rashida Forest is located in the upper valley and on the right hand of the dam. This forest will be drowned in water upon the inundation of the Bumbuna Hydroelectric Project. Radio Hill is smaller and located on the route to the residential quarters. In the savannah habitats, the vegetation is predominantly grass and small-scale farming is practiced. The disturbed habitats are farm bush habitats with about 3–5 years fallow period, which has decreased by 1 year over this period. Even when recovering from activities like large-scale slash-and-burn farming and logging, small-scale farming for the cultivation of vegetables by local people continues.

Fruit trapping

Traps used were basically as described by DeVries (1987, 1988) and Mühlenberg (1993) and the bait was a fermented banana. Species from 4 out of the 7 African Nymphalidae subfamilies (Libytheinae, Danainae, Satyrinae, Charaxinae, Apaturinae, Nymphalinae and Acraeinae) use fermenting fruit as a resource, including Nymphalinae, Satyrinae, Charaxinae and Apaturinae. In Africa, the following Nymphalinae genera feed consistently on fermenting fruit: *Euphaedra*, *Bebearia*, *Euriphene*, *Euryphura*, *Cymothoe*, *Pseudacraea*, *Euptera* and *Pseudathyma* (Larsen 1994a). Some other Nymphalinae such as *Hypolimnys*, *Salamis* and *Antanartia* are found on both flowers and fruits. Most tropical Satyrinae, especially within the *Bicyclus* and *Gnophodes*, are exclusively fruit-feeders. The Charaxinae and Apturinae are attracted to both fruit and rotting animal matter and faeces. The baited traps were installed 1.0 m above the ground within the study site. Three traps were situated in each of 3 habitats: the understorey of mature forest patches, disturbed forest and savanna habitats. Traps were checked every 24 h and baited with rotting banana, mango and animal faeces. The traps were regularly moved to cover most of the collecting area.

Walk-and-capture

Walk-and-capture transect routes of 1 km each were surveyed during 2 weeks. Three transects were walked between 09:00 and 12:00 in the morning and between 15:00 and 17:00 in the afternoon under sunny weather conditions, each for a duration of 2 hours. Three transects were situated in each of the three habitats: undisturbed forest, disturbed forest and savannah. A total of

14 person-hours were obtained for each habitat. All butterflies seen 2.5m either side of the transect route and up to 5m in front were trapped or released after marking when positive identification was possible (Pollard 1977; Hill et al. 1995).

Butterfly identification and geographic range classification

Butterflies collected were identified using 'Butterflies of West Africa' (Larsen 2005). Habitat associations (preference for certain forest types) and geographic distributions were adopted from Belcastro and Larsen (2006).

Calculation of community parameter


Butterfly diversity was estimated using the following estimators: Shannon-Weiner (H') diversity index (Magurran 1988) and Evenness (J'). The mean number of individuals in each family was calculated and mean separation done using the Least Significant Difference (LSD) test (SAS 1998). Differences in species richness were tested among the forest, disturbed and savannah habitats comparing number of species (S), Shannon-Weiner (H') diversity and Evenness (J') with Kruskal-Wallis ANOVA. Kruskal's gamma rank correlation coefficient (γ) was used to analyze the relationship between geographic range and habitat specificity (Statsoft 1995).

Results and Discussion

Species richness

A total of 195 butterfly species were recorded within the Bumbuna Forest Reserve during the present study (see Appendix). The Lycaenidae (19.00 ± 1.45) and Hesperiiidae (13.67 ± 4.67) are under-represented, constituting only 20% and 14% of the total butterfly fauna. Due to the focus on the fruit-feeding butterfly community, the Nymphalidae family is significantly ($F_{1,4} = 18.48$, $P < 0.05$; 49.33 ± 9.60) represented and comprises 51% of the butterfly fauna listed: members of the subfamily Limenitidinae (35 species) with *Euphaedra*, *Neptis* and *Bebearia* as important forest under-storey genera, Charaxinae (16 species) with *Charaxes* and *Palla* as dominant genera and Satyrinae (14 species) with the genera *Bicyclus*. The Papilionidae (6.67 ± 2.19) and Pieridae (7.67 ± 2.16) make up 7% and 8%, respectively of the total records. No species of the family Riodinidae was recorded.

The total butterfly abundances differ significantly among the three areas (2-way ANOVA, $F_{1,2} = 3.83$, $p < 0.05$), with the highest number captured from the forest habitat. The Shannon-Weiner diversity indices calculated for each sample were significantly higher for both the forest and the disturbed habitats compared to the savannah habitat (Kruskal-Wallis ANOVA, $H = 27.02$ and $P < 0.05$) and no significant difference could be found for evenness (Table 1).



Other surveys of butterflies have been conducted in the Bumbuna Forest yielding totals different from this survey. Larsen in his recent survey in May 2006, recorded 313 butterfly species during one month (including the 195

included in the current survey), while Belcastro (1990b, 1986a,b) made collections, though not regularly, and recorded additional 131 species. Thus 444 species are known from the Bumbuna area. According to TB Larsen

Table 1. Summary of butterfly individuals captured by family and habitat type.

% Species				
	Forest habitat	Disturbed habitat	Savannah habitat	Mean ± SE
Papilionidae	11	5	4	6.67 ± 2.19 b
Pieridae	12	5	6	7.67 ± 2.16 b
Lycaenidae	20	22	16	19.00 ± 1.45 b
Nymphalidae	66	50	32	49.33 ± 9.60 a
Hesperiidae	15	21	5	13.67 ± 4.67 b
Total Families	5	5	5	
Total Individual numbers	124	103	63	
Mean ± SE	24.8 ± 14.4 a	20.4 ± 8.2 ab	12.6 ± 5.3 b	
Shannon (H')	1.33	1.3	1.29	
Evenness (J')	0.83	0.81	0.8	

Means followed by the same letters are not significantly different from each other ($p = 0.05$; LSD). Within habitats, total number of individual butterflies differ significantly (2-way ANOVA, $P < 0.05$, LSD).
Kruskal-Wallis ANOVA indicates significant different in Shannon diversity index.

(personal comm.) another 50 species or so should occur for a total of about 500 in all. According to this estimate, the Bumbuna Forest would comprise 50% of all West African species (West of the Dahomey Gap).

The estimated total species richness is comparable with that of the roughly 21,600 ha large Bossematié Forest (also with about 500 species in total) (Larsen 1994b, 1995a). The actual number of species recorded in this study represent only three-fourths of those recorded in Bossematié Forest.

Ecological composition

Most African butterflies tend to be restricted to one or a limited number of ecological zones and are found in specific habitats (Larsen 1995a). For example, there is a very large difference in total species between fauna of the forest and the northern Sudan savannas, which are separated by the Guinea savanna (Larsen and Mei 1998). The butterfly fauna of West Africa (west of the Dahomey Gap) consist of approximately 1000 species (Larsen 2005).

The species recorded in the Bumbuna Forest thus amount to 19.6% of all butterflies recorded in West Africa (Table 2). Although slash-and-burn agriculture has resulted in a mosaic of forest and disturbed habitats, the overall ecological conditions of the Bumbuna Forest still

appear to meet the habitat requirements of a large number of forest species.

However, the number of forest butterflies species recorded in this study accounts for only 18% of all West African forest species. By contrast, approximately 66% of all West African ubiquitous species were recorded. Only 18% of all savanna species were recorded, which is less than might have been expected. Both ubiquitous and savanna species constitute approximately one fourth of the total number of butterflies sampled in the Bumbuna Forest.

147 (75.4%) of the recorded butterfly species belong to the category of forest species (Table 2), species centered on closed forests that do not usually colonize savanna or other open habitats (Larsen 1994b, 1998; Emmel and Larsen 1997). Only a small fraction are either ubiquitous species (9.8%) or habitat specialists linked to swampy zones (1.5%) or belong to the savanna butterfly community (13.3%). Almost half of all true forest species are species found generally distributed in all forest types, whereas 26% are centered on the moist semi-deciduous forests. Despite the significant number of true forest species, only 18% of all West African forest species have been recorded in the present study.

Table 2. Number of butterfly species collected in the Bumbuna Forest (BF) in the present study by ecological category (Belcastro and Larsen 2006), compared with the total fauna of West Africa (WA) west of the Dahomey Gap (data on the West African butterfly fauna after Larsen and Mei 1998).

Ecological category	Species		% Species	
	BF	WA	BF	WA
Generally distributed in forests	72		36.9	
Centered on the drier forests	13		6.7	
Centered on moist semi-deciduous forests	50		25.6	
Centered on evergreen forest types	12		6.2	
Total forest species	147	821	75.4	82.4
Centered on the Guinea Savannah and forest fringes	24		12.3	
Centered on the Sudan Savannah	2		1	
Total savannah species	26	141	13.3	14.2
Ubiquitous species	19	29	9.8	2.9
Especially demanding species habitats	3	5	1.5	0.5
TOTAL	195	996	100	100

Habitat preference and geographic distribution

Tables 3 and 4 show the number of species recorded during this study in the Bumbuna Forest and do not include the additional species listed by Larsen (2006). As expected, 76% of species classified as forest (Larsen 2006) were collected in the Bumbuna Forest Reserve, while 48% of forest species were collected in farm bush. This suggests that the forest butterflies were largely “robust” species that can survive in farm-bush and small bits of forests and village fruit and sacred groves. Twenty-three percent (23%) of forest species were even found in savanna, mainly in small gallery forests along streams.

By contrast, the largest proportion of savanna species were found in disturbed habitats (69%). A comparable proportion was recorded in the savanna habitat (62%), which was not surprising. However, less than one fourth were recorded in forest habitats.

The proportion of species present in the forest, disturbed and savanna habitats within the Bumbuna Forest Reserve, classified according to their geographic range (Belcastro and Larsen 2006), also show a clear pattern (Table 4). A significant negative correlation between geographic range and habitat specificity was visible in the 195 species captured (gamma rank correlation for multiple ties, $\gamma = -0.2737$, $n = 195$ spp., $P < 0.001$). The

Table 3. Number of butterfly species per ecological category (Belcastro and Larsen 2006) recorded in Bumbuna during the present study, northern Sierra Leone, and percentage of species exclusively recorded in either forest, disturbed or savannah habitats within Bumbuna.

	% Species			
	Bumbuna	Forest habitat	Disturbed habitat	Savannah habitat
Forest species	147	75.5	47.6	23.1
Savannah species	26	15.4	69.2	61.5
Ubiquitous species	19	42.1	73.9	63.2
Especially Demanding Species Habitats	3	37.3	33.3	33.3
TOTAL	195	63.6	52.8	32.3

Table 4. Number of butterfly species by geographic area (Belcastro and Larsen 2006) recorded in Bumbuna, Northern Sierra Leone, and percentage of species exclusively recorded in either forest, disturbed or savannah habitats within Bumbuna.

	% Species			
	Bumbuna	Forest habitat	Disturbed habitat	Savannah habitat
Species found through practically throughout Africa	55	40	61.8	52.7
Species in both the Nigeria and Liberia subregions and extending to the coastal forests of East Africa	15	73.3	66.7	53.3
Species that are endemic to most of Africa west of the Dahomey Gap (Liberia and Ghana subregions)	9	88.9	22.2	11.1
Species in both the Liberia and Ghana subregions and extending to most of equatorial forest zone	89	72.7	51.1	20.2
Species found in both the Liberia and Ghana subregions and extending only to Cameroun, Gabon, Congo, CAR	16	75	43.8	25
Species found in both the Liberia and Ghana subregions and extending only to Niger River or western Cameroun	7	71.4	42.9	14.3
Species that are endemic to Liberia subregion only	1	100	-	-
Species found in both the Liberia and Ghana subregions and extending East Africa north of the Equator or just south thereof	2	-	100	100
No information of geographical distribution	1	100	-	-
TOTAL	195	63.6	52.8	32.3

proportion of species present in forest appears to increase with decreasing geographic range. Only 40% of the species recorded in forest belong to the most widespread group of species, as compared to 61.8% in disturbed and 52.7% in savanna habitats.

Overall species richness in the forest is comparatively the high. It might therefore, be expected that species with a smaller geographic range will thrive in restricted habitats having specific requirements within the Bumbuna Forest Reserve. This emphasizes their importance for maintaining biodiversity on a regional scale. Similar patterns have been found for other West African (Fermon et al. 2000) and South East Asian (Hamer et al. 1997; Hill et al. 1995; Spitzer et al. 1993) forest butterflies. Species with a restricted geographic distribution appear to be more sensitive to human disturbance and forest structure changes than widespread species.

Acknowledgments

We would particularly like to thank TB Larsen, London, for his assistance in the field, sharing his knowledge and experience of the butterflies of West Africa. He was also very helpful with identifications.

References

- Barrie AH. 2002. *Post-conflict conservation status of large mammals in the western area forest reserve, Sierra Leone*. Postgraduate thesis, Njala University College, University of Sierra Leone.
- Beck J, Schulze C. 2000. Diversity of fruit-feeding butterflies (Nymphalidae) along a gradient of tropical rainforest succession in Borneo with some remarks on the problem of 'pseudoreplicates'. *Transactions of the Lepidoptera Society of Japan* 51: 89-98.
- Belcastro C, Larsen TB. 2006. *Butterflies as an indicator group for the conservation value of the Gola forests in Sierra Leone*. Report to the Gola Forest Conservation Concession Project.
- Belcastro C. 1990b. Nymphalidae of Sierra Leone. Ricerche biologiche in Sierra Leone; Problem attuali de scienza e di cultura. *Accademia nazionale dei Lincei* 265: 79-100.
- Belcastro C. 1986a. A new Euriphene (Lepidoptera: Nymphalidae) from Loma mountains, Sierra Leone. Ricerche biologiche in Sierra Leone; Problem attuali de scienza e di cultura. *Accademia nazionale dei Lincei* 260: 195-196.
- Belcastro C. 1986b. A preliminary list of Hesperidae (Lepidoptera) from Loma mountains, Sierra Leone. Ricerche biologiche in Sierra Leone; Problem attuali de scienza e di cultura. *Accademia nazionale dei Lincei* 260: 165-193.
- Brown KS. In: Collins NM, Thomas JA, editors. 1991. Conservation of Neotropical environments: insects as indicators. The conservation of insects and their habitats. 15th Symposium of the Royal Entomological Society of London September 1989 405-424. London. Academic Press.

- Brown KS. 1997. Diversity, disturbance, and sustainable use of Neotropical forests: insects as indicators for conservation monitoring. *Journal of Insect Conservation* 1: 25-42.
- Connell JH. 1978. Diversity in tropical rain forests and coral reefs: high diversity of trees and corals maintained only in a non-equilibrium state. *Science* 199: 1302-10.
- DeVries PJ. 1988. Stratification of fruit-feeding nymphalid butterflies in a Costa Rican rainforest. *Journal of Research on the Lepidoptera* 26: 98-108.
- DeVries PJ. 1987. *The Butterflies of Costa Rica and their natural history. Papilionidae, Pieridae, Nymphalidae*. Princeton University Press.
- DeVries PJ, Murray D, Lande R. 1997. Species diversity in vertical, horizontal, and temporal dimensions of a fruit-feeding butterfly community in an Ecuadorian rainforest. *Biological Journal of the Linnean Society* 62: 343-64.
- Dobson AP. 1995. *Conservation and biodiversity*. Scientific American Libraries, Freeman.
- Eggleton P, Bignell DE, Sands WA, Waite B, Wood TG, Lawton JH. 1995. The species richness of termites (Isoptera) under differing levels of forest disturbance in the Mbalmayo Forests Reserve, southern Cameroon. *Journal of Tropical Ecology* 11: 85-98.
- Emmel TC, Larsen TB. 1997. Butterfly diversity in Ghana, West Africa. *Tropical Lepidoptera* 8: 1-13.
- Fermon H, Waltert M, Larsen TB, Dall'Asta U, Muhlenberg M. 2000. Effects of forest management on diversity and abundance of fruit-feeding nymphalid butterflies in south-eastern Cote d'Ivoire. *Journal of Insect Conservation* 4: 173-185.
- Fermon H, Schulze CH, Waltert M, Muhlenberg M. 2001. The butterfly community of the Noyau Central, Lama Forest (Republic of Benin), with notes on its ecological composition and geographic distribution. *African Entomology* 9: 177-185.
- Hamer KC, Hill JK, Lace L A, Langan AM. 1997. Ecological and biogeographical effects of forest disturbance on tropical butterflies of Sumba, Indonesia. *Journal of Biogeography* 24: 67-75.
- Hamer KC, Hill JK. 2000. Scale-dependent effects of habitat disturbance on species richness in tropical forests. *Conservation Biology* 14: 1435-1440.
- Holloway JD, Kirk-Spriggs AH, Khen CV. 1992. The response of some rain forest groups to logging and conversion to plantation. *Philosophical Transactions of the Royal Society of London B* 335: 425-436.
- Hill JK, Kramer KC, Lace LA, Banham WMT. 1995. Effects of selective logging on tropical forest butterflies on Buru, Indonesia. *Journal of Applied Ecology* 32: 754-760.
- Hunter ML. 1999. *Maintaining biodiversity in forest ecosystems*. Cambridge University Press.
- Klein BC. 1989. Effects of forest fragmentation on dung and carrion beetle communities in Central Amazonia. *Ecology* 70: 1715-1725.
- Kremen C. 1992. Assessing the indicator properties of species assemblages for natural areas monitoring. *Ecological Applications* 2: 203-17.
- Kremen C. 1994. Biological inventory using target taxa: a case study of the butterflies of Madagascar. *Ecological Applications* 4: 407-422.
- Larsen TB, Riley J, Cornes MA. 1979. The butterfly fauna of a secondary bush locality in Nigeria. *Journal of Research on the Lepidoptera* 18: 4-23.
- Larsen TB. 1994a. Fruit-feeding butterflies in large numbers on flowers. *Entomologists' Record and Journal of Variation* 106: 157-8.
- Larsen TB. 1994b. *The butterflies of Ghana- their implications for conservation and sustainable use*. Report to IUCN and Dept. of Game and Wildlife, Ghana.
- Larsen TB. 1995a. Butterfly biodiversity and conservation in the Afro-tropical region. In: Pullin AS, editor. *Ecology and Conservation of Butterflies*, pp. 290-303. Chapman and Hall.
- Larsen TB, Mei M. 1998. Butterflies from a Guinea transition area, the Parc National de Haut Niger (Republic of Guinea), with description of a new subspecies of *Platylesches robusta* (Lepidoptera: Hesperidae). *Bollettino della Societa Entomologica Italiana* 130: 255-272.
- Larsen TB. 2005. *Butterflies of West Africa* 2 vols., 596pp., 125 plates. Apollo Books.
- Larsen TB. 2006. *The Ghana Butterfly Fauna and its Contribution to the Objectives of the Protected Areas System*. WDSP Report no. 63. Wildlife Division (Forestry Commission) and IUCN (World Conservation Union) 207
- Lewis OT. 2002. Effects of experimental selective logging on tropical butterflies. *Conservation Biology* 15: 389-400.
- Lovejoy TE, Bierregaard RO, Ryland AB. 1986. Edge and other effects of isolation on Amazon forest fragments. In: Soulé, editor. *Conservation biology: the science of scarcity and diversity*, pp. 257-285. Sinauer Associates Inc.
- Magurran AE. 1988. *Ecological diversity and its measurement*. Princeton University Press.
- May RM. 1990. How many species? *Philosophical Transactions of the Royal Society of London B* 330: 293-304.
- Mühlenberg M. 1993. *Freilandökologie*, 3rd edition. Quelle and Meyer.
- Pollard E. 1977. A method for assessing changes in the abundance of butterflies. *Biological Conservation* 12: 16-134.
- SAS Institute 1998. *StatView 5.0*. SAS Institute.
- Statsoft 1995. *Statistica 5.1* Vol. 1 available online at www.statsoft.com
- Sparrow HR, Sisk TD, Ehrlich PR, Murphy DD. 1994. Techniques and guidelines for monitoring neotropical butterflies. *Conservation Biology* 8: 800-809.
- Spitzer K, Novotný V, Tonner M, Lepš J. 1993. Habitat preferences, distribution and seasonality of the butterflies (Lepidoptera: Papilionidae) in a montane tropical rain forest, Vietnam. *Journal of Biogeography* 20: 109-121.
- Spitzer K, Jaroš J, Havelka J, Lepš J. 1996. Effects of small-scale disturbance on butterfly communities of an Indochinese montane rainforest. *Biological Conservation* 80: 9-15.

- Stork NE. 1988. Insect diversity: facts, fiction and speculation. *Biological Journal of the Linnean Society* 35: 321-337.
- Sutton SL, Collins NM. In: Collins NM, Thomas JA, editors. 1991. Insects and tropical forest conservation. The conservation of insects and their habitats. 15th Symposium of the Royal Entomological Society of London September 1989 405-24. Academic Press.
- Thomas CD. 1991. Habitat use and geographic ranges of butterflies from the wet lowlands of Costa Rica. *Biological Conservation* 55: 269-281.
- Terborgh J. 1999. *Requiem for Nature*. Island Press.
- Warren MS. 1985. The influence of shade on butterfly numbers in woodland rides to the wood white *Leptidea sinapis*. *Biological Conservation* 33: 147-164.
- Whitemore TC, Sayer JA. 1992. *Tropical deforestation and species extinction*. Chapman and Hall.
- Wilson EO. 1988. The current state of biological diversity. In: Wilson EO, editor. *Biodiversity*, pp. 3-18. Washington National Academic Press. , Washington National Academic Press.
- Wood B, Gillman MP. 1998. The effects of disturbance on forest butterflies using two methods of sampling in Trinidad. *Biodiversity and Conservation* 7: 597-616.
- World Conservation Monitoring Centre 1992. *Global Biodiversity: status of the Earth's living resources*. Chapman and Hall.

Appendix. The preliminary check-list of butterfly species recorded from the Bumbuna Forest in the present study. Authorities, ecological category, geographic distribution (Belcastro and Larsen 2006) and habitat of capture within Bumbuna are given.

			Habitat within-Bumbuna		
	Ecological category	Geographical distribution	FH	DH	SH
FAMILY PAPILIONIDAE Latreille					
Subfamily Papilioniae Latreille					
Genus Papilio Linnaeus					
<i>P. dardanus dardanus</i> Brown	ALF	WWT	x	-	-
<i>P. horribilis</i> Butler	WEF	ENW	x	-	-
<i>P. chrapkowskoides nurettini</i> Koçak	MEF	WWQ	x	-	-
<i>P. nireus nireus</i> Linnaeus	ALF	WWT	-	-	x
<i>P. menestheus menestheus</i> Drury	MEF	WWC	x	-	-
<i>P. demodocus demodocus</i> Esper	UBQ	AFR	x	x	-
Genus Graphium Scopoli					
<i>G. angolanus calabar</i> Hancock	GUI	AFR	x	x	-
<i>G. leonidas leonidas</i> Fabricius	UBQ	AFR	x	x	x
<i>G. illyris</i> Hewitson	WEF	WWQ	x	-	-
<i>G. polices</i> Cramer	ALF	WWT	x	x	x
<i>G. liponesco</i> Suffert	MEF	WWN	x	-	-
<i>G. antheus</i> Cramer	DRF	WWT	x	x	x
FAMILY PERIDAE Swainson					
Subfamily Coliadinae Swainson					
Genus Catopsilia Hübner					
<i>C. florella</i> Fabricius	UBQ	AFR	x	x	x
Genus Eurema Hübner					
<i>E. senegalensis</i> Boisduval	MEF	WWQ	x	-	-
<i>E. hecabe solifera</i> Butler	UBQ	AFR	x	x	x
<i>E. floricola leonis</i> Butler	GUI	AFR	x	x	x
<i>E. brigitta brigitta</i> Stoll	GUI	AFR	-	-	x
Subfamily Pierinae Swainson					
Genus Nephronia Butler					
<i>N. pharis pharis</i> Boisduval	ALF	WWQ	x	-	-
Genus Colotis Hübner					
<i>C. euipe euipe</i> Linnaeus	UBQ	AFR	-	-	x
Genus Belenois Hübner					
<i>B. calypso calypso</i> Drury	ALF	WWQ	x	x	-
Genus Appias Hübner					
<i>A. sylvia sylvia</i> Fabricius	ALF	WWQ	x	x	-
<i>A. sabina sabina</i> Felder & Felder	MEF	WWT	x	-	-
Genus Leptosia Hübner					
<i>L. alcesta alcesta</i> Stoll	ALF	AFR	x	-	x
<i>L.medusa</i> Cramer	MEF	WWN	x	-	-
Genus Mylothris					
<i>M. chloris chloris</i> Fabricius	UBQ	AFR	x	-	-
<i>M. schumanni schumanni</i> Suffert	MEF	WWQ	x	-	-

Appendix (con't.)

			Habitat within- Bumbuna		
	Ecological category	Geographical distribution	FH	DH	SH
FAMILY LYCAENIDAE Leach					
Subfamily Miletinae Reuter					
Genus <i>Euliphyra</i> Holland					
<i>E. hewitsoni</i> Aurivillius	MEF	WWC	x	-	-
Genus <i>Spalgis</i> Moore					
<i>S. lemolea pilos</i> Druce	DRF	AFR	x	-	-
Genus <i>Lachnocnema</i> Trimen					
<i>L. emperamus</i> Snellen	DRF	WWQ	x	-	x
Subfamily Lipteninae					
Genus <i>Ptelina</i> Clench					
<i>P. carnuta</i> Hewitson	MEF	WWQ	x	-	-
Genus <i>Pentila</i> Westwood					
<i>P. pauli pauli</i> Staudinger	DRF	AFR	x	-	-
<i>P. cf. condamini</i> Stempffer	MEF	ENL	x	-	-
Genus <i>Citrinophila</i> Kirby					
<i>C. marginalis</i> Kirby	ALF	WWN	x	-	-
Genus <i>Tetrarhanis</i> Karsch					
<i>T. baralingam</i> Larsen	WEF	ENW	x	-	-
Genus <i>Epitolina</i> Aurivillius					
<i>E. dispar</i> Kirby	MEF	WWQ	x	-	-
Subfamily Theclinae Swainson					
Genus <i>Myrina</i> Fabricius					
<i>M. silenus silenus</i> Fabricius	GUI	AFR	-	x	-
Genus <i>Oxylides</i> Hübner					
<i>O. faunus faunus</i> Drury	MEF	WWQ	x	-	-
Genus <i>Dapidodigma</i> Karsch					
<i>D. hymen</i> Fabricius	MEF	WWN	x	-	x
Genus <i>Axiocerses</i> Hübner					
<i>A. harpax harpax</i> Fabricius	GUI	WWE	-	x	x
Genus <i>Iolaus</i> Hübner					
<i>I. eurisus</i> Cramer	ALF	WWC	-	x	-
Genus <i>Hypolycaena</i> Felder					
<i>H. philippus philippus</i> Fabricius	GUI	AFR	-	x	x
<i>H. lebona lebona</i> Hewitson	WEF	WWQ	x	-	-
<i>H. scintillans</i> Stempffer	MEF	WWC	-	x	-
<i>H. antifaunus antifaunus</i> Westwood	MEF	WWQ	-	x	-
<i>H. hatita habita</i> Hewitson	MEF	WWQ	-	x	-
Genus <i>Pilodeudorix</i> Libert					
<i>P. cameroni cameroni</i> Plötz	MEF	WWQ	x	-	-
Subfamily Polyommatinae Swainson					
Genus <i>Anthene</i> Doubleday					
<i>A. liodes</i> Hewitson	ALF	AFR	x	-	-

Appendix (con't.)

			Habitat within-Bumbuna		
			FH	DH	SH
<i>A. amarah</i> Guérin-Ménéville	SUD	AFR	-	-	x
<i>A. lunulata</i> Trimen	GUI	AFR	-	x	x
<i>A. larydas</i> Cramer	ALF	WWQ	x	x	x
<i>A. lachares lachares</i> Hewitson	WEF	WWC	x	-	-
Genus <i>Neurypexina</i> Bethune-Baker					
<i>N. lyzanius</i> Hewitson	MEF	WWQ	x	x	-
Genus <i>Cupidesthes</i> Aurivillius					
<i>C. leonina</i> Bethune-Baker	MEF	WWQ	-	x	-
Genus <i>Pseudonacaduba</i>					
<i>P. sichela sichela</i> Wallengren	UBQ	AFR	--	x	-
Genus <i>Uranothauma</i> Butler					
<i>U. falkensteini</i> Dewitz	ALF	WWQ	-	x	x
Genus <i>Leptotes</i> Scudder					
<i>L. pirithous</i> Linné	UBQ	AFR	-	x	-
Genus <i>Eicochrysops</i> Bethune-Baker					
<i>E. hippocrates</i> Fabricius	SPE	AFR	x	-	-
Genus <i>Cupidopsis</i> Karsch					
<i>C. cissus cissus</i> Godart	GUI	AFR	-	-	x
Genus <i>Euchrysops</i> Butler					
<i>E. albistriata greenwoodi</i> D'Abrera	GUI	WWQ	-	-	x
Genus <i>Thermoniphas</i> Karsch					
<i>T. micylus</i> Cramer	MEF	WWN	-	x	-
Genus <i>Oboronia</i> Karsch					
<i>O. guessfeldti</i> Dewitz	DRF	WWQ	-	x	x
<i>O. ornata ornata</i> Mabille	ALF	WWQ	x	x	x
Genus <i>Azanus</i> Moore					
<i>A. moriqua</i> Wallengren	SUD	AFR	-	x	x
<i>A. mirza</i> Plötz	UBQ	AFR	-	x	x
<i>A. isis</i> Drury	DRF	WWQ	x	x	x
Genus <i>Zizeeria</i> Chapman					
<i>Z. knysna</i> Trimen	UBQ	AFR	-	-	x
Genus <i>zizina</i> Chapman					
<i>Z. antanossa</i> Mabille	GUI	AFR	-	x	-
FAMILY NYMPHALIDAE Swainson					
Subfamily Danainae Boisduval					
Genus <i>Danaus</i> Kluk					
<i>D. chrysippus</i> Linnaeus	UBQ	AFR	-	x	x
Genus <i>Amauris</i> Hübner					
<i>A. niavius niavius</i> Linnaeus	GUI	AFR	x	-	-
Subfamily Satyrinae Boisduval					
Genus <i>Gnophodes</i> Westwood					
<i>G. betsimena parmeno</i> Doubleday	ALF	AFR	x	-	-

Appendix (con't.)

			Habitat within- Bumbuna		
	Ecological category	Geographical distribution	FH	DH	SH
Genus <i>Melanitis</i> Fabricius					
<i>M. leda helena</i> Linnaeus	UBQ	AFR	x	-	-
Genus <i>Elymniopsis</i> Fruhstorfer					
<i>E. bammakoo bammakoo</i> Westwood	MEF	WWQ	x	-	-
Genus <i>Bicyclus</i> Kirby					
<i>B. procora</i> Karsch	MEF	WWQ	x	-	-
<i>B. taenias</i> Hewitson	ALF	WWC	x	-	-
<i>B. vulgaris</i> Butler	ALF	WWQ	-	x	x
<i>B. dorothea dorothea</i> Cramer	ALF	WWC	-	x	x
<i>B. sandace</i> Hewitson	ALF	WWQ	x	-	-
<i>B. funebris</i> Guérin-Ménéville	DRF	WWQ	-	x	-
<i>B. madetes</i> Hewitson	MEF	WWQ	x	-	-
<i>B. martius melas</i> Condamin	MEF	WWQ	x	x	-
Genus <i>Ypthima</i> Van Son					
<i>Y. doleta</i> Kirby	ALF	WWQ	-	x	x
<i>Y. impura impure</i> Elwes & Edwards	GUI	AFR	-	x	x
Genus <i>Ypthimomorpha</i> Van Son					
<i>Y. itonia</i> Hewitson	SPE	AFR	-	-	x
Subfamily Charaxinae Ochsenheimer					
Genus <i>Charaxes</i> Ochsenheimer					
<i>C. fulvenscens</i> Aurivillius	ALF	WWT	-	x	-
<i>C. cynthia cynthia</i> Butler	ALF	WWQ	x	x	-
<i>C. lucretius lucretius</i> Cramer	ALF	WWQ	-	x	-
<i>C. castor castor</i> Cramer	DRF	WWT	x	x	x
<i>C. tiridates tiridates</i> Cramer	ALF	WWQ	x	x	-
<i>C. numenes numenes</i> Hewitson	ALF	WWQ	x	-	-
<i>C. nobilis claudaei</i> Druce	WEF	WWQ	-	-	x
<i>C. zingha</i> Stoll	MEF	WWQ	x	x	-
<i>C. etesipe etesipe</i> Godart	DRF	AFR	x	x	x
<i>C. achaemenes Atlanta</i> van Someren	GUI	AFR	-	x	x
<i>C. eupale eupale</i> Drury	ALF	WWQ	x	x	x
<i>C. anticlea anticlea</i> Drury	ALF	WWQ	x	x	-
<i>C. virilis virilis</i> Van Sommeren & Jacks	MEF	WWQ	-	-	x
Genus <i>Euxanthe</i> Hübner					
<i>E. eurinome eurinome</i> Cramer	ALF	WWQ	-	x	-
Genus <i>Palla</i> Hübner					
<i>P. violinitens violinitens</i> Crowley	MEF	WWQ	x	-	-
<i>P. decius</i> Cramer	MEF	WWQ	x	--	-
Subfamily Nymphalinae Swainson					
Genus <i>Kallimoides</i> Shirôzu-Nakanishi					
<i>K. rumia rumia</i> Doubleday	ALF	WWQ	x	-	-
Genus <i>Précis</i> Hübner					

Appendix (con't.)

			Habitat within-Bumbuna		
			FH	DH	SH
<i>P. pelarga</i> Fabricius	ALF	WWQ	-	x	-
Genus Hypolimnas Linnaeus					
<i>H. misippus</i> Linnaeus	UBQ	AFR	-	x	x
<i>H. salmacis salmacis</i> Drury	ALF	WWQ	x	-	-
Genus Salamis Boisduval					
<i>S. cacta cacta</i> Fabricius	ALF	WWT	x	x	x
Genus Protogoniomorpha Wallengren					
<i>P. parhassus</i> Drury	ALF	AFR	x	x	-
Genus Junonia Hübner					
<i>J. oenone oenone</i> Linnaeus	UBQ	AFR	x	x	x
<i>J. sophia sophia</i> Fabricius	ALF	WWQ	x	x	x
<i>J. stygia stygia</i> Aurivillius	ALF	WWQ	x	x	x
<i>J. chorimene</i> Guérin-Ménéville	GUI	WWE	-	x	x
<i>J. terea terea</i> Drury	ALF	WWQ	x	x	x
Subfamily Cyrestinae Guenee					
Genus Cyrestini Boisduval					
<i>C. camillus camillus</i> Fabricius	ALF	WWT	x	x	x
Subfamily Biblidinae Boisduval					
Genus Byblia Hübner					
<i>B. anvata crameri</i> Aurivillius	UBQ	AFR	-	x	-
Genus Ariadne Horsfield					
<i>A. albifasca</i> Joicey & Talbot	ALF	WWQ	x	x	-
Genus Eurytela Boisduval					
<i>E. dryope dryope</i> Cramer	DRF	WWT	x	-	-
Subfamily Limenitidinae Behr					
Genus Harma Doubleday					
<i>H. theobene theobene</i> Doubleday	MEF	WWQ	x	-	-
Genus Cymothoe Hübner					
<i>C. egesta egesta</i> Cramer	MEF	WWQ	x	-	-
<i>C. mabillei</i> Overlaet	MEF	ENW	x	-	-
<i>C. sangaris</i> Godart	MEF	WWQ	x	x	x
Genus Pseudoneptis Snellen					
<i>P. bugandensis ianthe</i> Hemming	ALF	WWQ	x	-	-
Genus Pseudacraea Westwood					
<i>P. lucretia lucretia</i> Cramer	ALF	AFR	x	-	-
<i>P. semire</i> Cramer	ALF	WWQ	x	-	-
Genus Neptis Fabricius					
<i>N. nemetes nemetes</i> Hewitson	ALF	WWQ	-	x	-
<i>N. metella metalla</i> Doubleday & Hewitson	ALF	WWQ	x	-	-
<i>N. kiriakoffi</i> Overlaet	GUI	AFR	-	x	-
<i>N. morosa</i> Overlaet	GUI	AFR	-	-	x
<i>N. trigonophora melicertula</i> Staudinger	MEF	WWT	x	-	-

Appendix (con't.)

			Habitat within-Bumbuna		
	Ecological category	Geographical distribution	FH	DH	SH
<i>N. agouale agouale</i> Pierre-Baltus	ALF	WWQ	x	-	-
<i>N. melicerta melicerta</i> Drury	ALF	WWQ	x	-	-
<i>N. troundi</i> Pierre-Baltus	MEF	WWQ	x	-	-
Genus <i>Catuna</i> Kirby					
<i>C. crithea crithea</i> Drury	ALF	WWQ	x	-	-
<i>C. angustatum</i> Felder	MEF	WWQ			
Genus <i>Euryphura</i> Staudinger					
<i>E. chalcis chalcis</i> Felder	ALF	WWQ	x	-	-
Genus <i>Hamanumida</i> Linnaeus					
<i>H. daedalus</i> Fabricius	GUI	AFR	-	x	x
Genus <i>Aterica</i> Boiduval					
<i>A. galena</i> Brown	ALF	WWT	x	x	x
Genus <i>Euriphene</i> Boisduval					
<i>E. aridatha</i> Staudinger	MEF	WWC	x	-	-
<i>E. coerulea</i> Boisduval	WEF	WWN	-	x	-
Genus <i>Bebearia</i> Hemming					
<i>B. osyris</i> Schultze	WEF	ENW	x	-	-
<i>B. absolon absolon</i> Fabricius	ALF	WWQ	x	-	-
<i>B. cocalia</i> Felder & Felder	ALF	WWQ	x	-	-
<i>B. sophus phreone</i> Feisthamel	ALF	WWQ	x	x	-
Genus <i>Euphaedra</i> Hübner					
<i>E. medon</i> Linnaeus	ALF	WWQ	x	x	-
<i>E. xypete</i> Hewitson	MEF	WWC	x	x	-
<i>E. cyparissa</i> Cramer	DRF	WWC	x	x	x
<i>E. themis</i> Hübner	DRF	WWN	x	x	-
<i>E. phaethusa</i> Butler	ALF	ENW	x	x	x
<i>E. inanum</i> Butler	MEF	ENW	-	x	-
<i>E. edwardsii</i> Van der Hoeven	ALF	WWQ	x	x	-
<i>E. harpalyce</i> Cramer	ALF	WWQ	x	x	-
<i>E. eupalus</i> Fabricius	WEF	ENW	x	-	-
Subfamily <i>Heliconiinae</i> Swainson					
Genus <i>Acraea</i> Fabricius					
<i>A. serena</i> Fabricius	UBQ	AFR	x	x	x
<i>A. bonasia bonasia</i> Fabricius	ALF	WWQ	x	x	-
<i>A. egina egina</i> Cramer	ALF	WWT	-	x	-
<i>A. pseudEGINA</i> Westwood	UBQ	WWQ	-	x	x
<i>A. endoscota</i> Le Doux	ALF	WWQ	x	-	-
<i>A. quirina quirina</i> Fabricius	ALF	WWT	x	x	x
<i>A. macaria</i> Fabricius	MEF	ENW	x	-	-
<i>A. alcinoe alcinoe</i> Felder & Felder	MEF	WWQ	x	-	-
<i>A. epaea epaea</i> Cramer	ALF	WWQ	-	-	x
Genus <i>Phalanta</i> Horsfield					

Appendix (con't.)

			Habitat within-Bumbuna		
			FH	DH	SH
<i>P. phalantha aethiopica</i> Roth and Jordan	UBQ	AFR	-	-	x
<i>P. eurytis eurytis</i> Doubleday	ALF	AFR	x	x	x
FAMILY HESPERIIDAE Latreille					
Subfamily Coeliadinae Evans					
Genus Coeliades Hübner					
<i>C. forestan forestan</i> Stoll	UBQ	AFR	x	-	-
<i>C. pisistratus</i> Fabricius	ALF	AFR	x	-	-
Subfamily Pyrginae Swainson					
Genus Celaenorrhinus Hübner					
<i>C. leona</i> Berger	WEF	ENW	x	-	-
<i>C. galenus galenus</i> Fabricius	WEF	???	x	-	-
Genus Tagiades Hübner					
<i>T. flesus</i> Fabricius	ALF	AFR	-	x	-
Genus Eagris Guenée					
<i>E. denuba</i>	ALF	WWN	x	-	-
Genus Eretis Mabille					
<i>E. lugens</i> Rogenhofer	GUI	AFR	x	-	-
<i>E. melania</i> Mabille	DRF	AFR	-	-	x
Genus Sarangesa Moore					
<i>S. tertullianus</i> Fabricius	MEF	WWQ	x	x	-
<i>S. brigida brigida</i> Plötz	MEF	WWQ	x	-	-
Genus Spialia Swinhoe					
<i>S. ploetzi occidentalis</i> de Jong	ALF	WWQ	-	x	-
Subfamily Hesperinae Latreille					
Genus Astictopterus Felder & Felder					
<i>A. abjecta</i> Snellen	GUI	WWQ	-	x	-
Genus Gorgyra Holland					
<i>G. bina</i> Evans	MEF	WWQ	x	-	-
<i>G. afikpo</i> Druce	MEF	WWQ	x	-	-
<i>G. sara</i> Evans	ALF	WWC	x	-	x
Genus Teniorhinus Holland					
<i>T. watsoni</i> Holland	MEF	WWC	x	-	-
Genus Pardaleodes Butler					
<i>P. incerta</i> Murcia	GUI	WWT	-	x	-
<i>P. edipus</i> Stoll	ALF	WWQ	-	x	-
Genus Hypoleucis Mabille					
<i>H. ophiura ophiura</i> Hewitson	ALF	WWQ	x	x	-
Genus Meza Hemming					
<i>M. meza</i> Hewitson	ALF	WWQ	x	-	-
<i>M. mabea</i> Holland	MEF	WWQ	x	-	-
<i>M. leucophaea leucophaea</i> Holland	MEF	WWC	x	-	-

Appendix (con't.)

			Habitat within-Bumbuna		
	Ecological category	Geographical distribution	FH	DH	SH
Genus <i>Andronymus</i> Holland					
<i>A. caesar caesar</i> Fabricius	ALF	AFR	-	x	-
<i>A. helles</i> Evans	MEF	WWQ	-	x	-
<i>A. evander evander</i> Mabilie	MEF	WWC	x	x	x
Genus <i>Gretna</i> Evans					
<i>G. waga</i> Plötz	ALF	WWQ	-	x	