

Entomopathogenic Activity of a Variety of the Fungus, *Colletotrichum acutatum*, Recovered from the Elongate Hemlock Scale, *Fiorinia externa*

Authors: Marcelino, José A. P., Gouli, Svetlana, Parker, Bruce L, Skinner, Margaret, and Giordano, Rosanna

Source: Journal of Insect Science, 9(13) : 1-9

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.009.1301>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Entomopathogenic activity of a variety of the fungus, *Colletotrichum acutatum*, recovered from the elongate hemlock scale, *Fiorinia externa*

José A. P. Marcelino^{1,a}, Svetlana Gouli¹, Bruce L. Parker¹, Margaret Skinner¹ and Rosanna Giordano²

¹Department of Plant and Soil Science, Entomology Research Laboratory, The University of Vermont, Burlington, VT 05405-0105 USA

²Illinois Natural History Survey, Division of Biodiversity and Ecological Entomology, Champaign, IL 61820 USA

Abstract

A fungal epizootic in populations of *Fiorinia externa* Ferris (Hemiptera: Diaspididae) infesting hemlock trees, *Tsuga canadensis* (L.) Carrière (Pinales: Pinaceae) in forests of the Northeastern US has been recently detected. The current known distribution of the epizootic spans 36 sites in New York, Pennsylvania, New Jersey and Connecticut. *Colletotrichum acutatum* Simmonds var. *fiorinae* Marcelino and Gouli var. nov. inedit. (Phyllachorales: Phyllachoraceae) was the most prevalent fungus recovered from infected scales. Bioassays indicated that this *C. acutatum* variety is highly pathogenic to *F. externa*. Mortality rates of >90 and >55% were obtained for *F. externa* crawlers and settlers, respectively. Significantly lower mortality levels, ≤ 22%, were obtained when three other species of insects were assayed. *C. gleosporioides* has also been shown to have pathogenic activity towards a scale insect. The data suggest that *C. acutatum* var. *fiorinae* from *F. externa* epizootics in the US, and the previously reported *C. gleosporioides* f. sp. *ortheziidae* causing *Orthezia praelonga* epizootics in Brazil, may constitute distinct biotypes of *Colletotrichum* that have attained the ability to infect insects in addition to the commonly reported plant hosts.

Keywords: fungal epizootic, *Tsuga canadensis*, *Orthezia praelonga*

Correspondence: ^ajmarcelino@uac.pt

Received: 5 November 2007 | **Accepted:** 17 January 2008 | **Published:** 16 April 2009

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 9, Number 13

Cite this paper as:

Marcelino JAP, Gouli S, Parker BL, Skinner M, Giordano R. 2009. Entomopathogenic activity of a variety of the fungus, *Colletotrichum acutatum*, recovered from the elongate hemlock scale, *Fiorinia externa*. 9pp. *Journal of Insect Science* 9:13, available online: insectscience.org/9.13

Introduction

The eastern hemlock, *Tsuga canadensis* (L.) Carrière (Pinales: Pinaceae), a common species in forests of the Northeastern United States is in decline (Orwig et al. 2002). The invasive elongate hemlock scale (EHS), *Fiorinia externa* Ferris (Hemiptera: Diaspididae), has been identified as one of the causal agents of this decline (Lambdin et al. 2005). Attempts to control this pest have not been successful. The unique shield-like cover of the scale provides protection from contact insecticides, natural enemies and adverse climatic conditions. Because of its high reproductive rate, even when mortality exceeding 90% occurs, populations quickly rebound (Baranyovits 1953; Johnson and Lyon 1988).

In 2002 a fungal epizootic, whose geographic point of origin was unknown, was reported within the population of *F. externa* in the Mianus River Gorge Preserve in Bedford, NY (McClure 2002). Sclerotia were found concealing the bodies of adult mummified scales. Evidence of this infection was found among scales in 36 different sites in New York, Pennsylvania, Connecticut and New Jersey. A complex of entomopathogenic, phytopathogenic and saprophytic fungi was morphologically and molecularly identified as being associated with the diseased insects (Marcelino et al. 2009a). One species, *Colletotrichum acutatum* var. *fiorinae* var. nov. inedit. (Marcelino et al. 2008), was dominant in this complex and consistently recovered in *F. externa* populations in most of the epizootic localities.

Members of the genus *Colletotrichum* are known as cosmopolitan plant pathogens (Sreenivasaprasad and Talhinhos 2005), many of which cause anthracnose in several commercially important crops (Byrne et al. 1997; Lardner et al. 1999; Khan and Hsiang 2003; Horowitz et al. 2004; Xiao et al. 2004). Literature on the phytopathogenic genus *Colletotrichum* (Bailey and Jeger 1992; Prusky et al. 2000) includes a single report of the species *C. gloeosporioides* causing significant epizootics in the scale *Orthezia praelonga* Douglas 1891 (Hemiptera: Ortheziidae), a major pest of citrus in Brazil. This fungus, *C. gloeosporioides* f. sp. *ortheziidae* is under commercial development for management of *O. praelonga* (Cesnik and Ferraz 2000). The epizootic caused by *C. acutatum* var. *fiorinae* is the second report of a member of this genus infecting a scale insect. To understand the role of this fungus in the *F. externa* epizootic, the virulence of *C. acutatum* var. *fiorinae* to four insect species from three orders (Hemiptera, Lepidoptera and Thysanoptera) was evaluated.

Materials and Methods

Isolates

The virulence of five *C. acutatum* var. *fiorinae* isolates from different areas of the *F. externa* epizootic in the Northeast

U.S. was tested (Table 1). In addition, the following other isolates were also assayed: *C. gloeosporioides* f. sp. *ortheziidae* from the Brazilian epizootic in *O. praelonga* (ARSEF4360) (obtained from the Agricultural Research Service Entomopathogenic Fungal Collection, Cornell University, Ithaca, NY), two phytopathogenic *C. acutatum*, one isolated from blueberry (ERL1379) and one from tomato (ERL1380), *Lecanicillium lecanii* (Zimmerman) Gams & Zare (EHS132), an entomopathogenic fungal strain isolated from *F. externa* and one *Metarhizium anisopliae* (Metschn) Sorokin (CA-1), recovered from litter in a California avocado orchard (used only in the *Frankliniella occidentalis* bioassays because of its virulence to this insect). The initial isolation of the fungi was obtained by growth on potato dextrose agar medium (39 g/l) supplemented with penicillin (5 ml/l) and streptomycin (12.5 ml/l). Fungal isolates have been deposited at the University of Vermont Entomology Research Laboratory (UVM ERL) Worldwide Collection of Entomopathogenic Fungi, Burlington, VT. Isolates have been stored as mature mycelium (2 weeks old) in potato dextrose agar cubes (1 cm²) in cryogenic vials (8 replicates) containing 10% glycerol and held at -80° C.

Fungi used in the bioassays were grown in potato dextrose agar (39 g/l) for 10–12 days before being harvested with sterile Pasteur pipettes to obtain inoculum suspensions, in sterile distilled water. Calibration of conidial spore suspensions to 10⁶ and 10⁷ conidia/ml⁻¹, concentrations commonly used in insect inoculation bioassays (Abe and Ikegami 2005; Fransen 1987; Klingen et al. 2002), was done using an Improved Neubauer haemocytometer (Propper®) according to the protocol of Goettel and Inglis (1997).

Insects

The virulence of the above fungal isolates was tested for the two increasing doses of inoculum against several insect species representing three orders, Hemiptera, Thysanoptera and Lepidoptera, to understand their comparative infectiveness and dose-mortality response.

The silverleaf whitefly, *Bemisia argentifolii* Bellows and Perring (Hemiptera: Aleroydidae), was reared on poinsettia, *Euphorbia pulcherrima* Wild.ex Klotsch (Malpighiales: Euphorbiaceae) at the UVM ERL according to the protocol of Negasi et al. (1998). For these bioassays, terminal leaves of 2–3 week old bean plants, *Phaseolus vulgaris* L. (Fabales: Fabaceae) var. Royal Burgundy, were excised and the petiole placed in an Oasis® rooting cube (Smithers-Oasis, www.smithersoasis.com) held in place by absorbent cotton wool. Cubes were placed in tap water in 9 mm diameter Petri dishes until roots formed (~4 days), then 18 mating pairs of whiteflies were removed from poinsettia, anesthetized for 2 seconds with carbon dioxide and placed on each bean leaf. Infested leaves were held in vented plastic boxes (8.7 cm wide x 9.5 cm

Table 1. Fungal isolates tested in the insect bioassays

Fungus type	Species	Code	Host	Geographic origin	Year of collection
Entomopathogenic fungi	<i>Colletotrichum acutatum</i> var. <i>fioriniae</i> var. nov. inedit. (<i>C. a. fioriniae</i>)	EHS41	<i>Fiorinia externa</i>	Mohonk, NY	2005
	<i>C. acutatum</i> var. <i>fioriniae</i>	EHS48	<i>Fiorinia externa</i>	Bayberry Lane, NY	2005
	<i>C. acutatum</i> var. <i>fioriniae</i>	EHS51	<i>Fiorinia externa</i>	Esopus, NY	2005
	<i>C. acutatum</i> var. <i>fioriniae</i>	EHS58	<i>Fiorinia externa</i>	Ward Pound Ridge Reservation, NY	2005
	<i>C. acutatum</i> var. <i>fioriniae</i>	EHS61	<i>Fiorinia externa</i>	Ward Pound Ridge Reservation, NY	2005
	<i>Lecanicilium lecanii</i>	EHS132	<i>Fiorinia externa</i>	South Salem, NY	2005
	<i>Metarhizium anisopliae</i> ^a	CA-1	Soil from avocado plantation	Temecula, CA	2001
	<i>C. gloeosporioides</i> f. sp. <i>ortheziidae</i>	ARSEF4360	<i>Orthezia praelonga</i>	Jaguariuna, Sao Paulo, Brazil	1994
Phytopathogenic fungi	<i>C. acutatum</i>	ERL1379	Blueberry fruit	NJ	2005
	<i>C. acutatum</i>	ERL1380	Tomato fruit	Burlington, VT	2005

^aUsed in the *F. occidentalis* bioassays only

long) at 16:8 LD, 75% RH and 24°C. Adults were removed after 24 h to ensure age homogeneity of the progeny. On each leaf 40–180 1st instars were produced.

Elongate hemlock scales, *Fiorinia externa* Ferris (Hemiptera: Diaspididae), were field-collected from understory eastern hemlock, *T. canadensis* trees at the Mount Tom Forest Preserve, Holyoke, MA, which is located outside the known area of the *F. externa* epizootic. One day prior to the bioassay, 30 branches (50 cm long) with new growth and naturally infested with a healthy population of *F. externa* crawlers (i.e. 1st instar mobile nymph stage emerged from the 3rd instar adult female exuvia) and settlers (i.e. 2nd instar immobile nymph stage after inserting stylets in the epidermal cells of hemlock leaves, losing their legs and remaining anchored for life) were randomly sampled. Branches were kept cool during transport and held at 4° C prior to treatment. Eighty 10 cm long terminal twigs with new growth were clipped from the branches for the assay. On each infested twig 10–200 settlers and 1–46 crawlers were counted. Freshly pruned branches were gathered for each assay repetition.

Western flower thrips, *Frankliniella occidentalis* Pergande (Thysanoptera: Thripidae), were reared at the UVM ERL on bean leaves (var. Royal Burgundy) according to the protocol of Doane et al. (1998). Two-day-old 2nd instars were used for testing due to the natural high mortality of 1st instars. Each replicate of the assay had 10 thrips/leaf.

Eggs of the beet armyworm, *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae), were purchased from Benzon Research, Inc. (www.benzonresearch.com). Upon delivery, eggs were allowed to hatch in a glass container (12

cm diam. x 20 cm high) containing cabbage leaves. The containers were wiped with an antistatic tissue before introduction of eggs. The glass containers were held for 3–4 days at 22° C and 16:8 LD. After eclosion 20 1st instars were randomly selected for each replicate.

Bioassays

For all bioassays, fungal concentrations of 10⁶ and 10⁷ conidia/ml⁻¹ were suspended in sterile distilled water with 0.02% Silwet (Momentive, www.gesilicones.com) as a surfactant. A 0.02% solution of Silwet was used for the controls, with the exception of *F. externa* trials where blank controls were used. Each insect bioassay was repeated three times with four replicates for each treatment.

For the *B. argentifolii* assays, each treatment consisted of four leaves, each with 40–180 2 day old 1st instars. A Potter Precision Laboratory Spray Tower (Burkard Manufacturing Co. Ltd., www.burkard.com.uk) operating at 0.84 kg/cm with a 0.25 mm diameter nozzle was used to spray 2.5 ml of the fungal suspension. Mortality was assessed after 30 days. Individual whole insects were first inspected for morphological changes in the cuticle or body, i.e. changes in color or body turgor. In cases where mortality could not be confirmed in this manner, insects were squashed on a glass slide and checked for the presence of hyphae and fungal spores in the hemolymph, as well as, hyphae penetrating through the cuticle.

For the *F. externa* assays, isolates were tested against settlers and crawlers using modified protocols of Rose (1990) and Butt & Goettel (2000). For each treatment (i.e. 10⁶ and 10⁷ conidia/ml⁻¹), four twigs, each containing ≥ 10 crawlers or settlers, were held vertically in a metal test

tube rack and individually sprayed at a distance of 38 cm with 250 μl of a microdroplet mist of a conidial suspension using a hand-held plastic spray bottle. Homogeneous distribution, density and size of sprayed droplets were visually assessed. Twigs were allowed to air dry for 2 min, and were then placed individually in sterile graduated 50 ml conical plastic tubes (www.fishersci.com) containing 16 g of sterilized sand (Quikrete, www.quikrete.com) and 7 ml sterile distilled water. Each tube was covered loosely with a cap to allow ventilation. Tubes were placed in plastic bags to prevent desiccation and held at 22° C with 16:8 LD. Mortality of crawlers and settlers was determined 21 days after treatment as described above for the whitefly assay.

For the *F. occidentalis* assays, bean leaf discs (3.3 cm diameter) were placed on moist filter paper in 3.5 cm diameter Petri dishes, to which 10–2nd instars were added. Each Petri dish assembly was sprayed with 2 ml of the fungal suspension using a Potter Spray Tower, as described previously. After being air dried for 2 min, Petri dishes were covered and sealed with Parafilm, and held in the dark at 22 ± 1°C. Mortality was assessed 7 days after treatment. Insects that exhibited obvious signs of fungal infection, i.e. displayed an abnormal body color or lacked turgor, and those that did not respond when gently probed with a small insect pin, were considered dead.

S. exigua were assayed in well plates with 5 x 4 cells (13 mm diameter/cell) (Model #BIO-BA-128, Color-Dec, www.color-dec.it). A 10 mm diameter disc of moist filter paper was placed in the bottom of each well, followed by 10 mm diameter cabbage leaf disc and one 1st instar beet armyworm. Each 20-cell unit was sprayed with 2 ml of the test suspension with a Potter Spray Tower. Cell units were air dried for 2 min, covered with clear plastic wrap and held in an incubator in the dark at 22 ± 1° C. Mortality was assessed after 7 days as described for the *F. occidentalis* assays.

To test Koch's postulates (i.e. re-isolation of the test fungi from a diseased host after treatment of a healthy individual), a random subsample of 10 insects from each bioassay was taken. Each species of insect was surface sterilized in 0.01% NaOCl, rinsed in sterile distilled water and placed in a Petri dish on potato dextrose agar with 5 ml/l penicillin and 12.5 ml/l streptomycin. Petri dishes were held in the dark at 22 ± 1° C for 7 days, and then cadavers were examined for the presence of *C. acutatum* var. *fioriniae* or the other fungi tested.

Statistical analyses

Protocols differed among insect species assayed due to differences in sample sizes of *F. externa* and *B. argentifolii* that varied according to female fecundity. Therefore, a statistical treatment adjusted for an unbalanced design was used. Variances were not homogeneous (using

Levene's test), hence, a Welch's one way ANOVA (unpooled variances) was carried out. Transformation of the data was not required since they were normally distributed as observed by plotting the residuals from the ANOVA. An adjusted pairwise comparison between fungal isolates within a test insect species was made using a post-hoc Tukey-Kramer test. The effect of suspension concentration (10⁶ or 10⁷ conidia/ml⁻¹) was determined with an adjusted least square means (LS means). *P* < 0.05 was considered statistically significant. All statistical analyses were performed using SAS[®] (SAS Institute 1990) and plotted using SPSS[®] (SPSS Inc. 2005).

Results

Definite signs of infection were observed among insects treated with *C. acutatum* var. *fioriniae* var. nov. inedit., demonstrating its entomopathogenic capacity (Figure 1A-E). In the control treatments and those sprayed with *C. gloeosporioides* f. sp. *orthesiidae* *F. externa* underwent normal development and reached maturity (Figure 1F-G). However, normal development was halted among *F. externa* treated with *C. acutatum* var. *fioriniae* isolates and phytopathogenic *C. acutatum* isolates. Infected *F. externa* settlers did not attain maturity (Figure 1E) and both settlers and crawlers showed symptoms of mycosis (Figure 1D-E). Koch's postulate was successfully achieved for the *C. acutatum* var. *fioriniae* isolates in *F. externa*, silverleaf whitefly and western flower thrips. All 10 cadavers of *F. externa*, *S. exigua* and *B. argentifolii*, and two individuals of *F. occidentalis* showed evidence of infection with *C. acutatum* var. *fioriniae* (i.e. pink mycelia growing outward from the body of the insects to the media). Infection by *C. acutatum* var. *fioriniae* was confirmed by visual examination of spores using a stereomicroscope.

Insect mortality varied depending on isolate, conidial concentration and insect species tested. Differences in mortality between the two conidial concentrations tested (10⁶ and 10⁷ conidia/ml⁻¹) were not statistically significant for any of the isolates tested against *F. externa* crawlers and settlers whereas for other species tested, *S. exigua*, *B. argentifolii* and *F. occidentalis*, significant differences were observed between the two concentrations tested (Tables 2 and 3).

Although a statistical comparison of mortality across species could not be done due to differences in protocols and sample sizes used in the experiments conducted with the four insect species, the mortality caused by *C. acutatum* var. *fioriniae* isolates in crawlers and settlers of *F. externa* was higher than in the other species tested (Figures 2 and 3). Crawlers of *F. externa* were highly susceptible to *C. acutatum* var. *fioriniae* isolates (Figure 1D) with a maximum *F. externa* crawler mortality rate of 92.64% for isolate EHS48 at 10⁶ conidia/ml⁻¹ concentration and 93.44% for isolate EHS61 at 10⁷ conidia/ml⁻¹. When *F. externa*




Figure 1. Fungal structures from bioassays of *C. acutatum* var. *fiorinae* isolates and symptomatic and non-symptomatic *F. externa*. A) acervuli arising from head of an infected whitefly; B) crawler with septicemia from EHS51; C) septicemia in *F. externa* adult from EHS41; D) generalized septicemia in crawlers from EHS51 treatment; E) arrested development in diseased *F. externa*; F) normal development of the scale cover in *F. externa* from controls; G) normal development of the scale cover in nature; H) mycelium arising from adult *F. externa*. Bars: A = 50µm; B = 0.1 mm; C = 0.5 mm; D, E = 1 mm; F, G = 0.25 mm; H = 50 µm.

Table 2. Results of statistical analysis for bioassays with *Fiorinia externa* immatures

		<i>F. externa</i> settlers			<i>F. externa</i> crawlers		
		(N=12,343)			(N=989)		
Main effect		F	P	df	F	P	df
(Welch's one way ANOVA)		33.19	<0.0001*	9	41.72	<0.0001*	9
	Test isolate						
Mortality (Isolate * concentration) (LS means)	EHS41	0.75	0.38	1.00	0.07	0.79	1.00
	EHS48	0.00	0.97	1.00	0.66	0.41	1.00
	EHS51	0.51	0.47	1.00	0.00	0.94	1.00
	EHS58	1.12	0.29	1.00	1.01	0.31	1.00
	EHS61	0.03	0.86	1.00	0.58	0.44	1.00
	ERL1379	1.20	0.27	1.00	0.61	0.43	1.00
	ERL1380	0.01	0.92	1.00	1.04	0.30	1.00
	EHS132	0.01	0.93	1.00	0.14	0.70	1.00
	ARSEF4360	0.07	0.79	1.00	0.18	0.67	1.00
	Blank	0.29	0.59	1.00	0.00	0.98	1.00

*Significant differences in percent mortality

settlers were inoculated with isolate EHS58 a maximum mortality rate of 59.27% and 50.28% was observed with 10⁷ conidia/ml⁻¹ and 10⁶ conidia/ml⁻¹ respectively. In general mortality from *C. acutatum* var. *fiorinae* isolates was about 35% greater among crawlers than settlers. For the crawler bioassay, mortality caused by the two other entomopathogens tested, *C. gloeosporioides* f. sp. *orthosporiae* ARSEF4360 and *L. lecanii* EHS132, was significantly different from all other fungi tested, with the exception of the plant pathogen ERL1379. In contrast, mortality of *F.*

externa infected with the plant pathogen ERL1380 was significantly greater than the two entomopathogenic fungi ARSEF4360 and EHS132 but always statistically below levels attained with *C. acutatum* var. *fiorinae* isolates. Mortality of 22% or less was obtained for the other three insect species tested (*P* < 0.05) with *F. occidentalis* and *S. exigua* at <10%, and *B. argentifolii* between 15 to 22%. However three exceptions were observed: *L. lecanii* EHS132 at 10⁷ conidia/ml⁻¹ caused 25% mortality in *B.*

Table 3. Results of statistical analysis for bioassays with *Spodoptera exigua*, *Bemisia argentifolii* and *Frankliniella occidentalis*

		<i>S. exigua</i>			<i>B. argentifolii</i>			<i>F. occidentalis</i>		
		(N=1,200)			(N=11,492)			(N=1,320)		
Test isolates		F	P	df	F	P	df	F	P	df
Main effect ^a		3.07	0.0029*	9	3.79	0.0006*	8	1.33	0.24	10
Mortality ^b	EHS41	10.91	<0.001*	1.00	3.87	0.05	1.00	1.44	0.23	1.00
	EHS48	17.67	<0.001*	1.00	2.21	<0.13	1.00	0.00	1.00	1.00
	EHS51	20.28	<0.001*	1.00	1.19	<0.27	1.00	0.36	0.54	1.00
	EHS58	0.81	0.368	1.00	6.49	<0.01*	1.00	3.24	0.07	1.00
	EHS61	0.36	0.543	1.00	6.45	<0.01*	1.00	1.44	0.23	1.00
	ERLI379	1.44	0.231	1.00	1.56	<0.21	1.00	1.44	0.23	1.00
	ERLI380	3.24	0.073	1.00	4.32	<0.03*	1.00	5.76	0.01*	1.00
	EHS132	70.67	<0.001*	1.00	7.41	<0.007*	1.00	0.00	1.00	1.00
	ARSEF4360	0.00	1.000	1.00	•	•	•	0.00	1.00	1.00
	CA-1	•	•	•	•	•	•	81.03	<0.0001*	1.00
	SDW	3.24	0.073	1.00	0.01	0.93	1.00	0.36	0.54	1.00

^aWelch's one way ANOVA
^bIsolate x concentration, LS means
*Significant differences in percent mortality
•Not tested

argentifolii and 14.5% in *S. exigua*. *M. anisopliae* CA-1 at 10⁷ conidia/ml⁻¹ caused 12.5% mortality in *F. occidentalis* (Figure 3). Greater mortality was obtained at the higher conidia concentration. With the exception of *F. externa*, whiteflies appeared to be the most susceptible to infection at the higher concentration.

Discussion

This research indicates that the fungus *C. acutatum* var. *fiorinae* var.nov. inedit., isolated from infected *F. externa* adults recovered from several localities in the North East US hemlock forests where a fungal epizootic occurred was highly pathogenic to this insect host, particularly in the crawler stage.

High mortality rates were caused in both developmental stages of *F. externa* infected with inoculations of *C. acutatum* var. *fiorinae* var. nov. inedit. Mortality, but at significant lower levels, was also obtained when *F. externa* crawlers and settlers were treated with different *C. acutatum* isolates known to be phytopathogenic. Mortality in the other three insect species tested with the *C. acutatum* var. *fiorinae* and the other fungi was lower: *Bemisia argentifolii* had levels of ≤ 22%, and both *Spodoptera exigua* and *Frankliniella occidentalis* had levels <10%. These results indicate that the latter three species displayed lower susceptibility to both entomopathogenic and phytopathogenic *Colletotrichum* isolates.

This report of virulence of *C. acutatum* var. *fiorinae* to *F. externa* along with the demonstrated virulence and

biocontrol activity of *C. gloeosporioides* f. sp. *ortheziae* to scales of citrus for the last 20 years (Cesnik and Ferraz 2000) supports the generally held hypothesis that members of the genus *Colletotrichum* have a broader host range and inhabit niches other than those currently reported (Guerber and Correll 2001, Peres et al. 2005). The pattern of infection of *F. externa* with *C. acutatum* var. *fiorinae* has a patchy but widespread geographic distribution. The infection has been detected in the northeastern states of Connecticut, New York, New Jersey and Pennsylvania, suggesting that this infection is established in *F. externa* populations in the Northeastern U.S. While the number of insect species tested was limited it appears that the entomopathogenic activity of *C. acutatum* var. *fiorinae* was higher against *F. externa* than against the other insects tested, suggesting that this variety of *C. acutatum* may preferentially infect scale insects. Arthropods from additional orders (Araneae, Hymenoptera, Lepidoptera, Orthoptera) collected from the trunks of hemlocks occurring within the epizootic areas showed no evidence systemic infection with *Colletotrichum* spp. Specimens belonging to the above orders were surface sterilized (following the procedure mentioned herein) and placed on potato dextrose agar did not show evidence of *Colletotrichum* spp. mycelia growth (Marcelino et al. unpublished data).

C. acutatum var. *fiorinae* has been found growing endophytically in over 28 different species of plants within the epizootic areas. In addition, we have conducted laboratory tests to access the pathogenicity of this fungus to several plants, i.e. strawberries, beans, peppers, tomato, barley and hemlock (Marcelino et al. 2009b). Since this




Figure 2. Percent mortality among *F. externa* settlers and crawlers in fungal bioassays. ($\pm$ 95% CI of the mean). Bars with the same letters are not significantly different (data for both concentrations combined) ($P < 0.05$) using the post-hoc multiple comparison Tukey-Kramer test.

fungus is commonly found in a variety of plants within the Northeast hemlock forest it is possible that this fungus is a native species and that the invasive *F. externa* became infected subsequent to its arrival. An alternative hypothesis may be that introduced specimens of *F. externa* were infected with this variety of fungus at the time of introduction and that both scale and fungus are invasive species.

Mutualist or commensal endophytic associations between plants and members of the genus *Colletotrichum* have been reported (Zulfiqar et al. 1996; Makowski and Mortensen 1998; Tsrar and Johnson 2000; Lu et al. 2004). Several similar reports have also been published regarding *C. acutatum* (Zulfiqar et al. 1996; Freeman et al. 2001; Horowitz et al. 2002). It has been suggested that the life strategies adapted by *Colletotrichum* spp. (i.e. mutualism, parasitism or commensalism) are controlled in part by the host plant genotype (Redman et al. 2001) and that both chemical and physical factors between the fungus and plant host, direct gene expression of the fungus (Memmott et al. 2002; Dickman 2000, 2004). Similar relationships may have occurred between *F. externa* and *C. acutatum fiorinae*. Studies with *C. magna* endophytic mutants showed that strains with nonpathogenic life strategies had a broader host range than the parental pathogenic *C. magna* strains (Redman et al. 2001). *C. acutatum* also displays endophytic activity, in most plants we have sampled in the field and tested in the laboratory. This non-pathogenic behavior may have facilitated the shifting of hosts, of this *C. acutatum* variety, from plants to insects.

It is apparent from these reports that members of the genus *Colletotrichum* display a great degree of plasticity in host choice, however, to date, most of this work has focused solely on plants. Our data as well as that of Cesnik and Ferraz (2000) provides strong support that members of this genus can also be effective primary pathogens of insects.

Acknowledgments

We thank Tom Doubleday, Cheryl Frank, Teri Hata and Dr. Robert Jones for their help. This work was funded in part through a grant awarded by the Northeastern Area State and Private Forestry, USDA Forest Service (#04-CA-11244225286) and is in partial fulfillment of requirements for the Ph.D. degree of J.M. at the University of Vermont.

References

- Abe M, Ikegami T. 2005. Susceptibility of five species of thrips to different strains of the entomopathogenic fungus, *Beauveria bassiana*. *Applied Entomology and Zoology* 40: 667-674.
- Baranyovits F. 1953. Some aspects of the biology of armored scales. *Endavour* 12: 202-209.
- Bailey JA, Jeger MJ. 1992. *Colletotrichum: Biology, Pathology and Control*. CAB International.
- Butt TM, Goettel MS. 2000. Bioassays of entomogenous fungi. In: Navon A, Ascher KRS, editors. *Bioassays of Entomopathogenic Microbes and Nematodes*, pp. 141-195. CABI.




Figure 3. Percent mortality among different test insect species in fungal bioassays. ($\pm 95\%$ CI of the mean). Bars with the same letters are not significantly different (data for both concentrations combined) ($P < 0.05$) using the post-hoc multiple comparison Tukey-Kramer test. • Statistical significance was not estimated for this isolate because data for only one concentration was collected.

Byrne JM, Hausbeck MK, Hammerschmidt R. 1997. Conidial germination and apressorium formation of *Colletotrichum coccodes* on tomato foliage. *Plant Disease* 81: 715-718.

Dickman MB. 2000. Signal exchange during *Colletotrichum trifolii*-Alfalfa interactions. In: Prusky D, Freeman S, Dickman MB, editors. *Colletotrichum – Host specificity, pathology and host-pathogen interaction*, pp. 205-231. APS Press.

Dickman MB. 2004. Apressorium development in *Colletotrichum trifolii*. *Inoculum* 55: 13.

Doane EN, Parker BL, LaRosa S, Skinner M, Boone J, Pivot Y. 1998. *Mass rearing of western flower thrips, Frankliniella occidentalis (Thysanoptera: Thripidae) on beans*. Agricultural Experimental Station, University of Vermont Technical Note 4.

Frasen JJ. 1987. *Aschersonia aleyrodis* as a microbial control agent of greenhouse whitefly. *Ph.D. thesis*. Agricultural Univ. of Wageningen, The Netherlands. 167 .

Freeman S, Horowitz S, Sharon A. 2001. Pathogenic and nonpathogenic lifestyles in *Colletotrichum acutatum* from strawberry and other plants. *Phytopathology* 91: 986-992.

- Goettel MS, Inglis GD. 1997. Fungi: Hyphomycetes. In: Lacey LA, editor. *Manual of Techniques in Insect Pathology*, pp. 213-249. Academic Press.
- Guerber JC, Correll JC. 2001. Morphological description of *Glomerella acutata*, the teleomorph of *Colletotrichum acutatum*. *Mycologia* 93: 216-229.
- Horowitz S, Freeman S, Sharon A. 2002. Use of green fluorescent protein-transgenic strains to study pathogenic and non-pathogenic lifestyles in *Colletotrichum acutatum*. *Phytopathology* 92: 743-749.
- Horowitz S, Yarden O, Zveibil A, Freeman S. 2004. Development of a robust screening method for pathogenicity of *Colletotrichum* spp. on strawberry seedlings enabling forward genetic studies. *Plant Disease* 88: 845-851.
- Johnson WT, Lyon HH. 1988. *Insects that feed on trees and shrubs*, 2nd edition. Cornell University Press. 556 p.
- Khan A, Hsiang T. 2003. The infection process of *Colletotrichum graminicola* and relative aggressiveness on four turfgrass species. *Canadian Journal of Microbiology* 49: 433-442.
- Klingen I, Meadow R, Aandal T. 2002. Mortality of *Delia floralis*, *Galleria mellonella* and *Mamestra brassicae* treated with insect pathogenic hyphomycetous fungi. *Journal of Applied Entomology* 126: 231-237.
- Lambdin P, Lynch C, Grant J, Reardon R, Onken B, Rhea R. In: Onken B, Reardon R, editors. 2005. *Elongate hemlock scale and its natural enemies in the southern Appalachians, 3rd Symposium on Hemlock Woolly Adelgid in the Eastern United States*. Asheville, NC February 1-3 2005 pp. 145-154.
- Lardner R, Johnston PR, Plummer KM, Pearson MN. 1999. Morphological and molecular analysis of *Colletotrichum acutatum sensu lato*. *Mycological Research* 103: 275-285.
- Lu G, Cannon PF, Reid A, Simmons CM. 2004. Diversity and molecular relationships of endophytic *Colletotrichum* isolates from the Iwokrama Forest Reserve, Guyana. *Mycological Research* 108: 53-63.
- McClure MS. In: Onken B, Reardon R, Lashomb J, editors. 2002. The elongate hemlock scale, *Fiorinia externa* Ferris (Hemiptera: Diaspididae): A new look at an old nemesis. *Proceedings of the Hemlock Woolly Adelgid in the Eastern United States Symposium*. East Brunswick, NJ February 5-7 2005 pp. 248-253.
- Makowski RMD, Mortensen K. 1998. Latent infections and penetration of the bioherbicide agent *Colletotrichum gloeosporioides* f. sp. *malvae* in non-target field crops under controlled environmental conditions. *Mycological Research* 102: 1545-1552.
- Marcelino J, Gouli S, Giordano R, Gouli VV, Parker BL, Skinner M. 2009a. Fungi associated with a natural epizootic in *Fiorinia externa* Ferris (Hemiptera: Diaspididae) populations. *Journal of applied entomology* 133: 82-89.
- Marcelino J, Giordano R, Gouli S, Gouli VV, Parker BL, Skinner M, Cesnik R, TeBeest D. 2008. *Colletotrichum acutatum* var. *fiorinae* (teleomorph: *Glomerella acutata* var. *fiorinae* var. nov.) infection of a scale insect. *Mycologia* 100: 353-374.
- Marcelino J, Gouli S, Parker BL, Skinner M, Schwarzberg L, Giordano R. 2009b. Host plant associations of an entomopathogenic variety of the fungus, *Colletotrichum acutatum*, recovered from *Fiorinia externa*. *Journal of Insect Science*. In press.
- Memmott SD, Ha Y, Dickman MB. 2002. Proline reverses the abnormal phenotypes of *Colletotrichum trifolii* associated with expression of endogenous constitutively active Ras. *Applied and Environmental Microbiology* 68: 1647-1651.
- Negasi A, Parker BL, Brownbridge M. 1998. Screening and bioassay of entomopathogenic fungi for the control of silverleaf whitefly, *Bemisia argentifolii*. *Insect Science and its Applications* 18: 37-44.
- Mausel DL. 2002. Landscape patterns of hemlock decline in New England due to the introduced hemlock woolly adelgid. *Journal of Biogeography* 29: 1475-1487.
- Peres NA, Timmer LW, Adaskaveg JE, Correll JC. 2005. Lifestyles of *Colletotrichum acutatum*. *Plant Disease* 89: 784-796.
- Prusky D, Freeman S, Dickman MB. 2000. *Colletotrichum – Host specificity, Pathology and Host-Pathogen Interaction*. APS Press.
- Redman RS, Dunigan DD, Rodriguez RJ. 2001. Fungal symbiosis from mutualism to parasitism: who controls the outcome, host or invader? *New Phytologist* 151: 705-716.
- Rose M. 1990. Rearing and mass rearing. In: Rosen D, editor. *Armored scale insects. Their Biology, Natural Enemies and Control* [Series title: World crop pests, Vol. 4A: The Armored Scale Insects], pp. 357-365. Elsevier.
- SAS Institute 1990. *SAS/STAT user's guide*, 4th edition. Cary, NC. SAS Institute.
- SPSS Inc 2005. *SPSS Base 14.0 for Windows User's Guide*. Chicago, IL. SPSS Inc.
- Sreenivasaprasad S, Talhinhas P. 2005. Genotypic and phenotypic diversity in *Colletotrichum acutatum*, a cosmopolitan pathogen causing anthracnose on a wide range of hosts. *Molecular Plant Pathology* 6: 361-378.
- Tsrer L, Johnson D. 2000. *Colletotrichum coccodes* on potato. In: Prusky D, Freeman S, Dickman MB, editors. *Host specificity, pathology and host pathogen interaction of Colletotrichum*, pp. 362-373. APS Press.
- Xiao CL, MacKenzie SJ, Legard DE. 2004. Genetic and pathogenic analyses of *Colletotrichum gloeosporioides* isolates from strawberry and noncultivated hosts. *Phytopathology* 94: 446-453.
- Zulfiqar M, Brlansky RH, Timmer LW. 1996. Infection of flower and vegetative tissues of citrus by *Colletotrichum acutatum* and *C. gloeosporioides*. *Mycologia* 88(1): 121-128.