

Silkworm Thermal Biology: A Review of Heat Shock Response, Heat Shock Proteins and Heat Acclimation in the Domesticated Silkworm, Bombyx mori

Authors: Manjunatha, H. B., Rajesh, R. K., and Aparna, H. S.

Source: Journal of Insect Science, 10(204): 1-16

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.010.20401

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Silkworm thermal biology: A review of heat shock response, heat shock proteins and heat acclimation in the domesticated silkworm, Bombyx mori

H. B. Manjunatha^{1*}, R. K. Rajesh², and H. S. Aparna³

¹Department of Sericulture University of Mysore, Mysore 570 006, Karnataka, India ²Department of Biochemistry, Karnatak University, Dharwad 580 003, Karnataka, India ³Department of Biotechnology, University of Mysore, Mysore 570 006, Karnataka, India

Abstract

Heat shock proteins (HSPs) are known to play ecological and evolutionary roles in this postgenomic era. Recent research suggests that HSPs are implicated in cardiovascular biology and disease development, proliferation and regulation of cancer cells, cell death via apoptosis, and several other key cellular functions. These activities have generated great interest amongst cell and molecular biologists, and these biologists are keen to unravel other hitherto unknown potential functions of this group of proteins. Consequently, the biological significance of HSPs has led to cloning and characterization of genes encoding HSPs in many organisms including the silkworm, Bombyx mori L. (Lepidoptera: Bombycidae). However, most of the past investigations in *B. mori* were confined to expression of HSPs in tissues and cell lines, whereas information on their specific functional roles in biological, physiological, and molecular processes is scarce. Naturally occurring or domesticated polyvoltines (known to be the tropical race) are more resistant to high temperatures and diseases than bi- or univoltines (temperate races). The mechanism of ecological or evolutionary modification of HSPs during the course of domestication of *B. mori* - particularly in relation to thermotolerance in geographically distinct races/strains - is still unclear. In addition, the heat shock response, thermal acclimation, and hardening have not been studied extensively in *B. mori* compared to other organisms. Towards this, recent investigations on differential expression of HSPs at various stages of development, considering the concept of the whole organism, open ample scope to evaluate their biological and commercial importance in *B. mori* which has not been addressed in any of the representative organisms studied so far. Comparatively, heat shock response among different silkworm races/strains of poly-, bi-, and univoltines varies significantly and thermotolerance increases as the larval development proceeds. Hence, this being the first review in this area, an attempt has been made to collate all available information on the heat shock response, HSPs expression, associated genes, amino acid sequences, and acquired/unacquired thermotolerance. The aim is to present this as a valuable resource for addressing the gap in knowledge and understanding evolutionary significance of HSPs between domesticated (*B. mori*) and non-domesticated insects. It is believed that the information presented here will also help researchers/breeders to design appropriate strategies for developing novel strains for the tropics.

Keywords: acclimation, commercial traits, thermotolerance Correspondence: * manjunathahb@yahoo.com Editor: Brad Coates was editor of this paper. Received: 15 October 2009, Accepted: 9 April 2010 Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed. ISSN: 1536-2442 | Vol. 10, Number 204 Cite this paper as:

Manjunatha HB, Rajesh RK, Aparna HS. 2010. Silkworm thermal biology: A review of heat shock response, heat shock proteins and heat acclimation in the domesticated silkworm, *Bombyx mori. Journal of Insect Science* 10:204 available online: insectscience.org/10.204

Introduction

In recent years, the processes of heat shock responses and the role of heat shock proteins (HSPs) have not been confined merely to "molecular chaperons" (Parsell and Lindquist 1991), but spread over to determine their ecological and evolutionary role in the post genomic era (Sorensen and Loeschcke 2007). It is well known that both prokaryotic and eukaryotic cells respond to unfavourable environmental conditions bv increased synthesis of stress proteins such as HSPs. It is a universal phenomenon that most of the HSPs have conserved sequences ranging from bacteria to human, but certain features of the response do vary from organism to organism (Craig 1985). Unlike humans, plants and insects have a narrow range of tolerance to elevated temperatures and hence struggle to cope with these conditions. Consequently, the organisms that adapt over a period of time thrive whilst the others become extinct. For example, although Bombyx mori L (Lepidoptera: Bombycidae) is derived from a wild progenitor Bombyx mandarina (Arunkumar et al. 2006), it has lost its temperature-tolerance due to continuous domestication over 5000 years. As a result of such a long period of domestication silkworm races/strains diverged and the strains (polyvoltines) grown in tropical environments became resistant to high temperatures and diseases, while the strains reared in temperate conditions (bivoltines) remained susceptible.

Current research points to the fact that the loss of tolerance to environmental insults in *B. mori*, unlike *B. mandarina*, is due to prolonged domestication, which offers opportunities for systematic reinvestigation of this phenomenon while substantial diversity remains among various silkworm strains/races. Concerted efforts have been made during the past two decades resulting in the evolution of heat-tolerant silkworm strains, in response to conventional breeding strategies. This process has been successful, to some degree, in the tropical environment of the Indian subcontinent. Among several breeds developed, the only bivoltine silkworm breed that performed better all through the year over three decades in the field was NB4D2. Other temperate breeds were seasondependent. This observation poses the obvious questions regarding whether or not this adaptation is due to thermal acclimation, or if it is a process of hardening. A systematic investigation followed not only with reference to NB4D2, but also other geographically distinct silkworm strains such as Diazo (p50), Nistari etc. Consequently, many qualitatively and quantitatively superior productive/robust (thermotolerant) breeds were developed using Japanese commercial hybrids (temperate origin) as genetic resource material. Due to their low tolerance to the fluctuating environmental conditions in tropical climate they become unsuitable for growing year round (Nazia et al. 2005). Thus, the efforts made in the previous three decades were futile, and the spread and success of silkworm rearing was mainly due to the introduction of F_1 hybrids of native multivoltine as female parent (for resistance) and bivoltine as male parent (for high quality silk). Even now it is a challenging task to develop not only stressand disease-resistant strains, but also to provide high yielding silkworm strains with improved stress tolerance.

The cellular stress responses were described in *Drosophila melanogaster* for the first time by Ritossa (1962) and the term "heat shock protein" was introduced by Tissieres et al. (1974) as these proteins increased in synthesis

due to sudden increases in temperature. HSPs are identified based on their molecular mass ranges from 19 to 110 kDa in size and are broadly classified as large (major) HSPs and small HSPs (SmHSPs). The large HSPs are involved in major physiological processes such as cell division, transcription, protein membrane folding. transport, functions (Alique et al. 1994; Chen et al. 1996), and cytoprotective functions (Bakau and Horwich 1988; Chirico et al. 1988; Deshaies et al. 1988; Mizzen and Welch 1988; Palleros et al. 1991; Garrido et al. 2001; Kregel 2002). They can also form as large oligomeric complexes (Bentley et al. 1992; Leroux et al. 1997; Haslbeck et al. 1999) playing important roles in thermotolerance in mammalian cells (Landry et al. 1989), Drosophila (Landry and Huot 1995), house fly (Tiwari et al. 1997), and Lucilia cuprina (Tiwari et al. 1995) cells; but not in yeast cells (Nicholl and Quainlan SmHSPs bind specifically 1994). to cytoskeletal elements such as actin, and to intermediate filaments such as desmin, vimentin, and glial fibrillay acidic protein (Bennardine et al. 1992; Nicholl and Quainlan 1994). It has also been reported that SmHSPs modulate apoptosis (Arrigo 1998, 2005) and are involved in cell growth and differentiation (Mehlen et al. 1997). Recent approaches in genome wide identification of HSF (heat shock factor) – targeting genes provide novel insights into the complex metabolic reprogramming that occurs in cells in response to stress (Hahn et al. 2004).

Even to date, although some information is available, applicable data are insufficient to envisage the biological importance of HSPs in *B. mori*. To understand the complex phenomena governing silkworm thermal biology, integrative genomic, proteomic, and biotechnological approaches are required. Hence, the present review aims to collate research findings accrued over the last 16 years on the heat shock response, HSPs expression, associated genes, amino acid sequences, and thermotolerance in *B. mori*. It is believed that this review will help to uncover gaps in knowledge in this area that have not been documented to date in any other organisms. We also aim to explore the possibility of using heat shock proteins as molecular markers in conventional-molecular breeding for improvement of silkworm strains.

Heat shock and thermotolerance

The terms 'heat shock', 'acclimation', and 'hardening' are commonly used to describe the changes in an organism's living state caused by external environmental conditions and treatments (Bowler 2005; Loeschcke and Sorensen 2005; Lagerspetz 2006). The usage of these terms in the silkworm thermal biology literature has not been well-defined and requires systematic study to draw a line between them. The thermal tolerance of economically important organisms to environmental fluctuations attains significance in field-rearing conditions as performance in field/nature mainly depends on native varied adaptability to environmental conditions which is governed by molecular mechanisms of the cell. Notably, the polyvoltine silkworm strains exhibit better survivability over bivoltine strains, which might be due to their adaptation to thermal stress. For example, a polyvoltine strain, C. nichi, proved to be more tolerant than the bivoltine strain NB4D2 (Joy and Gopinathan Interestingly, in India 1995). among bivoltines, NB4D2 exhibited better tolerance to environmental fluctuation both in laboratory and field conditions compared to other newer bivoltine hybrids (CSR2, CSR4, NP2, KSO1, etc.), (Vasudha et al. 2006), an observation also confirmed by the recent

study of Firdose and Reddy (2009). However, the Chinese race, Feng, was the most tolerant strain followed by Japanese races, Kuo and J-09, while another Chinese race, C-54, was most susceptible (Hsieh et al. 1995). Since the range and significance of individual adaptive reactions differ in various species under different environmental conditions, the level of tolerance in elevated temperature varies between polyvoltine and bivoltine strains/races of B. mori. This diversity could be due to the fact that the races (species) living in hot and desert conditions for many thousands of years altered the molecularbiological mechanisms of adaptation, which facilitated their normal life-cycle even under extreme conditions (Evgen'ev et al. 2005).

Furthermore, the researchers' earlier study (Vasudha et al., 2006) demonstrated for the first time that the heat shock response in five bivoltine breeds (NB4D2, NP2, KSO1, CSR2, CSR4) varied during different and developmental stages. Of the five instars young silkworms, including first, second, and third instars, were relatively sensitive to any given heat shock temperature while older silkworms tolerated high temperatures for relatively longer periods of time. Similar observations were also reported by Joy and Gopinathan (1995). Notably, thermotolerance increased as larval development proceeded, sequentially in the order of first instar > second instar > third instar > fourth instar > fifth instar (Vasudha et al. 2006). The highest mortality (21%) was observed in the first instar and 100% survival in the fifth instar larvae of CSR4, while NP2 exhibited relatively lower mortality in the first instar and also 100% survival in the fifth instar. Comparatively, another lepidopteran model species. Manduca sexta. exhibited 100% survival at 42° C, but mortality increased as the heat shock temperature was raised to 48°

C in the fifth instar (Fittingoff and Riddiford 1990). In the case of pure mysore, a tropical multivoltine strain of B. mori in India, no mortality occurred at 42° C (for one hour) and 100% mortality was noticed at 46° C (Lohmann and Riddiford 1992). Interestingly, a few Japanese and Chinese silkworm strains also exhibited 100% mortality at 46° C after one hour of induced heat shock (Hsieh et al. 1995). In comparison with other insects, the threshold temperature that induced 100% mortality was 40° C in D. melanogater (Lindquist 1986); 45° C in different strains of silkworm, viz., NB4D2, NP2, KSO1, CSR2, and CSR4 (Vasudha et al. 2006); 46° C in Chinese, Japanese (Hsieh et al. 1995), and Indian silkworm strains of B. mori (pure mysore) (Lohmann and Riddiford 1992; Joy and Gopinathan 1995); 46° C in Musca domestica (Tiwari et al. 1997); 48° C in Manduca sexta (Fittingoff and Riddiford 1990); 48° C in Lucilia cuprina (Tiwari et al. 1995); and 50° C in Locusta migrotoria (Qin et al. 2003). However, no such information is available for wild silkmoths (Tasar -Antheraea mylitta; Muga – A. assamensis; Eri - Samia cynthia ricini). Hypothetically, all these studies imply that the heat shock treatment could be employed to determine the level of thermotolerance based on mortality (Loeschcke and Sorensen 2005), and they reveal that thermotolerance varies in different strains/races of silkworms and other insects.

The high thermotolerance in fifth instar larvae of *B. mori* reflects its adaptation to high temperatures that are encountered in the course of their normal life. However, in existing rearing practices, young silkworm larvae are recommended to be reared at high temperature (28° C) and high relative humidity (RH 80%); whereas older silkworm larvae are reared at lower temperature (24° C) and humidity (RH 65%). These practices thus leave ambiguity over the impact of heat shock on larval biological and commercial traits. In our estimation, the reason why farmers lose cocoon crops during the summer is likely due to elevated cell stress caused by high temperatures during rearing of young silkworm larvae.

Acclimation and hardening

As pointed out above, there is a need to differentiate acclimation from heat shock responses with special reference to B. mori. As per Lagerspetz (2006), there are three or more definitions of thermal acclimation; and an infinite number of possible combinations can be derived between acclimation. hardening, and heat shock responses (for details see Bowler 2005; Loeschcke and Lagerspetz Sorensen 2005; 2006). Importantly, the suggested definition of acclimation, from the Commission of Thermal Physiology of the International Union of Physiological Sciences, includes hardening and heat shock (Bowler 2005; Loeschcke and Sorensen 2005). Prosser (1955) used the term phenotypic acclimation for adaptive alterations exhibited by individual organisms. A very good example of developmental plasticity, which is known as seasonal polyphenism, is that of adult butterflies belonging to the genus Bicyclus that exhibit different wing patterns and variation in egg size representing alternating generations between a wet season form and a dry season form (Brakefield et al. 2007). Furthermore, species and populations adapt through natural operating on generations selection. of individuals and their hereditary property. In the light of these observations, we speculate that the polyvoltine and bi-/univoltines of *B*. mori likely adapted to dry (tropical polyvoltine type) and wet (temperate-bivoltine type) seasons, respectively, during the course of domestication. In B. mori, this adaptation did not exhibit any developmental plasticity in adult phenotype beyond changes in egg sizes, which are small in polyvoltine compared to bi- and/or univoltines and diapause in bi-/univoltine eggs.

More precisely, the term acclimation may be used to describe longer-term treatments, which may or may not be beneficial, but that is dependent on the exact conditions of treatment and the trait tested (Bowler 2005; Loeschcke and Sorensen 2005). Whether this phenomenon rightly fits with the treatment given for evaluation of thermotolerant (Robust) bivoltine breeds developed utilising Japanese thermotolerant hybrids as one of the parents needs to be analysed appropriately. However, comparison of the resultant robust bivoltine hybrids (CSR18, CSR19, HT1, etc.) subjected to thermal treatment revealed more tolerance to high temperature treatments than productive breeds affecting not only the survivability, but also other cocoon traits of the insect (Suresh et al. 1999). Unfortunately, performance of the thermotolerant the bivoltine breeds under fluctuated environment was very poor in the field. Thus, the question regarding the role of stress responses in thermal adaptation in nature still remains unanswered in B. mori as well as other organisms with different geographical origins. Additionally, some related questions, which were asked 10 years ago, are still valid and remain unresolved. Perhaps, cross-disciplinary approaches integrating proteomic, genomic, evolutionary, biological, and physiological methods might help to address these questions.

Proteome approach - *expression of HSP in tissues and whole organism*

The expression, regulation, localization, and functions of heat shock proteins have been studied extensively in different organisms.

The kinetics of HSP synthesis revealed distinct and reproducible differences between cell cultures of *B. mori* and the gypsy moth *Lymantria dispar*. While mulberry silkworm cells synthesize all HSP classes at temperature reaching 48° C, the gypsy moth cells synthesize no proteins at a 40° C and above and die under extreme conditions (Evgen'ev et al. 1987). In view of this, Evgen'ev et al. (1987) proposed to investigate whether high thermo-resistance was inherent only in the cultured cells, or if cells also behave in a similar way *in vivo*.

The differential expression of heat shock proteins in newly evolved bivoltine strains, NP2, KSO1, CSR2, and CSR4, was compared with that of the NB4D2 strain, which exhibited acclimation in the field over three decades (Vasudha et al. 2006). Interestingly, expression of only one set of HSPs with a molecular mass of 90 kDa in first, second (Figure 1), and third instars, and an 84 kDa HSP in the fourth instar was confirmed by Vasudha et al. (2006). Surprisingly, five different sets of 84, 62, 60, 47, and 33 kDa HSPs were also observed in the fifth instar larvae of NB4D2, KSO1, and CSR2 strains (Figure 2). Whereas, in the other two bivoltine strains expression of three HSPs (84, 47, and 33 kDa) in the NP2 and only two HSPs (84 and 47 kDa) in the CSR4 strains were reported (at 35 and 40° C for 2 h, Vasudha et al. 2006). In a multivoltine silkworm strain, pure mysore, 84, 70, 31, 30, and 29 kDa HSPs at 42° C (1 h, Lohmann and Riddiford 1992) and 83, 80, 74, 70, 68, 25, and 23 kDa at 48° C were found expressed in cells and organs (for 1 h, Evgen'ev et al. 1987). Between two multivoltines, 93, 46, and 28 kDa HSPs from pure mysore and 93, 70, 46, and 28 kDa HSPs from C. nichi were reported (Joy and Gopinathan 1995). This clearly indicated that different sets of HSPs were being expressed at



various heat shock temperatures, in different breeds of *B. mori* of which 90 and 84 kDa HSPs were ubiquitous (Table 1). In addition, expression of HSPs in different tissues varied depending on the stage of development, the temperature, and/or at which stage exposure was performed (Joy and Gopinathan 1995). Notably, concentration of HSPs and their distribution to specific sub-cellular sites is an important factor in acquisition of thermotolerance (Kampinga 1993).

Most of these studies were carried out following single dimensional electrophoresis (1-DE) and blotting techniques that made it possible to interrogate underlying mechanisms with greater certainty, albeit higher resolution of the proteins could not be achieved. To



mori larvae of CSR2 strain, heat shocked (HT) at 40° C and untreated control (C). Arrows indicate expression of 84, 60, 62, 47, 42, and 33 kDa heat shock proteins. M indicates molecular weight marker. High quality figures are available online.

Journal of Insect Science | <u>www.insectscience.org</u>

resolve these constraints in *B. mori*, advanced proteomic tools and techniques were for employed, which paved the way understanding differentiation and identification of different HSPs in the whole body of B. mori. A small number of protein spots were excised from the sample and two-dimensional separated bv gel electrophoresis (2-DE). After analysis of the resultant mass peptide finger prints with search engine Protein prospector, they were identified as the protein HSP70 (Rajesh et al. 2008). In addition, a comparative analysis of silk gland proteins in 2-DE gels of heat shock induced and normal silkworm larvae of NB4D2 revealed discrete differences with new and over expressed protein spots (Rajesh et al. 2009). Thus, application of advanced proteome techniques proved to be a promising approach in identification of different HSPs and opened new avenues to uncover more HSPs in *B. mori*.

Genomic approach - *hsp* genes and amino acid sequence of *B. mori*

The HSP family consists of ubiquitous phylogenetically proteins, which are conserved from bacteria to mammals and plants (Craig 1985). They have been divided into sub-families such as HSP110, HSP100, HSP90, HSP70, HSP60, HSP40, and HSP20 on the basis of their molecular weights (Nover and Scharf 1997; Gething 1998). Although, expression of HSPs has been reported from different silkworm strains (Table 1), only a few have been characterized in B. mori. Recently, Landais et al. (2001) characterized a cDNA encoding a 90 kDa HSP in *B. mori* and compared it with Spodoptera frugiperda (both lepidopteran insects). These two cDNAs encode 716 aa (amino acid) and 717 aa proteins in *B. mori* and *S. frugiperda*, respectively, with calculated molecular mass of 83 kDa which is similar to Drosophila.

Unlike in vertebrates, hsp90 does not contain introns and is a unique gene both in the B. mori and S. frugiperda genomes. Comparison of aa sequences of *B. mori* and *S. frugiperda* with that of *D. melanogaster*, Homo sapiens, and S. cerevisiae revealed a high percentage of similarity and phylogenetic relationships (for details see Landais et al. 2001). Apparently, extensive study is required to determine their expression at different developmental stages of different silkworm strains as the HSP90 expression is found rather in early instars than late instars (Vasudha et al. 2006) and expression of some hsp genes changes during development (Craig 1985). In D. melanogaster, hsc70-4 (constitutive *hsp* gene family) was expressed at a high level in embryos, larvae, and adults. whereas the *hsc70-1* and *hsc70-2* expression

was highest in adults but not detected in larvae. The *hsc70-1* was expressed at a low level while no expression of *hsc70-2* was observed in the embryo. In *Chironomus tentans*, *hsc70* expression was evident at all developmental stages but slightly lower in the embryo than older stages (Karouna-Renier et al. 2003).

Small heat shock proteins (smHSPs or sHSPs) belong to a family of genes that are seemingly less conserved compared with those of major *hsp* gene families, but occur ubiquitously in a variety of organisms. These proteins are involved in apoptosis as well as protection against heat stress (Arrigo 2005; Feder and Hofmann 1999). In *B. mori* (strain p50) six genes encoding sHSP19.9, sHSP20.1, sHSP20.4, sHSP20.8, sHSP21.4, and

HSPs	Silkworm race/breeds/strains	Tissues	References	
93kDa	C.Nichi, Pure mysore, NB4D2	Heamolymph	Joy Omana & Gopinathan, 1995	
3kDa	C.Nichi, NB4D2	Fat bodies	Joy Omana & Gopinathan, 1995	
90kDa	NB4D2, KSO1, NP2, CSR2, CSR4	Whole larvae	Vasudha et al., 2006	
		(I, II & III instars)		
90kDa	Strain Y;C	Silkgland	Zhang et al., 2006	
34kDa	Pure mysore	Epidermis	Lohmann & Riddiford, 1992	
84kDa	NB4D2, KSO1, NP2, CSR2, CSR4	Whole larvae	Vasudha et al., 2006	
		(IV and V instars)		
33kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	
32kDa	C.Nichi, NB4D2	Fat bodies	Joy Omana & Gopinathan, 1995	
30kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	
74kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	
70kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	
70kDa	Pure mysore	Epidermis	Lohmann & Riddiford, 1992	
70kDa	Chinese and Japanese races	Heamocytes	Hsieh et al., 1995	
70kDa	Chinese and Japanese races	Fat bodies	Hsieh et al., 1995	
70kDa	NB4D2, NP2, KSOI	Whole larvae	Vasudha et al., 2006	
70kDa	C.Nichi	Heamocytes	Joy Omana & Gopinathan, 1995	
70kDa	C.Nichi, NB4D2	Fat bodies	Joy Omana & Gopinathan, 1995	
70kDa	Strain Y;C	Silkgland	Zhang et al., 2006	
68kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	
62kDa	NB4D2, KSO1	Whole larvae	Vasudha et al., 2006	
		(V instar)		
60kDa	NB4D2, KSO1, CSR2	Whole larvae	Vasudha et al., 2006	
		(V instar)		
60kDa	Strain Y;C	Silkgland	Zhang et al., 2006	
47kDa	NB4D2, KSO1, NP2, CSR2, CSR4	Whole larvae	Vasudha et al., 2006	
		(V instar)		
33kDa	NB4D2, KSO1, NP2	Whole larvae	Vasudha et al., 2006	
		(V instar)		
31kDa	Pure mysore	Epidermis	Lohmann & Riddiford, 1992	
30kDa	Pure mysore	Epidermis	Lohmann& Riddiford, 1992	
29kDa	Pure mysore	Epidermis	Lohmann & Riddiford, 1992	
26kDa	Pure mysore	Epidermis	Lohmann & Riddiford, 1992	
25kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	
23kDa	Pure mysore	Cell line	Evgen'ev et al., 1987	

sHSP23.7 were reported (Sakano et al. 2006) although their biological and commercial roles remain unknown. The deduced amino acid residues of these sHSPs (Table 2) are quite similar to each other. CLUSTALW multiple alignments indicated 82, 80, and 80% identity between Pia25 and sHSP20.8, sHSP20.8, sHSP20.4, sHSP20.4, and sHSP19.9, respectively. Besides the α -crystallin domain, the N-terminal XXLXDQXFG motifs are commonly conserved in the sequences of these HSPs (Sakano et al. 2006). Further, transcriptase-polymerase reverse chain reaction (RT-PCR) analysis showed no difference in expression levels of *smHSP* genes in different organs (Sakano et al. 2006), but indicated an increased amount of transcripts following heat shock in B. mori strains p50 (Sakano et al. 2006), Nistari and NB4D2 (Velu et al. 2008), which was found to be strain specific. BmHSPs (B. mori HSPs) with other organisms was computed using available data in National Center for Biotechnology Information (NCBI) data bank (http://www.ncbi.nlm.nih.gov) and presented in Table 3.

Can HSP help in acquired thermotolerance?

The response to heat shock is an integral part of survival in the environment, as it is for

domestic silkworms that are derived from geographically distinct regions but reared under standard conditions. Hence, a new strategy was adopted wherein the whole egg (Manjunatha et al. 2005) and larvae (Vasudha et al. 2006) of B. mori were subjected to heat shock at various temperatures during different developmental stages, to determine the importance of HSPs in acquired thermotolerance. These studies revealed that different sets of HSPs expressed in different developmental stages have a profound influence not only on the performance of larvae (rated in terms of mortality), but also to complete life cycle under natural environmental conditions. The well-defined role of HSPs (expressed either individually or collectively) in acquired thermotolerance in the silkworm and other insects is not known. In order to derive more accurate and novel hypotheses, the expression of heat shock proteins should be correlated with currently available information on the tolerance of silkworm strains reared in tropical environments.

Role of HSP in relation to commercial traits

To date, the greatest emphasis has been given to HSP70 and HSP90 as molecular chaperons that help organisms to cope with stresses of

mbyx mori heat sh	ock protei	ns (BmHSPs) accession	numbers, protein ID	s and their deduced a	
HSI	Ps	Accession no.	Protein ID	Total no. of	
			no.	amino acids	
Hsp	90	AB060275	BAB41209	716	
Hsp	70	DQ311189	ABD36134.1	676	
Chaper (Hspe	ronin 60)	NM_001079879	NP_001073348	545	
Hsp	40	AB206400	BAD90846.1	351	
Hsp2	3.7	AB195973	BAD74198.1	209	
Hsp2	1.4	AB195972	BAD74197.1	187	
Hsp2	0.8	AF315317	AAG30944.1	186	
Hsp20	.8A	AF315319	AAG30946	186	
Hsp2	0.4	AF315318	AAG30945.2	181	
Hsp2	0.1	AB195971	BAD74196.1	178	
Hspl	9.9	AB195970	BAD74195.1	177	
Hsp		DQ443370.1	ABF51459	198	
α-cryst	allin I	AF309497.1	AAK06407	122	
α-cryst	allin2	AF309499.1	AAK06409	90	

internal and external nature. Recent approaches not only revealed the importance of HSP90 in normal growth and development of eukaryotes, and parasite (Plasmodium falciparum) growth in human erythrocytes (Banumathy et al. 2003), but also elucidated the relationship between HSPs and life history focusing on the ecological and traits evolutionary relevance (Sorensen et al. 2003; Sorensen and Loeschcke 2007). Concomitantly, the relationship between heat shock, HSPs expression, and commercial traits was studied in great detail in the case of B. mori (Vasudha et al. 2006). Notably, an increased cocoon weight of 17.52 vs. 13.48%, and increase in shell weight of 19.44 vs. 13.45% in NB4D2 over its control was observed following heat shock at 35 and 40° C, respectively. Concurrently, CSR2 also exhibited a 13.11 vs. 6.44% increase in cocoon weight and 16.26 vs. 5.03% increase in shell weight at 35 and 40° C heat shock over their respective controls. The increased cocoon and shell weight observed in heat shock induced bivoltine silkworm strains compared to controls would be due to expression of HSPs at larval stage. While Joy and Gopinathan (1995) did not observe any heat shock effects on commercial traits, Lohmann and Riddiford (1992) reported that of the nine animals heat shocked at 44° C for 1 h, only 5 resumed feeding, while 3 spun cocoons. Commercial traits of these animals were not evaluated and compared with that of controls. Consequently, as a novel strategy. heat shocked larvae (whole organisms) were allowed to grow under natural environmental conditions and they spun better quality cocoons than the non heat shocked larvae reared in natural environmental conditions (Vasudha et al. 2006). These investigations highlighted the fact that knowledge obtained from model organisms under normal laboratory conditions does not always reflect what happens out in the field, where conditions are continuously changing and hostile. Interestingly, unpredictably the increased cocoon weight and shell weight over control, reflects the positive correlation between heat shock responses and silk protein content in the cocoon. Abramova et al. (1991) reported suppression of fibroin synthesis in the silk gland following heat shock, but recently Zhang et al. (2006) identified HSP90, HSP70, and HSP60 in the silk glands of B. mori, offering the opportunity for further systematic investigation in different breeds of silkworm. None of the larvae recovered from heat shock at 45° C (Vasudha et al. 2006) and 46° C (Lohmann and Riddiford 1992), were able to spin cocoons. However, the observed differences between cocoon weight, shell weight, and shell ratio among various silkworm will require strains further investigations to determine species-specific responses to heat shock. Altogether, these observations clearly indicate that mild heat shock between 35 and 40° C for 2 h facilitates bivoltine silkworm larvae to respond and fluctuating overcome the natural environmental conditions in succeeding instars. The practical application of this phenomenon will need to be explored systematically positively and (using multivoltine and bivoltine silkworm strains) in laboratory and field conditions in order to achieve stabilized sericulture farming in tropical countries like India.

Hypothetical view on the heat shock and acclimation in the Silkworm, *B. mori*

Based on published scientific reports, and our own experimental observations (Manjunatha et al. 2005; Vasudha et al. 2006; Rajesh et al. 2008, 2009), it is suggested that silkworm researchers should more thoroughly delineate heat shock and thermal acclimation phenomena in *B. mori*. These studies would aid our understanding of the silkworm thermal biology much better until bio-molecular evidences further substantiate its relevance.

First, the heat shock treatment given to B. mori larvae is to determine built in thermotolerance based on mortality and differential expression of heat shock proteins. HSP expression patterns dictate different levels of thermotolerance in individual silkworm strains at varied heat shock temperatures. The hypothetical interpretation of thermotolerance (Figure 3) refers to the state of silkworm larvae that can perform better and withstand threshold heat shock temperatures for a fixed (short) period in a particular stage or generation. Obviously, this physical state is supported by expression of one or many HSPs in a given generation, but whether the same rate of expression appears under deleterious environmental condition in subsequent generations remain unclear and will require further investigation. Thermotolerance varies among silkworm races/strains, and it is categorized by tolerance levels 1, 2, 3, and 4 (Table 3). At level-1, any

polyvoltine (many generations per year, nondiapause type) or bivoltine (two generations per year, diapause at egg stage) or univoltine (one generation per year, diapause at egg stage) races/strains exhibiting better performance through several generations under fluctuating environmental conditions adapt completely. However, the survival rate varies in accordance with genomic organisation and ultimately leads to different levels of tolerance as levels 2, 3, and 4 exhibit 75%. 50%. and 25% survivability, respectively, compared to the insects that ones which did not respond to acclimation or exhibit adaptability in the natural environment. For instance, the pure mysore and nistari strains (native polyvoltines) have inbuilt adaptability to high temperatures (level 1), whereas the NB4D2 bivoltine strain (known as temperate race) exhibited better acclimation (level 2) during continuous rearing in the field round the year for three decades. As a consequence, NB4D2 acquired better adaptability than other bivoltine race/breeds/stains. Comparatively, of the new bivoltine strains, CSR2 showed better





response to heat shock (Manjunatha et al. 2005; Vasudha et al. 2006) and performance in the field (level 3) than other strains (level 4) (Nazia et al. 2005). Furthermore, between polyvoltine and bivoltine strains, the Nistari exhibited higher expression of HSP70 and HSP40 genes than the NB4D2 strain (Velu et al. 2008).

Second, thermal acclimation is a longer-term treatment (ranging from days to weeks), which results in increased resistance to temperature. It is not likely to be related to HSP production as acclimation occurs within the normal temperature range experienced, and its effect lasts as long as the new acclimation conditions persist (for proposed definitions refer to Bowler 2005 and Loeschcke and Sorensen 2005). To define it more precisely in *B. mori*, the information available is insufficient and warrants further investigation in poly-, bi-, and/or univoltine strains.

Third, expression of nearly 18 different HSPs has been reported so far from various tissues and whole body of different *B. mori* strains (Table 1). Interestingly, although differential expression of HSPs is noticed during different developmental stages, their role in altering biological, physiological, and commercial traits remains enigmatic.

documented here on *B. mori* would be useful for comparative genomic and proteomic research for other lepidopterans and other organisms. The genome wide analysis of hsp genes (Hahn et al. 2004) and their regulatory factors provide novel insights into the complex metabolic reprogramming that occurs within cells in response to stress. The domesticated silkworm *B. mori*, together with its wild progenitor, Bombyx mandarina, and non-mulberry silkworms (Tasar - Antheraea mylitta; Muga – A. assamensis; Eri - Samia cynthia ricini), which are reared in nature, open ample scope to investigate the ecological and evolutionary modification of HSPs and identify the candidate gene(s). The individual or collective role of HSPs in relation to biological, commercial, physiological, and immunological features among different silkworm races/breeds/strains (including nonmulberry silkworms) is important for understanding the factors that govern thermotolerance and acclimation in insects. Knowledge of HSPs and their use as molecular markers would facilitate conventional breeders to select better parents, with a reduction in laborious crosses for development of suitable silkworm strains, important for tropical countries under silkworm race improvement programmes.

Organisms	HSP90	HSP70	HSP23.7	HSP21.4	HSP20.8	HSP20.4	HSP20.I	HSP19.9	α - crystallin
A. thaliana	70.69	75.19	-	-	-	-	-	27.78	-
C. elegans	77.73	78.36	42.17	50.75	39.78	40.45	35.83	27.78	40.58
D. melanogaster	83.8	89.55	48.52	39.81	49.43	62.77	56.77	-	62.67
E. coli	37.67	52.92	-	-	-	-	-	-	-
H. sapiens	82.67	80.6	42.57	41.38	50.33	55.06	48.05	46.75	55.07
M. musculus	81.88	80.6	42.57	41.38	50.33	50	48.05	60.29	55.07
R. norvegicus	81.54	80.6	42.57	41.38	50 .00	47.75	48.7	-	54.41
S. cerevisiae	62.06	76.49	-	-	-	-	-	-	-

In conclusion, the research resources

Acknowledgements

This paper was made possible with the support of project No. F.30-210/2004, UGC, New Delhi. The authors wish to thank Dr. Ramnath Elaswarapu, Science Leader – Functional Genomics, LGC Ltd. (Queen's Road, Teddington, TW11 0LY) for critically editing the manuscript coupled with suggestions.

References

Abramova IYU, Ulmasov KHA, Akopov SB, Karaev KK, Babaeva AKH, Evgene'v MB. 1991. Dynamics of the synthesis of fibroin and heat shock proteins in different organs and at different stages of silkworm, *Bombyx mori* ontogenesis. *Prikladnya Biokhimiya Mikrobiologiya* 27: 147-156.

Alique R, Akhavan NH, Russell P. 1994. A role for hsp90 in cell cycle control: Weel tyrosine kinase activity requires interaction with hsp90. *EMBO Journal* 13: 6099-6160.

Arrigo AP. 1998. Small heat shock proteins: Chaperones that act as regulators of intracellular redox state and programmed cell death. *Journal of Biological Chemistry* 379: 19-26.

Arrigo AP. 2005. In search of the molecular mechanism by which small stress proteins counteract apoptosis during cellular differentiation. *Journal of Cellular Biochemistry* 94: 241-246.

Arunkumar KP, Metta M, Nagaraju J. 2006. Molecular phylogeny of silkmoths reveals the origin of domesticated silkmoth, *Bombyx mori* from Chinese *Bombyx mandarina* and paternal inheritance of *Antheraea proylei* mitochondrial DNA. *Molecular Phylogenetics and Evolution* 40: 419-427. Bakau B, Horwich AL. 1988. The Hsp70 and Hsp 60 chaperone machines. *Cell* 92: 351-366.

Banumathy G, Singh V, Pavithra SR, Utpal Tatu. 2003. Heat shock protein 90 function is essential for *Plasmodium falciparum* growth in human erythrocytes. *Journal of Biological Chemistry* 278: 18336-18345.

Bennardine F, Wrzosek A, Chiesi M. 1992. α , β -crystallin in cardiac tissue. Association with actin and desmin filaments. *Circulation Research* 71: 288-294.

Bentley NJ, Fitch IT, Tuite MF. 1992. The small heat shock protein hsp26 of *Saccharomyces cerevisiae* assembles into a high molecular weight aggregate. *Yeast* 8: 95-106.

Bowler K. 2005. Acclimation, heat shock and hardening. *Journal of Thermal Biology* 30: 125-130.

Brakefield PM, Pijpe J, Zwaan BJ. 2007. Developmental plasticity and acclimation both contribute to adaptive responses to alternating seasons of plenty and of stress in *Bicyclus* butterflies. *Journal of Bioscience* 32: 465-475.

Chen CF, Chen Y, Dai K, Chen PL, Riley DL, Lee WH. 1996. A new member of the hsp90 family of molecular chaperone interacts with the retinoblastoma protein during mitosis and after heat shock. *Molecular Cell Biology* 16: 4691-4699.

Chirico WJ, Waters MG, Blobel G. 1988. 70K heat shock related proteins stimulate protein translocation into microsomes. *Nature* 332: 805-810.

Craig CA. 1985. The heat shock response. *CRC Critical Reviews in Biochemistry* 18: 239-280.

Deshaies RJ, Koch BD, Werner-Washburne M, Craig EA, Schekman R. 1988. A subfamily of stress proteins facilitates translocation of secretary and mitochondria precursor polypeptides. *Nature* 332: 800-805.

Evgen'ev MB, Garbuz DG, Zatsepina OG. 2005. Heat shock proteins: Functions and role in adaptation to hyperthermia. *Russian Journal of Developmental Biology* 36: 218-224.

Evgen'ev MB, Sheinker VS, Levin AV, Braude ZTY, Titarenko EA, Shuppe NG, Karaev KK, Ul'masov KA, Zolotareva TY. 1987. Molecular mechanisms of adaptation to hyperthermia in higher organisms. (I). Synthesis of heat-shock proteins in cell cultures of different species of silkworms and in caterpillars. *Molecular Biology* 21: 410-419.

Feder ME, Hofmann GE. 1999. Heat shock proteins, molecular chaperons and the stress response: evolutionary and ecological physiology. *Annual Review of Physiology* 61: 243-282.

Firdose AM, Reddy YS. 2009. Racial features in the heat tolerance limits in the silkworm, *Bombyx mori. Acta Entomologica Sinica* 52: 478-486.

Fittingoff CM, Riddiford LM. 1990. Heat sensitivity and protein synthesis during heatshock in the tobacco hornworm, *Manduca Sexta. Journal of Comparative Physiology and Biochemistry* 160: 349-356.

Gething M. 1998. *Guidebook to molecular chaperons and protein folding catalyses*. Oxford University Press, Oxford.

Garrido C, Gurbuxani S, Ravagnan L, Kroemer G. 2001. Heat shock proteins: endogenous modulators of apoptotic cell death. *Biochemical Biophysical Research Communications* 286: 433-442.

Hahn JS, Hu Z, Thiele DJ, Iyer VR. 2004. Genome-Wide analysis of the biology of stress responses through Heat Shock Transcription Factor. *Molecular and Cellular Biology* 24: 5249-5256.

Haslbeck M, Walke S, Stromer T, Ehrnsperger M, White HE, Chen S, Saibil HR, Buchner J. 1999. Hsp26: A temperatureregulated chaperon. *EMBO Journal* 18: 6744-6751.

Hsieh FK, Yu SJ, Su SY, Peng SJ. 1995. Studies on the thermotolerance of the Silkworm, *Bombyx mori*. *Chinese Journal of Entomology* 15: 91-101.

Joy O, Gopinathan KP. 1995. Heat shock response in mulberry silkworm races with different thermotolerance. *Journal of Bioscience* 20: 499-513.

Kampinga HH. 1993. Thermotolerance in mammalian cells: Protein denaturation and aggregation and stress proteins. *Journal of Cell Science* 104: 11-17.

Karouna-Renier NK, Yang WJ, Rao KR. 2003. Cloning and characterization of a 70 kDa heat shock cognate gene (HSF70) from two species of *Chironomus*. *Insect Molecular Biology* 12: 19-26.

Kregel KC. 2002. Heat shock proteins: modifying factors in physiological stress responses and acquired thermotolerance. *Journal of Applied Physiology* 92: 2177-2186.

Landais I, Pommet J, Mita K, Nohata J, Gimenez S, Fournier P, Devauchelle G, Duonor-Cerutti M, Ogliastro M. 2001. Characterization of the cDNA encoding the 90 kDa heat shock protein in the Lepidoptera

Bombyx mori and *Spodoptera frugiperda*. *Gene* 271: 223-231.

Landry J, Huot J. 1995. Modulation of actin dynamics during stress and physiological stimulation by a signaling pathway involving p38 MAP kinase and heat shock ptotein27. *Biochemistry and Cell Biology-Biochimie et Biologie Cellulaire* 73: 703-707.

Landry J, Chretien P, Lambert H, Hickey E, Weber LA. 1989. Heat shock resistance conferred by expression of the human hsp27 gene in rodent cells. *Journal of Cell Biology* 109: 7-15.

Lagerspetz KYH. 2006. What is thermal acclimation? *Journal of Thermal Biology* 31: 332-336.

Leroux M, Melki R, Gordan B, Batelier G, Candido EP. 1997. Structure-function studies on small heat shock protein oligomeric assembly and interaction with unfolded polypeptides. *Journal of Biological Chemistry* 272: 24646-24656.

Lindquist S. 1986. The heat shock response. Annual Review of Biochemistry 55: 1151-1191.

Loeschcke V, Sorensen JG. 2005. Acclimation, heat shock and hardening – a response from evolutionary biology. *Journal of Thermal Biology* 30: 255-257.

Lohmann CMF, Riddiford LM. 1992. The heat shock response and heat sensitivity of *Bombyx mori. Sericologia* 32: 533-537.

Manjunatha HB, Zamood A, Vasudha BC, Aparna HS. 2005. Heat shock response and analysis of egg proteins in new bivoltine strains of *Bombyx mori. Sericologia* 45: 403-408. Mehlen P, Mehlen A, Godet J, Arrigo AP. 1997. hsp27 as a switch between differentiation and apoptosis in murine embryonic stem cells. *Journal of Biological Chemistry* 72: 31657-31665.

Mizzen LA, Welch WJ. 1988. Characterization of the thermotolerant cell. I. Effects on protein synthesis activity and the regulation of heat shock protein 70 expression. *Journal of Cell Biology* 106: 1105-1116.

Nazia C, Mal Reddy N, Basavaraja HK, Dandin SB. 2005. Alteration in qualitative and quantitative chararacters in bivoltine silkworm, *Bombyx mori* due to the thermal stress. *Indian Journal of Sericulture* 44: 100-106.

Nicholl ID, Quainlan RA. 1994. Chaperon activity of α -crystallins modulates intermediate filament assembly. *EMBO Journal* 13: 945-953.

Nover L, Scharf KD. 1997. Heat stress proteins and transcription factors. *Cellular and Molecular Life Sciences* 53: 80-103.

Palleros DR, Welch WJ, Fink AL. 1991. Interaction of hsp 70 with unfolded proteins: effects of temperature and nucleotides on the kinetics of binding. *Proceedings of National Academy of Sciences USA* 88: 5719-5723.

Parsell DA, Lindquist S. 1991. The function of heat shock proteins in stress tolerance: Degradation and reactivation of damaged proteins. *Annual Review of Biochemistry* 60: 321-347.

Prosser CL. 1955. Physiological variation in animals. *Biological Review* 30: 229-262.

Qin W, Tyshenko MG, Wu BS, Walker VK, Robertson RM. 2003. Cloning and

characterization of a member of the hsp 70 gene family from *Locusta migratoria*, a highly thermotolerant insect. *Cell Stress and Chaperones* 8: 39-53.

Rajesh RK, Aparna HS, Manjunatha HB. 2008. Molecular identification and analysis of heat shock proteins in *Bombyx mori*. *Scenario of Seri-biotechnological Research in India*, 183-191.

Rajesh RK, Aparna HS, Manjunatha HB. 2009. Comparative analysis of silk gland proteins of both heat shocked and normal silkworm larvae of NB4D2 strain by 2-DE. *International Journal of Applied Agricultural Research* 4: 125-130.

Ritossa F. 1962. New puffing pattern induced by temperature shock and DNP in *Drosophila*. *Experientia* 18: 571-573.

Sakano D, Li B, Xia Q, Yamamto K, Hiroshi F, Aso Y. 2006. Genes encoding small heat shock proteins of the silkworm *Bombyx mori*. *Bioscience Biotechnology and Biochemistry* 70: 2443-2450.

Sorensen JG, Loeschcke V. 2007. Studying stress responses in the post genomic era: its ecological and evolutionary role. *Journal of Bioscience* 32: 447-456.

Suresh KN, Kishor KCM, Basavaraja HK, Mal Reddy N, Ramesh Babu M, Datta RK. 1999. Comparative performance of robust and productive bivoltine hybrids of *Bombyx mori* L. under high temperature conditions. *Sericologia* 39: 567-571.

Tissieres A, Mitchell HK, Tracey UM. 1974. Protein synthesis in salivary glands of *D. melanogaster* relation to chromosome puffs. *Journal of Molecular Biology* 84: 389-398. Tiwari PK, Joshi A, Mohan DRK. 1997. Thermotolerance and heat shock response in *Musca domestica*. *Current Science* 72: 501-506.

Tiwari PK, Mohan DRK, Joshi A. 1995. Developmental study of thermotolerance and the heat shock response in *Lucilia cuprina* (Weidemann). *Journal of Bioscience* 20: 341-354.

Vasudha BC, Aparna HS, Manjunatha HB. 2006. Impact of heat shock on heat shock proteins expression, biological and commercial traits of *Bombyx mori. Insect Science* 13: 243-250.

Velu D, Ponnuvel KM, Qadri, HSM. 2008. Expression of the heat shock protein genes in response to thermal stress in the silkworm *Bombyx mori. International Journal of Industrial Entomology* 16: 21-27.

Zhang P, Aso Y, Yamamato K, Banno Y, Wang Y, Tsuchida K, Kawaguchi Y, Fujii H. 2006. Proteome analysis of silk gland proteins from the silkworm, *Bombyx mori. Proteomics* 6: 2586-2599.