

## **Differential Consumption of Four Aphid Species by Four Lady Beetle Species**

Authors: Finlayson, Christy, Alyokhin, Andrei, Gross, Serena, and Porter, Erin

Source: Journal of Insect Science, 10(31) : 1-10

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.010.3101>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Differential consumption of four aphid species by four lady beetle species

Christy Finlayson<sup>a</sup>, Andrei Alyokhin, Serena Gross and Erin Porter

The University of Maine School of Biology and Ecology, 5722 Deering Hall, Room 202, Orono, ME 04469-5722

### Abstract

The acceptability of four different aphid species *Macrosiphum albifrons* (Essig), *Macrosiphum euphorbiae* (Thomas), *Macrosiphum pseudorosae* Patch, and *Myzus persicae* (Sulzer) (Hemiptera: Aphididae), as prey for four lady beetle species, one native species *Coccinella trifasciata* L, and three non-native *Coccinella septempunctata* L, *Harmonia axyridis* Pallas, *Propylea quatuordecimpunctata* L (Coleoptera: Coccinellidae) were tested in the laboratory. The relative field abundance of adults of the same lady beetle species on host vegetation, *Lupinus polyphyllus* Lindley (Fabales: Fabaceae), *Solanum tuberosum* L (Solanales: Solanaceae), and *Rosa multiflora* Thunberg (Rosales: Rosaceae), both with and without aphids present was also observed. In the laboratory, *H. axyridis* generally consumed the most aphids, while *P. quatuordecimpunctata* consumed the fewest. The exception was *P. quatuordecimpunctata*, which consumed a greater number of *M. albifrons* nymphs, and *C. trifasciata*, which consumed a greater number of *M. albifrons* nymphs and adults, compared with the other two beetle species. Lady beetles consumed fewer *M. albifrons* compared with the other three aphid species, likely because of deterrent compounds sequestered by this species from its host plant. In the field, *P. quatuordecimpunctata* was the most abundant species found on *L. polyphyllus* and *S. tuberosum*.

**Keywords:** predation, biological control, competition, non-native species

**Correspondence:** <sup>a</sup> christy.finlayson@umit.maine.edu

**Associate Editor:** J.P. Michaud was editor of this paper.

**Received:** 9 November 2008, **Accepted:** 9 January 2009

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 10, Number 31

#### Cite this paper as:

Finlayson C, Alyokhin A, Gross S, Porter E. 2010. Differential consumption of four aphid species by four lady beetle species. *Journal of Insect Science* 10:31 available online: [insectscience.org/10.31](http://insectscience.org/10.31)

## Introduction

Lady beetles (Coleoptera: Coccinellidae) are known to be voracious predators of plant pests such as aphids (Hemiptera: Aphididae) (Hodek 1973; Gordon 1985). It is often assumed that aphidophagous lady beetles are highly polyphagous, consuming most (if not all) aphid species that they encounter (Pedigo and Rice 2006). However, there is evidence that not every aphid species is equally suitable for every lady beetle species (Obrycki and Orr 1990; Phoofolo and Obrycki 1997; Kalushkov 1998; Michaud 2000; Kalushkov and Hodek 2004; Mignault et al. 2006). For example, Michaud (2000) conducted choice tests with seven lady beetle species and two aphid species, *Toxoptera citricida* and *Aphis spiraecola*. Although all lady beetles tested consumed both aphid species, four species *Coccinella septempunctata*, *Coleomegilla maculata fuscilabris*, *Coelophora inaequalis*, and *Olla v-nigrum*, were not able to complete their developmental cycle with either aphid species. Depending on the aphid species consumed and the addition of supplements (pollen) to the diet, the other three species, *Hippodamia convergens*, *Cycloneda sanguinea*, and *Harmonia axyridis*, varied considerably in the number of eggs laid, egg viability, larval development time, and adult weight.

Lady beetles are commonly released to combat a diverse range of pests (Gordon 1985; Koch 2003), despite the fact that little is known about specific prey preferences of different species. The success of such pest control measures depends, in part, upon the willingness of the lady beetles to consume the pest in question. Releases of non-native species may supplement pest control by native species when their prey species do not overlap

or when prey is plentiful. Native lady beetle abundance, however, may be reduced through competition with non-native species with overlapping prey preferences. Additionally, non-native lady beetles may alter aphid community structure. Determining differences in prey consumption by different lady beetle species may provide insight into changes that occur in systems where non-native species become established. In the laboratory, one native and three non-native lady beetle species were provided four different species of aphid prey and their consumption was recorded. To determine if any differences documented in the laboratory were reflected in the field, lady beetle species were observed for their association with these aphids under field conditions.

## Materials and Methods

### Study species

The four lady beetle species chosen for this study are aphidophagous (Gordon 1985) and abundant in Maine in the same habitats (Finlayson et al. 2008). The native lady beetle species used was *Coccinella trifasciata perplexa* Mulsant, which is found from Labrador south to New Jersey and west to California and Alaska (Gordon 1985). The non-native lady beetle species used were *Coccinella septempunctata* L., *Harmonia axyridis* Pallas, and *Propylea quatuordecimpunctata* L. These three species are Palearctic in origin and were intentionally and inadvertently introduced in North America. *C. septempunctata* has been established in North America since 1973 (Angalet and Jacques 1975), *H. axyridis* since 1988 (Chapin and Brou 1991; Tedders and Schaefer 1994), and *P. quatuordecimpunctata* since 1968 (Wheeler 1990).

Four aphid species that are abundant and readily available in the region were chosen to serve as the prey for the selected lady beetle species. The potato aphid, *Macrosiphum euphorbiae* (Thomas), feeds on over 200 plant species (Blackman and Eastop 1984). The green peach aphid, *Myzus persicae* (Sulzer), feeds on over 40 different plant families (Blackman and Eastop 1984). The hosts of the rose aphid, *Macrosiphum pseudorosae* (Patch), include the genus *Rosa* and a variety of herbaceous plants (Footitt and Maw 1997). The lupine aphid, *Macrosiphum albifrons* Essig, is a specialist, feeding exclusively on plants in the genus *Lupinus* (Blackman and Eastop 1984). While *M. persicae* is believed to be Palearctic in origin (Blackman and Eastop 1984), the other three aphid species are Nearctic (Stroyan 1981; Blackman and Eastop 1984).

### Laboratory trials

Lady beetles were collected from the field 48–72 hours before test initiation, maintained on a 50/50 diet of honey/egg yolk, then provided with water, but no food, for 48 hours before test initiation. Lady beetles were collected from a variety of locations and plants in Orono, Maine (44.8835° N, 68.6721° W), that included mixed shrub (*Solidago* sp., *Rubus* sp., *Prunus* sp., *Rosa* sp., *Cornus sericea*, *Alnus* sp.), apple (*Malus* sp.), grain (*Hordeum* sp., *Avena* sp.), mixed organic crops (*Solanum lycopersicon*, *Allium* sp., *Brassica* sp., *Pisum* sp., *Phaseolus* sp.) and fallow fields (*Phleum pratense*, *Trifolium* sp., *Cirsium* sp., *Vicia* sp., *Fragaria* sp.).

Potato aphids and green peach aphids were obtained from colonies maintained in the laboratory. The colonies were originally founded by aphids collected from potato, *Solanum tuberosum* (Solanales: Solanaceae), in Presque Isle, Maine, and then maintained

for at least 20 generations on excised potato foliage in the laboratory. Rose and lupine aphids were collected in the field from host vegetation including multi-flora rose, *Rosa multiflora* Thunberg (Rosales: Rosaceae), and lupine, *Lupinus polyphyllus* Lindley (Fabales: Fabaceae), respectively, and then maintained in the laboratory on excised host vegetation for up to 3 days before use in trials.

For each experiment, 10 aphids of the same species were placed, using a paintbrush, on an excised leaflet held within a 100 x 15 mm polystyrene Petri dish. Leaves used in trials were from the host plants from which aphids were collected in the field, as previously stated. Each trial was initiated when a single lady beetle previously housed in a separate Petri dish was added to the Petri dish containing the aphids by quickly exchanging lids between the two Petri dishes when the lady beetle was on the lid. After 24 hours, the beetle was removed and the number of aphids remaining in the dish was recorded. When a partial aphid remained, it was estimated to the 0.25 aphid. The experiment was conducted separately with adult apterae and with first to second instars. Sixty trials were conducted with each lady beetle species/aphid species pairing: 30 replicates with adult aphids and 30 replicates with the nymphs.

Lady beetles, aphid colonies, and test dishes were housed in Percival I-33VL Intellus environmental chambers at a 16:8 L:D photoperiod and 20! C. Trials with *M. euphorbiae* and *M. albifrons* were conducted in 2005, from June 16 to August 12 and from June 2 to August 12, respectively. Trials with *M. persicae* and *M. pseudorosae* were conducted in 2006, from May 24 to August 16 and from August 10 to August 24, respectively. Trials were conducted continuously throughout the range of dates

and in random order with respect to beetle species, aphid species, and choice of aphid nymph or adult.

### Field observations

Plots of *L. polyphyllus*, *S. tuberosum*, and *R. multiflora* were observed for 30 minutes each in Orono, ME (44.8974°N, 68.6873°W). Observations were made between 10:00 am and 2:00 pm in plots at least 0.1 ha in size where the vegetation of interest was dominant ( $\geq 50\%$ ). The number of adult lady beetles on host vegetation where aphids were absent (designated “absent”) or where *M. albifrons*, *M. euphorbiae*, or *M. pseudorosae* were present (designated “present”) was recorded. Because aphid populations were fairly contiguous where present, with no break in distribution greater than approximately one meter, each of the plots observed was designated as either “absent” or “present.” Forty observation trials were conducted for each of the three species. *M. persicae* were not found in the field in numbers sufficient to conduct observations. *M. albifrons* colonies were observed from June 2 to July 12, 2005, *M. euphorbiae* colonies were observed from June 17 to July 30, 2005, and *M. pseudorosae* colonies were observed from June 20 to August 24, 2006.

### Statistical analyses

Normality of laboratory-generated data was tested using the Wilk-Shapiro test (PROC UNIVARIATE; SAS Institute Inc. 2002). The data were transformed using rank transformations (Conover and Iman 1989). Means and standard errors reported in this paper were calculated from the untransformed data. Differences between lady beetle species were analyzed separately for each aphid species using one-way ANOVA followed by Tukey’s multiple comparison test (PROC GLM; SAS Institute Inc. 2002). Analyses

were conducted separately for aphid nymphs and adults.

Poisson regression (PROC GENMOD; SAS Institute Inc. 2002; SAS Institute Inc. 2005) was used to analyze lady beetle count data generated during field observations. Each plant species observed was analyzed separately, with the number of lady beetles as the response variable and lady beetle species and aphid presence/absence as the predictor variables. Overdispersion for *M. albifrons* and *M. pseudorosae* was corrected using a multiplicative overdispersion factor (Pearson chi-square divided by degrees of freedom) (Cox 1983; Allison 1999; SAS Institute Inc. 2005).

## Results

### Laboratory trials

There were always significant differences in the numbers of aphids consumed by different lady beetle species (Figure 1). *H. axyridis* consumed the most nymphs and adults of *M. persicae* (nymphs:  $F_{3,116} = 6.27$ ,  $p < 0.0006$ ; adults:  $F_{3,116} = 37.37$ ,  $p < 0.0001$ ), *M. euphorbiae* (nymphs:  $F_{3,116} = 11.98$ ,  $p < 0.0001$ ; adults:  $F_{3,116} = 20.67$ ,  $p < 0.0001$ ), and *M. pseudorosae* (nymphs:  $F_{3,116} = 32.59$ ,  $p < 0.0001$ ; adults:  $F_{3,116} = 48.47$ ,  $p < 0.0001$ ) compared with the other three lady beetle species, while *P. quatuordecimpunctata* consumed the fewest adults of these three aphid species and the fewest nymphs of *M. persicae* and *M. euphorbiae*. *C. septempunctata* consumed the lowest numbers of *M. pseudorosae* nymphs compared with the other three beetle species.

Lady beetles generally consumed fewer *M. albifrons* (Figure 1) compared with the other three aphid species. *C. trifasciata* and *P. quatuordecimpunctata* consumed a greater

number of *M. albifrons* nymphs compared with the other two beetle species ( $F_{3,116} = 11.86$ ,  $p < 0.0004$ ); *C. trifasciata* also consumed the greatest number of lupine aphid adults ( $F_{3,116} = 6.46$ ,  $p < 0.0006$ ).


### Field observations

All four lady beetle species were found on *S. tuberosum*, while only *H. axyridis* and *P. quatuordecimpunctata* were found on *R. multiflora* and only *C. trifasciata* and *P. quatuordecimpunctata* were found on *L. polyphyllus* (Table 1). There were significant differences in mean numbers of lady beetle species documented in two of the three vegetation types observed. The most abundant species in *S. tuberosum* was *P. quatuordecimpunctata*, followed by *C. septempunctata* ( $X^2 = 18.17$ ,  $p < 0.0001$ ), *H. axyridis* ( $X^2 = 22.02$ ,  $p < 0.0001$ ), and *C. trifasciata* ( $X^2 = 18.84$ ,  $p < 0.0001$ ). On *L. polyphyllus*, *P. quatuordecimpunctata* was more abundant than *C. trifasciata* ( $X^2 = 5.52$ ,  $p = 0.0188$ ). However, there was no

difference in the relative abundance of *P. quatuordecimpunctata* and *H. axyridis* on *R. multiflora*. Although mean lady beetle numbers were higher in six out of the eight occasions where aphids were present compared to absent (Table 1), those differences were not significant.

### Discussion

Consumption rates of the four aphid species differed among the four lady beetle species. With the exception of *M. albifrons*, *H. axyridis* was the most voracious predator, while *P. quatuordecimpunctata* removed the least prey. There may be a number of reasons for these differences. First, consumption rates may have been affected by the size of the beetles and/or the size of the prey. *C. septempunctata* is the largest of the lady beetles studied, followed by *H. axyridis*, *C. trifasciata*, and *P. quatuordecimpunctata* (Finlayson unpublished data). Being the



**Figure 1.** Mean ( $\pm$  standard error) consumption of aphids by different lady beetle species (Ct = *Coccinella trifasciata*, Cs = *Coccinella septempunctata*, Ha = *Harmonia axyridis*, Pq = *Propylea quatuordecimpunctata*). For each aphid species, nymphs and adults were analyzed separately; means with the same letter are not significantly different. High quality figures are available online.

smallest in size, *P. quatuordecimpunctata* may be satiated with fewer aphids compared with the other species. *M. albifrons* is larger than the other aphid species, thus fewer *M. albifrons* may satiate a beetle compared with the other species offered. Consumption rate may also be affected by differences in handling (Pervez and Omkar 2005), nutritional suitability of prey (Houck 1991; Roger et al. 2001; Gagné et al. 2002), or chemical deterrence (Pasteels et al. 1983; Nishida and Fukami 1989).

Observations of adult beetles on field plots were generally consistent with expectations based on their consumption of aphids in the laboratory. *H. axyridis* consumed the most *M. pseudorosae* in laboratory trials and was one of two species found in the field with *M. pseudorosae*. *C. trifasciata* consumed the most *M. albifrons* in laboratory trials and was one of two species found in the field with *M. albifrons*. The other beetle species found with *M. pseudorosae* and *M. albifrons* was *P. quatuordecimpunctata*, the species that consumed the second largest number of *M. pseudorosae* and *M. albifrons*, although this difference was only statistically significant for *M. albifrons* nymphs. It is also not surprising to find *P. quatuordecimpunctata* in all observations because this species is probably the most abundant lady beetle in Maine

(Finlayson et al. 2008).

Three of the species tested in this study, *H. axyridis*, *C. septempunctata*, and *P. quatuordecimpunctata*, are not native to Maine (Gordon 1985). Because lady beetle species differ in their prey consumption, decreases in the relative abundance of native species following the establishment of non-native lady beetle species that has been reported in a number of studies (Elliot et al. 1996; Brown and Miller 1998; Colunga-Garcia and Gage 1998; Michaud 2002; Brown 2003; Turnock et al. 2003; Alyokhin and Sewell 2004) may favor some aphid species over others. For example, Alyokhin et al. (2005) observed a significant reduction in both density and the amplitude of annual oscillations of populations of *M. persicae* and *Aphis nasturtii* following the establishment of *H. axyridis* and *P. quatuordecimpunctata*.

When compared with other aphidophagous coccinellid species, *H. axyridis* has been shown to have superior competitive abilities regarding its feeding rate (Michaud 2002), intraguild predation (Hironori and Katsuhiko 1997; Yasuda et al. 2001; Yasuda et al. 2004), and interactions with natural enemies (Dutcher et al. 1999; Saito and Bjørnson 2006; Finlayson et al. 2009). Similarly, in this study, *H. axyridis* exhibited greater prey

**Table 1.** Mean (± standard error) number of lady beetles documented during field observations. Mean beetles documented where aphids were present on vegetation are presented alongside mean beetles that were documented where aphids were absent.

Lady Beetle Species		Plant/Aphid Species					
		potato		rose		lupine	
		present	absent	present	absent	present	absent
N		22	18	26	14	36	4
<i>C. trifasciata</i>	Mean	0.05	0.06 d	0	0	0.39	0 b
	SE	0.045	0.055			0.151	
<i>C. septempunctata</i>	Mean	0.36	0.17 b	0	0	0	0
	SE	0.105	0.090				
<i>H. axyridis</i>	Mean	0.14	0.17 c	0.81	0.14 a	0	0
	SE	0.075	0.090	0.236	0.097		
<i>P. quatuordecimpunctata</i>	Mean	1.36	0.89 a	0.46	0.36 a	1.25	0.5 a
	SE	0.242	0.196	0.149	0.169	0.377	0.289

N = the number of observations, out of 40, where aphids were either present or absent.  
For each plant/aphid species, lady beetle species with the same letter are not significantly different.



consumption of three of the four aphid species tested compared with the other lady beetle species tested. The true voracity of *H. axyridis*, however, may have been underestimated because it consumed close to the upper limit of what was made available in trials. Providing more than ten aphids may have improved the resolution of species differences.

*M. albifrons* is native to the study area (Stroyan 1981) and is known to sequester toxic compounds from its host plant that have been shown to cause a “narcotizing effect” on *C. septempunctata* (Gruppe and Roemer 1988). It is thus notable that *H. axyridis* and *C. septempunctata*, both introduced species without historical exposure to *M. albifrons*, consumed the lowest numbers of this species. In contrast, *C. trifasciata*, which is native to the area, consumed the most *M. albifrons* adults. It would seem that *C. trifasciata* may have evolved the ability to tolerate these compounds, whereas the recently introduced non-native species have yet to do so. By virtue of being able to exploit lupine aphids, *C. trifasciata* may enjoy a refuge from prey competition with the non-native species. These differences in prey consumption suggest that different lady beetle species should not be considered equal consumers of aphids.

## Acknowledgments

The authors would like to thank the following for their assistance in the field and laboratory: Lauren Little, David Ginsberg, Allison Fleming, and Todd Finlayson. We would also like to thank Joseph Cannon, John Jemison, Black Bear Food Guild, and Orono Land Trust for providing access and guidance on their land, and Frank Drummond and Malcolm Hunter, Jr., J.P. Michaud, and two

anonymous reviewers for providing comments on the manuscript. This research was supported by the Maine Agricultural and Forest Experiment Station (Hatch ME08466-01) and the National Science Foundation’s GK-12 Teaching Fellows Program (Grant # DGE – 0231642 to S. Brawley et al.). This is Publication No. 3034 of the Maine Agricultural and Forest Experiment Station.

## References

- 
- Allison PD. 1999. *Logistic Regression Using the SAS System: Theory and Application*. SAS Institute Inc.
- Alyokhin A, Sewell G. 2004. Changes in lady beetle community following the establishment of three alien species. *Biological Invasions* 6: 463-471.
- Alyokhin A, Drummond FA, Sewell G. 2005. Density-dependent regulation in populations of potato-colonizing aphids. *Population Ecology* 47: 257-266.
- Angalet GW, Jacques JR. 1975. The establishment of *Coccinella septempunctata* L. in the continental United States. United States Department of Agriculture Cooperative Economic Insect Report 25: 883-884.
- Blackman RL, Eastop VF. 1984. *Aphids on the World's Crops: An Identification and Information Guide*. John Wiley & Sons.
- Brown MW, Miller SS. 1998. Coccinellidae (Coleoptera) in apple orchards of eastern West Virginia and the impact of invasion by *Harmonia axyridis*. *Entomological News* 109: 136-142.
- Brown MW. 2003. Intraguild responses of aphid predators on apple to the invasion of an



exotic species, *Harmonia axyridis*. *BioControl* 48: 141-153.

Chapin EA. 1966. A new species of Myrmecophilous Coccinellidae, with notes on other Hyperspini (Coleoptera). *Psyche* 73: 278-283.

Colunga-Garcia M, Gage SH. 1998. Arrival, establishment, and habitat use of the multicolored Asian lady beetle (Coleoptera: Coccinellidae) in a Michigan landscape. *Environmental Entomology* 27: 1574-1580.

Cox DR. 1983. Some remarks on overdispersion. *Biometrika* 70: 269-274.

Dutcher JD, Estes PM, Dutcher MJ. 1999. Interaction in Entomology: Aphids, aphidophaga and ants in pecan orchards. *Journal of Entomological Science* 34: 40-56.

Elliot N, Kieckhefer R, Kauffman W. 1996. Effects of an invading coccinellid on native coccinellids in an agricultural landscape. *Oecologia* 105: 537-544.

Finlayson CJ, Landry KM, Alyokhin AV. 2008. Abundance of native and non-native lady beetles (Coleoptera: Coccinellidae) in different habitats in Maine. *Annals of the Entomological Society of America* 101: 1078-1087.

Finlayson CJ, Alyokhin AV, Porter EW. 2009. Interactions of native and non-native lady beetle species (Coleoptera: Coccinellidae) with aphid-tending ants in laboratory arenas. *Environmental Entomology*, 38: 846-855.

Footitt RG, Maw E. 1997. Biological survey of Canada (terrestrial arthropods). In: Danks HV, Downes JA, editors. *Insects of the Yukon*. Canadian Museum of Nature.

Gagné I, Coderre D, Mauffette Y. 2002. Egg cannibalism by *Coleomegilla maculata lengi* neonates: preference even in the presence of essential prey. *Ecological Entomology* 27: 285-291.

Gordon RD. 1985. The Coccinellidae (Coleoptera) of America north of Mexico. *Journal of New York Entomological Society* 93: 1-912.

Gruppe A, Roemer P. 1988. The lupin aphid (*Macrosiphum albifrons* Essig, 1911) (Hom., Aphididae) in West Germany: Its occurrence, host plants and natural enemies. *Journal of Applied Entomology* 106: 135-143.

Hironori Y, Katsuhiko S. 1997. Cannibalism and interspecific predation in two predatory ladybirds in relation to prey abundance in the field. *Entomophaga* 42: 53-163.

Hodek I. 1973. *Biology of Coccinellidae*. W Junk Publisher.

Houck MA. 1991. Time and resource partitioning in *Stethorus punctum* (Coleoptera: Coccinellidae). *Environmental Entomology* 20: 494-497.

Kalushkov P. 1998. Ten aphid species (Sternorrhyncha: Aphididae) as prey for *Adalia bipunctata* (Coleoptera: Coccinellidae). *European Journal of Entomology* 95: 343-349.

Kalushkov P, Hodek I. 2004. The effects of thirteen species of aphids on some life history parameters of the ladybird *Coccinella septempunctata*. *BioControl* 49: 21-32.

Koch RL. 2003. The multicolored Asian lady beetle, *Harmonia axyridis*: A review of its biology, uses in biological control, and non-target impacts. *Journal of Insect Science* 3:32, Available online:

<http://insectscience.org/3.32>.

Michaud JP. 2000. Development and reproduction of ladybeetles (Coleoptera: Coccinellidae) on the citrus aphids *Aphis spiraecola* Patch and *Toxoptera citricida* (Kirkaldy) (Homoptera: Aphididae). *Biological Control* 18: 287-297.

Michaud JP. 2002. Invasion of the Florida citrus ecosystem by *Harmonia axyridis* (Coleoptera: Coccinellidae) and asymmetric competition with a native species, *Cycloneda sanguinea*. *Environmental Entomology* 31: 827-835.

Mignault M-P, Roy M, Brodeur J. 2006. Soybean aphid predators in Quebec and the suitability of *Aphis glycines* as prey for three Coccinellidae. *BioControl* 51: 89-106.

Nishida R, Fukami H. 1989. Host plant iridoid-based chemical defense of an aphid, *Acyrtosiphon nipponicus*, against ladybird beetles. *Journal of Chemical Ecology* 15: 1837-1845.

Obrycki JJ, Orr CJ. 1990. Suitability of three prey species for Nearctic populations of *Coccinella septempunctata*, *Hippodamia variegata*, and *Propylea quatuordecimpunctata* (Coleoptera: Coccinellidae). *Journal of Economic Entomology* 83: 1292-1297.

Pasteels JM, Grégoire JC, Rowell-Rahier M. 1983. The chemical ecology of defense in arthropods. *Annual Review of Entomology* 28: 263-289.

Pedigo LP, Rice ME. 2006. *Entomology and Pest Management*. Pearson Prentice Hall.

Pervez A, Omkar. 2005. Functional responses of coccinellid predators: An illustration of a logistic approach. *Journal of Insect Science*

5:5, Available online:  
<http://insectscience.org/5.5>.

Phoofolo MW, Obrycki JJ. 1997. Comparative prey suitability of *Ostrinia nubilalis* eggs and *Acyrtosiphon pisum* for *Coleomegilla maculata*. *Biological Control* 9: 167-172.

Roger C, Coderre D, Vigneault C, Boivin G. 2001. Prey discrimination by a generalist coccinellid predator: Effect of prey age or parasitism? *Ecological Entomology* 26: 163-172.

SAS Institute Inc. 2002. SAS Institute Inc.

SAS Institute Inc. 2005. Fitting Poisson Regression Models Using the GENMOD Procedure Course Notes. SAS Institute Inc.

Saito T, Bjørnson S. 2006. Horizontal transmission of a microsporidium from the convergent lady beetle, *Hippodamia convergens* Guérin-Ménéville (Coleoptera: Coccinellidae), to three coccinellid species of Nova Scotia. *Biological Control* 39: 427-433.

Stroyan HLG. 1981. A North American lupin aphid found in Britain. *Plant Pathology* 30: 253.

Tedders WL, Schaefer PW. 1994. Release and establishment of *Harmonia axyridis* (Coleoptera: Coccinellidae) in the Southeastern United States. *Entomological News* 105: 228-243.

Turnock WJ, Wise IL, Matheson FO. 2003. Abundance of some native coccinellids (Coleoptera: Coccinellidae) before and after the appearance of *Coccinella septempunctata*. *Canadian Entomologist* 135: 391-404.

Wheeler AG. 1990. *Propylea quatuordecimpunctata*: Additional records of

an adventive lady beetle (Coleoptera: Coccinellidae). *Entomological News* 101: 164-166.

Yasuda H, Kikuchi T, Kindlmann P, Sato S. 2001. Relationships between attack and escape rates, cannibalism, and intraguild predation in larvae of two predatory ladybirds. *Journal of Insect Behavior* 14: 373-384.

Yasuda H, Evans EW, Kajita Y, Urakawa K, Takizawa T. 2004. Asymmetric larval interactions between introduced and indigenous ladybirds in North America. *Oecologia* 141: 1432-1939.