

A Review of the Implications of Heterozygosity and Inbreeding on Germplasm Biodiversity and Its Conservation in the Silkworm, *Bombyx mori*

Authors: Jingade, A.H., Vijayan, K., Somasundaram, P., Srivasababu, G.K., and Kamble, C.K.

Source: Journal of Insect Science, 11(8) : 1-16

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.011.0108>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



A review of the implications of heterozygosity and inbreeding on germplasm biodiversity and its conservation in the silkworm, *Bombyx mori*

A.H. Jingade^{1a*}, K. Vijayan^{2b}, P. Somasundaram^{1c}, G.K. Srinivasababu^{1d}, C.K. Kamble^{1e}

¹Central Sericultural Germplasm Resources Centre, Central Silk Board, Thally Road, Hosur, Tamil Nadu-635 109, India

²Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan-115, ROC.

Abstract

Silkworm genebanks assume paramount importance as the reservoirs of biodiversity and source of alleles that can be easily retrieved for genetic enhancement of popular breeds. More than 4000 *Bombyx mori* L (Lepidoptera: Bombycidae) strains are currently available and these strains are maintained through continuous sibling mating. This repeated sibling mating makes the populations of each strain more homozygous, but leads to loss of unique and valuable genes through the process of inbreeding depression. Hence, it is essential to maintain a minimal degree of heterozygosity within the population of each silkworm strain, especially in the traditional geographic strains, to avoid such loss. As a result, accurate estimation of genetic diversity is becoming more important in silkworm genetic resources conservation. Application of molecular markers help estimate genetic diversity much more accurately than that of morphological traits. Since a minimal amount of heterozygosity in each silkworm strain is essential for better conservation by avoiding inbreeding depression, this article overviews both theoretical and practical importance of heterozygosity together with impacts of inbreeding depression and the merits and demerits of neutral molecular markers for measurements of both heterozygosity and inbreeding depression in the silkworm *Bombyx mori*.

Keywords: conservation, genetic diversity, inbreeding depression, molecular markers

Abbreviations: HFC, heterozygosity fitness correlations; ID, inbreeding depression; SNP, single nucleotide polymorphisms; SSR, microsatellites

Correspondence: ^{a*} anujnh@yahoo.co.in, ^b kvijayan01@yahoo.com, ^c drpsunder@rediffmail.com, ^d babu_jica@yahoo.co.in, ^e director_csgrc@yahoo.com, *Corresponding author

Editor: Brad Coates was editor of this paper.

Received: 31 October 2009, **Accepted:** 26 March 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 11, Number 8

Cite this paper as:

Jingade AH, Vijayan K, Somasundaram P, Srinivasababu GK, Kamble CK. 2011. A review of the implications of heterozygosity and inbreeding on germplasm biodiversity and its conservation in the silkworm, *Bombyx mori*. *Journal of Insect Science* 11:8 available online: insectscience.org/11.8

Introduction

The mulberry silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae) is a completely domesticated lepidopteran insect that has been used for silk production for more than 5000 years (International Silkworm Genome Consortium 2008). Its production base is spread over 60 countries with Asian countries producing over 90% of mulberry silk production (Datta and Nanavathy 2007). In addition to being used in silk industry, *B. mori* is also used for production of recombinant proteins (Tamura et al. 2000; Tomita et al. 2003) and for basic research in microbiology, physiology, and genetics (Wills et al. 1995). Considering the great economic importance attached to *B. mori*, the traditional and leading silk producing countries such as China, Japan, India, Russia, Korea, and Bulgaria have collected a number of silkworm races suitable for a wide range of agro-climatic conditions (Tables 1, 2). Currently more than 4000 strains are available in the germplasm of *B. mori* (Nagaraju 2002; Kumaresan et al. 2004a, 2004b), which includes uni-, bi-, and

polyvoltine varieties. These silkworm strains have been maintained under *ex situ* conditions through years of continuous sibling mating and selection. Consequently, these strains have possibly undergone many evolutionary changes through mutations (natural as well as man-made selections), thereby creating a wide genetic diversity among the strains; but this diversity becomes more homogenous within the strain. This repeated inbreeding in each traditional strain results in the loss of valuable genes through a process called inbreeding depression. Therefore, it is imperative to prevent this genetic erosion due to excessive inbreeding to conserve the genetic resources of geographical strains for their use in divergent breeding programmes. In this context it is worth bearing in mind that the process of conservation of the genetic resources of these strains does not only necessitate acquisition and physical possession of them, it is equally important to ensure that the original genetic characters are maintained and the rare and endemic strains are well protected from extinction. Therefore, all possible measures need to be taken to

Table 1. Silkworm, *Bombyx mori* genetic resources available in different countries

Sl. No.	Country	Multivoltine	Bivoltine	Total
1	Brazil	10	65	75
2	Bulgaria	0	183	183
3	China	20	580	600
4	France	0	53	53
5	India	150	450	600
6	Iran	0	50	50
7	Italy	0	30	30
8	Japan	30	1542	1572
9	N. Korea	5	281	286
10	Russia	0	500	500
11	S. Korea	6	300	306
12	Thailand	5	25	30
13	Vietnam	5	20	25
	Total	231	4079	4310

(Source: FAO Manual, 2003)

Table 2. Country-wide silkworm germplasm maintained at Central Sericulture Resources Centre, Hosur, Tamil Nadu, India

Sl.No	Country	Bivoltine	Multivoltine	Total
1	Bangladesh	0	3	3
2	Brazil	3	0	3
3	China	40	4	44
4	France	11	0	11
5	India	207	63	270
6	Indonesia	1	0	1
7	Iraq	1	0	1
8	Japan	64	3	67
9	Poland	3	0	3
10	Russia	19	0	19
11	South Korea	1	0	1
12	Thailand	4	0	4
13	Ukraine	2	0	2
14	Vietnam	3	0	3
	Total	359	73	432

preserve the originality of the *B. mori* races as the gene banks are reservoirs of gene assemblies consisting of geographically isolated races, genetic stocks, and breeds collected from indigenous and exotic regions. Frankel (1984) developed the concept of core collections to preserve germplasm accessions with minimum repetitiveness and at the same time retain most of its original genetic diversity. This concept was further elaborated by Frankel and Brown (1984). Rapid replacement of traditional races by high yielding breeds/hybrids also threatens the very survival of a number of old races, which are the result of conscious or subconscious selection over a long period of time. Some of the old Indian traditional silkworm races such as Chotopolu and ancient breeds viz., Nistid, Nismo, Ichat, and Itan are either extinct or on the verge of extinction due to the absence of a proper conservation system (Thangavelu 1998). It is, therefore, high time for silkworm conservationists and geneticists to take appropriate measures to protect their valuable resources from further degradation and extinction.

Genetic diversity in the silkworm *Bombyx mori*

Genetic diversity is the genetic variation within species, both among geographically separated populations and among individuals within a single population. Genetic diversity is an essential aspect in conservation biology because a fundamental concept of natural selection states that the rate of evolutionary change in a population is proportional to the amount of genetic diversity present in it (Fisher 1930). Decreasing genetic diversity increases the extinction risk of populations due to a decline in fitness. Therefore, both biochemical and molecular markers have recently been employed to estimate the extent of genetic diversity present among various types of silkworm strains such as mono-, bi- and multivoltines present in China, Japan, Korea, India, and several other countries. For instance, Nagaraja and Nagaraju (1995) estimated genetic diversity among thirteen silkworm strains from tropical and temperate regions. The study placed the silkworms into two distinct groups consisting of six diapausing and seven non-diapausing

genotypes. The genotype, Pure Mysore, was found to be the most divergent genotype among the genotypes studied. Likewise, Chatterjee and Pradeep (2003), Dalirsefat et al (2009), Li et al. (2005), Lu et al. (2002), Nagaraju et al. (2001), Prasad et al. (2005), Reddy et al. (1999a, b), and Xia et al. (1998) used AFLP, RFLP, ISSR, and SSR markers to estimate the genetic distance among silkworm strains that differed in several morphological traits. Pairwise genetic diversity estimation among 20 mutant strains by Velu et al. (2008) revealed 0.045 to 0.553 genetic diversity and consequent grouping among the silkworms. Recently Xia et al (2009) constructed a single-base pair resolution genetic map using genomic information from 40 domesticated and wild silkworm strains. With the help of 16 million single nucleotide polymorphism (SNP) markers, identified from the total genomic information, the domestication events and subsequent genetic differentiation in *B. mori* have been worked out. The study revealed significant genetic divergence between wild and domesticated strains of *B. mori*. The genetic variation was measured using the population size scaled mutation rate which was significantly smaller in domesticated strains (0.011) when compared to the wild strains (0.013). The rate of heterozygosity in domesticated strains was two times lower than that in wild varieties (0.003 and 0.008, respectively), which is most likely attributed to inbreeding experienced by domesticated lines. The study also shows clear genetic separation between domesticated and wild varieties and relatively less gene flow between them.

Significance of heterozygosity and inbreeding depression in silkworm race maintainance

Heterozygosity is the state of having different alleles with regard to a given character. Allelic diversity, on the other hand, determines a population's ability to respond to long-term selection over many generations, and ultimately the survival of the population (perhaps even of the species). Individual heterozygosity is expected to be correlated between parents and their offspring. The expected correlation is maximal, approaching $r = 0.50$, when allelic frequencies are highly asymmetric, and it is zero when the allelic frequencies are equal to 0.50 (Mitton et al. 1993). Inbreeding is the process of mating of two closely related individuals. The inbreeding coefficient of an individual refers to how closely related its parents are. The major effect of inbreeding is the reduction in mean performance of the population. Inbreeding tends to reduce the level of the characters connected with fitness and consequently can lead to loss of general vigour and fertility. This reduced fitness of populations caused by the manifestation of deleterious recessive genes is termed inbreeding depression. ID usually reduces reproductive performance and survival (Falconer 1989), and has been documented in many populations ever since Darwin (1876) described it (Wright 1977; Charlesworth and Charlesworth 1987; Ralls et al. 1988; Falconer 1989).

In *B. mori*, it is speculated that the wild gene pool has undergone considerable selection in every generation during the course of its domestication, though it might be unintentional (<http://unp.un.org/>). Consequently the pool reached its current level of diversity of characteristics within the species. Genetic principles and breeding techniques of modern genetic science were utilised for cross breeding and exploitation of the phenomenon of hybrid vigour in the

silkworm. A high level of heterosis/hybrid vigour for various quantitative characters was found to occur in the crosses of highly inbred lines and geographically divergent populations of the silkworm (Harada 1957; Harada 1961; Krishnaswami et al. 1964; Hirobe 1968; Yokoyama 1979; Gamo and Hirabayashi 1983; Nagaraju 1990). Low heritability traits, such as survival rate, displays extensive heterosis and have become the focal point for exploitation of heterosis in the silkworm (Nagaraju and Pavan Kumar 1995). In silkworms, when an initial choice of parents has to be made to obtain heterosis, it is important to ascertain the level of parental divergence (Jolly et al. 1989; Chatterjee et al. 1993; Mukerjee et al. 1999; Kumaresan et al. 2003a, 2003b). Parental homozygosity has a distinct influence on the degree of manifestation of heterosis (Nagaraju and Goldsmith 2002). Hybrid vigour arises when the crossing of two inbred lines (or an outbred line with an inbred line) produces individuals that are heterozygous at loci that were previously homozygous for deleterious alleles (Falconer 1981). Petkov et al. (1999) has attempted to solve the problem of applying inbreeding to the silkworm. However the evidence of effects of inbreeding on inbreeding coefficients in silkworm populations is limited.

Estimation of inbreeding depression

Based on the Hardy and Weinberg law, which states that in large random mating populations, all the genotypes have equal fitness, i.e. there is no selection, therefore both allelic and genotypic frequencies in the population remain constant as long as other disturbing factors such as non-random mating, mutations, selection, limited population size, random genetic drift, and gene flow do not act (Hardy 1908). Wright (1922) formulated F-statistics to measure the inbreeding coefficient

'F' at a locus using the formula $2pq(l - F)$ where p and q are the allelic frequencies at a diallelic locus and F is the inbreeding coefficient. Individuals with the same expected inbreeding co-efficients vary in their degree of homozygosity hence there is opportunity for natural selection to modify the array of genotypic frequencies, particularly to favour the more heterozygous individuals. Inbreeding depression for a trait is a linear function of the inbreeding coefficient (Lynch and Walsh 1998). Inbreeding depression has been measured at the population level by comparing the mean phenotypes of inbred and outbred individuals. Inbreeding depression is estimated from the difference in mean fitness between inbred and outbred individuals within the same family, often the progeny of a single maternal parent. Different techniques are available to determine the level of inbreeding. The advantages and disadvantages of different approaches for measuring inbreeding and inbreeding depression is given in Table 3. Genetic drift (F_{ST}), quantified by effective population size, is another parameter with which inbreeding depression within a population is measured. F_{ST} measures the decrease in heterozygosity due to the finite size of the population and the effective size. In fact F_{ST} gives a measure of the genetic drift among conceptual replicates of the population in an infinite nonrandom mating population, $F_{ST} = 0$ and $F_{IT} = F_{\sigma}$, rising toward a value which depends on the proportion of inbred matings. Thus, genetic drift (F_{ST}) and decline in heterozygosity (F_{IT}) will have the same rate of change except that the drift runs one generation ahead of the decline in heterozygosity.

In countries that practice sericulture, several pure races belonging to different voltinism groups are maintained in germplasm centres by selection and inbreeding. Sibling mating of the progenies derived from a single brood is

Table 3. Advantages and disadvantages of approaches for measuring inbreeding and inbreeding depression

Sl.No.	Method	Advantages	Disadvantages
1	Pedigree analysis	Natural mating patterns	Requires accurate information on paternity and long term monitoring to establish pedigrees
		Natural levels of environmental variation	
		Accurate estimates of individual inbreeding and its effects	
		Easy comparisons among studies	
2	Genetic analysis	Natural mating patterns	Tracking of changes in F so far limited to partially selfing plants and animals
		Natural levels of environmental variation	Many techniques do not allow comparisons of results among studies
		Parentage assignment using genetic markers	
3	Experimentally inbred mating	Most precise control of inbreeding and potentially confounding variables	Levels of inbreeding studied usually exceed those present in natural populations, thus the results might not show how much inbreeding depression occurs in nature, if high levels of inbreeding bring synergistic effects
		True replication providing rigorous and accurate estimates of inbreeding depression	

(Source: Keller and Wallar, 2002)

preferred for pure stocks so that the original traits of the races are maintained through generations. The homozygous races maintained in the germplasm are an important source of breeding new silkworm breeds through hybridization (Subramanya and Bishop 2009). Generally breeders try to maintain the original characters of the races through selection with care to avoid inbreeding depression. Although, much work has been carried out on quantitative traits such as yield and biochemical parameters (Chatterjee et al. 1993; Natcheva et al. 2001; Nagaraju 2002) not much work has been done on the impacts of inbreeding on quantitative traits. Two statistical models, viz. general mixed model and a fixed model, are available for estimating the impact of inbreeding on quantitative traits (Subramanya and Bishop 2009).

Inbreeding depression in silkworm races

The general procedure of silkworm breeding is the judicious selection of parents and their subsequent matings to meet definite purposes. Genetic improvement of silkworm

races is generally achieved by conventional breeding methods of hybridization between selected parents combined with selection for desired traits. The improvement of silkworm strains has been achieved over a long period by pure line selections. The advantage of using F1 hybrids for commercial exploitation was emphasized by Toyama as early as in 1906. Inbred lines are developed by crossing between two strains showing excellent characteristics and subsequent sibling-mating and progeny selections for seven or more generations. Silkworm strains maintained at germplasms serve as genetic stocks. Genetic stock refers to an inbred stock that carries one or several qualitative characters. A parental pure line refers to an inbred stock that can be crossed with other purelines in order to produce a hybrid variety. This is done in order to exploit heterosis or the hybrid vigour phenomena of quantitative characters and qualitative traits that meet the needs and interests of breeders. Parents that are complementary for desired characters in one line are selected and a cross is made. Sibling-mating is carried out within an egg

batch of the F1, and the individual F2 silkworm pairs are selected with each maintained as a separate family through F5 or F6. General combining ability with a topcross tester is tested when uniformity is found to be necessary. The best general combiners are selected to identify strains used recurrently in a series of breeding programmes and specific crosses are made and tested for combining ability.

The most intense form of inbreeding in the silkworm is full-sibmating. Less intense forms of inbreeding occurs between cousins, second cousins, and other types of family matings. The relative effects can be seen in the inbreeding coefficient, F , in consecutive generations. For the mating type of full-sib, inbreeding coefficients increase from 0.25, 0.37, 0.50, 0.59, and up to 0.67 as inbred generations proceed from generation 1, 2, 3, 4, and go up to 5, respectively. In practical terms, the breeding expectation is that inbreeding leads to isolation of homozygous lines with reduced vigor. The reduction in vigor is not in a direct relationship with generation, but generally parallels the reduction in heterozygosity. Inbreeding can be expedited by selecting for the ultimate inbred characters, if they are defined. That is, if the inbred characteristic is round cocoon and long larval duration, selection of such individuals will give a better chance for achieving a relatively high degree of homozygosity. If characters are undefined, the breeder should select against heterotic properties (<http://unp.un.org/>). The success and spread of silkworm rearing was mostly due to the use of F1 silkworm hybrids of exotic temperate silkworm male parental strains and native female tropical silkworm strains (Nagaraju 2002). Continuous inbreeding results in the accumulation of many deleterious genes leading to inbreeding depression, thereby

resulting in the deterioration of some commercial characters. However, the overall combination of beneficial traits could be achieved in a reasonable way by employing inbreeding techniques coupled with selection as the silkworm breeder has at his disposal a diversified array of gene combination to manipulate and isolate new silkworm breeds having desirable qualities for commercial exploitation (Raju and Krishnamurthy 1993). The improvement of indigenous breed could be achieved through hybridizations utilizing exotic breeds (Kovalov 1970). Inbreeding the hybrids to isolate and stabilize silkworm breeds for the desired traits has been well documented by many breeders (Hirobe 1968; Kovalov 1970; Gamo 1976). Artificial selection has been widely utilised in the breeding programmes concerning this commercially important insect. Selection increases the frequency of homozygotes and makes homozygous effects stronger (Whitlock 2002). In *B. mori*, qualitative changes in the genetic basis, induced by artificial selection, resulted in genetically divergent populations with different phenotypic values of larval duration. In the Nistari strain of *B. mori*, where short larval duration is part of fitness component, when artificial selection was imposed the selection pressure acted against this fitness component and caused longer larval duration inbred lines by favouring variation at the multilocus level thus causing genetic divergence (Pradeep et al. 2005).

Molecular markers for heterozygosity estimation

Heterozygosity fitness correlations (HFC) historically appeared as an opportunity to develop marker-based genetics, an approach boosted by the discovery of allozyme polymorphism (Lewontin and Hubby 1966). Geneticists had known since as early as 1910–1920 that manipulations of genomic

heterozygosity (using inbred lines) could affect fitness components and crop yields (Shull 1952). HFC provided a potential 'microscopic' basis of the 'macroscopic' effects observed using inbred strains. Although the relationship between heterozygosity at the genome level and fitness was not in doubt, the question came in identifying the particular genes involved and how they affected fitness. Genes showing HFC were obviously good candidates. HFC is quantified by the variance in a fitness trait explained by its regression on heterozygosity (r^2). Most estimates yield an order of magnitude of 0.01-0.05. However, sample size (N) must be considered as, even if the true correlation is zero, an average value of $1/(N-1)$ is expected by chance (Sokal & Rohlf 1995). As most samples are of the order 10^2 , $1/(N-1)$ is of the same magnitude as the r^2 values observed. For multiple regressions (L loci as separate 'heterozygosity' variables) the null expectation for R^2 is $L/(N-1)$, a serious problem if L is not $<N$ (Bush et al. 1987). However, the overall lack of significant correlations in natural populations of such a model organism as *Drosophila* (Houle 1989) and in many other species, even among bivalves or pines (Britten 1996), strongly suggests that HFC is not universal. For a long time allozymes have been the most reliable and convenient genetic markers. This situation is now changing with the development of PCR-based codominant markers such as microsatellites, anonymous RFLP and intron length polymorphisms (Mitton 1994). Heterozygosity may be underestimated due to lack of resolution (alleles with similar mobility appear as the same allele), somatic aneuploidy, or null alleles. Most PCR-based markers also have null alleles. Most molecular markers have the same inherent defects as allozymes. However, PCR markers, unlike allozymes, allow

scoring of very small organisms (such as bivalve larvae) at many poly-morphic loci. Polymorphic markers (microsatellites) seem unsuitable as almost all individuals will be heterozygous. However, microsatellites provide estimates of the quantity of divergence between two alleles (specifically, the difference in repeat number, under a stepwise mutation model), an interesting alternative to the traditional heterozygosity (in which alleles are classified as either identical or different). Moreover molecular markers can be used in conjunction with allozymes, in order to compare the latter with *a priori* neutral loci. Such experiments allow evaluation of possible causes of HFC other than null alleles or aneuploidy.

Assessment of heterozygosity in each silkworm race is important for efficient management and conservation of genetic resources (Graner et al. 2004). Genotyping-based approaches (Table 4) can be used to quantify levels of inbreeding and to gain a better understanding of the genetic architecture of inbreeding depression through the detection of genes that affect it. Although not much effort has been made to measure the heterozygosity in different strains of *B. mori*, the few attempts which made recently have revealed that within strains genetic diversity is much less than it is among the strains. Staykova (2008) used isozyme patterns of 480 individuals of eight silkworm strains to assess the heterozygosity among the silkworm populations. The heterozygosity among the strains varied from 0.099 in China to 0.238 in Kinshu. Further, the study also revealed that the observed heterozygosity was much less than the expected value, which was attributed to the effect of the inbreeding. Similarly, using molecular markers such as AFLP in 23 silkworm strains composed of 12 Japanese and 11 Chinese strains, Gavria et al (2006)

Table 4. Characteristics of different molecular markers for assessing genetic diversity

Sl.No.	Marker	Dominance	Genome mapping	Reproducibility	Null alleles
1	Isozymes	Co-Dominant	Limited	Very high	Rare
2	RFLP	Co-Dominant	Good	Very high	Rare
3	ISSR	Dominant	Limited	High	Rare
4	RAPD	Dominant	V.good	Medium	NA
5	AFLP	Dominant	V.good	High	NA
6	SSR	Co-Dominant	Good	High	Common
7	SSCP	Co-Dominant	Good	High	Common
8	SNP	Co-Dominant	Good	High	Common
9	STS/EST	Co-Dominant	Good	High	Rare

showed that Japanese strains are more heterogenous than the Chinese strains. However, a recent study with three Iranian and three Japanese silkworm strains revealed low within-strain genetic variation (26.6%) and high genetic diversity between strains (75.4%). Among the strains, within strain variation was least in the Japanese strain P107 (0.08), and highest in the Iranian Khorasan Orange strain (0.18) (Dalirsefat et al. 2007). Since these studies are with isozyme (less polymorphism) and dominant molecular markers (AFLP), better depth of the heterozygosity cannot be revealed. Co-dominant markers that can reveal both homozygous and heterozygous loci have to be used. The most commonly used PCR based co-dominant markers are microsatellites (SSR) and single nucleotide polymorphisms (SNPs). Since a number of SSR primers have been developed from silkworm genome (Reddy et al. 1999a) they can be used for assessing the loss of heterozygosity in subsequent generations. SNPs is a new generation marker system, and represents stable, diallelic nucleotide variations that are distributed throughout the genome and are the most abundant resource for analysis of genetic variation (Kwok et al. 1994). Generally, microsatellites provide a higher level of resolution because of the large number of alleles per locus. But SNPs provide large data

sets because of high abundance in the genome and high throughput genotyping methods. To detect inbreeding depression, the level of heterozygosity is correlated with one or more measures of fitness. Modern genomic technologies enable screening of a large genome-wide set of molecular markers, which give a reliable estimate of F1. In combination with fitness measures, the genomic data also enable detection of genes associated with inbreeding depression (Kristensen et al. 2010). The results of SNPs can provide a fundamental description of the nucleotide diversity of the silkworm genome (Cheng et al. 2004), which is evident from the results reported by Xia et al (2009). Using information from 16 million SNPs identified in genomes of 40 silkworm strains, they estimated the heterozygosity in wild and domesticated silkworm strains. The rate of heterozygosity in the wild silkworm (0.0080) was more than double of that present in domesticated strains (0.0032). This significant reduction in the heterozygosity of the domesticated strains was attributed to inbreeding or the population-size bottle neck experience by the domesticated strains, as a 90% reduction in the effective population size occurred due to domestication.

Restoration of genetic diversity in silkworm races

B. mori is a completely domesticated insect and has undergone intense artificial selection, (Goldsmith 2005). Therefore, it has to depend on humans for its survival. Years of continuous domestication have made *B. mori* unable to survive in the wild. Therefore, natural selection induced genetic diversity is limited to few traits such as voltinism. Hence, it is essential to conserve and utilize the wild relatives of *B. mori* to broaden the genetic base of the domesticated races apart from utilizing the geographically divergent races, mutants, sex-limited races, evolved breeds, breeder's genetic stocks, and exotic resources. The wild relatives of *Bombyx* are ecologically very vulnerable and the vulnerability at different spatial and temporal scales is not known. The design of biodiversity network in sericulture involving the complementaries of wild relatives and domesticated *B. mori* is also not well established. Therefore, conservation of wild as well as domesticated seribiodiversity resources is very essential for sustainable development of sericulture since loss of genetic resources of domesticated and wild relatives of *Bombyx* species along with their unique genes may be disadvantage for future generations (FAO Manual 2003).

Summary

With the advent of co-dominant DNA markers that are superior to allozymes, the identification of heterozygosity among the populations of races, variety, or breeds of a species became easier. Use of SSR and SNP markers to identify the heterozygosity levels for conservation of genetic material for true-to-type genotype maintenance among germplasm stocks has become a regular practice. The use of DNA markers to relate genetic homozygosity and genetic distance of the parental strains to the manifestation of hybrid vigour, and understanding the genetic

basis of heterosis, will be quite useful to select suitable parental lines for hybridization programmes. Considering the large number of silkworm inbred lines that are maintained in the silkworm breeding stations, it is prudent to use molecular markers to identify parental lines that are most suitable for developing heterotic hybrids. Silkworm strain-specific markers could also be used for authentication of races, protection of breeders' rights, and periodical monitoring of the purity of varieties in the conservation centres. So far, evidence of effects of inbreeding on inbreeding coefficients in silkworm populations is limited. Hence understanding the effects of inbreeding for various traits can be very crucial points in the management of germplasm. Another important objective of silkworm genetic resource conservation is to protect the available seri-biodiversity from extinction. At present only few old Indian indigenous races survive, viz. Barapolu, Chotapolu, Nistari, Sarupat, and Moria; several indigenous races like Ichot, Nishmo, Itan, and the univoltine Kashmiri races are extinct (FAO Manual 2003). Therefore, it is essential to develop appropriate strategies for the conservation of biodiversity not only for the vulnerable populations, but also for sustainable utilisation of silkworm genetic resources. Effective conservation plans need to be developed.

References

-
- Britten HB. 1996. Meta-analyses of the association between multilocus heterozygosity and fitness. *Bulletin of US Department Agriculture* 1090:1-36
- Bush RM, Smouse PE, Ledig FT. 1987. The fitness consequences of multiple locus heterozygosity: The relation-ship between

heterozygosity and growth rate in pitch pine (*Pinus rigida* Mill.). *Evolution* 41: 787-798.

Charlesworth D, Charlesworth B. 1987. Inbreeding depression and its evolutionary consequences. *Annual Review of Ecology and Systematics* 18: 237-268.

Chatterjee S N, Rao CGP, Chatterjee GK, Aswath SK, Patnaik AK. 1993. Correlations between yield and biochemical parameters in the mulberry silkworm *Bombyx mori* L. *Theoretical and Applied Genetics* 87: 385-391.

Chatterjee SN, Pradeep AR. 2003. Molecular markers (RAPD) associated with growth, yield and origin of the silkworm, *Bombyx mori* in India. *Russian Journal of Genetics* 39: 1612-24.

Cheng TC, Xia QY, Qian JF, Liu C, Lin Y, Zha XF, Xiang ZH .2004. Mining single nucleotide polymorphisms from EST data of silkworm, *Bombyx mori*, inbred strain Dazao. *Insect Biochemistry and Molecular Biology* 34: 523-530.

Dalirsefat SB, Mirhoseini S Z. 2007. Assessing genetic diversity in Iranian native silkworm (*Bombyx mori* L.) strains and Japanese commercial lines using AFLP markers. *Iranian Journal of Biotechnology* 5: 25-33.

Dalirsefat S, Meyer A, Mirhoseini S. 2009. Comparison of similarity coefficients used for cluster analysis with amplified fragment

length polymorphism markers in the silkworm, *Bombyx mori*. *Journal of Insect Science* 9:71. Available online: <http://insectscience.org/9.71>.

Darwin C. 1876. *The Effects of Cross and Self Fertilisation in the Vegetable Kingdom*. John Murray.

Datta RK, Nanavaty M. 2007. Global Silk Industry. A.P.H. Publishing Corp.

Falconer DS. 1981. *Introduction to Quantitative Genetics*. Longman Scientific and Technical.

Falconer DS. 1989. *Introduction to Quantitative Genetics*, Longman/Wiley.

FAO 2003. Conservation Status of Sericultural Germplasm Resources in the World-1. Food and Agricultural Organization of the United Nation (FAO), FAO Corporate Document Repository. Available online: <http://www.fao.org/>.

Fisher RA. 1930. *The Genetic Theory of Natural Selection*. Clarendon Press.

Frankel OH. 1984. Genetic perceptiveness of germplasm conservation. In: Arber WK, Llimense K, Peacock WJ, Strarlinger P, editors. *Genetic manipulation: Impact on man and society*, pp-161-170. Cambridge University Press.

Frankel OH, Brown AHD. 1984. Current plant genetic resources- a critical appraisal. In:

Genetics: New frontiers. Oxford and IBH Publication Co.

Gamo T. 1976. Recent concepts and trends in silkworm breeding. *Farming Japan* 10: 11-12

Gamo T, Hirabayashi T. 1983. Genetic analysis of growth rate, pupation rate and some quantitative characters by diallel crosses in silkworm, *Bombyx mori*. L. *Japanese Journal of Breeding* 3: 178-190.

Gaviria DA, Aguilar E, Serrano HJ, Alegria AH. 2006. DNA fingerprinting using AFLP markers to search for markers associated with yield attributes in the silkworm, *Bombyx mori*. *Journal of Insect Science* 6: 15 Available online: <http://insectscience.org/6.15>.

Graner A, Dehmer KJ, Thiel T, Borner A. 2004. Plant genetic resources: benefits and implications of using molecular markers, In: de Vicente MC, editor. *Issues in Genetic Resources* No. 11, pp. 26-32. IPGRI.

Goldsmith M.R, Shimada T, Abe H. 2005. The Genetics and Genomics of the Silkworm, *Bombyx mori*. *Annual Reviews Entomology*, 50: 71-100.

Harada C. 1957. On the relation between commercial characters and their Fi hybrids *Bombyx mori*. *Proceedings of International Genetics Symposium* pp. 352-356.

Harada C. 1961. On the heterosis of quantitative characters in the silkworm.

Bulletin of Sericulture Experimental Station 17: 50-52.

Hardy, G. H. 1908. Mendelian proportions in a mixed population, *Science* 28: 49–50.

Hirobe T. 1968. Evolution, differentiation and breeding of the silkworm the silk road, past and present genetics in Asian countries. pp. 25-36. *XII International Congress Genetics, Tokyo*

Houle D. 1989. Allozyme-associated heterosis in *Drosophila melanogaster*. *Genetics* 123: 789-801.

Jolly MS, Datta RK, Noamani MKR, Iyengar MNS, Nagaraj CS, Basavaraj HK, Kshamarani G, Rao PR. 1989. Studies on the genetic divergence in mulberry silkworm *Bombyx mori* L. *Sericologia* 29: 545-559.

Kovalov PA. 1970. *Silkworm breeding technique*. Translated and published by Central Silk Board. Bangalore, India.

Krishnaswami S, Jolly MS, Subba Rao S. 1964. Diallel analysis of quantitative characters in multivoltine races of silkworm. *Indian Journal of Genetics* 24: 213-222.

Kristensen TN, Pedersen KS, Vermeulen CJ. 2010. Research on inbreeding in the ‘Omic’era. *Trends in ecology and evolution* 25(1): 44-52.

Kumaresan P, Sinha RK, Thangavelu K. 2003a. Heterosis studies in some elite multivoltine silkworm (*Bombyx mori* L.) races

with popular bivoltine NBD2. *International Journal of Industrial Entomology* 7: 221-229

Kumaresan P, Mahadevamurthy TS, Thangavelu K, Sinha RK. 2003b. Further studies on the genetic divergence of multivoltine silkworm (*Bombyx mori* L.) genotypes based on economic characters. *Entomon* 28: 193-198.

Kumaresan P, Mohan B, Koundinya PR, Sinha RK, Thangavelu K. 2004a. Silkworm germplasm conservation-A perspective. *Indian Silk* 44: 4-9.

Kumaresan P, Sinha RK, Mohan B, Thangavelu K. 2004b. Conservation of multivoltine silkworm (*Bombyx mori* L.) germplasm in India-An overview. *International Journal of Industrial Entomology* 9: 1-13.

Kwok PY, Carlson C, Yager TD, Ankener W, Nickerson DA. 1994. Comparative analysis of human DNA variations by fluorescence-based sequencing of PCR products. *Genomics* 23: 138-144.

Lewontin RC, Hubby JL. 1966. A molecular approach to the study of heterozygosity in natural populations II. Amount of variation and degree of heterozygosity in natural populations of *Drosophila pseudoobscura*. *Genetics* 54: 595-609.

Li MW, Shen L, Xu AY, Miao XX, Hou CX, Sun PJ, Zang YH, Huang YP. 2005. Genetic diversity among the silkworm (*Bombyx mori*

L., Lep, Bombycidae) germplasm revealed by microsatellites. *Genome* 48: 802-810.

Lynch M and Walsh B. 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer Associates.

Mitton JB, Schuster WSF, Cothran EG, De Fries JC. 1993. Correlation between the individual heterozygosity of parents and their offspring. *Heredity* 71: 59-63

Mitton JB. 1994. Molecular approaches to population biology. *Annual Review Ecology and Systematics* 25: 45-69.

Mukharjee P, Mukharjee S, Kumaresan P. 1999. An analysis of genetic divergence in Indian (multivoltine silkworm *Bombyx mori*) germplasm. *Sericologia* 39: 337-347.

Nagaraja GM, Nagaraju J. 1995. Genome fingerprinting in silkworm *Bombyx mori* using random arbitrary primers. *Electrophoresis* 16: 1633-1638.

Nagaraju J. 1990. Studies on some genetic aspects of quantitative characters in tropical silkworm *Bombyx mori*. Ph.D. Thesis, University of Mysore.

Nagaraju J. 2002. Application of genetic principles for improving silk production. *Current Science* 83: 409-414.

Nagaraju J, Kumar P. 1995. Effects of selection on cocoon filament length in divergently selected lines of the silkworm,

Bombyx mori. *Journal of Sericulture Science of Japan* 64: 103-109

Nagaraju J, Reddy KD, Nagaraja GM, Sethuraman BN. 2001. Comparison of multilocus RFLPs and PCR-based marker systems for genetic analysis of the silkworm *Bombyx mori*. *Heredity* 86: 588-597.

Nagaraju J, Goldsmith MR. 2002. Silkworm genomics - Progress and Prospects. *Current Science* 83: 415-425.

Natcheva Y, Petkov N, Ignatova L, Tzenov P, Braslavskii M, Stostkii M, Liushenko J. 2001. Correlations between some characters in silkworm outbred populations. *Sericulture (Ukraine)* 23: 29-34.

Petkov N, Natcheva Y, Tzenov P. 1999. About the problem of applying inbreeding to silkworm *Bombyx mori* L. *Agricultural Science* 1: 37-39.

Pradeep AR, Chatterjee SN, Sarathchandra B, Raje Urs S. 2005. Allelic variants of a juvenile hormone responsive gene which connote genetic differentiation in strains of the silkworm *Bombyx mori*. *Journal of Genetics and Breeding* 59: 213-224.

Prasad MD, Muthulakshmi M, Madhu M, et al. 2005. Survey and analysis of microsatellites in the silkworm, *Bombyx mori*: Frequency, distribution, mutation, marker potential and their conservation in heterologous species. *Genetics* 169: 197-214.

Raju PJ, Krishnamurthy NB. 1993. Breeding of two bivoltines, MG 511 and MG512, of silkworm *Bombyx mori* L. for higher viability and silk productivity. *Sericologia* 33: 577-587.

Ralls K, Ballou JD, Templeton A. 1988. Estimation of lethal equivalents and the cost of inbreeding in mammals. *Conservation Biology* 2: 185-193.

Reddy KD, Abraham EG, Nagaraju J. 1999a. Microsatellites in the silkworm *Bombyx mori*: Abundance, polymorphism, and strain characterisation. *Genome* 42: 1057-1065.

Reddy KD, Nagaraju J, Abraham EG. 1999b. Genetic characterization of the silkworm *Bombyx mori* by simple sequence repeat (SSR) anchored PCR. *Heredity* 83: 681-87.

Shull GH. 1952. Beginnings of the heterosis concept. In: Gowen JW, editor. *Heterosis*, pp. 14-48. Iowa State College Press.

Sokal RR, Rohlf FJ. 1995. *Biometry*, 3rd edition. W.H. Freeman.

Staykova T. 2008. Genetically-determined polymorphism of nonspecific esterases and phosphoglucomutase in eight introduced breeds of the silkworm, *Bombyx mori*, raised in Bulgaria. *Journal of Insect Science* 8:18. Available online: <http://insectscience.org/8.18>.

Subramanya G, Bishop S. 2009. A novel procedure for estimating inbreeding coefficient (F) in the inbred populations of mulberry silkworm *Bombyx mori* L.

Proceedings of National workshop on Seribiodiversity Conservation, 7-8 March 2009.

Tamura T, Thibert C, Royer C, Kanda T, Abraham E, et al. 2000. Germline transformation of the silkworm *Bombyx mori* L. using a *piggyBac* transposon-derived vector. *National Biotechnology* 18: 81–84.

Thangavelu K. 1998. Silkworm breeding in India-at a cross road. In: Subramanya G, editor. *Silkworm Breeding*. Oxford & IBH Publishing Co. Pvt, Ltd.

The International Silkworm Genome Consortium/Insect Biochemistry and Molecular Biology, 38, 2008, 1036-1045.

Tomita M, Munetsuna H, Sato T, Adachi T, Hino R, et al. 2003. Transgenic silkworms produce recombinant human type III procollagen in cocoons. *National Biotechnology* 21: 52–56.

Toyama. 1906. Breeding method of silkworms (Jap). *Sangyo-Shimpo* 158: 283-286.

United Nations Publication website, ESCAP. 1993. Principles and Techniques of Silkworm Breeding. Available online: <http://unp.un.org/>.

Velu D, Ponnuvel KM, Muthulakshmi M, Sinha RK, Qadri SMH. 2008. Analysis of genetic relationship in mutant silkworm strains of *Bombyx mori* using inter-simple sequence repeat (ISSR) markers. *Journal of Genetics and Genomics* 35: 1-7.

Whitlock MC. 2002. Selection, load, and inbreeding depression in a large metapopulation. *Genetics* 160: 1191-1202.

Willis JH, Wilkins AS, Goldsmith MR. 1995. A brief history of Lepidoptera as model systems. In: Goldsmith MR, Wilkins AS, editors. *Molecular Model Systems in the Lepidoptera*, pp. 1–20. Cambridge University Press.

Wright S. 1922. Coefficients of inbreeding and relationship. *American Naturalist* 56: 330–338.

Wright S. 1977. Evolution and Genetics of populations. Vol. 3. *Experimental Results and Evolutionary Deductions*. University of Chicago Press.

Xia Q, Guo Y, Zhang Z, Li D, Xuan Z, Li Z, Dai F. et al. 2009. Complete resequencing of 40 genomes reveals domestication events and genes in silkworm (*Bombyx*)., *Science* 326: 433-436. Available online: www.sciencemag.org. [DOI:10.1126/science.1176620]

Xia QY, Zhou ZY, Lu C, Xiang ZH. 1998. Molecular Phylogenetic study on the racial differentiation of *Bombyx mori* by random amplified polymorphic DNA (RAPD) markers. *Acta Entomologica Sinica* 41: 32-40.

Yokoyama T. 1979. In: Hoy MA, McKelvey JJ Jr., editors. *Genetics in Relation to Insect*

Mangement, pp. 71-83. The Rockefeller
Foundation Press.