

Utility of Multi-Gene Loci for Forensic Species Diagnosis of Blowflies

Authors: Zaidi, Farrah, Wei, Shu-jun, Shi, Min, and Chen, Xue-xin

Source: Journal of Insect Science, 11(59) : 1-12

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.011.5901>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Utility of multi-gene loci for forensic species diagnosis of blowflies

Farrah Zaidi^{1, 2a}, Shu-jun Wei^{1b}, Min Shi^{1c}, Xue-xin Chen^{1d*}

¹Institute of Insect Sciences, Zhejiang University, Hangzhou 310029, China

²Department of Zoology, University of Peshawar, Peshawar 25120, Pakistan

Abstract

Contemporary studies in forensic entomology exhaustively evaluate gene sequences because these constitute the fastest and most accurate method of species identification. For this purpose single gene segments, cytochrome oxidase subunit I (COI) in particular, are commonly used. However, the limitation of such sequences in identification, especially of closely related species and populations, demand a multi-gene approach. But this raises the question of which group of genes can best fulfill the identification task? In this context the utility of five gene segments was explored among blowfly species from two distinct geographic regions, China and Pakistan. COI, cytochrome b (CYTB), NADH dehydrogenase 5 (ND5), nuclear internal transcribed spacers (ITS1 and ITS2), were sequenced for eight blowfly species including *Chrysomya megacephala* F. (Diptera: Calliphoridae), *Ch. pinguis* Walker, *Lucilia sericata* Meigen, *L. porphyrina* Walker, *L. illustris* Meigen, *Hemipyrellia ligurriens* Wiedemann, *Aldrichina grahamsi* Aldrich, and the housefly, *Musca domestica* L. (Muscidae), from Hangzhou, China; while COI, CYTB, and ITS2 were sequenced for four species, i.e. *Ch. megacephala*, *Ch. rufifacies*, *L. cuprina*, and the flesh fly, *Sarcophaga albiceps* Meigen (Sarcophagidae), from Dera Ismail Khan Pakistan. The results demonstrate a universal utility of these gene segments in the molecular identification of flies of forensic importance.

Keywords: Calliphoridae, cytochrome b, molecular identification, NADH dehydrogenase 5, nuclear internal transcribed spacers

Abbreviations: **COI**, cytochrome oxidase subunit I; **CYTB**, cytochrome b; **MP**, maximum parsimony; **ML**, maximum likelihood; **ITS1**, **ITS2**, nuclear internal transcribed spacers 1 and 2; **ND5**, NADH dehydrogenase 5

Correspondence: ^a zaidi_farah@yahoo.com, ^b shujun268@163.com, ^c shimin0623@zju.edu.cn, ^{d*} xxchen@zju.edu.cn, *Corresponding author

Editor: Zhijian Jake Tu was editor of this paper

Received: 12 February 2010, **Accepted:** 24 September 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 11, Number 59

Cite this paper as:

Zaidi F, Wei S-j, Shi M, Chen X-x. 2011. Utility of multi-gene loci for forensic species diagnosis of blowflies. *Journal of Insect Science* 11:59 available online: insectscience.org/11.59

Introduction

Blowflies are the first insects to arrive at the scene of death, often seen ovipositing on the cadaver during the first few hours after death (Smith 1986). The growth periods, developmental rates (van Laerhoven 2008), and diapause responses (Ash and Greenberg 1975) are all substantially different for closely related species. These peculiar features make blowflies the primary and most accurate indicators of the post mortem interval (Grassberger et al. 2003). However, the immature stages pose a great identification challenge for forensic entomologists due to their lack of species-specific anatomical characters (McDonagh et al. 2009). Identification of these stages using traditional methods (Mendonca et al. 2008) and/ or advanced techniques (McDonagh et al. 2009) is therefore the focus of many forensic studies. The most popular method of identification in recent years includes molecular taxonomy (Harvey et al. 2003; Smith and Baker 2008). In this regard several loci have been explored for their phylogenetic utility, but mitochondrial cytochrome oxidase subunit I (COI) (Harvey et al. 2003; Ratcliffe et al. 2003; Nelson et al. 2007; Ying et al. 2007; Smith and Baker 2008) is often used, sometimes along with subunit II (COII) (Ratcliffe et al. 2003; Ying et al.,2007). The 5'

end of COI is also the site of the proposed universal animal DNA “barcode” (Hebert et al. 2003). COI barcodes have been successfully utilized in the identification of many blowfly species (Nelson et al. 2007). However, the barcoding approach has its own limitations (Moritz and Cicero 2004) as is seen with its failure to identify some closely related species of blowflies (Nelson et al. 2007; Whitworth et al. 2007). Thus, COI barcodes, or for that matter any single gene, seems unlikely to resolve the identities of all calliphorid species of forensic importance because such phylogenies only infer the evolutionary relationships for the particular gene used. Consequently, a switch to multi-gene approach becomes necessary (McDonagh et al. 2009). To date only a handful studies have utilized such an approach (Wallman et al. 2005).

Therefore, a multi-gene approach was employed for the identification of blowfly species of forensic importance from Hangzhou, China and Dera Ismail Khan, Pakistan. Gene data was freshly generated for a novel combination of five loci, including the barcode region of mitochondrial cytochrome oxidase subunit I (COI) (Hebert et al. 2003), NADH dehydrogenase 5 (ND5), cytochrome b (CYTB), and nuclear internal transcribed spacers 1 and 2 (ITS1 and ITS2).

Table 1. Species collected from Hangzhou, China and Dera Ismail Khan, Pakistan

Species	Family	Location
<i>Chrysomya megacephala</i> (Fabricius)	Calliphoridae	Hangzhou, China
<i>Chrysomya pinguis</i> (Walker)	Calliphoridae	Hangzhou, China
<i>Aldrichina grahami</i> (Aldrich)	Calliphoridae	Hangzhou, China
<i>Lucilia sericata</i> (Meigen)	Calliphoridae	Hangzhou, China
<i>Lucilia porphyrina</i> (Walker)	Calliphoridae	Hangzhou, China
<i>Lucilia illustris</i> (Meigen)	Calliphoridae	Hangzhou, China
<i>Hemipyrellia liguriensis</i> (Wiedemann)	Calliphoridae	Hangzhou, China
<i>Musca domestica</i> (Linnaeus)	Muscidae	Hangzhou, China
<i>Chrysomya megacephala</i> (Fabricius)	Calliphoridae	Dera Ismail Khan, Pakistan
<i>Chrysomya ruffiacis</i> (Macquart)	Calliphoridae	Dera Ismail Khan, Pakistan
<i>Lucilia cuprina</i> (Wiedemann)	Calliphoridae	Dera Ismail Khan, Pakistan
<i>Sarcophaga albiceps</i> Meigen	Sarcophagidae	Dera Ismail Khan, Pakistan

Materials and Methods

Sample collection and Identification

Samples of blowflies were collected from Huajiachi, Hangzhou over the period March 2007 to June 2008, while in Dera Ismail Khan sampling was carried out in December 2008. Identification was performed using morphological keys (Fan 1992; Fan 1997;

Table 2. New sequences of flies from Hangzhou, China and Dera Ismail Khan, Pakistan

Species/individuals	GenBank accession numbers				
	ITS1	COI	CYTB	ITS2	ND5
Hangzhou, China					
<i>Chrysomya megacephala</i> 1	FJ790779	FJ614818	FJ614835	FJ614850	FJ614869
<i>Chrysomya megacephala</i> 2	FJ790780	FJ614817	FJ614834	FJ614851	FJ614868
<i>Chrysomya megacephala</i> 3	FJ790781	FJ614816	FJ614833	FJ614852	FJ614867
<i>Chrysomya pinguis</i> 1	-	FJ614820	FJ614837	FJ614853	FJ614871
<i>Chrysomya pinguis</i> 2	-	FJ614819	FJ614836	FJ614854	FJ614870
<i>Hemipyrellia ligurriens</i> 1	FJ790782	FJ614823	FJ614840	FJ614856	FJ614874
<i>Hemipyrellia ligurriens</i> 2	-	FJ614822	FJ614839	FJ614855	FJ614873
<i>Hemipyrellia ligurriens</i> 3	-	FJ614821	FJ614838	FJ614857	FJ614872
<i>Lucilia sericata</i> 1	-	FJ614824	FJ614841	FJ614858	FJ614875
<i>Lucilia sericata</i> 2	-	FJ614825	FJ614842	FJ614859	FJ614876
<i>Lucilia sericata</i> 3	-	FJ614826	FJ614843	FJ614860	FJ614877
<i>Lucilia illustris</i> 1	FJ790783	FJ614827	FJ614844	FJ614861	FJ614878
<i>Lucilia illustris</i> 2	-	FJ614828	FJ614845	FJ614862	FJ614879
<i>Lucilia porphyrina</i> 1	FJ790784	FJ614829	FJ614846	FJ614863	FJ614880
<i>Lucilia porphyrina</i> 2	-	FJ614830	FJ614847	FJ614864	FJ614881
<i>Aldrichina grahami</i> 1	FJ790785	FJ614831	FJ614848	FJ614866	FJ614882
<i>Aldrichina grahami</i> 2	-	FJ614832	FJ614849	FJ614865	FJ614883
Dera Ismail Khan, Pakistan					
<i>Sarcophaga albiceps</i>	-	GQ912668	GQ912671	GQ912697	-
<i>Lucilia cuprina</i>	-	GQ912667	GQ912670	GQ912696	-
<i>Chrysomya megacephala</i>	-	GQ912669	GQ912672	GQ912698	-
<i>Chrysomya ruffifacies</i> 1	-	GQ912645	GQ912673	GQ912674	-

Senior-White et al. 1940). The identified species are listed in Table 1.

DNA extraction, amplification and sequencing

Twenty-six individuals including 22 blowflies (7 species) from Hangzhou and 4 individuals (including 4 species) from Dera Ismail Khan (Table 1) were used for DNA extraction, amplification, and sequencing. Total DNA was extracted from leg and thorax regions of each adult fly using the DNeasy Tissue Kit (QIAGEN, www.qiagen.com) following the manufacturer's protocol. Previously reported primers along with protocols of polymerase chain reaction (PCR) conditions were used to amplify the COI, ITS2 (Nelson et al. 2007), ND5 (Zehner et al., 2004), ITS1, and CYTB (Kengne et al. 2007). Thermal cycling was performed in a Mastercycler (Eppendorf, www.eppendorf.com). Amplified products were purified and sequenced by Shanghai Invitrogen (www.invitrogen.com). The newly

sequenced blowfly specimens from Hangzhou and Dera Ismail Khan with their related GenBank accession numbers for ITS1, COI, ITS2, ND5, and CYTB are listed in Table 2. The previously published sequences utilized during the present study are presented in Table 3.

Sequence analysis

All sequences were partial except for ITS2. Each COI, CYTB, and ND5 sequence represents part of the corresponding genes of

Table 3. Previously published COI sequences used in our phylogenetic analysis

Species	GenBank accession numbers
<i>Chrysomya pinguis</i>	EU661327, EU661325, EU661326
<i>Chrysomya ruffifacies</i> (Macquart)	-
<i>Chrysomya megacephala</i>	AJ426041, EU661323, EU661322, EU661321
<i>Hemipyrellia ligurriens</i>	EU661328
<i>Hemipyrellia pulchra</i> (Wiedemann)	-
<i>Lucilia porphyrina</i>	EU661329, EU661330, EU661331
<i>Lucilia illustris</i>	-
<i>Lucilia bazini</i> (Seguy)	-
<i>Lucilia sericata</i>	AJ422212
<i>Lucilia cuprina</i> (Wiedemann)	-
<i>Aldrichina grahami</i>	EU661320
<i>Calliphora vicina</i> (Robineau-Desvoidy)	-

M. domestica (GenBank accession EU154477) with COI pertaining to positions 1385-2040 (655 bp), CYTB to 10722-11192 (470 bp), and ND5 to 6340-6785 (445 bp) while ITS-II with its 422 bp length was completely sequenced. ITS1 sequences were not used in the phylogenetic analysis due to the difficulties in alignment as well as an inadequate data set. Nevertheless, ITS1 was sequenced for most of the species from Hangzhou with the legible length ranging between 400 to 700 bp.

The software Dnadist (Felsenstein 1989) was used to compute distance matrices for nucleotide sequences of different gene segments and graphs were plotted using the same data.

Chromas Pro 34-Version 1.33 (available online, www.technelysium.com.au/ChromasPro.html) was used for manual editing of sequences that were subsequently aligned in CLUSTAL X using default parameter settings.

Phylogenetic analyses were performed using maximum parsimony (MP) with PAUP* 4.0b10 (Swofford 2002) and maximum likelihood (ML) with PhyML (Guindon et al. 2005). The MP analyses were run with default

heuristic search options except that 100 replicates of random stepwise additions were used. Models of DNA substitution were estimated in Modeltest 3.7 (Posada and Crandall 1998). For ML we used GTR + G model for COI, Cytb and ND5, and F81+G model for ITSII nucleotide sequences. Bootstrap proportions were obtained after 1000 replicates by using 10 replicates of random stepwise additions of taxa.

Results and Discussion

Sequence data were generated for 12 species, including 10 blowflies, from two geographic zones of Pakistan and China.

Sequence variability for ITS

The transcribed spacer ITS2 has been sequenced recently and successfully utilized in the molecular identification of blowflies (Nelson et al. 2007; Song et al. 2008). However, ITS1 has not been sequenced before for blowflies. This region was found to be highly variable and was amplified with great difficulty. The species *L. sericata*, *L. cuprina*, *Ch. rufifacies*, and *S. albiceps* posed difficulties in amplification. Additionally, the baseline noise in chromatograms made the ITS1 sequences unreliable for *Ch. pinguis*. In contrast, ITS2 was sequenced without any problems.

Intra-specific sequence variation

The ITS 1 intra-specific sequence divergence was computed only for *Ch. megacephala*. The variation was in a reasonable range, i.e. 0.29 - 1.16 %, and was comparable to the mitochondrial segments of COI and CYTB genes (Table 4). However, previous studies usually showed no intra-specific variation in the Drosophilidae for ITS1 region (Baffi and Ceron 2002).

Table 4. Intra-specific ITS1, COI and CYTB divergence among *Chrysomya megacephala* individuals

ITS1 Sequence Divergence			
Species/ individuals	<i>Ch. megacephala</i> 1	<i>Ch. megacephala</i> 2	<i>Ch. megacephala</i> 3
<i>Chrysomya megacephala</i> 1	0	-	-
<i>Chrysomya megacephala</i> 2	1.20%	0	-
<i>Chrysomya megacephala</i> 3	0.90%	0.30%	0
COI Sequence Divergence			
Species/ individuals	<i>Ch. Megacephala</i> 1	<i>Ch. Megacephala</i> 2	<i>Ch. Megacephala</i> 3
<i>Chrysomya megacephala</i> 1	0	-	-
<i>Chrysomya megacephala</i> 2	0	0	-
<i>Chrysomya megacephala</i> 3	0.16%	0.16%	0
CYTB Sequence Divergence			
Species/ individuals	<i>Ch. Megacephala</i> 1	<i>Ch. Megacephala</i> 2	<i>Ch. Megacephala</i> 3
<i>Chrysomya megacephala</i> 1	0	-	-
<i>Chrysomya megacephala</i> 2	0	0	-
<i>Chrysomya megacephala</i> 3	0.24%	0.24%	0


The ITS2 region showed no intra-specific variation for most of the blowfly species except for *L. sericata*. This result seems to be consistent with an earlier study that suggested that ITS2 cannot be utilized in differentiation of geographical populations of some blowfly species (Song et al. 2008).

ITS1 may be a good tool for identification at the population level. The most important reason is the high mutation rate that cause deletions or insertions. A prominent indel characteristic (a 184 bp long deletion) was detected in one of the *Ch. megacephala* sequences (FJ790781). The size and location suggested a loop deletion. Loop deletions, when present in coding regions, might have deleterious effects on the organism like any other mutation (Stefanovic et al. 2007), but their exact impact on non-coding regions such as ITS is unknown. In any case, use of indels (i.e. insertions and or deletions) as genetic markers has been recommended in

phylogenetic studies of natural populations (Väli et al. 2008). Thus a high expectancy of insertion/deletions among ITS1 sequences might make them useful in identification of geographic populations of blowflies of forensic importance.

Inter-specific sequence variation

Inter-specific variation was explored for a wide array of paired species (blowfly-blowfly; blowfly-fleshfly, and blowfly-housefly). Both of the ITS regions were found highly variable as compared to mtDNA. The ITS1 segment was abnormally divergent among all gene segments (Supplement Table S1), ranging between 7.3-7.8 % (Figure 1). The results of the present study are consistent with previous studies of Dipteran families, such as Tephritidae, which also revealed little evidence of similarity between ITS1 sequences of species, especially those belonging to different genera (Douglas and Haymer 2001). The most likely reason seems to be the rapidly evolving and mutating nature of non-coding regions, such as ITS (Haymer 1994). On the other hand, ITS1 showed little variation among members of the same species as shown by *Ch. megacephala* individuals (Table 4). These results also showed no overlapping between intra- and inter-specific variations for ITS1. The ITS1 intra-specific variation as recorded for *Ch. megacephala* during the present study was 0.24 -1.16 % (Table 4), while inter-specific variation between different species pairs was in the range of 8- 60 % (Supplement Table S1). However, intra- and inter-specific overlapping seems plausible for ITS2 when the smallest inter-specific ITS2 value of divergence is recorded at 1.2 % for *Ch. megacephala* and *Ch. pinguis* (Supplement Table S1). Similar overlapping is previously observed among blowflies (Song et al. 2008), though without any taxonomic impact in that study,






Figure 2. Intra-specific ND5 sequence variation between Hangzhou and other regions. Note: Transitions are the most prominent type of mutations among ND5 sequences. High quality figures are available online.

nevertheless overlapping can negatively influence the identification process. However, at the family level ITS2 possessed strong resolution power as it showed higher values of variation between blowfly-flesh fly pairs (14.7-21.1 %) and blowfly-housefly pairs (30-35.2 %), as compared to blowfly-blowfly pairs (Supplement Table S1). This result is also consistent with previous studies (Song et al. 2008). Another important finding of the present study was abnormally high ITS1 divergence values for sister species *L. porphyrina* and *L. illustris*, i.e. 22 % (Figure 1). At other gene loci, including ITS2, the divergence values between these two species were comparatively low, but adequate for differentiation (4-6 %, Figure 1).

Sequence variability for mt DNA

Mitochondrial gene segments showed less sequence variability as compared to the ITS regions although sequence divergence values were almost comparable between CYTB and ITS2 (Figure 1). Insertion and deletions were comparatively fewer in mtDNA genes.

Intra-specific sequence variation

These results showed a mean of 0.15% COI intra-specific variation for *Ch. megacephala*.

The results are similar to those for *Chrysomya* species of blowflies (0.097 %) reported by Nelson et al. (2007). The mt gene CYTB showed variation of 0.24 % among both *Ch. megacephala* and *L. sericata* individuals. On the other hand, ND5 sequences of blowflies were highly conserved with no intra-specific variation. However, comparisons with ND5 sequences from other regions of China that recently became publicly available (Table 3) showed significant variation among the transition/transversion ratios and the indel characteristics between species from Hangzhou and their counterparts in other Chinese cities (Figure 2). These results show the presence of distinct geographic populations of blowflies or perhaps cryptic species. Mutations such as insertions, deletions, transitions, and transversions (Figure 2) are the result of, or an indicator of, divergence of natural populations, making them an important tool in the population identification studies (Väli et al. 2008).

Inter-specific sequence variation

The inter-specific variation for mtDNA segments, including COI, CYTB, and ND5, was found suitable for blowfly species diagnosis. However, for some sister species




Figure 3. Maximum-parsimony phylogram based on COI (A), CYTB (B), and ITS2 (C) sequences of blowflies from Dera Ismail Khan, Pakistan and Hangzhou, China. High quality figures are available online.

this variation was considerably low, e.g. 0.78 % of COI (*L. sericata* vs *L. cuprina*) and 0.5 % of CYTB (*L. sericata* vs *L. cuprina*). Even ITS2 that provided a least variation of 3.5 % for the *L. sericata* and *L. cuprina* pair showed low variation for another blowfly species pair, i.e. *Ch. megacephala* and *Ch. pinguis* (1.2 %). These results justify the need of a novel molecular marker for sister species diagnosis.

Phylogenetic analyses

In an attempt to compare blowfly species from two distinct climatic zones, single (Figures 3A, B, C) and multi-gene trees (Figures 4A, B; Figure 5) based on COI, CYTB, ND5, and ITS2 sequences were constructed utilizing maximum parsimony and maximum likelihood methods. Both approaches identified the 11 fly species of forensic




Figure 4. Maximum parsimony (A) and maximum likelihood (B) phylograms based on COI, CYTB, and ITS2 sequences of blowflies from Dera Ismail Khan, Pakistan and Hangzhou, China. High quality figures are available online.

importance from China and Pakistan. This is the first time that the molecular data from the regions of Dera Ismail Khan, Pakistan and Hangzhou, China are presented and compared for blowflies.

Single vs. multi-gene approach

The significance of single genes in molecular forensics is undeniable (Nelson et al. 2007; Wells et al. 2007; Song et al. 2008). Nonetheless, additional genes become essential in challenging identifications (Nelson et al. 2007). A reasonable group of genes not only clarify doubts and provides validation, but also shed light on new evolutionary relationships. A unique instance is that of *A. grahamsi*. Grouped together with




Figure 5. Maximum-likelihood phylogram based on COI, CYTB, ND5, and ITS2 sequences of blowflies. High quality figures are available online.

Lucilini tribe in the COI gene tree (Figure 3A) this species presented a deviation from traditional taxonomy because this species (Calliphorinae) is identified as sisters with Chrysomyiinae rather than Luciliinae (Rognes 1997). Contrary to COI, the CYTB and ITS2 trees (Figures 3b, c) followed this classification. These two distinct patterns of evolution were also observed previously for blowfly phylogenies based on 28rRNA sequences (Stevens and Walls 2001). The

multi-gene trees validated both views. The 4-gene ML tree (Figure 5) agreed with the classical taxonomy while the 3-gene MP and ML trees (Figures 4A, B) displayed the alternative scheme.

However, each of the single gene trees, although they adequately identified blowflies, they also showed some limitations in terms of evolutionary relationships between species. For example, the ITS2 tree incorrectly put *H. ligurriens* within genus *Lucilia* (Figure 3C). Previously another single gene phylogeny based on the COI locus also grouped this species with *L. cuprina* (Wells et al. 2007). The multi-gene trees made amendments by supporting the monophyletic status of genus *Hemipyrellia* (Figures 4A, B). *Chrysomya rufifacies* provided another example of misplacement by single gene tree, CYTB in this case (Figure 3B). The 3-gene ML tree clarified this ambiguity by rightly placing this species along with other species of genus *Chrysomya* (Figure 4B).

These results demonstrate a universal utility of the respective five gene segments. Though single genes exhibited easy identification, the wrong placing of some species illustrates the possibility of future misdiagnosis. Therefore a switch to a multi-gene tactic was made with future work aiming at the examination of more variable segments and species from distinct and multiple geographic regions.

Acknowledgements

We thank two anonymous reviewers for their valuable comments and suggestions. The study has been supported jointly by the grants from National Science Fund for Distinguished Young Scholars (No. 30625006), the National Basic Research Program of China (973 Program) (No. 2006CB102005), and National

S&T Infrastructure Platforms (Nos. 2005DKA21105, 2005DKA21402).

References

Ash N, Greenberg B. 1975. Developmental temperature responses of the sibling species *Phaenicia sericata* and *Phaenicia pallescens*. *Annals of the Entomological Society of America* 68: 197.

Baffi MA, Ceron CR. 2002. Molecular analysis of the rDNA ITS-1 Intergenic Spacer in *Drosophila mulleri*, *D. arizonae*, and their hybrids. *Biochemical Genetics* 40: 411-421.

Caenazzo L, Ceola F, Ponzano E, Novelli E. 2008. Human identification analysis to forensic purposes with two mitochondrial markers in polyacrilamide mini gel. *Forensic Science International: Genetics Supplement Series* 1(1): 266-268.

Douglas LJ, Haymer DS. 2001. Ribosomal ITS1 polymorphisms in *Ceratitis capitata* and *Ceratitis rosa* (Diptera: Tephritidae). *Annals of the Entomological Society of America* 94: 726-731.

Fan ZD. 1992. *Key to the common flies of China*. Academia Sinica, Shanghai Institute of Entomology.

Fan ZD. 1997. *Fauna Sinica, Insecta, Diptera: Calliphoridae*, Volume 6. Science Press Co. Ltd, Chinese Academy of Sciences.

Grassberger M, Friedrich E, Reiter C. 2003. The blowfly *Chrysomya albiceps* (Wiedemann) (Diptera: Calliphoridae) as a new forensic indicator in Central Europe. *International Journal of Legal Medicine* 117: 75-81.

Guindon S, Lethiec F, Duroux P, Gascuel O. 2005. PHYML Online—a web server for fast maximum likelihood-based phylogenetic inference. *Nucleic Acids Research*, 33(suppl 2): W557-W559.

Harvey ML, Mansell MW, Villet MH et al. 2003. Molecular identification of some forensically important blowflies of southern Africa and Australia. *Medical and Veterinary Entomology* 7: 363-369.

Haymer DS. 1994. Random amplified polymorphic DNAs and Micro satellites: What are they, and can they tell us anything we don't already know? *Annals of the Entomological Society of America* 87: 717-722.

Hebert PD, Cywinska A, Ball SL et al. 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B* 270: 313-321.

Kengne P, Antonio-Nkondjio C, Awono-Ambene HP et al. 2007. Molecular differentiation of three closely related members of the mosquito species complex, *Anopheles moucheti*, by mitochondrial and ribosomal DNA polymorphism. *Medical and Veterinary Entomology* 21: 177-182.

McDonagh L, Thornton C, Wallman JF et al. 2009. Development of an antigen-based rapid diagnostic test for the identification of blowfly (Calliphoridae) species of forensic significance. *Forensic Science International: Genetics* 3(3): 162-165.

Mendonca PM, Santos-Mallet JR, Mello RP et al. 2008. Identification of fly eggs using scanning electron microscopy for forensic investigations. *Micron* 39: 802-807.

Moritz C, Cicero C. 2004. DNA barcoding: Promise and pitfalls. *PLoS Biology* 2: 1529-1531.

Nelson LA, Wallman JF, Dowton M. 2007. Using Coi Barcodes to identify forensically and medically important blowflies. *Medical and Veterinary Entomology* 21: 44-52.

Posada D, Crandall KA. 1998. Modeltest: Testing the Model of DNA Substitution, *Bioinformatics* 14: 817-818.

Ratcliffe ST, Webb DW, Weinzievl RA et al. 2003. PCR-RFLP identification of Diptera (Calliphoridae, Muscidae and Sarcophagidae)- a generally applicable method. *Journal of Forensic Science* 48: 783-785.

Senior-White R, Aubertin D, Smart J. 1940. *The fauna of British India, including remainder of the Oriental region. Diptera VI. Family Calliphoridae.* Taylor and Francis.

Smith JA, Baker NC. 2008. Molecular genetic identification of forensically important flies in the UK. *Forensic science International: Genetics Supplement Series* 1(1): 620-622.

Smith KGV. 1986. *A manual of forensic entomology.* The British Museum (Natural History) and Cornell University Press.

Song Z, Wang X, Liang G. 2008. Species identification of some common necrophagous flies in Guangdong province, southern China based on the rDNA internal transcribed spacer 2 (ITS2). *Forensic Science International* 175(1): 17-22.

Stefanovic M, Markham NO, Parry EM et al. 2007. An 11-amino acid β -hairpin loop in the cytoplasmic domain of band 3 is responsible

for ankyrin binding in mouse erythrocytes, *Cell Biology. Proceedings of the National Academy of Sciences USA* 104: 13972-13977.

Stevens J, Wall RL. 2001. Genetic relationships between blowflies (Calliphoridae) of forensic importance. *Forensic Science International* 120(1): 116-123.

Swofford DL. 2002. PAUP*, *Phylogenetic Analysis Using Parsimony (*and Other Methods)*, Version 4, Sinauer Associates, Sunderland, Massachusetts.

Väli U, Brandström M, Johansson M et al. 2008. "Insertion-deletion polymorphisms (indels) as genetic markers in natural populations". *BMC Genetics* 9: 8.

van Laerhoven SL. 2008. Blind validation of postmortem interval estimates using developmental rates of blow flies. *Forensic Science International* 180: 76-80.

Wallman JF, Leys R, Hogendoorn K. 2005. Molecular systematics of Australian carrion-breeding blowflies (Diptera: Calliphoridae) based on mitochondrial DNA. *Invertebrate Systematics* 19: 1-15.

Wells JD, Wall R, Stevens JR. 2007. Phylogenetic analysis of forensically important *Lucilia* flies based on cytochrome oxidase I sequence: a cautionary tale for forensic species determination. *International Journal of Legal Medicine* 121: 229-233.

Whitworth TL, Dawson RD, Magalon H et al. 2007. DNA barcoding cannot reliably identify species of the blowfly genus *Protophthora* (Diptera: Calliphoridae). *Proceedings of Biological Sciences* 274: 1731-1739.

Ying BW, Liu TT, Fan H et al. 2007. The application of mitochondrial DNA cytochrome oxidase II gene for the identification of forensically important blowflies in western China. *The American Journal of Forensic Medicine and Pathology* 28: 308-313.

Zehner R, Amendt J, Schutt S et al. 2004. Genetic identification of forensically important flesh flies (Diptera: Sarcophagidae). *International Journal of Legal Medicine* 118: 245-247.

Supplement Table S1. Pair-wise comparison of inter-specific variation among different gene segments

Species	% sequence variation				
	ITS1	NDS	ITS2	COI	9:33 PM
<i>Ch. megocephala</i> vs <i>Ch. pinguis</i>	-	5	1.2	2.2-2.5	5.2- 5.5
<i>Ch. megocephala</i> vs <i>L. illustris</i>	50-35	7.7	14.6-14.9	8.3-8.5	12.2-12.5
<i>Ch. megocephala</i> vs <i>L. porphyrina</i>	25-27.2	9.1	15.4-15.7	9.8- 9.9	12-12.2
<i>Ch. megocephala</i> vs <i>L. sericata</i>	-	10.3	15.8- 16.4	9.1-9.2	14-14.5
<i>Ch. megocephala</i> vs <i>H. ligurriens</i>	28-31.2	10.9	17	8.3-8.65	15.1- 15.4
<i>Ch. megocephala</i> vs <i>A. grahami</i>	51-57.3	10.4	8.7	8.5-8.6	12.7-13
<i>Ch. megocephala</i> vs <i>M. domestica</i>	-	16.3	30.2-30.4	10.3-10.5	15.1- 15.4
<i>Ch. pinguis</i> vs <i>L. illustris</i>	-	9.7	14.4	9.4-9.5	13.4
<i>Ch. pinguis</i> vs <i>L. porphyrina</i>	-	9.9	15.7	10.3-10.4	11.7
<i>Ch. pinguis</i> vs <i>L. sericata</i>	-	10.8	15.6-16	10-10.1	13-13.1
<i>Ch. pinguis</i> vs <i>H. ligurriens</i>	-	12.1	16.3	8.8- 9.2	14
<i>Ch. pinguis</i> vs <i>A. grahami</i>	-	10.6	8.8	8.8-9	12.5
<i>Ch. pinguis</i> vs <i>M. domestica</i>	-	17	30.2	11-11.2	16.7
<i>Ch. ruffiacis</i> vs <i>Ch. megocephala</i>	-	-	7.5-7.7	6.5- 6.8	13.7-13.9
<i>Ch. ruffiacis</i> vs <i>Ch. pinguis</i>	-	-	6.8	8-8.2	11.9
<i>Ch. ruffiacis</i> vs <i>L. sericata</i>	-	-	16.2-16.4	11.6	12.7-13
<i>Ch. ruffiacis</i> vs <i>L. illustris</i>	-	-	17	10.8	11.2
<i>Ch. ruffiacis</i> vs <i>L. porphyrina</i>	-	-	15.8	11.95-12.1	14.7
<i>Ch. ruffiacis</i> vs <i>H. ligurriens</i>	-	-	18.9	10.4-10.6	9
<i>Ch. ruffiacis</i> vs <i>A. grahami</i>	-	-	8.7	9.4	11.7
<i>Ch. ruffiacis</i> vs <i>S. albiceps</i>	-	-	15.4	10.3	12.4
<i>Ch. ruffiacis</i> vs <i>M. domestica</i>	-	-	31	12.4	15
<i>L. illustris</i> vs <i>L. porphyrina</i>	22	4.5	5.1	4.6 (4)	6.1
<i>L. illustris</i> vs <i>L. sericata</i>	-	7.9	11.1-11.4	4.7	9.1
<i>L. illustris</i> vs <i>H. ligurriens</i>	17.7	10.1	10.6	6-6.2	9
<i>L. illustris</i> vs <i>A. grahami</i>	26.6	7.8	19	8.5	10.6
<i>L. illustris</i> vs <i>M. domestica</i>	-	16.2	30	10.9	12.5
<i>L. porphyrina</i> vs <i>L. sericata</i>	-	8.1	11.6-12	5.8-6	8
<i>L. porphyrina</i> vs <i>H. ligurriens</i>	7.8	10.1	11.4	7.2-7.6	10.4
<i>L. porphyrina</i> vs <i>A. grahami</i>	41.3	8.1	17.7	8.8-9	11.8
<i>L. porphyrina</i> vs <i>M. domestica</i>	-	16.5	33	12.5	13.7
<i>L. sericata</i> vs <i>L. cuprina</i>	-	-	3.5-3.7	0.78	0.5-0.7
<i>L. sericata</i> vs <i>H. ligurriens</i>	-	9.5	13.7-14	5.2-5.4	9.5-9.8
<i>L. sericata</i> vs <i>A. grahami</i>	-	10.4	18.4-18.7	7.6	13.2-13.5
<i>L. sericata</i> vs <i>M. domestica</i>	-	16.3	33.4-33.8	12.4	16.4
<i>L. cuprina</i> vs <i>L. illustris</i>	-	-	11.3	4.8	9.1
<i>L. cuprina</i> vs <i>L. porphyrina</i>	-	-	12.2	5.8	8.5
<i>L. cuprina</i> vs <i>H. ligurriens</i>	-	-	14	5.5-5.7	9.6
<i>L. cuprina</i> vs <i>A. grahami</i>	-	-	16.2	7.2	13.2
<i>L. cuprina</i> vs <i>Ch. megocephala</i>	-	-	14-14.6	8.8-9.2	14.6-14.9
<i>L. cuprina</i> vs <i>Ch. pinguis</i>	-	-	14.4	10.1-10.3	13.7
<i>L. cuprina</i> vs <i>Ch. ruffiacis</i>	-	-	15	10.9-11.1	12.2
<i>L. cuprina</i> vs <i>S. albiceps</i>	-	-	19.5	10.1	16
<i>L. cuprina</i> vs <i>M. domestica</i>	-	-	32.5	12.6	16.5
<i>H. ligurriens</i> vs <i>A. grahami</i>	42.8	12.4	18.7	7.4-7.6	10.4
<i>H. ligurriens</i> vs <i>M. domestica</i>	-	16	30	12.1-12.4	13.5
<i>A. grahami</i> vs <i>M. domestica</i>	-	16.3	35.2	13.5	15.5
<i>S. albiceps</i> vs <i>Ch. megocephala</i>	-	-	14.7-15	9.9-10.3	15.9-16.2
<i>S. albiceps</i> vs <i>Ch. pinguis</i>	-	-	14.8	10.7-10.8	16.2
<i>S. albiceps</i> vs <i>L. sericata</i>	-	-	19-19.3	10.5	16
<i>S. albiceps</i> vs <i>L. illustris</i>	-	-	18.1	10.3	14.5
<i>S. albiceps</i> vs <i>L. porphyrina</i>	-	-	18.3	12.3-12.5	15.4
<i>S. albiceps</i> vs <i>H. ligurriens</i>	-	-	21.1	11.8-12	14
<i>S. albiceps</i> vs <i>A. grahami</i>	-	-	15.8	11.6	17 (2)
<i>S. albiceps</i> vs <i>M. domestica</i>	-	-	29.5	13.5	12.7
Variation at species level (blowflies vs. blowflies)	7.8– 57.3 %	4.5– 12.4 %	1.2– 19 %	0.78– 10.4 %	0.5– 15.4 %
Variation at family level (blowflies vs. houseflies & blowflies vs. flesh flies)	-	16– 17 %	14.7– 35.2 %	9.9– 13.5 %	12.5– 16.7 %