

## **Insecticide Activity of Essential Oils of *Mentha longifolia*, *Pulicaria gnaphalodes* and *Achillea wilhelmsii* Against Two Stored Product Pests, the Flour Beetle, *Tribolium castaneum*, and the Cowpea Weevil, *Callosobruchus maculatus***

Authors: Khani, Abbas, and Asghari, Javad

Source: Journal of Insect Science, 12(73) : 1-10

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.7301>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## **Insecticide activity of essential oils of *Mentha longifolia*, *Pulicaria gnaphalodes* and *Achillea wilhelmsii* against two stored product pests, the flour beetle, *Tribolium castaneum*, and the cowpea weevil, *Callosobruchus maculatus***

Abbas Khani<sup>1a\*</sup> and Javad Asghari<sup>1, 2b</sup>

<sup>1</sup>Department of Plant Protection, Faculty of Agriculture, University of Zabol, Zabol, Iran

<sup>2</sup>Jihad and Agriculture Organization of South Khorasan, Birjand, Iran

### **Abstract**

Essential oils extracted from the foliage of *Mentha longifolia* (L.) (Lamiales: Lamiaceae) and *Pulicaria gnaphalodes* Ventenat (Asterales: Asteraceae), and flowers of *Achillea wilhelmsii* C. Koch (Asterales: Asteraceae) were tested in the laboratory for volatile toxicity against two stored-product insects, the flour beetle, *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) and the cowpea weevil, *Callosobruchus maculatus* F. (Coleoptera: Bruchidae). The chemical composition of the isolated oils was examined by gas chromatography-mass spectrometry. In *M. longifolia*, the major compounds were piperitenon (43.9%), tripal (14.3%), oxathiane (9.3%), piperiton oxide (5.9%), and d-limonene (4.3%). In *P. gnaphalodes*, the major compounds were chrysanthenyl acetate (22.38%), 2L -4L-dihydroxy eicosane (18.5%), verbenol (16.59%), dehydroaromadendrene (12.54%),  $\beta$ -pinen (6.43%), and 1,8 cineol (5.6%). In *A. wilhelmsii*, the major compounds were 1,8 cineole (13.03%), caranol (8.26%), alpha pinene (6%), farnesyl acetate (6%), and p-cymene (6%). *C. maculatus* was more susceptible to the tested plant products than *T. castaneum*. The oils of the three plants displayed the same insecticidal activity against *C. maculatus* based on LC<sub>50</sub> values (between 1.54  $\mu$ l/L air in *P. gnaphalodes*, and 2.65  $\mu$ l/L air in *A. wilhelmsii*). While the oils of *A. wilhelmsii* and *M. longifolia* showed the same strong insecticidal activity against *T. castaneum* (LC<sub>50</sub> = 10.02 and 13.05  $\mu$ l/L air, respectively), the oil of *P. gnaphalodes* revealed poor activity against the insect (LC<sub>50</sub> = 297.9  $\mu$ l/L air). These results suggested that essential oils from the tested plants could be used as potential control agents for stored-product insects.

**Keywords:** fumigant toxicity, gas chromatography-mass spectrometry, mono terpenoids

**Correspondence:** <sup>a</sup> [abbkhani@uoz.ac.ir](mailto:abbkhani@uoz.ac.ir), <sup>b</sup> [jaas542@yahoo.com](mailto:jaas542@yahoo.com), \* Corresponding author

**Received:** 26 June 2011, **Accepted:** 26 September 2011

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 12, Number 73

#### **Cite this paper as:**

Khani A, Asghari J. 2012. Insecticide activity of essential oils of *Mentha longifolia*, *Pulicaria gnaphalodes* and *Achillea wilhelmsii* against two stored product pests, the flour beetle, *Tribolium castaneum*, and the cowpea weevil, *Callosobruchus maculatus*. *Journal of Insect Science* 12:73 available online: [insectscience.org/12.73](http://insectscience.org/12.73)

## Introduction

The global pest-harvest grain losses by insect damage and other bio-agents range from 10% to 40% (Papachristos and Stamopoulos 2002). Chemicals largely used as pesticides in crop protection could have undesirable effects such as ozone depletion, environmental pollution, toxicity to non-target organisms, pest resistance, pesticide residues, and direct toxicity to users (Isman 2006). With heightened concern for environmental problems and human health, the search for readily biodegradable and environmentally friendly insecticides is of interest among scientists (Shaaya et al. 1997; Isman 2000). Plants offer an alternative source of insect-control agents because they contain a range of bioactive chemicals, many of which are selective and have little or no harmful effect on non-target organisms and the environment (Shaaya et al. 1997; Rajendran and Sriranjini 2008).

The genus *Mentha* belongs to the family Lamiaceae (Labiatae), and consists of about 25-30 species, most of which are found in temperate regions of Eurasia, Australia and South Africa (Lange and Croteau 1999). *Mentha longifolia* (L.) (Lamiaceae), commonly known as wild mint, is a perennial herb that can grow 1-2 m high. Various biological activities have been reported for some species of *Mentha*, such as antibacterial (Oyedeki and Afolayan 2006; Hajlaoui et al. 2008), antifungal (Bouchra et al. 2003), and insecticidal properties (Franzios et al. 1997; Lamiri et al. 2001; Pavela 2005; Saljoqi et al. 2006). The oils of *M. longifolia* are known to contain numerous monoterpenoids with piperitone oxide, piperitone, piperitenone, pulegone, d-limonene, carvone, menthone,  $\beta$ -caryophyllene, 1,8-Cineole, and menthol as

dominating compounds; however, there have been some variations in the constituents of this oil from different countries, and a chemo-geographical variation has been observed in essential oil composition of this species (Oyedeki and Afolayan 2006).

The *Pulicaria* genus belongs to the family Compositae (Asteraceae), tribe Inuleae, which contains more than 77 species that are widely distributed throughout Asia, Europe and Africa (Anderberg 1991). The chemical investigation of the genus showed the presence of terpenes such as monoterpenes and oxygenated monoterpenes (Weyerstahl et al. 1999), diterpenes (Muhammad et al. 1992), and sesquiterpenes (Weyerstahl et al. 1999; Dendougui et al. 2000). Various biological activities have been reported for some species of *Pulicaria*, such as antibacterial, antifungal (El-Kamali et al. 1998; Bahman et al. 2002; Liu et al. 2010), and insecticidal properties (Ross et al. 1997; Dubaie and El-Khulaidi 2005).

The herb *Achillea*, which belongs to the family Compositae (Asteraceae), is a genus with more than 100 species around the world. These plants are medicinal perennial rhizomous herbs, native to Europe and Western Asia, but also found in Australia, New Zealand, and North America (Chevallier 1996). Previous research has investigated the chemical composition of the essential oil of *Achillea*, such as its antibacterial (Barel et al. 1991; Magiatis et al. 2002) and insecticidal properties (Calmasur et al. 2006; Jovanovic et al. 2007; Magdy and Samir 2008). Previous work showed that the essential oil extracted from *Achillea wilhelmsii* C. Koch (Asteraceae) leaves exhibited volatile toxicity to *Sitophilus granarius* and *Tribolium confusum* (Calmasur et al. 2006).

In the present study, the chemical components of essential oils from *M. longifolia* L. (Lamiaceae) and *P. gnaphalodes* vent. (Asteraceae) aerial parts, and *A. wilhelmsii* C. Koch (Asteraceae) flowers, were determined, and the insecticidal activity of them was tested against the adult stages of the stored-products pests, *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) and *Callosobruchus maculatus* F. (Coleoptera: Bruchidae). No study has been reported concerning the activity of the three test oils as fumigants against these stored product insects.

## Materials and Methods

### Insect material

*C. maculatus* and *T. castaneum* were reared in plastic containers (20 cm length, 14 cm width, and 8 cm height, covered by a fine mesh cloth for ventilation) containing bean grain and wheat flour mixed with yeast (10:1, w/w), respectively. The culture was maintained in the dark, in a growth chamber set at 27±1°C and 65±5 relative humidity. All experiments were carried out under the same environmental conditions.

### Plants and essential oils

Aerial parts (foliage) of *M. longifolia* and *P. gnaphalodes*, and flowers of *A. wilhelmsii*, were collected respectively in Masabi, Sarayan (33° 51' N, 58° 31' E; 1500 m a.s.l.), Birjand suburbs (32° 50' N, 59° 13' E; 1500 m asl) and Sade, Ghaenat (33° 19' N, 59° 14' E; 1900 m asl), located in South Khorasan province, Iran, from May to July, 2009. The plant material was dried naturally on laboratory benches at room temperature (23-27 °C) until crisp. The dried material was stored at -24 °C, and then hydrodistilled to extract its essential oil. Essential oil was extracted from the plant samples using a

Clevenger-type apparatus where the plant material is subjected to hydrodistillation. Conditions of extraction were 50 g of samples, 1:10 plant material/water volume ratio, and a four-hour distillation. The oil was dehydrated with anhydrous sodium sulphate (10 min), and immediately stored in airtight glassware in a refrigerator at 4 °C.

### Gas chromatography-mass spectrometry

The essential oils were analyzed on a gas chromatograph mass spectrometer (Shimadzu -17A-QP5050, Japan). The gas chromatography column was a super CP-Sil 5CB capillary column (50 m × 0.32 mm ID, 0.25 µm film thickness). The column oven temperature was set at 70 °C for 1 min, increased to 100 °C at a rate of 1.5 °C/min, increased to 180 °C at a rate of 4 °C/min, and held at 180 °C for 1 min. Next, it was increased to 200 °C at a rate of 10 °C/min, increased to 250 °C at a rate of 2.5 °C/min, and held at 250 °C for 5 min. Injector and detector temperatures were 280 °C and 300 °C respectively. The gas chromatography mass analysis was carried out with the same characteristics as used in gas chromatography. The ionization energy was 70 eV, with a scan time of 1 sec, and a mass range of 40-300 amu. Unknown essential oil was identified by comparing its gas chromatography retention time to that of known compounds, and its mass spectra to known compounds or published spectra.

### Fumigant toxicity

To determine the fumigant toxicity of the oil, glass vials (volume 70 mL) were prepared, each containing 10 adults (1-7 days old of undefined sex) of each species. Filter papers (2 cm diameter) were prepared by adding 5, 15, or 30 µl of oil to individual papers (without using any solvent). Then, each filter paper was attached to the under-surface of a

screw cap and the cap was screwed tightly on the vial in order to generate concentrations of 71.43, 214.29, and 428.57  $\mu\text{L/L}$  air, respectively. Each concentration and control was replicated five times. Mortality was determined 3, 6, 9, 12, and 24 hours after exposure. When no signs of leg or antennal movement were observed, insects were considered dead.

Another experiment was designed to assess insect mortality using 50% lethal doses ( $\text{LC}_{50}$ ). The concentrations of the essential oils were chosen based on range-finding tests (to cause mortality between 5 and 90%). For each bioassay, five different concentrations, each with five replicates and ten individuals per replicate, were used. The volumes of the glass vials were 300 mL and 500 mL for *T. castaneum* and *C. maculatus*, respectively. The dead and living insects in each bottle were counted 24 hours after initial exposure to the essential oil. The mortality was determined as described in the previous experiment. The treatment bottles were monitored for 48 hours after recording the data, and no affected insect recovered. Data obtained from each dose response bioassay were subjected to probit analysis.  $\text{LC}_{50}$  values were determined by log-probit regression using SPSS 16.0 for Windows.

## Results and Discussion


### Chemical composition of essential oils

The chemical composition of the essential oils of the studied plants, *M. longifolia* and *P. gnaphalodes* aerial parts, and *A. wilhelmsii* flowers, are presented in Table 1.

In *M. longifolia*, the major compounds were piperitenone (43.9%), triphal (14.3%), oxathiane (9.3%), piperitone oxide (5.9%), and d-limonene (4.3%). The major compounds

of the Iranian *M. longifolia* oil were piperitone (43.9%), limonene (13.5%), and trans-piperitol (12.9%) (Rasooli and Rezaei 2002). However, identification of piperitone as the major compound in the *M. longifolia* oil is in sharp contrast to other reports where the oil had carvone (Monfared et al. 2002) or cis-carveol (Zeinali et al. 2005) as the major component.

The analysis of *P. gnaphalodes* essential oil revealed that chrysanthenyl acetate (22.38%), 2L-4L-dihydroxy eicosane (18.5%), verbenol (16.59%), dehydroaromadendrene (12.54%),  $\beta$ -pinene (6.43%), and 1,8 cineol (5.6%) were the main products. Weyerstahl et al. (1999) reported the oil prepared from aerial parts of *P. gnaphalodes* collected in the Elbrus mountains, Tehran province, Iran contained about 65% monoterpenes, with  $\alpha$ -pinene (34%) and 1,8-cineole (12%) as main compounds, and  $\beta$ -pinene (0.6%), allo-aromadendrene (0.4%) and trans-verbenol (0.2%) as minor compounds. Also, cis-chrysanthenol (oxidized monoterpenes) (2.3%) and its esters cis-chrysanthenyl formate (2.9%), cis-chrysanthenyl acetate (0.2%), chrysanthenone (monoterpene) (2%), and its related product isochrysanthenone (0.7%) were identified in the oil of this plant. Chrysanthenyl acetate was reported as the main component in the oil of *Artemisia vulgaris* L. collected from some localities in North Lithuania (Judžentienė and Buzelytė 2006), *Tanacetum balsamita* subsp. *balsamita* (Asteraceae) from Turkey (Bagci et al. 2008b), and *Tanacetum parthenium* (L.) from England and the Netherlands (Hendriks et al. 1996; Christensen et al. 1999). However, there are no data in the literature on the prevalence of chrysanthenyl acetate, 2L-4L-dihydroxy eicosane, verbenol, and dehydroaromadendrene in *P. gnaphalodes* essential oil.



**Figure 1.** Mean (5 replications, 10 individuals each) cumulative percentage mortality of *Tribolium castaneum* and *Callosobruchus maculatus* exposed to various concentrations of essential oils from *Mentha longifolia* and *Pulicaria gnaphalodes* aerial parts and *A. wilhelmsii* flowers at various periods of time. High quality figures are available online.

Gas chromatography-mass spectrometry analysis indicated that there are 20 major compounds in the oil of *A. wilhelmsii* flowers, comprising 88% of the total weight. 1,8-cineole was the most abundant compound (13.03%), followed by caranol (8.26%),  $\alpha$  pinene (6%), farnesyl acetate (6%), p-cymene (6%), together with lesser amounts of the other important insecticidal compounds, including camphor (4.2%), carvacrol (3.7%), and terpineol (3.1%). The results of our analysis are in agreement with previous reports that have also reported carvacrol, 1,8-cineol, camphor, and  $\alpha$ -pinene as the major components of the oil of *A. wilhelmsii* from Iran (Afsharypour et al. 1996; Javidnia et al. 2004; Ghani et al. 2008) and Turkey (Bagci et al. 2008a).

### Fumigant toxicity

In all cases, considerable differences in insect mortality due to essential oil vapor were observed using different concentrations and exposure times. The mortality increased with rising concentrations and exposure time (Figure 1). Results indicated that the oils of all three plants were significantly more toxic against *C. maculatus* than *T. castaneum*, as inferred by the confidence intervals of LC<sub>50</sub> (Figure 1). Furthermore, a difference in the response of the insect species to the essential oils has previously been reported for stored-product insects (Lee et al. 2003; Negahban et al. 2007).

Based on LC<sub>50</sub> (Table 1) and fumigant toxicity experiments (Figure 1), the oils of the three

plants displayed the same strong insecticidal activity against *C. maculatus* (between 1.54  $\mu\text{L/L}$  air in *P. gnaphalodes*, and 2.65  $\mu\text{L/L}$  air in *A. wilhelmsii*). While the oils of *A. wilhelmsii* and *M. longifolia* showed the same strong insecticidal activity against *T. castaneum* ( $\text{LC}_{50}$ =10.02 and 13.05  $\mu\text{L/L}$  air, respectively), the oil of *P. gnaphalodes* showed poor activity against the insect ( $\text{LC}_{50}$ =297.9  $\mu\text{L/Lair}$ ) (Table 2).

No study has been previously reported on the insecticidal activities of the oils of three tested plants against *C. maculatus* and *T. castaneum*; however, it has been reported that the oils and extracts of the tested plants had insecticidal activity against other insects. For example, the ethanol extracts from the leaves of *M. longifolia* revealed insecticidal activity with a maximum of 70% mortality six days after grain pollution (5 grams of plant powder extracted with 500 mL ethanol, diluted to 10%, and used at an amount of 4 mL per 20 grams of grains in plastic vials with 100 mL capacity) (Saljoqi et al. 2006). Contact and fumigant insecticidal actions of *Achillea wilhelmsii* (Calmasur et al. 2006) and other *Achillea* species (Jovanovic et al. 2007; Magdy and Samir 2008) have been demonstrated against stored product pests. One hundred percent mortality was achieved with 2  $\mu\text{L/L}$  air doses of the essential oils extracted from the leaves of *A. wilhelmsii* against *Sitophilus granarius* after an exposure time of 48 hours. However, mortality of *Tribolium confusum* at the same condition was about 90% (Calmasur et al. 2006).

Monoterpenes have been well documented as active fumigants, repellents, and insecticides toward stored-product insects (Papachristos et al. 2004). The insecticidal activity of the essential oils investigated in our study may be attributed to their having major monoterpenes

components, because some major compounds of the test oils, such as carvacrol, camphor, 1,8-cineole,  $\alpha$ -pinene, p-cymene, piperitenone oxide, and terpineol possessed insecticidal effects against the test insects (Traboulsi et al. 2002; Papachristos et al. 2004; Miresmailli et al. 2006; Cetin et al. 2007; Kordali et al. 2008; Magdy and Samir 2008).

The essential oils from these plants could become a viable alternative to conventional chemical control strategies. However, further studies need to be conducted in order to evaluate the safety of these oils before practical use in stored-product insect control.

### Acknowledgements

We wish to thank Dr. V.A. Mozafarian, member of the scientific board at Research Institute of Forests and Rangelands, for plant species identification, and M.A. Allah Resani, expert of basic science faculty of Birjand University, for gas chromatography-mass spectrometry analysis. This research was supported by the University of Zabol, which is greatly appreciated.

### References

- Afsharypour S, Asgary S, Lockwood B. 1996. Constituents of the essential oil of *Achillea wilhelmsii* from Iran. *Planta Medica* 62: 77-78.
- Anderberg AA. 1991. Taxonomy and phylogeny of the tribe Inuleae (Asteraceae). *Plant Systematic and Evolution* 176: 75-123.
- Bagci E, Koçak A, Yüce E. 2008a. The Composition of the essential oils of two *Achillea* L. (*Achillea wilhelmsii* C. Koch. ve *Achillea schischkinii* Sosn.) species. *Science*

and Engineering Journal of Firat University 20 (2): 251-255.

Bagci E, Kursat M, Kocak A, Gur S. 2008b. Composition and Antimicrobial Activity of the Essential Oils of *Tanacetum balsamita* L. subsp. *balsamita* and *T. chiliophyllum* (Fisch. et Mey.) Schultz Bip. var. *chiliophyllum* (Asteraceae) from Turkey. *Journal of Essential Oil Bearing Plants* 11(5): 476-484.

Barel S, Segal R, Yashphe J. 1991. The antimicrobial activity of the essential oil from *Achillea fragrantissima*. *Journal of Ethnopharmacology* 33: 187-191.

Bouchra C, Achouri M, Idrissi Hassani LM, Hmamouchi M 2003. Chemical composition and antifungal activity of essential oils of seven Moroccan Labiatae against *Botrytis cinerea* Pers. *Journal of Ethnopharmacology* 89: 165-169.

Calmasur O, Kordali S, Kaya O, Aslan I. 2006. Toxicity of essential oil vapours obtained from *Achillea* spp. to *Sitophilus granarius* (L.) and *Tribolium confusum*. *Journal of Plant Diseases and Protection* 113(1): 37-41.

Cetin H, Erler F, Yanikoglu A. 2007. Comparative evaluation of *Origanum onites* essential oil and its four major components as larvicides against the pine processionary moth, *Thaumetopoea wilkinsoni* Tams. *Pest Management Science* 63(8): 830-833.

Chevallier A. 1996. *The Encyclopedia of Medicinal Plants*. Dorling Kindersley Publishing.

Christensen LP, Jakobsen HB, Paulsen E, Hodal L, Andersen KE. 1999. Airborne compositae dermatitis: Monoterpenes and no

parthenolide are released from flowering *Tanacetum parthenium* (feverfew) plants. *Archives of Dermatological Research* 291: 425-431.

Dendougui H, Benayache S, Benayache F, Connolly JD. 2000. Sesquiterpene lactones from *Pulicaria crispa*. *Fitoterapia* 71: 373-378.

Dubaie AS, El-Khulaidi AA. 2005. *Medicinal and aromatic plants in Yemen, deployment-components of effective-uses*. Ebadi Center for Studies and Publishing.

El-Kamali HH, Ahmed AH, Mohammed AS, Yahia AAM, El-Tayeb IH, Ali AA. 1998. Antibacterial properties of essential oils from *Nigella sativa* seeds, *Cymbopogon citratus* leaves, and *Pulicaria undulata* aerial parts. *Fitoterapia* 69 (1): 77-78.

Franzios G, Mirotsou M, Hatziapostolou E, Kral J, Scouras ZG, Mavragani-Tsipidou P. 1997. Insecticidal and genotoxic activities of mint essential oils. *Journal of Agricultural and Food Chemistry* 45: 2690-2694.

Ghani A, Azizi M, Hassanzadeh-Khayyat M, Pahlavanpour AA. 2008. Essential Oil Composition of *Achillea eriophora*, *A. nobilis*, *A. biebersteinii* and *A. wilhelmsii* from Iran. *Journal of Essential Oil Bearing Plants* 11(5): 460-467.

Hajlaoui H, Snoussi M, Ben Jannet H, Mighri Z, Bakhrouf A. 2008. Comparison of chemical composition and antimicrobial activities of *Mentha longifolia* L. ssp. *longifolia* essential oil from two Tunisian localities (Gabes and Sidi Bouzid). *Annals of Microbiology* 58(3): 103-110.



- Hendriks H, Bos R, Woerdenbag J. 1996. The essential oil of *Tanacetum parthenium* (L.) Schultz-Bip. *Flavour and Fragrance Journal* 11: 367-371.
- Isman MB. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annual Review of Entomology* 51: 45-66.
- Isman MB. 2000. Plant essential oils for pest and disease management. *Crop Protection* 19: 603-608.
- Javidnia K, Miri R, Sadeghpour H. 2004. Composition of the volatile oil of *Achillea wilhelmsii* C. Koch from Iran. *Daru* 12(2): 63-66.
- Jovanovic Z, Kosti M, Popovic Z. 2007. Grain-protective properties of herbal extract against the bean weevil *Acanthoscelides obtectus* Say. *Industrial Crops and Products* 26: 100-104.
- Judžentienė A, Buzelytė J. 2006. Chemical composition of essential oils of *Artemisia vulgaris* L. (mugwort) from North Lithuania. *CHEMIJA* 17(1): 12-15.
- Kordali S, Cakir A, Ozer H, Cakmakci R, Kesdek M, Mete E. 2008. Antifungal, phytotoxic and insecticidal properties of essential oil isolated from Turkish *Origanum acutidens* and its three components, carvacrol, thymol and p-cymene. *Bioresource Technology* 99(18): 8788-8795.
- Lamiri A, Lhaloui S, Benjilali B, Berrada M. 2001. Insecticidal effects of essential oils against Hessian fly, *Mayetiola destructor* (Say). *Field Crops Research* 71: 9-15.
- Lange BM, Croteau R. 1999. Genetic engineering of essential oil production in mint. *Current Opinion in Plant Biotechnology* 2: 139-144.
- Lee S, Peterson CJ, Coats JR. 2003. Fumigation toxicity of monoterpenoids to several stored product insects. *Journal of Stored Product Research* 39: 77-85.
- Liu LL, Yang JL, Shi YP. 2010. Phytochemicals and Biological Activities of *Pulicaria* Species. *Chemistry and Biodiversity* 7(2): 327-349.
- Magdy IEM, Samir AMA. 2008. Chemical composition and insecticidal potential of essential oils from Egyptian plants against *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) and *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Applied Entomology and Zoology* 43(4): 599-607.
- Magiatis P, Skaltsounis AL, Chinov I, Haroutounian SA. 2002. Chemical composition and in vitro antimicrobial activity of the essential oils of three greek *Achillea* species. *Zeitschrift für Naturforschung* 57: 287-290.
- Miresmailli S, Bradbury R, Isman MB. 2006. Comparative toxicity of *Rosmarinus officinalis* L. essential oil and blends of its major constituents against *Tetranychus urticae* Koch (Acari: Tetranychidae) on two different host plants. *Pest Management Science* 62(4): 366-371.
- Monfared A, Rustaiyan A, Nabid M. 2002. Composition of a carvone chemotype of *Mentha longifolia* (L.) Huds. from Iran. *Journal of Essential Oil Research* 14: 51-52.

- Muhammad I, El-Feraly FS, Mossa JS, Ramadan AF 1992. Terpenoids from *Pulicaria glutinosa*. *Phytochemistry* 31(12): 4245-4248.
- Negahban M, Moharramipour S, Sefidkon F. 2007. Fumigant toxicity of essential oil from *Artemisia sieberi* Besser against three stored product insects. *Journal of Stored Product Research* 43: 123-128.
- Nickavar B, Amin GR, Ghavamian P 2002. Antimicrobial activity of *Pulicaria dysenterica*. *Iranian Journal of Pharmaceutical Research* 1: 31-32.
- Oyedeki OA, Afolayan AJ. 2006. Chemical composition and antibacterial activity of the essential oil isolated from South African *Mentha longifolia* (L.) subsp. *capensis* (Thunb.) Briq. *Journal of Essential Oil Research* 18: 57-59.
- Papachristos DP, Karamanoli KI, Stamopoulos DC, Menkissoglu-Spiroudi U. 2004. The relationship between the chemical composition of three essential oils and their insecticidal activity against *Acanthoscelides obtectus* (Say). *Pest Management Science* 60(5): 514-520.
- Papachristos DP, Stamopoulos DC. 2002. Repellent, toxic and reproduction inhibitory effects of essential oil vapors on *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). *Journal of Stored Product Research* 38(2): 117-128.
- Pavela R. 2005. Insecticidal activity of some essential oils against larvae of *Spodoptera littoralis*. *Fitoterapia* 76: 691-696.
- Rajendran S, Sriranjini V. 2008. Plant products as fumigant for stored-product insect control. *Journal of Stored Product Research* 44: 126-135.
- Rasooli I, Rezaei MB. 2002. Bioactivity and chemical properties of essential oils from *Zataria multiflora* Boiss and *Mentha longifolia* (L.) Huds. *Journal of Essential Oil Research* 14: 141-146.
- Ross SA, El-Sayed KA, El-Sohly MA, Hamann MT, Abdel-Halim OB, Ahmed AF, Ahmed MM. 1997. Phytochemical Analysis of *Geigeria alata* and *Francoeuria crispa* essential oils. *Planta Medica* 63: 479-482.
- Saljoqi AUR, Afridi MK, Khan SA, Rehman S. 2006. Effects of six plant extracts on rice weevil *Sitophilus oryzae* L. in the stored wheat grains. *Journal of Agricultural and Biological Science* 1 (4): 1-5.
- Shaaya E, Kostyukovsky M, Eilberg J, Sukprakarn C. 1997. Plant oils as fumigants and contact insecticides for the control of stored-product insect. *Journal of Stored Product Research* 33: 7-15.
- Traboulsi AF, Taoubi K, El-Haj S, Bessiere JM, Rammal S. 2002. Insecticidal properties of essential plant oils against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Management Science* 58(5): 491-495.
- Weyerstahl P, Marschall H, Wahlburg HC, Christiansen C, Rustaiyan A, Mirdjalili F. 1999. Constituents of the essential oil of *Pulicaria gnaphalodes* (Vent.) Boiss. from Iran. *Flavour and Fragrance Journal* 14(2): 121-130.
- Zeinali H, Arzani A, Razmjoo K, Rezaee M. 2005. Evaluation of oil compositions of Iranian mints (*Mentha* spp.). *Journal of Essential Oil Research* 17: 156-159.

**Table 1.** Chemical composition of essential oils of *Mentha longifolia* and *Pulicaria gnaphalodes* vegetative parts and *Achillea wilhelmsii* flowers.

Compound <sup>a, b</sup>	Kovats index	Concentration (%)		
		<i>M. longifolia</i>	<i>P. gnaphalodes</i>	<i>A. wilhelmsii</i>
P Hydroxy cyclohexanone	—	—	4.5	—
2L-4L-Dihydroxy Eicosane	199	—	18.5	—
Camphene	946	—	—	4
$\alpha$ pinene	949	—	—	8.5
Delta 3 caren	951	—	—	5.8
Sabinen	957	1.6	—	—
$\beta$ pinene	959	1.3	6.4	1
Linalool	964	2.7	—	—
1,8 Cineol	981	—	5.6	13
DI lilmonen	983	4.3	—	—
Trans orth - cimen	984	1.5	—	—
Carene	988	1.5	—	—
p-cymene	994	—	—	6
Eucalyptol	995	—	—	3.2
Verbenol	1014	—	16.6	—
Camphor	1021	—	1.2	4.2
Terpineol-4	1036	—	2.2	—
Terpinene-4-ol	1037	—	2.1	—
Terpineol	1052	—	—	3.1
$\alpha$ Terpineol	1055	—	—	1.5
Oxaspiro	1068	3.3	—	—
Chrysanthenyl Acetate	1069	—	22.4	—
Carvacrol	1082	—	—	3.7
Piperiton oxide	1086	5.9	—	—
Tripal	1108	14.3	—	—
Piperitenon	1112	43.9	—	—
Caranol	1149	—	—	8.3
Oxathiane	1150	9.3	—	—
Cyclohexane	1154	—	—	1.3
Delta elemene	1231	1.4	—	—
Caryophyllene oxide	1344	—	—	1.4
Germacrene	1465	1.3	—	—
Caryophyllene oxide	1528	1.1	—	—
Aromadendrene oxide	1535	—	—	5.7
Guaiol	1577	1	—	—
$\alpha$ humulene	1612	—	—	1.3
Dehydroaromadendrene	1633	—	12.54	—
Farnesyl acetate	1691	—	—	6.1
Palmitic acid	1834	—	—	2.3
Nerolidol epoxy acetate	1871	—	—	2.8
Di hydroxy eicosane	1983	—	—	5.1

<sup>a</sup> Compounds are listed in the order of their elution.<sup>b</sup> Identification based on authentic standards, Wiley libraries spectra, and literature.**Table 2.** Efficiency of essential oil extracted from three plants against *Tribolium castaneum* and *Callosobruchus maculatus* adults.

Plants	Insects*	95% CL	$\chi^2$ (df)	Probability	LC <sub>50</sub>	Slope $\pm$ SE
<i>M. longifolia</i>	<i>T. castaneum</i>	(10.9-15.1)	2.31(3)	0.51	13.05	3.25 $\pm$ 0.44
	<i>C. maculatus</i>	(1.29-2.78)	5.66 (3)	0.13	2.05	3.54 $\pm$ 0.45
<i>P. gnaphalodes</i>	<i>T. castaneum</i>	(255.1-362.0)	4.65 (3)	0.33	297.9	2.58 $\pm$ 0.40
	<i>C. maculatus</i>	(0.5-3.0)	7.14 (3)	0.07	1.54	1.88 $\pm$ 0.28
<i>A. wilhelmsii</i>	<i>T. castaneum</i>	(8.7-11.7)	3.98 (3)	0.26	10.02	3.06 $\pm$ 0.46
	<i>C. maculatus</i>	(2.3-3.0)	4.74 (3)	0.93	2.65	0.61 $\pm$ 0.08

\* Ten individuals per replicate, five replicates per concentration, five concentrations per assay for *T. castaneum* and *C. maculatus*, respectively; LC: lethal concentration  $\mu$ L/L air, CL: confidence limits