

First Record of the Leafhopper Genus Varicopsella Hamilton, 1980 (Hemiptera: Cicadellidae: Macropsinae) in China, with Descriptions of a New Subgenus and New Species, a Checklist, and a Key to Species

Authors: Li, Hu, Dai, Ren-Huai, and Li, Zi-Zhong

Source: Journal of Insect Science, 14(125) : 1-10

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.014.125

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



First record of the leafhopper genus Varicopsella Hamilton, 1980 (Hemiptera: Cicadellidae: Macropsinae) in China, with descriptions of a new subgenus and new species, a checklist, and a key to species

Hu Li^{1,2a}, Ren-Huai Dai,^{1b*} and Zi-Zhong Li^{1c}

¹Institute of Entomology, Guizhou University, Guiyang, Guizhou, and the Provincial Key Laboratory for Agricultural Pest Management of Mountainous Region, Guiyang, Guizhou 550025 P.R. China ²School of Biological Sciences & Engineering, Shaanxi University of Technology, Hanzhong, Shaanxi, and Bio-resources Key Laboratory of Shaanxi Province, Shaanxi University of Technology, Hanzhong, Shaanxi 723000 P.R. China

Abstract

A new monobasic leafhopper subgenus, *Varicopsella (Multispinulosa)* Li, Dai, and Li, **subgen. nov.**, of the subfamily Macropsinae (Hemiptera: Auchenorrhyncha: Membracoidea: Cicadellidae) is proposed to accommodate *Varicopsella (Multispinulosa) hamiltoni* Li, Dai, and Li, **sp. nov.** from Guangxi province of China. The new subgenus and new species are described and illustrated. They can be distinguished mainly by characteristics of the fore wings with two anteapical cells; weak dorsoventrally flattened body; aedeagal shaft with paired apical processes on ventral margin; and the shape of the dorsal connective. An updated checklist and an illustrated key for identification of the species of *Varicopsella* along with geographical distributions of the species are given.

摘要

文章描述了采自中国广西的叶蝉 1 新亚属及 1 新种,汉密尔顿合板叶蝉 Varicopsella (Multispinulosa) hamiltoni Li, Dai, and Li, subgen. nov. and sp. nov., 新种的主要区别特征是:前翅具 2 个端前室;背腹向扁平体型不甚明显;阳茎干腹缘具一对端突;背连索的形状也不尽相同。同时给出该属最新的物种名录、附插图的检索表和地理分布图。

Keywords: Auchenorrhyncha, morphology, distribution Correspondence: ^a lihu@snut.edu.cn, ^b rhdai69@163.com, ^c lizizhong38@163.com, *Corresponding author Editor: Peter Kerr and Daniela Takiya were editors of this paper. Received: 28 October 2012 Accepted: 25 April 2013 Published: 1 September 2014 Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed. ISSN: 1536-2442 | Vol. 14, Number 125 Cite this paper as: Li H, Dai R-H, Li Z-Z. 2014. First record of the leafhopper genus *Varicopsella* Hamilton, 1980 (Hemiptera: Cicadellidae:

Li H, Dai R-H, Li Z-Z. 2014. First record of the leafhopper genus *Varicopsella* Hamilton, 1980 (Hemiptera: Cicadellidae: Macropsinae) in China, with descriptions of a new subgenus and new species, a checklist, and a key to species. *Journal of Insect Science* 14(125). Available online: <u>http://www.insectscience.org/14.125</u>

Journal of Insect Science | http://www.insectscience.org

Introduction

The oriental leafhopper genus Varicopsella Hamilton, 1980 (Hemiptera: Auchenorrhyn-Membracoidea: Cicadellidae: cha: Macropsinae) was established by Hamilton (1980) for six species from the Oriental region (Philippines and Borneo), with Macropsis breakevi Merino, 1936 as its type species by original designation. One year later, Viraktamath (1981), described a new species from India. Up to now, no other species was described in this genus. Considering the vast land and abundant resources, macropsine fauna are likely to be richer in southern China, including the genus Varicopsella.

Deposited in our collection from Guangxi Province of China (included in the Oriental Region), two specimens were found that bear the distinct generic characteristics of *Varicopsella*: the fused sclerites between lora and frons on face, and the independently evolved dorsal connectives and pygofer processes in males. These specimens have been identified as a new species of *Varicopsella*, and on the basis of the distinctive feature that the fore wings of the new species have two anteapical cells, we propose to place it in a new subgenus. The identification was verified by K. G. A. Hamilton, pers. comm. (Research Branch, Agriculture and Agri-Food Canada).

In this paper, the genus *Varicopsella* is reported for the first time from China based on the new species. The new taxa (*Varicopsella* (*Multispinulosa*) hamiltoni Li, Dai, and Li, **subgen. nov.** and **sp. nov.** from Guangxi province (China) are described and illustrated. Photographs of imago and illustrations of the male genitalia are provided, and an illustrated key to the species of the genus based on the original descriptions and illustrations, with a map of distribution of the genus based on the primary, data are given.

Materials and Methods

Morphological terminology used in this work follows Hamilton (1980) and of the rows of setae on the legs follows Rakitov (1998). The type specimens examined here are deposited in the Institute of Entomology, Guizhou University, Guiyang, China (GUGC).

Nomenclature

This paper and the nomenclature it contains have been registered in ZooBank. The LSID number is:

urn:lsid:zoobank.org:pub:F8C13A75-E433-4DA8-8534-1144943E3686

Systematics

Subfamily Macropsinae Evans, 1938 Genus Varicopsella Hamilton, 1980

Subgenus *Varicopsella* (*Varicopsella*) Hamilton, 1980 *Varicopsella* Hamilton, 1980: 900.

Type species: *Macropsis breakeyi* Merino, 1936.

Remarks. The genus *Varicopsella* can be distinguished from other Macropsinae mostly by the following characteristics: fused sclerites between lora and frons on face, and the unique male dorsal connectives each formed by two articulating parts with ventral pair appressed or fused with each other and dorsal ones produced as variously-shaped projections among species. The unique distinctive feature of the nominate subgeneric *Varicopsella* (*Varicopsella*) is fore wings with three anteapical cells.

Distribution. Oriental, distributed in Philippines, Borneo, and India (Hamilton 1980).

Subgenus *Varicopsella* (*Multispinulosa*) Li, Dai, and Li, **subgen. nov.**

Type species: *Varicopsella (Multispinulosa) hamiltoni* Li, Dai and Li, **sp. nov.**

Description. Body (Fig. 1–4) typical wedgeshaped; head, face, pronotum, and scutellum strongly striated.

Head (Fig. 1), in dorsal view, clearly arcuate anteriorly, and width across eyes equal to pronotum, crown shorter medially than besides eyes. Face (Fig. 4), across eyes slightly wider than long, obviously co-planar, with lora fused to frons, lacking clear sutures between them, dorsal part with median carina, along it bilaterally with slant striations ended below ocelli; distance between ocelli about 5× longer than that of ocellus to adjacent eye. Pronotum (Fig. 1) $2.1 \times$ as long as wide, anterior margin strongly prominent, posterior margin slightly concave medially, almost straight, surface with strongly oblique striations along median carina. Scutellum (Fig. 1) nearly triangular, with $1.2 \times$ as long as pronotum, strongly striated except bilateral corners with relatively smooth surface, coalescent suture between mesonotum and secutellum distinct, arcuate. Forewings opaque (Fig. 1-3), with 2 anteapical cells, veins protruding. Hind femoral macrosetae (Fig. 6) 2+1; hind tibia (Fig. 5) with 8 macrosetae on AD row, 5 on AV row, 11 on PD row, dense and slender on PV row.

Male genitalia. Pygofer (Fig. 7–8) broad basally, ventral margins with some setae subapically and produced into triangular processes with small spines on surface. Subgenital plates (Fig. 7) slender with scattered setae. Styles (Fig. 9) slender with marginal setae, nearly angled on basal 0.45, slightly intumescent subapically, apex nearly rectangular on ventral margin. Connective (Fig. 10–11) similar to others of genus. Aedeagus (Fig. 12–13) simple, broad basally, shaft sinuated, shaft apex sharp, dorsally directed, with pair of processes on ventral margin, dorsal apodeme weakly developed, ventral margin concave in middle; gonopore apical on ventral margin. Dorsal connectives (Fig. 14–15) strongly developed, ventral portions appressed with each other, dorsal portions slender, sinuate, with dorsal end bilobed.

Distribution. Oriental, China (new record of genus), Guangxi Province.

Remarks. The new subgenus can be easily distinguished from the other *Varicopsella* subgenus by the fore wings with two anteapical cells; weakly flattened external form of the body; unique male aedeagal shaft with paired apical processes on ventral margin; and shape of the dorsal connective.

Etymology. The subgenus name is derived from the Latin words "*multi-*" and "*spinulosus*" because of the several spines on the ventral margin projections of the pygofer lobe. The gender of the subgenus is feminine.

Varicopsella (Multispinulosa) hamiltoni Li, Dai, and Li **sp. nov.** (Fig. 1–15, 21)

Measurements. Body length including tegmina, 3, 3.9–4.1 mm.

Body color. Background color (Fig. 1–4) yellowish-green, striations on head, face, and pronotum with same tint as background except posterior margin of pronotum with brown markings. Head, face, and pronotum (Fig. 1– 4) yellowish-green. Eyes brown, with some-

what slightly and occasionally red crown; ocelli brightly lucid; antennal fossa, scape, pedicel, and flagellum successively vary from yellowish green to brown; terminal anteclypeus, beak, and outer margins of both lora yellowish-brown to brown. Scutellum (Fig. 1) fully black except bilateral margins slightly yellowish-brown medially. Fore wings (Fig. 1–3) dark brown, except terminal regions of veins 1A, distal claval suture and outer anteapical part with brown maculae successively enlarged in size, veins clearly spotted with white. Legs (Fig. 5–6) brown with black maculae.

Structural morphology. As in generic description.

Male genitalia. Pygofer (Fig. 7–8), in lateral view, broad basally, truncate caudally, in ventral view, ventral margins with some setae subapically and elongate to nearly triangular processes slightly exceeding pygofer length, and with several small spines on surface. Subgenital plate (Fig. 7), in lateral view, rodlike, scattered with hair-like setae, apex with bunched long setae, slightly projecting beyond pygofer. Style (Fig. 9), slender, margined with some setae, nearly angled on basal 0.45, slightly inflated subapically, apex angular with dorsally twisted digitation. Connective (Fig. 10-11), in dorsal aspect, clearly longer than wide, with anterior margin strongly wider than posterior margin, anterior margin excavated medially, and with finger-like process in middle, both lateral arms slender and blend dorsally. Aedeagus (Fig. 12–13) tubular, broad basally, shaft apex sharpened, with pair of triangular processes on ventral margin; in lateral aspect, dorsal apodeme weakly developed, shaft sinuated, ventral margin concave medially, tip twisted to dorsum; in ventral aspect, shaft nearly parallel margined; gonopore on ventral margin, apical. Dorsal connective (Fig. 14–15) strongly developed, ventral portions appressed with each other, with dorsal end bearing two digitations; in lateral view, dorsal portions long, sinuate; in caudal view, ventral paired digitations ventrally directed to each other.

Female. Unknown.

Materials examined. Holotype, ♂, CHINA: Guangxi Province, Longsheng County, Huaping National Natural Reserve, 18. V. 2012, collected by Li Hu; Paratype, 1 ♂, CHINA: Guangxi Province, Baise City, Tianlin County, Langping Village, 23. IV. 2012, collected by Yang Weicheng.

Distribution. Guangxi prov. (Huaping, Langping), China (Fig. 21).

Etymology. This species is named in honor of K. G. A. Hamilton for excellent contributions to the Auchenorrhyncha systematics and invaluable help to the first author.

Updated checklist and distributions of the species of genus *Varicopsella*

Varicopsella (Multispinulosa) hamiltoni sp. nov.

Distribution. China (Huaping, Langping).

Varicopsella (Varicopsella) basilana (Merino) Macropsis basilana Merino, 1936: 323. Varicopsella basilana, Hamilton, 1980: 900. Distribution. Philippines (Basilan, Mindanao).

Varicopsella (Varicopsella) breakeyi (Merino) Macropsis breakeyi Merino, 1936: 320. Varicopsella breakeyi, Hamilton, 1980: 900. Distribution. Philippines (Mindanao).

Varicopsella (Varicopsella) davaoensis (Merino)

Macropsis davaoensis Merino, 1936: 325. Varicopsella davaoensis, Hamilton, 1980: 900.

Distribution. Philippines (Mindanao).

Varicopsella (Varicopsella) elegans Viraktamath

Varicopsella elegans Viraktamath, 1981: 306. Distribution. India (Himalayas).

Varicopsella (Varicopsella) luzonensis (Merino) Macropsis luzonensis Merino, 1936: 324. Varicopsella luzonensis, Hamilton, 1980: 900. Distribution. Philippines (Luzon).

Varicopsella (Varicopsella) obtusa Hamilton Varicopsella obtusa Hamilton, 1980: 919. Distribution. Borneo (Sandakan).

Varicopsella (Varicopsella) otanesi (Merino) Macropsis otanesi Merino, 1936: 323. Varicopsella otanesi, Hamilton, 1980: 900. Distribution. Philippines (Basilan, Mindanao).

Key to the Subgenera and Species of Male *Varicopsella*

The *Varicopsella* species are keyed based on the variations of male genitalia, therefore two species only known by the female, *V*. (*V*.) basilana (Merino, 1936) and *V*. (*V*.) davaoensis (Merino, 1936), are excluded from the key.

1. Body form not strongly flattened dorsoventrally (Fig. 3); fore wings with 2 anteapical cells (Fig. 3); aedeagal shaft with pair of triangular processes apically (Fig. 13).....V. (*Multispinulosa*) hamiltoni **subgen. nov.** and **sp. nov.**

– Body form strongly flattened dorsoventrally; fore wings with 3 anteapical cells; aedeagal shaft without pair of triangular processes apically (Fig. 16–20)..........V. (Varicopsella) 2

3. Dorsal connective short with tip bifid or trifid (Fig. 17, 19)......4
– Dorsal connective slender with tip sharp, not furcated (Fig. 18, 20)......5

5. Pygofer with two small spines on each lobe (Fig. 20); dorsal connective elongate process with dorsal end directed ventrally (Fig. 20)......V. (Varicopsella) otanesi – Pygofer with two extremely tiny spines on lobe (Fig. 18); dorsal connective elongate process with dorsal end directed dorsally (Fig. 18).....V. (Varicopsella) luzonensis

Acknowledgements

We sincerely thank K. G. A. Hamilton (Research Branch, Agriculture and Agri-Food Canada, Ottawa, ON) for the kind suggestions to study; also thank our colleague and friend Mr. Wei-Cheng Yang (Institute of Entomology, Guizhou University) for providing the material for study. This project was supported by the National Natural Science Foundation of China (No. 31000952).

References

Hamilton, K. G. A. 1980. Contributions to the study of the world Macropsini (Rhynchota: Homoptera: Cicadellidae). *Can. Entomol.* 112: 875–932.

Merino, G. 1936. Philippine Cicadellidae (Homoptera). *Philip. J. Sci.* 61: 307–400.

Rakitov, R. A. 1998. On differentiation of cicadellid leg chaetotaxy (Homoptera, Auchenorrhyncha, Membracoidea). *Rus. Entomol. J.* 6: 7–27.

Viraktamath, C. A. 1981. Indian Macropsinae (Homoptera: Cicadellidae). II. Species described by W. L. Distant and descriptions of new species from the Indian subcontinent. *Entomol. Scand.* 12: 295–310.







Journal of Insect Science | http://www.insectscience.org

