

Mites Inhabiting a Lepidopteran Egg

Authors: Mota, Luísa Lima E, Tacioli, André, and Sendoya, Sebastian Felipe

Source: The Journal of the Lepidopterists' Society, 68(2) : 141-142

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.v68i2.a7>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MITES INHABITING A LEPIDOPTERAN EGG

Additional key words: Acari, ecosystem engineering, Cerrado, Mimallonidae

Lepidopteran eggs are highly variable in morphology, showing different shapes and sizes according to family, genus and/or species (Stehr 1987). After hatching, many species eat their egg shells (see examples in Braby & Nishida 2007), while others open a hole through which they hatch and leave the rest of the chorion intact (see examples in Kaminski et al. 2013). In some cases, the empty egg shell can remain for days or weeks in a shape similar to the original (e.g. egg shells of *Parrhasius polibetes* (Stoll, 1781) (Lycaenidae) may endure on the inflorescences of their hostplants for several weeks; L. L. Mota, pers. obs)

In April 2013, a lepidopteran egg shell was found attached to the lateral of a Myrtaceae leaf, in an area of Cerrado savanna belonging to the Laboratório Nacional de Luz Síncrotron (22°48'S, 47°03'W), in Campinas, Southeast Brazil. The egg was elongated (1.3 mm long and 0.4 mm wide), with longitudinal ribs, and one extremity was open with irregular edges, suggestive of biting marks left by the caterpillar while hatching (Fig. 1). It was attached to the leaf longitudinally, so the structure resembled a tunnel with one side closed. Its shape and position in the leaf are similar to the described for the Mimallonidae genus *Lacosoma* Grote, 1864 (Dyar 1900, Peterson 1961; 1966). A caterpillar was found on a nearby leaf on the same branch, and was possibly the one which hatched from this egg. It was under a net constructed of silk and frass, also similar to that described for early and mid-instar larvae of a *Lacosoma* species (Dyar 1900). These evidences strongly suggest the egg belonged to the family Mimallonidae.

With stereomicroscopy observation, four unidentified mites (Acari) were found inside the egg, at the opposite extremity to the opening (Fig. 1a). Feces and at least six elliptical mite eggs (0.09 mm long) were also present at this region (Fig. 1b). The mites stayed in the same position even with intense hand and pin manipulation. Under stereomicroscope light they started to move and only one of them left the egg, walked around the chorion and the leaf and entered the egg shell a few minutes later (Fig. 1c–d). Previously, no mites were found walking in the leaf outside the egg shell.

The presence of the mites, their feces and eggs inside this lepidopteran egg shell suggests they were consistently using the chorionic structure as a shelter and reproduction site. Mites use many leaf structures as shelters, such as acaridomatias (O'Dowd & Willson 1989,

Willson 1991), former lepidopteran shelters (Lima et al. 2013), rolled leaves (Fournier et al. 2003), and even adhesive traps of carnivorous plants (Antor & García 1995). Some mites tend to establish preferentially near wall structures, especially in gaps between two walls (Kawasaki et al. 2009), so the egg shell of this Mimallonidae seems to be a perfect site for them, representing a long structure with two parallel walls. The benefits proportioned by the egg shell could be related to diminishing predation risk and maintaining microclimatic conditions (which could prevent them from desiccating), as found for other kinds of shelters used by mites (Kawasaki et al. 2009, Lima et al. 2013). The mites and eggs were positioned as far as possible from this aperture, where it would be most difficult for a predator larger than the opening of the egg to reach, and where they were in contact with a third wall structure, possibly providing optimal microclimate conditions. It is interesting to note that egg shells with other shapes, such as round or flat, or with hatching apertures positioned in a different place, could differ as well in the conditions provided for mites.

Even though this is a single record, it is, as far as we know, the first record of opportunistic use of a lepidopteran egg shell as a shelter, and offers evidence that empty eggs can increase the environmental complexity of leaves or other substrates. This effect would be possibly more intense in the case of eggs oviposited aggregately, or in plants that host a high abundance of eggs. Organisms that change the availability of resources to others are considered ecosystem engineers (Jones et al. 1994). It is known that some lepidopterans act this way by leaving leaf shelters (Vieira & Romero 2013) after caterpillar use, and by the vacant space left by mining species (Kagata & Ohgushi 2004). We suggest that another way through which lepidopterans—and other insects as well—could act as ecosystem engineers is through the differentiated microhabitat created by egg shells left empty after hatching. This would apply for systems in which some small arthropod species largely use these structures, and is an interesting, yet unrecorded, possibility.

ACKNOWLEDGEMENTS

We are grateful to the Laboratório Nacional de Luz Síncrotron for the permission to work within its property. We also thank Camila Vieira and an anonymous reviewer, Paulo Oliveira and André Freitas for encouraging this work, Thaís Postalí for helpful comments, Geoff Gallice for English corrections and Marina Ferraz




FIG. 1. **a)** Hatched Mimallonidae egg in dorso-lateral view, with mites (arrow) at the opposite extremity to the egg opening; **b)** Mite eggs (arrow) inside the Mimallonidae egg; **c)** Lateral view of the Mimallonidae egg after heating and hand manipulation, with a mite (arrow) that exited the egg interior; **d)** The same mite (arrow) close to the egg aperture.

de Camargo Barbosa for helping with the mite eggs identification. The authors thank Fapesp (2011/18580-8). L.L.M. thanks CNPq (132277/2012-3) for a graduate fellowship; M.C.V thanks Fapesp (2010/13619-0) and CAPES (1198-13-0) for graduate fellowships; S.F.S thanks Fapesp (2012/23399-3) for a post-graduate fellowship.

LITERATURE CITED

- ANTOR, R. J. & M. B. GARCÍA. 1995. A new mite-plant association: mites living amidst the adhesive traps of a carnivorous plant. *Oecologia*. 101(1): 51–54.
- BRABY, M. F. & K. NISHIDA. 2007. The immature stages, larval food plants and biology of neotropical mistletoe butterflies. I. The *Hesperocharis* group (Pieridae: Anthocharidini). *J. Lepid. Soc.* 61(4): 181–195.
- DYAR, H. G. 1900. Notes on the larval-cases of Lacosomidae (Peroporidae) and life-history of *Lacosoma chiridota* Grt. *J. New York Entom. Soc.* 8(3): 177–180.
- FOURNIER, V., J. A. ROSENHEIM, J. BRODEUR, L. O. LANEY, & M. W. JOHNSON. 2003. Herbivorous mites as ecological engineers: indirect effects on arthropods inhabiting papaya foliage. *Oecologia*. 135(3): 442–450.
- JONES, C. G., J. H. LAWTON, & M. SHACHAK. 1994. Organisms as ecosystem engineers. *Oikos* 69: 231–386.
- KAGATA, H. & T. OHGUSHI. 2004. Leaf miner as a physical ecosystem engineer: secondary use of vacant leaf mines by other arthropods. *Ann. Entomol. Soc. Am.* 97(5): 923–927.
- KAMINSKI, L. A., L. L. MOTA, A. V. L. FREITAS, & G. R. P. MOREIRA. 2013. Two ways to be a myrmecophilous butterfly: natural history and comparative immature-stage morphology of two species of *Theope* (Lepidoptera: Riodinidae). *Biol. J. Linn. Soc.* 108: 844–870.
- KAWASAKI, T., S. YANO, & M. H. OSKABE. 2009. Effects of wall structure and light intensity on the settlement of a predatory mite, *Euseius sojaensis* (Ehara) (Acari: Phytoseiidae). *App. Entom. Zool.* 44: 81–84.
- LIMA, V. O., P. R. DEMITE, C. VIEIRA, R. J. F. FERES, & G. Q. ROMERO. 2013. Contrasting engineering effects of leaf-rolling caterpillars on a tropical mite community. *Ecol. Entomol.* 38: 193–200.
- O'DOWD, D. J. & M. F. WILLSON. 1989. Leaf domatia and mites on Australasian plants: ecological and evolutionary implications. *Biol. J. Linn. Soc.* 37: 191–236.
- PETERSON, A. 1961. Some types of eggs deposited by moths, Heterocera-Lepidoptera. *Fla. Entom.* 44: 107–114.
- . 1966. Some eggs of moths among the Liparidae, Lasiocampidae, and Lacosomidae (Lepidoptera). *Fla. Entom.* 49: 35–42.
- STEHF, F. W. 1987. Order Lepidoptera, pp. 288–596. *In* Stehr, F. W. (ed.), *Immature insects*. Vol. 1. Kendall/Hunt Publishing Company, Dubuque, Iowa.
- VIEIRA, C. & G. Q. ROMERO. 2013. Ecosystem engineers on plants: indirect facilitation of arthropod communities by leaf-rollers at different scales. *Ecology*. 94: 1510–1518.
- WILLSON, M. R. 1991. Foliar shelters for mites in the Eastern Deciduous Forest. *Am. Midl. Nat.* 126: 111–117.

LUÍSA LIMA E MOTA*, ANDRÉ TACIOLI, SEBASTIAN FELIPE SENDOYA *Departamento de Biologia Animal, Universidade Estadual de Campinas, 13083-970 Campinas, São Paulo, Brazil* AND MAYRA CADORIN VIDAL *Department of Biological Sciences, University of Denver, Denver, CO 80208, United States of America.*

*Correspondence should be addressed to Luísa Lima e Mota, email: lulismota@yahoo.com.br

Submitted for publication 14 September 2013; revised and accepted 27 January 2014.