

Nomenclatural and faunistic notes on some Italian Dytiscidae (Insecta: Coleoptera)

Author: Nardi, Gianluca

Source: Integrative Systematics: Stuttgart Contributions to Natural History, 5(1) : 25-35

Published By: Stuttgart State Museum of Natural History

URL: <https://doi.org/10.18476/2022.827907>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

RESEARCH ARTICLE

Nomenclatural and faunistic notes on some Italian Dytiscidae (Insecta: Coleoptera)

GIANLUCA NARDI

Abstract

Unusual Italian records witnessing the active flight of two predaceous diving beetles, *Rhantus suturalis* (W. S. Macleay, 1825) and *Acilius (Acilius) sulcatus* (Linnaeus, 1758), are provided. The second species is recorded from the Aosta Valley region (N Italy) and from three northern Italian provinces (Asti, Padua, Sondrio) for the first time. The conservation status in Italy of both species is updated. *Deronectes silphoides* (Ponza, 1805), from the environs of Saluzzo in the Piedmont region (N Italy), was in the past treated as a *nomen dubium*. Later it was placed, as a *nomen oblitum*, in synonymy with *D. opatrinus* (Germar, 1824) (*nomen protectum*). However, it is herein newly declared a *nomen dubium* and removed from the above synonymy, since its type specimen is lost and its type locality is outside the area of distribution of the *D. opatrinus* species-group. At the same time, the identity of *Hydroporus silphoides* Villa & Villa, 1833 (*nomen nudum*), previously referred to *H. obscurus* Sturm, 1835, is declared as uncertain.

Keywords: Colymbetinae, Dytiscinae, Hydroporinae, nomenclature, chronogeonomy, conservation, flight, Italy.

Zusammenfassung

Der aktive Flug zweier Schwimmkäfer, *Rhantus suturalis* (W. S. Macleay, 1825) und *Acilius (Acilius) sulcatus* (Linnaeus, 1758), wird ungewöhnlicherweise aus Italien berichtet. Außerdem wird die letztgenannte Art zum ersten Mal aus der Region des Aosta-Tals (Norditalien) und aus drei norditalienischen Provinzen (Asti, Padua, Sondrio) nachgewiesen. Der Schutzstatus beider Arten für Italien wird aktualisiert. *Deronectes silphoides* (Ponza, 1805) aus der Umgebung von Saluzzo in der Region Piemont (Norditalien) wurde in der Vergangenheit als *nomen dubium* behandelt. Später wurde die Art als *nomen oblitum* als Synonym zu *D. opatrinus* (Germar, 1824) behandelt (*nomen protectum*). Sie wird hier erneut als *nomen dubium* deklariert und aus der obigen Synonymie entfernt, da ihr Typus-exemplar verloren gegangen ist und ihre Typuslokalität außerhalb des Verbreitungsgebietes der *D. opatrinus*-Arten-gruppe liegt. Zugleich wird die Identität von *Hydroporus silphoides* Villa & Villa, 1833 (*nomen nudum*), die bisher als *H. obscurus* Sturm, 1835 behandelt wurde, für unsicher erklärt.

Introduction

The aim of this short paper is to provide and discuss unusual Italian records of *Rhantus suturalis* (W. S. Macleay, 1825) and *Acilius (Acilius) sulcatus* (Linnaeus, 1758) related to their flight capacity, and update their chronogeonomy (sensu BRANDMAYR et al. 2007) and conservation status (BOLOGNA 2007; ROCCHI 2007). The status of *Deronectes silphoides* (Ponza, 1805) from the Piedmont region (N Italy) is also discussed.

Material and methods

The specimens examined herein were identified according to FRANCISCOLO (1979), DETTNER (1983) and NILSSON & HOLMEN (1995); the systematics and nomenclature follow NILSSON & HÁJEK (2022a). In the past (e.g. FRANCISCOLO 1979; ROCCHI 1982; RASTELLI & DELLA BEFFA 2007; ROCCHI 2007), *Rhantus suturalis*

(W. S. Macleay, 1825) was known as *R. pulverosus* (Stephens, 1828); for practical reasons, this latter name was not listed in the relative literature citations hereunder.

As stated also by COLONNELLI & NARDI (2019), the illustrations of the beetles in PONZA (1805) are on two plates, where single illustrations are aligned, the lines (“Lin.”) are numbered in the captions (Ponza 1805: 93–94), whereas individual illustrations in each plate are unnumbered. The names of the illustrated species are written in succession in the captions, whereas on the plates they are reported above each illustration. Hereunder, this information—the plate (pl.), the line (Lin.) and, in square brackets, the position of the illustration in the line—for the illustration of *Dytiscus silphoides* is reported as follows: “pl. 1, Lin. 4[.3]”.

The labels of the examined specimens are written in Italian; for practical reasons, the names of the administrative regions and provinces and collecting methods were translated into English hereunder, while other toponyms are written in Italian. The Italian regions are listed from North to South and from West to East, while other toponyms are listed alphabetically. The administrative provinces are listed according to the labels of the specimens, so provinces of recent institution (e.g., Prato in Tuscany,

Oristano in Sardinia, etc.) were not used. Possible interpolations are given in square brackets. Chorotypes were attributed according to VIGNA TAGLIANTI et al. (1999).

The conservation status in Italy of the involved species follows ROCCHI (2007) but was updated according to the IUCN (2022) criteria based on previously overlooked and subsequently published records (e.g., PONZA 1805; VON MARTENS 1824; CONTARINI 1843; SELLA 1864; DISCONZI 1865; RAGAZZI 1878; PAGLIA 1879; LAZZARINI 1895; IMPARATI 1897; RAGUSA 1921; HEYROVSKY 1968; MARCUZZI & TURCHETTO LAFISCA 1977; GIACHINO 1982; STOCH & DOLCE 1984; PILON 2002; SAPUPPO 2002; NARDI 2004, 2005; DETTNER 2006; ROCCHI & MASCAGNI 2006; TOLEDO 2006; RASTELLI & DELLA BEFFA 2007; MAZZA et al. 2008; PEDERZANI & PESARINI 2008; BOSI et al. 2009; CORRADI 2009; FERRARO 2009; MAZZOLDI 2009; ROMANO 2010; TERZANI et al. 2011; FORNACIARI 2011; SORGI et al. 2011; FABBRI et al. 2012; MAZZA et al. 2012; VIGLIOGLIA 2012; GUBELLINI 2013; MELANDRI 2013; PEZZI 2013; CIANFERONI et al. 2014; LUPI et al. 2014; LEO 2015; GATTI et al. 2016; ROCCHI & TERZANI 2016; ROCCHI et al. 2017; LENCIONI et al. 2019; TOLEDO & GROTTOLLO 2019; ANGELINI 2020; GROTTOLLO 2020; HENDRICH & BALKE 2020; ROCCHI & POGGI 2020; MOLA & TOLEDO 2021; ROCCHI et al. 2021a, 2021b, 2021c; STAUBLE 2021) and on the material examined.

All material examined for this study is deposited in Italian collections. The following abbreviations are used in the text: Coll. = Collection of; dint. = dintorni = environs; ex = specimen/s; Fraz. = Frazione = Hamlet; loc. = locality; hn = hand net; nc = no collector; ncd = no collection date; prov. = province; reg. = region; sdb = same data but. Acronyms of depositories are as follows:

ARDE	Coll. Associazione Romana di Entomologia c/o Museo Civico di Zoologia, Rome
CED	Coll. ERNESTO DE MAGGI c/o Museo civico di Zoologia, Rome
CEG	Coll. EMILIO GARAVAGLIA c/o Museo civico di Zoologia, Rome
CEM	Coll. ENRICO MIGLIACCIO, Rome
CEP	Coll. EUGENIO PACIERI c/o Museo di Storia Naturale dell'Università degli Studi di Firenze, Sezione di Zoologia "La Specola", Florence
CFB	Coll. FRANCESCO BIEGELEBEN c/o Museo civico di Zoologia, Rome
CGN	Author's private coll., Cisterna di Latina (Latina)
CGP	Coll. GIUSEPPE PACE, Rome
CGR	Coll. GIOVANNI EMILIO RASETTI c/o Museo civico di Zoologia, Rome
CMC	Coll. MARCELLO CERRUTI c/o Museo di Zoologia, Sapienza Università di Roma, Rome
CNCB	Centro Nazionale Carabinieri Biodiversità "Bosco Fontana", Marmirolo (Mantua)
CNS	Coll. NINO SANFILIPPO c/o Museo Civico di Storia Naturale "G. Doria", Genova (S. ROCCHI, pers. comm., 2022)
CPaC	Coll. PAOLO CORNACCHIA, Porto Mantovano (Mantua)
CPaM	Coll. PAOLO MALTZEFF, Rome
CPiC	Coll. PIETRO COSIMI c/o Museo di Storia Naturale dell'Università degli Studi di Firenze, Sezione di Zoologia "La Specola", Florence
CVS	Coll. VITTORIO ROSA c/o Museo di Zoologia, Sapienza Università degli Studi di Roma, Rome
MCC	Museo Civico di Storia Naturale di Carmagnola (Turin).

Results

Rhantus suturalis (W. S. Macleay, 1825)

Material examined

Piedmont reg., Cuneo prov.: Caramagna, 1973, nc, 2 ♂♂ (MCC); sdb 26.IV.1975, 1 ♀ (MCC); Caramagna, Bosco [del] Merlino, stagno, 28.III.1976, G. CURLETTI leg., 3 ♂♂, 3 ♀♀ (MCC). Piedmont reg., Torino prov.: Piobesi, 3.VI.1992, M. RASTELLI leg., 1 ♀ (MCC); Torino, Basse di Stura, 250 m a.s.l., 21.X.1971, D. GIANASSO leg., 1 ♂ (CPaC). Veneto reg., Venice prov.: [San Michele al Tagliamento, Fraz.] Bibione, [1 m a.s.l.] 23.VI.1983, COSTELLA leg., 1 ♂ (CGN). Veneto reg., Verona prov.: Valeggio sul Mincio, Via I Maggio, 45°20'57"N 10°43'41"E, 87 m a.s.l., 3.VI.2018, in letterbox, G. NARDI leg., 1 ex (CGN). Lombardy reg., Mantua prov.: Marmirolo, Riserva Naturale Statale "Bosco Fontana", Piazza Carbone, 45°12'09"N 10°45'00"E, 48 m a.s.l., 5.IX.2018, G. NARDI leg., at 90 W incandescent light, 20:00–21:00 hrs, 1 ♀ (CNCB). Lombardy reg., Pavia prov.: Torre d'Isola, 14.III.1990, G. PACE leg., puddle in a canal, 1 ♀ (CGN). Tuscany reg., Florence prov.: Prato, dint. Centrale ENEL, 30.III.1985, I. SCALI leg., 1 ♂, 2 ♀♀ (CGN). Tuscany reg., Livorno prov.: Tirrenia, VIII.1963, E. MIGLIACCIO leg., 1 ♀. (CEM). Abruzzo reg., Pescara prov.: Montesilvano, 22.VII.1975, [P. MALTZEFF leg.], 1 ♀ (CPaM). Lazio reg., Latina prov.: Cisterna di Latina, strada parallela al Fosso Teppia [= Via Tivera], 41°35'22"N 12°52'42", 73 m a.s.l., 23.IV.1989, G. NARDI & C. ESPOSITO leg., shallow cemented basin with rainwater in a large meadow with asphodels, 1 ♀ (CGN); sdb 31.III.1991, G. NARDI leg., sdb water temperature 14°5 C, 1 ♀ (CGN). Lazio reg., Rome prov.: [Roma], Maccarese, 12.V.1912, Luig. [= P. LUIGIONI leg.], 1 ♂ (CMC); Roma, Settecami, 1991, C. DE LIBERATO leg., 1 ♀ (CGN). Sardinia reg., Cagliari prov.: Domusnovas, 2.VI.1980, L. FANCELLO leg., 1 ♀ (CGN). Sardinia reg., Nuoro prov.: [Siniscola.] Case Capo Comino, 14–19.VI.2007, P. CORNACCHIA leg., attached to a wall in a house, attracted by artificial lights, 1 ♀ (CPaC); sdb 20–29.VI.2007, on the ground in a house, attracted by artificial lights, 1 ♀ (CPaC). Sardinia reg., Sassari prov.: Torre Grande di Oristano, 4–7.VII.1988, E. MIGLIACCIO leg., at lights, 1 ♂ (CEM).

Chorotype and distribution

Subcosmopolitan (cf. NILSSON & HÁJEK 2022a). *Rhantus suturalis*, originated in the highlands of New Guinea ca. 6.0–2.7 Mya (BALKE et al. 2009; TOUSSAINT et al. 2013), is a supertramp and is the most widely distributed dytiscid on earth, being widespread in the Palaearctic, Oriental, Pacific and Australian regions (BILTON 2014; HÁJEK 2017; HENDRICH & BALKE 2021; NILSSON & HÁJEK 2022a, 2022b). In Italy it is recorded from all regions except Aosta Valley (ROCCHI 2007; S. ROCCHI, pers. comm. 2022).

Ecology

This euryoecious pioneer species occurs chiefly in stagnant waters, from sea level to high mountains; it is an excellent flyer and is frequently collected at light (cf. FRANCISCOLO 1979; ROCCHI 1982; NILSSON & HOLMEN 1995; ANGELINI 1998; NARDI & MALTZEFF 2001; ŠUSTEK 2002; CASTRO et al. 2003; NARDI 2005; KEHL & DETTNER 2007; ROCCHI 2007; BURGHELEA et al. 2008; LÓPEZ-PÉREZ &

MILLÁN SÁNCHEZ 2012; BILTON 2014; LEO 2015; TOLEDO & GROTTOLLO 2019; MATSUSHIMA & YOKOI 2020; MOLA & TOLEDO 2021; RAMADAN & RAMADAN 2021) and in Malaise traps (MAJZLAN 2014, 2020). The diel flight activity of this species has one peak in the early/late evening, in summer and autumn (CSABAI et al. 2012). Adults are generalist predators and scavengers (BOSI 2001). An experiment demonstrated that *R. suturalis* females release sex pheromones to attract mating partners (HERBST et al. 2011). GROPPALI (1992: 37) recorded two specimens of “*Rhantus* [sic!] *notatus* (Fabr.)” found among the stomach contents of five *Glareola pratinicola* Linnaeus, 1758 (Aves: Charadriiformes: Glareolidae) in Apulia (SE Italy). *Rhantus notatus* sensu (Fabricius, 1781) is currently known as *R. frontalis* (Marsham, 1802) (cf. FRANCISCOLO 1979; NILSSON & HOLMEN 1995; NILSSON & HÁJEK 2022b). This species is known with certainty in Italy only from the Piedmont region (N Italy) and on the basis of very old specimens (ROCCHI 2007). In this framework, the record by GROPPALI (1992) probably refers to *R. suturalis*, which is the sole congeneric species occurring in the Apulia region (ROCCHI 2007).

Conservation status in Italy

This species is widespread and generally common in Italy. Based on the available data, it is hereunder evaluated as of “Least Concern (LC)” at the Italian scale, whereas previously it had been classified as “Data Deficient (DD)” (ROCCHI 2007).

Notes

The above localities in the Verona province (Veneto region) and Lombardy are in the Po Plain, where the species has been recorded from many sites, chiefly in the temporary, stagnant waters of ponds, rain puddles, flooded meadows, rice fields, etc. (cf. NARDI 2004; TOLEDO 2006; ROCCHI 2007; LUPI et al. 2014; TOLEDO & GROTTOLLO 2019). The above specimen from Valeggio sul Mincio (Verona prov.) was found in the author’s letterbox in a recently-built neighbourhood located at the foot of a small (120 m), wooded morainic hill and on the outskirts of town, not far from the cultivated fields of the Po Plain. The letterbox is attached to a gate in front of a road and some private gardens. There are no water bodies within a radius of 100 m except for a fish pond in a private garden and some swimming pools, but the Mincio river is only 480 m from the collecting site. The species was already recorded from another site in the same commune and from bordering areas (NARDI 2004; TOLEDO 2006; ROCCHI 2007; TOLEDO & GROTTOLLO 2019). The letterbox is a metallic, aluminium-coloured parallelepiped (31 x 22 x 10 cm) with a 17.5 x 3 cm slot at the top for inserting mail, covered by a small, liftable metal flap. Many species of water beetles, including *R. suturalis*, are attracted in flight by reflection-polariza-

tion signals emitted by the surface of, e.g., coloured car roofs (MUGGLETON 1966; JÄCH 1997; NILSSON 1997; VONDEL 1998; KRISKA et al. 2006; ROCCHI 2011; G. NARDI, unpublished data). This behaviour would also explain the above unusual find: the small flap of the letterbox was left open by the postman, after which the beetle landed in the small slot at the top of the letterbox and ended up trapped inside, where it died from heat exposure. The only other beetle trapped in the same letterbox during 2013–2022 was a specimen of *Mimela junii junii* (Duftschmid, 1805) (Scarabaeidae: Rutelinae) on 1.XI.2018.

The listed specimen of *Rhantus suturalis* from Mantua prov. (Lombardy) comes from a Nature Reserve where the species had been previously collected, but never at light (NARDI 2004). Malaise and window flight traps were heavily used in this reserve in the last two decades (cf. CERRETTI et al. 2004; BIRTELE & HARDERSEN 2012); however, differently from other areas (MAJZLAN 2014, 2020), the only dytiscid so far collected with these methods was a specimen of *Bidessus minutissimus* (Germar, 1824) (NARDI 2004; P. CORNACCHIA & G. NARDI, unpublished data).

The above records of *R. suturalis* from other regions are provided for a better knowledge of its Italian chronogeonomy or are related to the flight capacity of this species. Surprisingly, in the Cuneo province (Piedmont region, N Italy) it had so far been recorded from a single site only (RASTELLI & DELLA BEFFA 2007).

Acilius (Acilius) sulcatus (Linnaeus, 1758)

Material examined

Trentino-Alto Adige/Südtirol reg., Bolzano prov.: Aslago [= Bolzano, Oltrisarco-Aslago quarter], 11.III.1930, nc, 1 ♂ (CFB); Bolzano, Renon, VIII.[19]36, REANDA [leg.], 1 ♂ (CMC); Eppan [= Appiano sulla strada del vino], 15.V.1929, nc, 1 ♂ (CFB); sdb 21.V.1929, 1 ♀ (CFB); Eppan [= Appiano sulla strada del vino], illegible loc., 8.V.1929, nc, 2 ♂♂ (CFB); Eppan, Unter-rain [= Appiano sulla strada del vino, Fraz. Riva di Sotto], 27.IV.1929, nc, 2 ♀♀ (CFB). Piedmont reg., Asti prov.: Castelnovo Don Bosco, 6.VIII.1964, D. GIANASSO leg., 1 ♂ (CEM). Piedmont reg., Novara prov.: Fontaneto d’Agogna, 300 m a.s.l., 10.IX.1987, D. FONTANETO leg., 1 ♀ (CGN); sdb 16.X.1989, 1 ♀ (CGN); sdb 20.X.1989, 1 ♂ (CGN); Valenzani, 8.IX.1966, P. TESTA leg., 2 ex (CNS). Lombardy reg., Brescia prov.: Brescia dint., ncd, MARCHESI [leg.], 2 ♂♂ (CMC); sdb III.1938 2 ♀♀ (CMC). Lombardy reg., Sondrio prov.: Aprica, Pian Zembro, 21.VII.1957, [M.] BARAJON leg., 1 ♀ (CVS). Aosta Valley reg., Aosta prov.: Gran Villa, Lozoe, loc. Champloeg, 1650 m a.s.l., 14.V.1992, R. SINDACO leg., 1 ♀ (MCC). Veneto reg., Padua prov.: Colli Euganei, [Torreglia, Fraz.] Luvigliano, [about 30 m a.s.l.] X.1965, LOMAZZI leg., 1 ♂ (CPaC). Veneto reg., Treviso prov.: Monte Pizzo, 29.V.1989, nc, 1 ♂, 1 ♀ (CGN). Emilia-Romagna reg., [Forlì prov.]: [Cesenatico, Fraz.] Sala, 15.X.[19]32, nc, 1 ♂ (CMC). Emilia-Romagna reg., Piacenza prov.: App. Piacent. [= Piacentine Apennines], lago Cacciasù [untraced site], 9.VI.1956, [M.] BARAJON leg., 2 ♀♀ (CVS). Emilia-Romagna reg., Ravenna prov.: Ravenna, 28.V.1961, Senni leg., 1 ♂, 1 ♀

(ARDE). Umbria reg., Perugia prov.: Castel Ritaldi, Fraz. Bruna, summer 1949, nc, 1 ♂ (CPaM). [Lago] Trasimeno, IX. [no year], nc, 1 ♂ (CGR). Tuscany reg., Arezzo prov.: [Cavriglia, Fraz.] Montegonzi, 28.VII.1990, R. LISA leg., 1 ♂ (CGN); Cortona, 700 m a.s.l., VI.1931, nc, 1 ♀ (CEG); sdb no altitude, ncd, E. GARAVAGLIA leg., 1 ♂ (CEG); sdb GARAVAGLIA F.lli [= Fratelli = Brothers] [leg.], 2 ♀♀ (CEG); sdb 22.VII.[no year], E. GARAVAGLIA leg., 1 ♂ (CEG). Tuscany reg., Florence prov.: Massarella, Padule di Fucecchio, 4.II.1990, I. SCALI leg., 1 ♂ (CGN). Tuscany reg., Livorno prov.: Livorno dint., VIII.[19]36, D. C. [= ? leg.], 1 ♀ (CMC); Livorno, Tombolo, VII.[19]34, nc, 1 ♂ (CMC). Tuscany reg., Pisa prov.: Pisa, ncd, G. E. RASETTI leg., 1 ♂ (CGR); Pisa, Marina, ncd, nc, 1 ♀ (CGR); Tombolo, 5.VII.1936, nc, 1 ♀ (CGR). Lazio reg., Frosinone prov.: Monti Lepini, Morolo, Monte Alto, versante E [= E slope], 1300 m a.s.l., 7.V.2001, C. ESPOSITO leg., deep tank (diameter 75 m) in cement and fiberglass, 2 ♀♀ (CGN); Paliano, loc. La Selva, 25.VIII.1982, M. MEI leg., 1 ♀ (CGN). Lazio reg., Latina prov.: Cisterna di Latina, Via. E. Fermi, 41°35'21"N 12°50'18"E, 70 m a.s.l., 8.IV.1980, G. NARDI leg., in a small, almost dry puddle, in the excavation of the foundation of a house under construction, 1 ♀ (CNA); sdb 41°35'25"N 12°50'08"E. 15.IV.1984, 76 m a.s.l., R. PERGOLES leg., on small, low wall of vegetable garden, away from aquatic environments, 1 ♂ (CNA); Monti Lepini, Bassiano, loc. Antignana, 400 m a.s.l., 19.V.1991, G. NARDI leg., cattle pond devoid of vegetation, water temperature 19 °C, 1 ♀ (CGN); Monti Lepini, Norma, Monte Gorgoglione, Pozzo di Monte, 700 m a.s.l., 19.VI.1999, C. ESPOSITO leg., large pond, 1 ♀ (CGN). Lazio reg., Rome prov.: [Colli Albani,] Grottaferrata, [200–300 m] 7.X.1945, [E.] DE MAGGI leg., 1 ♂ (CED); Roma, Acilia, 6.IV.[19]31, nc, 1 ♀ (CMC); sdb 6.XI.1932, Coll. CASTELLANI, 1 ♀ (CMC); sdb 18.IV.1934, 1 ♂ (CMC); sdb 3.IV.[19]38, [M.] CERRUTI [leg.], 1 ♀ (CMC); Roma, Capocotta, 14.VIII.[19]39, [E.] DE MAGGI [leg.], 1 ♀ (CMC). Abruzzo reg., L'Aquila prov.: Gran Sasso, Campo Imperatore [1500–2100 m], IV.1977, E. PACIERI leg., 1 ♀ (CPiC); sdb 30.IV.1977, 1 ♀ (CEP); [Parco Nazionale d'Abruzzo,] Pescasseroli, VI.[19]36, [M.] CERRUTI leg. 1 ♂ (CMC).

Chorotype and distribution

Sibero-European, reaching Algeria (NILSSON & HÁJEK 2022b). A recent detailed record from the Oriental region (India, West Bengal) (CHANDRA et al. 2008) appears erroneous since it was later ignored (GHOSH & NILSSON 2012; BUKONTAITE et al. 2014; CHANDRA et al. 2017; HÁJEK 2017; CHANDRA et al. 2021; NILSSON & HÁJEK 2022a, 2022b). In Italy, *A. (A.) sulcatus* was previously recorded from all regions except Aosta Valley and Molise (ROCCHI 2007; ROCCHI et al. 2018; ROCCHI, pers. comm. 2021); it is here recorded from Aosta Valley (N Italy) for the first time.

Ecology

Acilius (Acilius) sulcatus is a pioneer species whose adults have high flight ability and can disperse from overwintering ponds to temporary pools in the spring, returning to permanent waters in the summer as the temporary pools dry up (cf. DAVY-BOWKER 2002; KEHL & DETTNER 2007; GIORIA 2014; IVERSEN et al. 2017). Its altitudinal range varies from sea level up to 1776 m (BERGSTEN & MILLER 2006; ROCCHI 2007; SORGI et al. 2011; ENKHNASAN &

BOLDGIV 2019). In Italy, this species is characteristic of muddy pools, often turbid and devoid of vegetation, typically in water basins for livestock, even of artificial origin, where it is often quite abundant; more rarely it lives in large ponds, as long as there are areas free from aquatic vegetation (TOLEDO 2016; TOLEDO & GROTTOLLO 2019), and is most common in the absence of fish; adults overwinter in the water, even when covered by ice (BERGSTEN & MILLER 2006), and can be preyed upon by the grey heron (*Ardea cinerea* Linnaeus, 1758) (Aves: Ciconiiformes: Ardeidae) (DRAULANS et al. 1987). Adults have also been collected by Malaise traps (MAJZLAN 2014, 2020) and in flight (MUGGLETON 1966; FORNACIARI 2011; MELANDRI 2013).

Conservation status in Italy

This species is relatively widespread but localized in Italy; it is here evaluated as of “Least Concern (LC)” at the Italian scale, whereas it had previously been classified as “Data Deficient (DD)” by ROCCHI (2007).

Notes

The above specimens from Cisterna di Latina (Latium, Latina prov.) confirm the high flight ability of this species. At the time of collection there were cultivated fields and natural meadows interspersed with houses and gardens. The only aquatic environments present were small roadside gutters for the collection of rainwater, large temporary puddles in the meadows and very few small, natural ditches. The female collected in 1980 was netted in a small, almost dry puddle in the foundations of a house under construction. A male of *Colymbetes fuscus* (Linnaeus, 1758) and some specimens of *Agabus (Gaurodytes) bipustulatus* (Linnaeus, 1767) were also netted (CGN) from a different part of the same foundations. The above specimen from 1984 was found attached to the small, low wall of a vegetable garden, away from aquatic environments. In the following decades, this site underwent intense building development and these species were no longer collected there (G. NARDI, unpublished data). Fortunately, in the same province, *Acilius (Acilius) sulcatus* was recorded also from a National Park (NARDI 2005) and is here recorded from the Lepini Mountains, an unprotected area of great conservation value (cf. CORSETTI et al. 2015; CORSETTI & MAROZZA 2020).

The other above records are provided for a better understanding of the Italian chorogeonomy of this localized species; it was previously (ROCCHI 2007) unrecorded from the Aosta Valley region. The elevation of the Aosta Valley site (1650 m) is also interesting, since the species ranges from 0 to 1776 m in Italy (ROCCHI 2007; SORGI et al. 2011), but most records come from elevations lower than 1300 m (cf. ROCCHI 2007; MAZZA et al. 2012; GATTI et al. 2016; LENCIONI et al. 2019; TOLEDO & GROTTOLLO 2019). The species had so far not been recorded (cf. ROCCHI 2007) from

the Asti (Piedmont), Padua (Veneto), and Sondrio (Lombardy) provinces, whereas it was only known from single sites in the provinces of Novara (Piedmont), Frosinone (Lazio) and L'Aquila (Abruzzo) (CONTARINI 1843; ROMANO 2010; VIGLIOGLIA 2012; IPLA 2017).

Deronectes silphoides (Ponza, 1805) (*nomen dubium*)

Dytiscus Silphoides Ponza, 1805. PONZA 1805: : 82, pl. 1, Lin. 4[.3]

Notes

The original description of this species is as follows: “Magnitudo *D. Fulvi* [= *Dytiscus fulvus*]: depressus, ovatus, niger; thorace utrique marginato, margine incrassato; antennis setaceis, femoribusque rufis.” (PONZA 1805: 82, as *Dytiscus silphoides*). No type locality is listed, but as stated in the title of the paper it is in the environs of Saluzzo (Cuneo province, Piedmont region, Italy).

The above “*D. Fulvi*”, is “*D. Fulvus*. *Ent. Paris*. I. 66. 5. *Geoffr. Paris*. I. 184. 5” (PONZA 1805: 81), and corresponds to *Dytiscus fulvus* Fabricius, 1801, currently *Haliphilus* (*Liaphilus*) *fulvus* (Halipilidae) (cf. VONDEL 1997: 66).

Although the type material of *Dytiscus silphoides* is lost (DUTTO et al. 2009) and the original description does not conform to modern standards, the illustration of its habitus (PONZA 1805: pl. 1, Lin. 4[.3]) is sufficiently diagnostic to be assigned to *Deronectes* Sharp, 1882, and not to *Haliphilus* Latreille, 1802.

Ponza's species was later listed in a few papers (e.g., ANONYMOUS 1806: 280, as *Dytiscus silphoides* Ponza; VON MARTENS 1824: 502, as *D. silphoides* [no author]; GEMMINGER & HAROLD 1868: 426, as *Haliphilus silphoides* Ponza; SHERBORN 1930: n. 5989, as *D. silphoides* Ponza; DUTTO et al. 2009: 429, as *D. Silphoides* Ponza), but it was overlooked in capital works (e.g., ZIMMERMAN 1920; FRANCISCOLO 1979).

Recently, NILSSON (2001: 151) treated *Dytiscus silphoides* as a *nomen dubium* in the genus *Deronectes* Sharp, 1882. Later, NILSSON & FERY (2006: 58) declared this species as a *nomen oblitum*, in order to ensure the continuous usage of its junior synonym *Deronectes opatrinus* (Germar, 1824) as a valid name (*nomen protectum*), and this is the current status of these two names (NILSSON & HÁJEK 2022b: 29).

Nevertheless, for three reasons, it is not possible that *Deronectes silphoides* (Ponza) is a synonym of *D. opatrinus*:

(1) No species of the *Deronectes opatrinus* species-group (FERY & BRANCUCCI 1997; FERY & RIBERA 2018) occurs in Italy (cf. ROCCHI 2007, 2021; NILSSON & HÁJEK 2022b).

(2) The old Italian literature records of *D. opatrinus* are based on misidentifications of *D. fairmairei* (Leprieur, 1876) (FRANCISCOLO 1979: 442) and/or *D. moestus*

inconspectus (Leprieur, 1876) [cf. ANGELINI 1984: 107, as *D. moestus* (Fairmaire, 1858)]. These species are not close to *D. opatrinus* but belong to two other species-groups (FERY & BRANCUCCI 1997; NILSSON 2001).

(3) NILSSON & FERY (2006: 58) attributed such synonymy to VILLA & VILLA (1833: 7), who under “*Hydroporus opatrinus* Ill. [sic!]” listed “*Hydroporus silphoides* Peirol. [= Peiroleri]”, not *Dytiscus silphoides* Ponza.

No paper by Peiroleri mentioning *H. silphoides* was found, so the authorship of this name is as follows: *Hydroporus silphoides* Villa & Villa, 1833 (*nomen nudum*). Later, the same authors (VILLA & VILLA 1844: 424, as “*Hydroporus silphoides* Peirol.”) listed it from the plain of Lombardy, with *obscurus* Jan (*nomen nudum*) [not *H. obscurus* Sturm, 1835] and *opatrinus* [sensu] Villa [& Villa], 1833 as its synonyms. *Hydroporus obscurus* Jan was already listed as a “synonym” of “*Hydroporus opatrinus* Ill. [sic!]” by VILLA & VILLA (1833); it corresponds to *H. obscurus* De Cristofori & Jan, 1832 (*nomen nudum*) from Sicily (DE CRISTOFORI & JAN 1832: 14, as *H. obscurus nob.* [= *nobis* = De Cristofori & Jan]). It must be underlined that the authorship of this *nomen nudum* must be assigned to both these authors, not only to JAN (VILLA & VILLA 1833, 1844) or DE CRISTOFORI (NILSSON & HÁJEK 2022b).

GHILIANI (1887: 227) recorded *Hydroporus* “*silphoides* Peiroleri, Ponza (*obscurus* Hen. [(sic!)])” from Borgomanero (Novara prov., Piedmont) and *H. “opatrinus* Villa, Germ. Aubè” from brooks of the Novara and Saluzzo areas (Piedmont). BAUDI (1890: 35) referred the former GHILIANI record to *H. “obscurus* St. [= Sturm, 1835] (*silphoides* Peirol., Ponza, p. 227)”, while the latter one to *Deronectes bombycinus* (Leprieur, 1876), which is a synonym of *D. fairmairei* (Leprieur, 1876) (NILSSON & HÁJEK 2022a, 2022b).

LUIGIONI (1929: 155), following BAUDI (1890), referred “*silphoides* Ponza” to *Hydroporus obscurus* Sturm, 1835.

ANGELINI (1984: 106), based on the records of GHILIANI (1887, as *H. silphoides* Peir.) and VILLA & VILLA (1844, as *H. silphoides* Peirol), listed *Hydroporus obscurus* Sturm, 1835 as doubtful for the Italian fauna, and this Holarctic species is currently unrecorded from Italy (ROCCHI 2021; NILSSON & HÁJEK 2022b). In this framework, the identity of *Hydroporus silphoides* Villa & Villa, 1833 (*nomen nudum*) remains uncertain.

Five species of *Deronectes* are recorded from the Piedmont region (cf. ROCCHI 2007): *D. angelinii* Fery & Brancucci, 1997, *D. aubei aubei* (Mulsant, 1843), *D. semirufus* (Germar, 1845), *D. fairmairei* and *D. moestus inconspectus*. Excluding *D. fairmairei*, which is known from Piedmont with doubt, the remaining taxa are known also from the Cuneo province (cf. ROCCHI 2007). *Deronectes silphoides* (Ponza, 1805) is therefore identical to one of these species; unfortunately, on the basis of its very short description (PONZA 1805), it is only possible to establish

that it is different from *D. angelinii* and *D. semirufus*. Therefore, *Deronectes silphoides* (Ponza, 1805) is here newly (see NILSSON 2001) declared as a *nomen dubium*. In this way, since *Deronectes silphoides* has priority over all five *Deronectes* taxa mentioned above as well as *D. moestus moestus*, no nomenclatural instability is created.

Discussion

The faunistic knowledge on Italian Dytiscidae was optimally summarized by ROCCHI (2007), which represents the basis for updating the conservation status of species in this family in Italy. Unfortunately, the conservation status of many species in this family appears to be clearly worsening, especially in lowland areas and agricultural landscapes where lentic aquatic environments are often destroyed and/or altered by human activities (cf. MINELLI 2001; BATTISTI 2004; STOCH 2005; STOCH & NASELLI-FLORES 2014). An additional threat is the Louisiana crayfish, *Procambarus clarkii* Girard, 1852 (Malacostraca: Decapoda: Cambaridae), an invasive predaceous species that is spreading throughout Italy (cf. BATTISTI & SCALICI 2020; LO PARRINO et al. 2020), causing the decrease and/or disappearance of many populations of water beetles (cf. PEDERZANI & FABBRI 2006; BAZZATO et al. 2016). In this framework, it is important and useful to provide recent faunistic data on water beetles allowing a correct evaluation of their conservation status; unfortunately, it is easy to predict that the conservation status of one of the above species, *Acilius (Acilius) sulcatus*, now classified as “Least concern”, will worsen, since its aquatic environments are often destroyed and/or altered by human activities.

Acknowledgements

Thanks to FERNANDO PEDERZANI (Ravenna) and SAVERIO ROCCHI (Museo di Storia Naturale dell'Università degli Studi di Firenze, Sezione di Zoologia “La Specola”, Florence) for the revision of the manuscript, and to PAOLO AUDISIO (Sapienza Università di Roma, Rome), LUCA BARTOLOZZI (Museo di Storia Naturale dell'Università degli Studi di Firenze, Sezione di Zoologia “La Specola”, Florence), FELICETTA COMANDINI, PAOLO CORNACCHIA (Porto Mantovano, Mantua), GIANFRANCO CURRETTI (Museo Civico di Storia Naturale di Carmagnola, Turin), CARMINE ESPOSITO (Velletri, Rome), FRANCO MASON (Verona), ANGELO NARDI (Cisterna di Latina, Latina), DINO PANICCIA (Frosinone), ROBERTO PERGOLES (Cisterna di Latina, Latina), EMANUELE PIATTELLA (Rome), ROBERTO POGGI (Museo Civico di Storia Naturale “G. Doria”, Genoa), SAVERIO ROCCHI (Museo di Storia Naturale dell'Università degli Studi di Firenze, Sezione di Zoologia “La Specola”, Florence) and LAURA SPADA (Centro Nazionale Carabinieri Biodiversità “Bosco Fontana”, Marmirolo, Mantua) for their support.

Last but not least, I thank the following, recently deceased Roman colleagues: PIERO COSIMI, GIANNI GOBBI, PAOLO MALTZEFF, EUGENIO PACIERI and AUGUSTO VIGNA TAGLIANTI.

References

- ANGELINI, F. (1984): Catalogo topografico dei Coleotteri Haliplidae, Hygrobiidae, Dytiscidae e Gyrinidae d'Italia. – *Memorie della Società Entomologica Italiana* **61A** (1982): 45–126.
- ANGELINI, F. (1998): Coleotterofauna reperita mediante trappola luminosa in due stazioni umide della Basilicata (Italia meridionale). – *Quaderni della Stazione di Ecologia del Civico Museo di Storia Naturale di Ferrara* **11**: 7–37.
- ANGELINI, F. (2020): Contribution to the knowledge of beetles (Insecta Coleoptera) of some protected areas of Apulia, Basilicata and Calabria (Italy). – *Biodiversity Journal* **11** (1): 85–254.
- ANONYMOUS (1806): LII. Proceedings of Learned Societies.. – *The Philosophic Magazine: comprehending the various branches of Science, the Liberal and Fine Arts, Agriculture, Manufactures, and Commerce* **24** (February, March, April, and May 1806): 271–281.
- BALKE, M., RIBERA, I., HENDRICH, L., MILLER, M. A., SAGATA, K., POSMAN, A., VOGLER, A.P. & MEIER, R. (2009): New Guinea highland origin of a widespread arthropod supertramp. – *Proceedings of the Royal Society B: Biological Sciences* **276** (1666): 2359–2367.
- BATTISTI, C. (2004): Frammentazione ambientale, connettività, reti ecologiche. Un contributo teorico e metodologico con particolare riferimento alla fauna selvatica, 248 pp.; Rome (Provincia di Roma, Assessorato alle Politiche Ambientali, Agricoltura e Protezione Civile).
- BATTISTI, C. & SCALICI, M. (2020): First records of the red swamp crayfish *Procambarus clarkii* (Girard, 1852) (Decapoda Cambaridae) from a small circum-Sardinian island (central Mediterranean Sea). – *BioInvasions Records* **9** (2): 333–339.
- BAUDI, F. (1890): Catalogo dei Coleotteri del Piemonte. – *Annali della Reale Accademia d'Agricoltura di Torino* **32** (1889): 1–274.
- BAZZATO, E., CILLO, D., NUOLI, M. T., FOIS, F. & SCARAVELLI, D. (2017): New distributional data of genus *Hydrophilus* Leach, 1815 in Sardinia (Coleoptera, Hydrophiloidea, Hydrophilidae). *Natura Rerum* **4** (2015–2016): 71–77.
- BERGSTEN, J. & MILLER, K. B. (2006): Taxonomic revision of the Holarctic diving beetles genus *Acilius* Leach (Coleoptera: Dytiscidae). – *Systematic Entomology* **31**: 145–197. <https://doi.org/10.1111/j.1365-3113.2005.00309.x>
- BILTON D. T. (2014): Chapter 9. Dispersal in Dytiscidae. – In: YEE, D. A. (ed.): *Ecology, systematics, and the natural history of predaceous diving beetles (Coleoptera: Dytiscidae)*, pp. 387–407; Berlin, Heidelberg (Springer Science + Business Media B.V.).
- BIRTELE, D. & HARDERSEN, S. (2012): Analysis of vertical stratification of Syrphidae (Diptera) in an oak-hornbeam forest in northern Italy. – *Ecological Research* **27** (4): 755–763. <https://doi.org/10.1007/s11284-012-0948-2>
- BOLOGNA, M. A. (2007) Criteria for the compilation of the first red list of species of national and regional interest. – In: RUFFO, S. & STOCH, F. (eds.): *Checklist and distribution of the Italian fauna*, pp. 63–65. – *Memorie del Museo Civico di Storia Naturale di Verona*, 2. Serie. Sezione Scienze della Vita **17** (2006).
- BOSI, G. (2001): Observations on colymbetina predation based on crop contents analysis in three species: *Agabus bipustulatus*, *Ilybius subaeneus*, *Rhantus suturalis* (Coleoptera Dytiscidae). – *Bollettino della Società Entomologica Italiana* **133** (1): 37–42.

- BOSI, G., BO, T. & FENOGLIO, S. (2009): Alcune considerazioni sulla distribuzione di Noteridae e Dytiscidae (Coleoptera) nella provincia di Alessandria. – *Rivista Piemontese di Storia Naturale* **30**: 79–93.
- BRANDMAYR, P., CASALE, A., PUZZO, F. & SCALERCIO, S. (2007): Cronogeonomy analysis: some examples regarding species of the Italian fauna. – In: RUFFO, S. & STOCH, F. (eds.): Checklist and distribution of the Italian fauna, pp. 41–45. – *Memorie del Museo Civico di Storia Naturale di Verona*, 2. Serie. Sezione Scienze della Vita **17** (2006).
- BUKONTAITE, R., MILLER, K. B. & BERGSTEN, J. (2014): The utility of CAD in recovering Gondwanan vicariance events and the evolutionary history of Aciliini (Coleoptera: Dytiscidae). – *BMC Evolutionary Biology* **14** (5), 1–18. <https://doi.org/10.1186/1471-2148-14-5>
- BURGHELEA, C., ZAHARESCU, D. & PALANCA, A. (2008): Spatial and temporal distribution of three aquatic beetles (Dytiscidae) in Monegros arid zone, NE Spain. – *Analele Științifice Ale Universității “Al. I. Cuza” Iași (Serie Nouă). Secțiunea I. Biologie Animală* **54**: 101–111.
- CASTRO, A., HIDALGO, J. M. & CÁRDENAS, A. M. (2003): Nuevos datos sobre los coleópteros acuáticos del Parque Nacional de Doñana (España): capturas realizadas mediante trampas de luz y técnicas de muestreo para fauna edáfica (Coleoptera). – *Boletín de la Sociedad Entomológica Aragonesa (S.E.A.)* **33**: 153–159.
- CERRETTI, P., HARDERSEN, S., MASON, F., NARDI, G., TISATO, M. & ZAPPAROLI, M. (eds.) (2004): Invertebrati di una foresta della Pianura Padana, Bosco della Fontana. Secondo contributo. Conservazione Habitat Invertebrati 3, 304 pp.; Verona (Cierre Grafica Editore).
- CHANDRA, G., MANDAL, S. K., GHOSH, A. K., DAS, D., BANERJEE, S. S. & CHAKRABORTY, S. (2008): Biocontrol of larval mosquitoes by *Acilius sulcatus* (Coleoptera: Dytiscidae). – *BMC Infectious Diseases* **8** (1): 1–8. <https://doi.org/10.1186/1471-2334-8-138>
- CHANDRA, K., DEEPA, J. & SHIVASHANKAR, C. (2021): Chapter 34. Insecta: Coleoptera (Aquatic). – In: CHANDRA, K., DEEPA, J., RAGHUNATHAN, C., JADHAV, S. S. & KARUTHAPANDI, M. (eds.): Current status of faunal diversity in Telangana, pp. 227–235; Kolkata (Zoological Survey of India).
- CHANDRA, K., JAISWAL, D. & GUPTA, D. (2017): Chapter 24: Insecta: Coleoptera (Aquatic Beetles). – In: CHANDRA, K., GOPUI, K. C., RAO, D. V., VALARMATHI, K. & ALFRED, J. R. B. (eds.): Current status of freshwater faunal diversity in India, pp. 379–400; Kolkata (Zoological Survey of India).
- CIANFERONI, F., ROCCHI, S., TERZANI, F. & MASCAGNI, A. (2014): Coleotterofauna acquatica e semiacquatica del bacino idrografico del Fiume Sieve (Toscana) (Insecta Coleoptera). *Bollettino della Società Entomologica Italiana* **146** (2): 51–81.
- COLONNELLI, E. & NARDI, G. (2019): The Cleridae, Anthribidae and Curculionidae (Coleoptera) described by Ponzá. – *Zootaxa* **4590** (1): 196–199. <https://doi.org/10.11646/zootaxa.4590.1.11>
- CONTARINI, N. (1843): Cataloghi degli uccelli e degli insetti delle provincie di Padova e Venezia, 42 pp.; Bassano (Tipografia Baseggio).
- CORRADI, L. (2009): *Acilius (Acilius) sulcatus* (Linnaeus, 1758) – Dytiscidae. Available from: https://www.naturamediterraneo.com/forum/topic.asp?TOPIC_ID=93149 (accessed 15 May 2022)
- CORSETTI, L., ANGELINI, C., COPIZ, R., MATTOCCIA, M. & NARDI, G. (eds.) (2015): Biodiversità dei Monti Lepini. Atti del convegno, Carpineto Romano (RM), 17 maggio 2014. *Le Scienze* (25), 352 pp.; Latina (Edizioni Belvedere).
- CORSETTI, L. & MAROZZA, L. (eds.) (2020): Atlante della biodiversità faunistica dei Monti Lepini. *Le Scienze* (33), 488 pp.; Latina (Edizioni Belvedere).
- CSABAI, Z., KÁLMÁN, Z., SZIVÁK, I. & BODA, P. (2012): Diel flight behaviour and dispersal patterns of aquatic Coleoptera and Heteroptera species with special emphasis on the importance of seasons. – *Naturwissenschaften* **99**: 751–765. <https://doi.org/10.1007/s00114-012-0957-6>
- DAVY-BOWKER, J. (2002): A mark and recapture study of water beetles (Coleoptera: Dytiscidae) in a group of semi-permanent and temporary ponds. – *Aquatic Ecology* **36** (3): 435–446. <https://doi.org/10.1023/A:1016550127986>
- DE CRISTOFORI, G. & JAN, G. (1832): Cataloghi sistematici e descrittivi degli oggetti di storia naturale esistenti nel museo di Giuseppe De Cristofori e Prof. Gioegio Jan. Contenenti il prodomo della fauna, della flora e della descrizione oritognostico-geognostica dell’Italia Superiore. Sectio III.^a Entomologia. Pars I.^a Conspectus Methodicus Insectorum. Fasc. I.^{us} Coleoptera, 3 + 111 + [2] + 16 pp.; Milan (Giovanni Pirotta)
- DETTNER, K. (1983): *Colymbetes schilcknechti* a new water beetle from Sardinia with a key to European species of the genus *Colymbetes* (Coleoptera, Dytiscidae). – *Aquatic Insects* **5** (1): 39–44. <https://doi.org/10.1080/01650428309361123>
- DETTNER, K. (2006): Adephagan water beetles of Elba Island (Tuscany) (Coleoptera Haliplidae, Dytiscidae, Noteridae, Gyrinidae). – In: NARDI, G. (ed.): Volume in memoria di Mario Enrico Franciscolo, pp. 85–122. – *Memorie della Società Entomologica Italiana* **85** (1).
- DISCONZI, F. (1865) Entomologia Vicentina; ossia catalogo sistematico degli insetti della provincia di Vicenza con osservazioni e descrizioni di moltissime specie degli insetti utili e dei nocivi particolarmente all’agricoltura colla giunta di un metodo pratico sulla caccia degli insetti e sul modo di apparecchiarli per le collezioni. Con disegni litografici, [2] + 316 pp. + 18 pls.; Padova (Tipografia G. B. Randi).
- DRAULANS, D., PERREMANS, K., VAN VESSEM, J. & POLLET, M. (1987): Analysis of pellets of the grey heron, *Ardea cinerea*, from colonies in Belgium. – *Journal of Zoology* **211** (4): 695–708. <https://doi.org/10.1111/j.1469-7998.1987.tb04480.x>
- DUTTO, M., POGGI, R. & CAVAZZUTI, P. (2009): I “Coleoptera Salutiensia” di Lorenzo Ponzá (1805), con dati bio-bibliografici sull’autore e alcune note. – *Annali del Museo Civico di Storia Naturale “G. Doria”* **100**: 347–458.
- ENKHNASAN, D. & BOLDGIV, B. (2019): Biogeography of predaceous diving beetles (Coleoptera, Dytiscidae) of Mongolia. – *ZooKeys* **853**: 87–108. <https://doi.org/10.3897/zookeys.853.33908>
- FABBRI, R., PEDERZANI, F., ROCCHI, S., MASCAGNI, A., CASALI, S. & BUSIGNANI, G. (2012): Coleotterofauna acquatica della Repubblica di San Marino (Insecta Coleoptera). – *Quaderno di Studi e Notizie di Storia Naturale della Romagna* **35**: 85–100.
- FERRARO, T. (2009): Catalogo della collezione coleotterologica del Dipartimento di Agronomia Ambientale e Produzioni Vegetali. Sottordine Adephaga: tassonomia–sinonimia–distribuzione. B.Sc thesis, 146 pp.; Padua (Università degli Studi di Padova).
- FERY, H. & BRANCUCCI, M. (1997): A taxonomic revision of *Deronectes* Sharp, 1882 (Insecta: Coleoptera: Dytiscidae)

- (Part I). – Annalen des Naturhistorischen Museums in Wien. Serie B (Botanik und Zoologie) **99**: 217–302.
- FERY, H. & RIBERA, I. (2018): Phylogeny and taxonomic revision of Deronectina Galewski, 1994 (Coleoptera: Dytiscidae: Hydroporinae: Hydroporini). – Zootaxa **4474** (1): 1–104. <https://doi.org/10.11646/zootaxa.4474.1.1>
- FORNACIARI, M. (2011): *Acilius (Acilius) sulcatus* (Linnaeus, 1758) – Dytiscidae. Available from: <http://www.entomologiitaliani.net/public/forum/phpBB3/viewtopic.php?f=11&t=26977> (accessed 14 April 2020)
- FRANCISCOLO, M. E. (1979): Fauna d'Italia. XIV. Coleoptera Halplidae, Hygrobiidae, Gyrinidae, Dytiscidae, I–VI + 1–804 pp.; Bologna (Edizioni Calderini).
- GATTI, E., DAL CORTIVO, M. & SOMMACAL, M. (2016): I Coleotteri delle riserve naturali gestite dall'U.T.B. di Belluno, 920 pp.; Feltre (Edizioni DBS).
- GEMMINGER, [M.] & HAROLD, B. DE (1868): Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Tom. II. Dytiscidae, Gyrinidae, Hydrophilidae, Staphylinidae, Pselaphidae, Gnostidae, Paussidae, Scydmaenidae, Silphidae, Trichopterygidae, Scaphididae, 425–752 pp. + [6 (Index Generum ad Tom. II.)]; Munich (E. H. Gummi).
- GHILLANI, V. (1887): Elenco delle specie di Coleotteri trovate in Piemonte. Opera postuma pubblicata per cura del Membro ordinario Dott. Lorenzo Camerano. – Annali della Reale Accademia d'Agricoltura di Torino **29** (1886): 195–381.
- GHOSH, S. K. & NILSSON, A. N. (2012): Catalogue of the diving beetles of India and adjacent countries (Coleoptera: Dytiscidae). – Skorvonnopparn Supplement **3**: 1–77.
- GIACHINO, P. M. (1982): Collezione coleotterologica di Massimiliano Spinola. – Cataloghi del Museo Regionale di Scienze Naturali di Torino **3**: 1–616.
- GIORIA, A. (2014): Chapter 7. Habitats. – In: YEE, D. A. (ed.): Ecology, systematics, and the natural history of predaceous diving beetles (Coleoptera: Dytiscidae), pp. 307–362; Berlin, Heidelberg (Springer Science + Business Media B.V.). https://doi.org/10.1007/978-94-017-9109-0_7
- GROPALI, R. (1992): Sull'alimentazione di sette specie di Charadriiformes in Italia. – Rivista Italiana di Ornitologia **62** (1–2): 35–40.
- GROTTOLO, M. (2020): Indagine sulla entomofauna della riserva naturale “della Rocca, del Sasso e Parco lacuale” (Manerba del Garda, Lombardia). – Natura Bresciana **43**: 73–117.
- GUBELLINI, L. (2013): Checklist dei Dytiscidae (Coleoptera) del bacino del Metauro [first edition; July 19, 2002]. – In: Associazione Naturalistica Argonauta e Comune di Fano, Fano (PU) (eds.): La Valle del Metauro. Banca dati sugli aspetti naturali e antropici del bacino del Metauro. Available from: <https://www.lavalledelmetauro.it/contenuti/funghi-flora-fauna/scheda/9274.html> (accessed April 24 2022)
- HÁJEK, J. (2017): Dytiscidae Régimbart, 1879. – In: LÖBL, I. & LÖBL, D. (eds.): Catalogue of Palaearctic Coleoptera. Volume 1, Archostemata – Myxophaga – Adephaga: Revised and updated edition, pp. 846–914; Leiden & Boston (Brill).
- HENDRICH, L. & BALKE, M. (2020): Ein weiterer Fund des Schwimmkäfers *Methles cribratellus* (Fairmaire, 1880) von Sardinien. – Nachrichtenblatt der Bayerischen Entomologen **69** (1): 16–19.
- HENDRICH, L. & BALKE, M. (2021): New records of the diving beetle *Rhantus simulans* Régimbart, 1908 in south-western Australia (Coleoptera, Dytiscidae, Colymbetinae). – Check List **17** (2): 643–648. <https://doi.org/10.15560/17.2.643>
- HERBST, C., BAIER, B., TOLASCH, T. & STEIDLE, J. L. (2011): Demonstration of sex pheromones in the predaceous diving beetle *Rhantus suturalis* (MacLeay 1825) (Dytiscidae). – Chemoecology, **21** (1): 19–23. <https://doi.org/10.1007/s00049-010-0061-3>
- HEYROVSKY, L. (1968): Contributo alla conoscenza della coleottero-fauna del Trentino-Alto Adige (Coleoptera) II. – Bollettino dell'Associazione Romana di Entomologia **23** (2): 41–44.
- IMPARATI, E. (1897): Coleotteri del Piacentino. – Rivista Italiana di Scienze Naturali e Bollettino del Naturalista Collettore, Allevatore, Coltivatore Bollettino del Naturalista **17** (4): 49–51.
- IPLA [Istituto Piante da Legno e l'Ambiente & Regione Piemonte] (2017): Zona Speciale di Conservazione IT1150007 – “Baraggia di Piano Rosa”. Piano di Gestione, 209 pp.; Turin (Istituto Piante da Legno e l'Ambiente & Regione Piemonte).
- IUCN [International Union for Conservation of Nature] (2022): Guidelines for Using the IUCN Red List Categories and Criteria. Version 15, 116 pp.; IUCN Standards and Petitions Committee. Available from: <https://www.iucnredlist.org/resources/redlistguidelines> (accessed 15 May 2022)
- IVERSEN, L. L., RANNAP, R., BRIGGS, L. & JENSEN, K. S. (2017): Time-restricted flight ability influences dispersal and colonization rates in a group of freshwater beetles. – Ecology and Evolution **7**: 824–830. <https://doi.org/10.1002/ece3.2680>
- JÄCH, M. (1997): Daytime swarming of rheophilic water beetles in Austria (Coleoptera: Elmidae, Hydraenidae, Halplidae). – Latissimus **9**: 10–11.
- KEHL, S. & DETTNER, K. (2007): Flugfähigkeit der in Deutschland vorkommenden adephagen Wasserkäfer (Coleoptera, Hydradephaga). – Entomologie Heute **19**: 141–161.
- KRISKA, G., CSABAI, Z., BODA, P., MALIK, P. & HORVATH, G. (2006): Why do red and dark-coloured cars lure aquatic insects? The attraction of water insects to car paintwork explained by reflection–polarization signals. – Proceedings of the Royal Society B, Biological Sciences **273**: 1667–1671. <https://doi.org/10.1098/rspb.2006.3500>
- LAZZARINI, C. (1895): Catalogo di coleotteri friuliani. – In Alto **6** (3): 44–46.
- LENCIONI, V., FATTORI, D., NARDI, G. & LATELLA, L. (2019): Insights into spatio-temporal dynamics of invertebrate communities from two alpine pasture ponds. – Aquatic Insects **40** (1): 30–52. <https://doi.org/10.1080/01650424.2018.1523434>
- LEO, P. (2015): *Rhantus (Rhantus) suturalis* (W.S. Macleay, 1825) – Dytiscidae. Available from: <http://www.entomologiitaliani.net/public/forum/phpBB3/viewtopic.php?t=64694> (accessed 22 February 2020)
- LO PARRINO, E., FICETOLA, G. F., MANENTI, R. & FALASCHI, M. (2020): Thirty years of invasion: the distribution of the invasive crayfish *Procambarus clarkii* in Italy. – Biogeographia – The Journal of Integrative Biogeography **35**: 43–50. <https://doi.org/10.21426/B635047157>
- LÓPEZ-PÉREZ, J. J. & MILLÁN SÁNCHEZ, A. (2012): Catálogo de los Adephaga acuáticos (Coleoptera: Dytiscidae, Gyrinidae, Halplidae, Hygrobiidae y Noteridae) de la provincia de Huelva (Sudoeste de Andalucía, España). – Boletín de la Sociedad Entomológica Aragonesa (S.E.A.) **51**: 251–260.
- LUIGIONI, P. (1929) I Coleotteri d'Italia. Catalogo sinonimico–topografico–bibliografico. – Memorie della Pontificia Accademia delle Scienze, I Nuovi Lincei (2) **13**: [4 +] 1–1159 [+ 1].

- LUPI, D., JUCKER, C. & ROCCO, A. (2014): Rice fields as a hot spot of water beetles (Coleoptera Adephegata and Polyphaga). – *Redia* **97**: 95–112.
- MAJZLAN, O. (2014): Chrobáky (Coleoptera) dvoch lokalít Závod-Šišuláky a Gajary na Záhorí. – *Entomofauna Carpathica* **26** (2): 12–62.
- MAJZLAN, O. (2020): Chrobáky (Coleoptera) pod pohorím Burda (Južné Slovensko). – *Naturae Tutela* **24** (1): 47–70.
- MARCUZZI, G. & TURCHETTO LAFISCA, M. (1977): Ricerche sui Coleotteri della Puglia raccolti da G. Marcuzzi (1960–63). I. Elenco delle Specie. – *Quaderni di Ecologia Ambientale* **9**: 1–186.
- MARTENS, G. VON (1824): Reise nach Venedig. Erster Theil. Von Stuttgart über Ulm, Wien und Triest nach Venedig, VIII + 472 pp.; Ulm (Stettin'sche Buchhandlung).
- MATSUSHIMA, R. & YOKOI, T. (2020): Flight capacities of three species of diving beetles (Coleoptera: Dytiscidae) estimated in a flight mill. – *Aquatic Insects* **41** (4): 1–7. <https://doi.org/10.1080/01650424.2020.1804065>
- MAZZA, G., AQUILONI, L., BELLAVITA, M., CIANFANELLI, S., CIANFERONI, F., ORIOLI, G., PALOMBI, A., PIAZZAI, M., ROCCHI, S., TERZANI, F. & GHERARDI, F. (2012): Ricerche faunistiche negli habitat stagnatili (trocce) della Riserva Naturale Monte Rufeno (Lazio, Italia Centrale). – *Studi Trentini di Scienze Naturali* **92**: 21–32.
- MAZZA, G., TERZANI, F. & ROCCHI, S. (2008): Ricerche florofaunistiche in alcune zone umide del Parco Nazionale delle Foreste Casentinesi, Monte Falterona e Campigna (Toscana, Emilia-Romagna). – *Quaderni della Stazione Ecologica del Civico Museo di Storia Naturale di Ferrara* **18**: 37–88.
- MAZZOLDI, P. (2009): La fauna a Idradeffagi del Pian di Spagna (Coleoptera: Halipilidae, Gyrinidae, Noteridae, Dytiscidae). – *Il Naturalista Valtellinese* **19** (2008): 65–81.
- MELANDRI, S. (2013): *Acilius (Acilius) sulcatus* (Linnaeus, 1758) – Dytiscidae. Available from: https://www.naturamediterraneo.com/forum/topic.asp?TOPIC_ID=210789 (accessed 24 April 2022)
- MINELLI, S. (ed.) (2001): Risorgive e Fontanili – Acque sorgenti di pianura nell'Italia Settentrionale. Quaderni Habitat 2, 157 pp.; Udine (Ministero dell'Ambiente e Museo Friulano di Storia Naturale).
- MOLA, L. & TOLEDO, M. (2021): Contributo alla conoscenza della coleotterofauna del Monte Netto (Lombardia, provincia di Brescia). I. Coleotterofauna acquatica (Coleoptera: Halipilidae, Noteridae, Dytiscidae, Helophoridae, Hydrochidae, Georissidae, Hydrophilidae, Hydraenidae, Heteroceridae, Dryopidae, Elmidae, Scirtidae). – *Studi Trentini di Scienze Naturali* **101**: 61–72.
- MUGGLETON, J. (1966): Diurnal flight in *Acilius sulcatus* (L.) (Col., Dytiscidae). – *Proceedings and Transactions of the South London Entomology and Natural History Society* **3**: 84.
- NARDI, G. (2004): Adephegous water beetles: faunistics, ecology and conservation (Coleoptera, Gyrinidae, Halipilidae, Noteridae, Dytiscidae). – In: CERRETTI, P., HARDERSEN, S., MASON, F., NARDI, G., TISATO, M. & ZAPPAROLI, M. (eds.): *Invertebrati di una foresta della Pianura Padana, Bosco della Fontana. Secondo contributo. Conservazione Habitat Invertebrati* 3, pp. 65–115; Verona (Cierre Grafica Editore).
- NARDI, G. (2005): Dati preliminari sui Coleotteri Idradeffagi del Parco Nazionale del Circeo. – In: ZERUNIAN, S. (ed.): *Habitat, flora e fauna del Parco Nazionale del Circeo*, pp. 151–178; Sabaudia (Corpo Foresatle dello Stato, Ufficio Gestione Beni ex ASFD di Sabaudia – Parco Nazionale del Circeo).
- NARDI, G. & MALTZEFF, P. (2001): Gli Idradeffagi della Tenuta Presidenziale di Castelporziano (Coleoptera, Gyrinidae, Halipilidae, Noteridae, Hygrobiidae, Dytiscidae). *Bollettino dell'Associazione Romana di Entomologia* **56** (1–4): 175–232.
- NILSSON, A. N. (1997): On flying *Hydroporus* and the attraction of *H. incognitus* to red car roofs. – *Latissimus* **9**: 12–16.
- NILSSON, A. N. (2001): Dytiscidae (Coleoptera). *World Catalogue of Insects*, 3, 395 pp.; Stenstrup (Apollo Books).
- NILSSON, A. N. & FERY, H. (2006): *World Catalogue of Dytiscidae – corrections and additions*, 3. (Coleoptera: Dytiscidae). – *Koleopterologische Rundschau* **76**: 55–74.
- NILSSON, A. N. & HÁJEK, J. (2022a): A world catalogue of the Family Dytiscidae, or the Diving Beetles (Coleoptera, Adephegata). Version 1.1.2022, 317 pp. Available from: http://www.waterbeetles.eu/documents/W_CAT_Dytiscidae_2022.pdf (accessed 15 May 2022)
- NILSSON, A. N. & HÁJEK, J. (2022b): *Catalogue of Palearctic Dytiscidae (Coleoptera)*. Internet version 2022-01-01, 89 pp. Available from: http://www.waterbeetles.eu/documents/PAL_CAT_Dytiscidae_2022.pdf (accessed 15 May 2022)
- NILSSON, A. N. & HOLMEN, M. (1995): *Fauna Entomologica Scandinavia* 32. The aquatic Adephegata (Coleoptera) of Fennoscandia and Denmark. II. Dytiscidae, 192 pp.; Leiden, New York, Köln (E. J. Brill).
- PAGLIA, E. (1879): *Saggio di storia naturale sul territorio mantovano*, 507 pp.; Mantova (V. Guastalla Tipografo-Editore).
- PEDERZANI, F. & FABBRI, R. (2006): Il quarto cavaliere dell'Apocalisse *Procambarus clarkii* (Girard, 1852). – *Quaderno di Studi e Notizie di Storia Naturale della Romagna* **23**: 199–212.
- PEDERZANI, F. & PESARINI, F. (2008): I Coleotteri Idradeffagi e Palpicorni della Collezione Campadelli (Coleoptera Caraboidea, partim, Hydrophiloidea) (Catalogo sistematico della Collezione Campadelli. IV contributo). – *Annali del Museo Civico di Storia Naturale di Ferrara* **9/10** (2006/2007): 29–42.
- PEZZI, G. (2013): L'entomofauna della Zona di Protezione Speciale (ZPS) “Bacini ex zuccherificio di Mezzano” (Ravenna). 5° contributo. Coleotteri Adephegati (escl. Carabidae sl) e Polifagi (escl. Scarabaeiformia) (Insecta Coleoptera). – *Quaderno di Studi e Notizie di Storia Naturale della Romagna* **37**: 105–122.
- PILON, N. (2002): Coleotteri – In: FURLANETTO, D. (ed.), *Atlante della biodiversità nel Parco Ticino*. Vol. 1. Elenchi sistematici, pp. 312–364; Corbetta (Il Guado).
- PONZA, L. (1805): *Coleoptera salutiensia sive enumeratio methodica Coleopterorum, quae in Agro Salutiensi reperiuntur, locorum natalium indicatione, observationibus, novisque aliquot speciebus aucta*. – *Mémoires de l'Académie Impériale des Sciences, Littérature et Beaux-Arts de Turin. Sciences Physiques et Mathématiques* **12–13** (1804/1805): 29–94 + pls. IV–V.
- RAGAZZI, V. (1878): Contribuzione alla fauna entomologica italiana. Catalogo metodico dei Coleotteri raccolti nella provincia Modenese nell'estate degli anni 1875–76. – *Bollettino della Società Entomologica Italiana* **10** (III): 179–188.
- RAGUSA, E. (1921): Coleotteri nuovi o poco conosciuti nella Sicilia. – *Bollettino della Società Entomologica Italiana* **56** (III): 85–100.
- RAMADAN, M. M. & RAMADAN, H. M. (2021): Phenology of predaceous diving beetles (Coleoptera: Dytiscidae) in a desert oasis. – *International Journal of Zoology and Animal Biology* **4** (2): 1–14.

- RASTELLI, M. & DELLA BEFFA, G. (2007): Le cenosi dei Coleotteri Idroade-fagi del tratto planiziale del Parco del Po Cuneese (Coleoptera: Dytiscidae, Haliplidae). Ricerche sugli ambienti acquatici del Po Cuneese. Risultati dell'Interreg IIIA "Aqua". – Memorie dell'Associazione Naturalistica Piemontese **8** (154): 111–120.
- ROCCHI, S. (1982): Ditisidi raccolti nel Nepal dal Dr. Enrico Migliaccio e dal Dr. Guido Sabatinelli (Coleoptera, Dytiscidae). – Bollettino dell'Associazione Romana di Entomologia **35** (1–4) (1980): 57–60.
- ROCCHI, S. (2007): Insecta Coleoptera Hydroade-phaga. – In: RUFFO, S. & STOCH, F. (eds.), Checklist and distribution of the Italian fauna, pp. 165–166 + CD-ROM. – Memorie del Museo Civico di Storia Naturale di Verona, 2. Serie. Sezione Scienze della Vita **17** (2006).
- ROCCHI, S. (2011): Contribution to the knowledge of the Hydrophiloidea of Sardinia (Coleoptera: Hydrochidae, Hydrophilidae, Sphaeridiidae, Spercheidae). – In: NARDI, G., WHITMORE, D., BARDIANI, M., BIRTELE, D., MASON, F., SPADA, L. & CERRETTI, P. (eds.): Biodiversity of Marganai and Montimannu (Sardinia). Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati 5, pp. 287–311; Verona (Cierre Grafica Editore).
- ROCCHI, S. (2021) Insecta Coleoptera Adephaga (aquatic). – In: BOLOGNA, M. A., ZAPPAROLI, M., OLIVERIO, M., MINELLI, A., BONATO, L., CIANFERONI, F. & STOCH, F. (eds.): Checklist of the Italian Fauna. Version 1.0. Last update: 2021-05-31. Available from: <https://www.lifewatchitaly.eu/en/initiatives/checklist-fauna-italia-en/checklist/> (accessed April 2022)
- ROCCHI, S. & MASCAGNI, A. (2006): Coleotterofauna acquatica e semiacquatica del Parco Nazionale delle Foreste Casentinesi, Monte Falterona e Campigna (Insecta Coleoptera Hydrade-phaga, Hydraenidae, Hydrophiloidea, Dryopoidea). – Quaderno di Studi e Notizie di Storia naturale della Romagna **21** (2005): 1–32.
- ROCCHI, S. & POGGI, R. (2020): La collezione Nino Sanfilippo di Coleotteri Idroade-fagi della regione paleartica, conservata nel Museo Civico di Storia Naturale "Giacomo Doria" di Genova. I. Ditisidi. (Coleoptera, Dytiscidae). – Annali del Museo Civico di Storia Naturale "G. Doria" **112** (2019): 167–228.
- ROCCHI, S. & TERZANI, F. (2016): Contributo alla conoscenza della coleotterofauna acquatica dell'Appennino piacentino (Insecta Coleoptera Gyrinidae, Haliplidae, Noteridae, Dytiscidae, Helophoridae, Georissidae, Hydrochidae, Hydrophilidae, Hydraenidae, Scirtidae, Elmidae, Dryopidae, Psephenidae, Chrysomelidae). – Quaderno di Studi e Notizie di Storia Naturale della Romagna **43**: 133–162.
- ROCCHI, S., TERZANI, F. & MASCAGNI, A. (2021a): Coleotterofauna acquatica e semiacquatica del bacino idrografico del fiume Cecina (Toscana). – Bollettino della Società Entomologica Italiana **153** (1): 3–22. <https://doi.org/10.4081/bollettinosei.2021.3>
- ROCCHI, S., TERZANI, F. & MASCAGNI, A. (2021b): Coleotterofauna acquatica e semiacquatica dell'Umbria (Insecta: Coleoptera: Gyrinidae, Haliplidae, Noteridae, Hygrobiidae, Dytiscidae, Helophoridae, Georissidae, Hydrochidae, Hydrophilidae, Hydraenidae, Scirtidae, Elmidae, Dryopidae, Limnichidae, Heteroceridae, Psephenidae, Chrysomelidae, Curculionidae). – Quaderno di Studi e Notizie di Storia Naturale della Romagna **53**: 65–153.
- ROCCHI, S., TERZANI, F., MASCAGNI, A. & CIANFERONI, F. (2021c): Gli Insetti di Firenze. La coleotterofauna acquatica (Coleoptera: Hydrade-phaga, Hydrophiloidea, Hydraenidae, Scirtidae, Elmidae, Dryopidae, Limnichidae, Heteroceridae). – Quaderni del Museo Civico di Storia Naturale di Ferrara **9**: 81–102.
- ROCCHI, S., TERZANI, F., CECCOLINI, F., PIZZOCARO, L. & CIANFERONI, F. (2018): Coleotterofauna acquatica e semiacquatica del Molise (Italia meridionale) (Insecta: Coleoptera: Gyrinidae, Haliplidae, Noteridae, Hygrobiidae, Dytiscidae, Helophoridae, Georissidae, Hydrochidae, Hydrophilidae, Hydraenidae, Scirtidae, Elmidae, Dryopidae, Limnichidae, Heteroceridae, Chrysomelidae, Curculionidae). – Quaderno di Studi e Notizie di Storia Naturale della Romagna **47**: 139–196.
- ROCCHI, S., TERZANI, F., CIANFERONI, F., FORBICIONI, L., PAPI, R. & PIZZOCARO, L. (2017): Aggiornamenti alla conoscenza della coleotterofauna acquatica dell'Arcipelago Toscano (Coleoptera). – Onychium **13**: 75–91. <http://dx.doi.org/10.5281/zenodo.495599>
- ROMANO, M. (2010): *Acilius (Acilius) sulcatus* (Linnaeus, 1758) – Dytiscidae. Available from: www.entomologiitaliani.net/public/forum/phpBB3/viewtopic.php?f=159&t=6000&hilit=Acilius (accessed 22 February 2020)
- SAPUPPO, G. (2002): Contributo alla conoscenza dei Coleotteri della zona etnea. Collezione Gulli-Sapuppo. – Atti e Memorie dell'Ente Fauna Siciliana **8** (2001): 35–280.
- SELLA, E. (1864) Sopra alcuni Coleotteri che s'incontrano nel Biellese. – Atti della Società Italiana di Scienze Naturali **7**: 105–148.
- SHERBORN, C. D. (1930): Index Animalium sive index nominum quae ab A.D. MDCCCLVIII generibus et speciebus animalium imposita sunt. Societatibus eruditorum adiuvantibus a Carolo Davies Sherborn confectus. Sectio secunda a kalendis ianuariis, MDCCCL usque ad finem decembris, MDCCCL. Part XXIV. Index *serratus-squamosus* [n. 5911–6118; 208 pp.]; London (British Museum).
- SORGI, G., DI PIETRO, R. & ALICATA, P. (2011): I macroinvertebrati degli stagni dei Nebrodi: diversità e problemi di conservazione. – Biogeographia – The Journal of Integrative Biogeography **30** (1): 551–565. <https://doi.org/10.21426/B630110593>
- STAUBLE, A. (2021): I Coleotteri Idroade-fagi nell'area urbana di Verona. – In: LATELLA, L. (ed.): Storia naturale della città di Verona. – Memorie del Museo Civico di Storia Naturale di Verona, 2. Serie. Monografie Naturalistiche **6**: 104–110.
- STOCH, F. (ed.) (2005): Pozze, stagni e paludi – Le piccole acque, oasi di biodiversità. Quaderni Habitat 11, 160 pp.; Udine (Ministero dell'Ambiente e Museo Friulano di Storia Naturale).
- STOCH, F. & DOLCE, S. (1984): Alimentazione e rapporti alimentari di *Triturus alpestris alpestris*, *Triturus cristatus cristatus* e *Triturus vulgaris meridionalis*. Quaderni dell'Ente Tutela Pesca – Udine. Rivista di Limnologia **9**: 17–28.
- STOCH, F. & NASELLI-FLORES, L. (2014): Acque temporanee: biodiversità, funzioni ecosistemiche, vulnerabilità e sensibilità ai cambiamenti climatici. – Biologia Ambientale **28** (2): 87–92.
- ŠUSTEK, Z. (2002): Flight of Dytiscidae, Hydrophilidae and Staphylinidae (Coleoptera) on lihgt [sic!] in centre of a great city. – Entomofauna Carpathica **14**: 59–63.
- TERZANI, F., CIANFERONI, F., FABIANO, F., MAZZA G., MONTE C., ZINETTI, F. & BALLERINI, C. (2011): Coleotterofauna della provincia di Prato (Toscana): elenco preliminare. – Onychium **8** (2010): 22–76.
- TOLEDO, M. (2006): Considerazioni preliminari sui cicli vitali e strategie riproduttive di alcune specie di Dytiscidae nella

- pianura padana (Coleoptera). – In: NARDI, G. (ed.): Volume in memoria di Mario Enrico Franciscolo, pp. 187–208. – Memorie della Società Entomologica Italiana **85** (1).
- TOLEDO, M. & GROTTOLLO, M. (2019): Contributo alla conoscenza dei Coleotteri acquatici nei bacini idrografici della provincia di Brescia (Lombardia) (Coleoptera: Gyrinidae, Haliplidae, Noteridae, Dytiscidae, Helophoridae, Hydrochidae, Hydrophilidae, Hydraenidae, Psephenidae, Heteroceridae, Dryopidae, Elmidae). – Memorie della Società Entomologica Italiana **96**: 3–288.
<https://doi.org/10.4081/memorieSEI.2019.3>
- TOUSSAINT, E. F., SAGATA, K., SURBAKTI, S., HENDRICH, L. & BALKE, M. (2013): Australasian sky islands act as a diversity pump facilitating peripheral speciation and complex reversal from narrow endemic to widespread ecological super-tramp. – Ecology and Evolution **3** (4): 1031–1049.
<https://doi.org/10.1002/ece3.517>
- VONDEL, B. J. VAN (1997): Insecta: Coleoptera: Haliplidae. – In: SCHWOERBEL, J. & ZWICK, P. (eds.): Süßwasserfauna von Mitteleuropa. Band 20/2, 3 und 4, pp. 1–95; Stuttgart (Gustav Fischer).
- VONDEL, B. [J.] VAN (1998): Another case of water beetles landing on a red car roof. – Latissimus **10**: 29.
- VIGLIOGLIA, V. (2012): *Acilius (Acilius) sulcatus* (Linnaeus, 1758) – Dytiscidae. Available from: <http://www.entomologiitaliani.net/public/forum/phpBB3/viewtopic.php?f=159&t=37731&hilit=ACILIUS> (accessed 15 April 2020)
- VIGNA TAGLIANTI, A., AUDISIO, P. A., BELFIORE, C., BIONDI, M., BOLOGNA, M. A., CARPANETO, G. M., DE BIASE, A., FATTORINI, S., PIATTELLA, E., SINDACO, R., VENCHI, A. & ZAPPAROLI, M. (1999): A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. – Biogeographia, Lavori Società Italiana di Biogeografia (n.s.) **20**: 31–59.
<https://doi.org/10.21426/B6110172>
- VILLA, A. & VILLA, G. B. (1833): Coleoptera Europae Dupleta in Collectione Villa, quae pro mutua commutatione offerri possunt, 36 pp.; Mediolani [= Milan].
- VILLA, A. & VILLA, G. B. (1844): Catalogo dei Coleopteri della Lombardia. – In: CATTANEO, C. (ed.): Notizie naturali e civili sulla Lombardia, vol. 1, 77 pp. [reprint]. Milan (Giuseppe Berdardoni di Giovanni).
- ZIMMERMANN, A. (1920): Dytiscidae, Haliplidae, Hygrobiidae, Amphizoidae. – In: SCHENKLING, S. (ed.): Coleopterorum Catalogus. Vol. 4, pars 71, 326 pp.; Berlin (W. Junk).

Author's address:

Centro Nazionale Carabinieri Biodiversità “Bosco Fontana”, Strada Mantova 29, I-46045 Marmirolo (MN), Italy;
l_nardi@hotmail.com;  <https://orcid.org/0000-0002-5542-296X>

ZooBank registration: <http://zoobank.org/References/7189977C-BACA-4105-BC32-A6920B554394>

Manuscript received: 03.V.2022; accepted: 19.V.2022.

