

# DiplesiocerasBuckman, 1920 – an enigmatic Middle Jurassic ammonite genus revisited

Authors: Dietze, Volker, Schweigert, Günter, and Chandler, Robert B.

Source: Palaeodiversity, 15(1): 111-117

Published By: Stuttgart State Museum of Natural History

URL: https://doi.org/10.18476/pale.v15.a8

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

### Diplesioceras BUCKMAN, 1920 – an enigmatic Middle Jurassic ammonite genus revisited

VOLKER DIETZE, GÜNTER SCHWEIGERT & ROBERT B. CHANDLER

#### Abstract

New specimens of the rare ammonite genus *Diplesioceras* BUCKMAN, 1920 from the Upper Bajocian of Normandy and from the Iberian Ranges of Spain provide additional information on the shell morphology and ontogeny of this previously monotypic genus. A new species of *Diplesioceras*, *D. fischeri*, is recorded by a single specimen probably originating from the Parkinsoni Zone. The new material clearly demonstrates the presence of a vertically undulating floored keel, a character previously only known from some Strigoceratidae BUCKMAN, 1924. This observation let us to discuss whether *Diplesioceras* should be assigned to Strigoceratidae BUCKMAN, 1924. Further characters such as ribbing style and conch ontogeny, intraspecific variation, biostratigraphic occurrence and palaeogeographic distribution were analyzed. Despite of a long stratigraphical gap to supposedly ancestral taxa, a systematic placement of *Diplesioceras* in Hammatoceratidae BUCKMAN, 1887 appears to be more convenient; however, we cannot exclude an origination in Strigoceratidae either.

K e y w o r d s : Hammatoceratidae, Strigoceratidae, morphology, ontogeny, phylogeny, ammonite systematics, Lazarus taxon.

#### 1. Introduction

Diplesioceras BUCKMAN, 1920 is a very rare ammonite genus occurring in the Upper Bajocian of the Middle Jurassic. In previous literature, only a few fragmentary specimens and nuclei of Diplesioceras have been illustrated, all of them originating from the western Tethyan realm (Buckman 1920; Dietl 1974; Fernández-López 1985). The only record mentioned from outside of this distribution area is a very questionable and fragmentary specimen from Kenva (GALÁCZ 1997). Since its introduction by BUCKMAN, the higher systematic placement of Diplesioceras has been controversially discussed. SPATH (1931), ROMAN (1938) and subsequently HAAS (1942) identified some morphological resemblance of Diplesioceras at that time only known by the tiny holotype – with the Cretaceous genus Dipoloceras as purely homoeomorphic. In the course of our studies of strigoceratids, a placement of Diplesioceras within this family was suspected but could not be confirmed due to the lack of adequately preserved new material (Schweigert et al. 2007). In the most recent attempt (HOWARTH 2013), Diplesioceras was still tentatively placed in «Incertae sedis» within Hildoceratoidea. The aim of this study is to present more complete material that adds significantly to our knowledge of the shell ontogeny of Diplesioceras and sheds some light upon the possible taxonomic relationships of this enigmatic genus.

### 2. The type locality and horizon of Diplesioceras diplesium

The genus Diplesioceras BUCKMAN, 1920 is based on the monotypic D. diplesium BUCKMAN, 1920 as type species. The type series consisted only of the holotype, a nucleus with a diameter of 16 mm (Fig. 1a). After BUCKMAN (1920), it originated from the "Shell Bed" (= Astarte-bed; cf. HUDLESTON 1887) of the Inferior Oolite of Vetney Cross near Bridport (Dorset). This location is synonymous with Vitney Cross (Hudleston 1887 [profile no. 2 = Vitney Cross Quarry] or Vinney Cross (PARSONS 1974 [= outcrop no. 14]; SENIOR 1977). HUDLESTON (1887) mentioned two small quarries at this locality, the stratigraphically more complete Limekiln Quarry [= probably HUDLESTON'S profile no. 2 and outcrop no. 13 in PARSONS 1974] and the Knight's Quarry. Knight's Quarry was lithostratigraphically studied in detail by RICHARDSON (1928-1929). His "Astarte-obligua Bed", which corresponds to the "Shell Bed" of BUCKMAN (1920), has a thickness of almost 60 cm and was said to be very fossiliferous. Interestingly, RICHARDSON (1928-1929) mentioned numerous heteromorphic ammonites from this bed. HUDLESTON (1887) divided the "Astarte-bed" into a lower part with the common bivalve Astarte obliqua (LAMARCK, 1819) and an upper part with the brachiopod Terebratula (today: Sphaeroidothyris) sphaeroidalis Sowerby, 1823. At present, this bed is officially termed "Astarte Bed". In the area



Downloaded From: https://complete.bioone.org/journals/Palaeodiversity on 23 May 2025 Terms of Use: https://complete.bioone.org/terms-of-use of the type locality of *D. diplesium*, within a few square kilometres between Bridport, Shipton Gorge and Uploders, the *Astarte* Bed belongs to the upper Garantiana Zone (PARSONS 1974), at least in its lower part. The road cutting "Loders Cross" located near outcrop no. 1 in PARSONS (1974) was exposed during the construction of the motorway A 35 in the 1980s and became well-known among amateur collectors for its abundant white-shelled specimens of *Prorsisphinctes*. In that outcrop, the lower part of the *Astarte* Bed (with *Prorsisphinctes*) belonged to the Garantiana Zone (Tetragona Subzone), whereas the upper part already contained ammonites of the Parkinsoni Zone (Acris Subzone). Similar observations were made in 2010 during an excavation in the *Astarte* Bed near Uploders (close to the outcrops no. 16 and 17 of PARSONS 1974).

Based on the descriptions of the lithological section by HUDLESTON (1887) and RICHARDSON (1928–1929) which include information on the associated perisphinctids and *Spiroceras*, we conclude that the holotype of *D. diplesium* most likely originates from the upper part of the Garantiana Zone (Tetragona Subzone) and not from the Acris Subzone of the Parkinsoni Zone. The information that *Diplesioceras* was a Bathonian genus (DACQUÉ 1934) is definitely erroneous; moreover, the latter misspelled the genus as "*Diplesiceras*".

### 3. Morphological observations in published *Diplesioceras* specimens

The three-dimensionally preserved holotype of *D. diplesium* (Fig. 1a) is a phragmocone with a diameter of only 16 mm preserved with its calcitic replacement shell. The coarse rectiradiate primary ribs do not diverge and curve ventrally towards the aperture but do not touch the keel. Every  $3^{rd}$ ,  $4^{th}$  or  $5^{th}$  rib is remarkably prominent and ventrolaterally flared. The floored keel is smooth and broadly rounded.

**Fig. 1. a, c**–e–*Diplesioceras diplesium* BUCKMAN, 1920. **a**: Holotype, British Geological Survey, coll. no. 32024, Inferior Oolite, Astarte Bed, Garantiana Zone (Tetragona Subzone), Vetney Cross near Bridport, Dorset, UK. **c**: SNSB-BSPG 2014 XXXIV 69269 (ex coll. H. KEUPP), Oolithe de Bayeux Formation, ? Garantiana Zone, Évrecy, Normandy, France. **d**: Specimen in private collection of DIETER BERGER, Wiesloch, Germany (plaster cast: SMNS 70656), Oolithe de Bayeux Formation, ? Garantiana Zone, Sainte-Honorine-des-Pertes, Normandy, France. **e**: SMNS 70657 (plaster cast), Garantiana Zone, Aldea del Pinar, near Salas de los Infantes, Burgos province, Spain (repository of original specimen unknown). **b** – *Diplesioceras fischeri* n. sp., holotype, SMNS 70658, Oolithe de Bayeux Formation, ? Parkinsoni Zone, Port-en-Bessin, Normandy, France. Scale bars equal 1 cm (Fig. 1a1–a3 in double size).

The fragmentary specimen from N Spain reported by DIETL (1974: pl. 3, fig. 3) and another small fragment illustrated by FERNÁNDEZ-LÓPEZ (1985: pl. 17, fig. 7) correspond well with the tiny holotype and do not provide further morphological information except for a juvenile suture line in the first one. The suture line at this early stage is very similarly developed in a wide range of Middle Jurassic hildoceratoid ammonoids and provided no hints for the systematic placement.

# 4. New records of *Diplesioceras diplesium* from Normandy and Spain

New and more complete specimens from the Upper Bajocian of Normandy and from northern Spain add to our knowledge of the morphological characters and ontogeny of *Diplesioceras*.

The first one is a D. diplesium (Fig. 1c) from the Oolithe de Bayeux Formation of Évrecy. It is a phragmocone with a diameter of c. 45 mm. The ventrally rounded keel exhibits a very weak vertical undulation. It seems that the onset of this undulating stage appears later than in the specimen of Diplesioceras fischeri n. sp. from Porten-Bessin described below. On the last preserved whorl, another short section of the keel is preserved; however, the undulation is still less prominent than in D. fischeri n. sp. The specimen from Évrecy shows an advanced ontogenetic stage, but its body chamber is broken off and thus we do not know the adult stage. After about one third of the outer whorl the ventrolateral nodes disappear, the ribs become weaker and shallower, especially on the inner flank. In some parts of the flank the shell is missing and the suture lines are discernible. The suture shows only few elements on the flank, a very deep and prominent lateral lobe and a single umbilical lobe. Despite of the relatively small size of the ammonite this suture line is very complex ("fissilobate") already at this early ontogenetic stage.

The exact finding level of this specimen within the highly condensed Oolithe de Bayeux Formation is unknown and cannot be reconstructed by lithological means alone (see PAVIA et al. 2013); however, typical ammonites of the upper Garantiana Zone are well recorded from Évrecy in an identical rock matrix.

A further specimen of *D. diplesium* (Fig. 1d) was collected from the Oolithe de Bayeux Formation at Sainte-Honorine-des-Pertes, Normandy. It is a phragmocone of c. 31 mm diameter preserved with its calcitic replacement shell. The ribbing is relatively coarse, exclusively consisting of rectiradiate to slightly falcoid primaries. Numerous flared ribs occur at irregular distances. The flares are clearly formed by parabolic nodes. In the falcoid stage present in the last half whorl, the kinking points of the ribs bear small nodes as well. On the venter, the ribs become proverse and widen up before they abruptly end

along the unsculptured floored keel. The keel is thick and ventrally rounded; parts of it were lost, however, in the latest part of the phragmocone there is an undulation.

The section at Sainte-Honorine-des-Pertes is the historical type section of D'ORBIGNY's Bajocian stage. A detailed analysis of the Oolithe de Bayeux section has shown that there is strong condensation and re-elaboration (PAVIA 1994). The preservation of ammonites allows to assign the *D. diplesium* of Fig. 1d to PAVIA's taphorecord 5/2 which corresponds to the late Garantiana to early Parkinsoni zones. We assume from the striking morphological similarity with other specimens that this specimen most likely originates from the late Garantiana Zone as well.

From the same area in the Burgos province of Spain where DIETL (1974) and FERNÁNDEZ-LÓPEZ (1985) previously had recorded very fragmentary specimens of Diplesioceras diplesium, two more complete specimens were collected decades ago by a German amateur fossil collector. Unfortunately, the whereabouts of these specimens are unknown, because the specimens were later purchased to an anonymous person; however, there exists a nice plaster cast (Fig. 1e) of the bigger one of the two specimens. The latter is a completely septate phragmocone of 40 mm diameter lacking the body chamber. The sharp undulating keel is partially preserved. The inner whorls of the specimen are poorly preserved and roughly prepared and thus cannot be studied. Flared ribs occur irregularly, some of which even directly follow one another, without any intercalated normal ribs. The latest flared rib occurs at a diameter of c. 30 mm. The last half of the outer whorl shows prorsiradiate falcoid primaries that arise from a steeply dipping umbilical edge. In the lower third of the flank, the kink of the falcoid ribs is pronounced by weak parabolic nodes.

Further records of *Diplesioceras* from the Garantiana Zone have been mentioned in faunal lists of various localities of the Sierra de Albarracín of eastern Spain (MELÉNDEZ et al. 2002); however, none of this material has been illustrated.

### 5. Description of a new species of *Diplesioceras* from Normandy

Genus Diplesioceras BUCKMAN, 1920

Туре species: *Diplesioceras diplesium* Вискман, 1920, by monotypy.

A d d i t i o n a l t a x a i n c l u d e d : *Diplesioceras fischeri* n. sp., herein.

#### Diplesioceras fischeri n. sp. Fig. 1b

1998 *Phlycticeras aenigmaticum.* – Schweigert & Dietze, p. 5 only.

Specimen	D	Wh	Ww	Uw	R
D. diplesium, holotype	16	7.5	9	?	?
D. diplesium, SMNS 70656	31	11	14.5	9.5	28
D. diplesium, SMNS 70657	40	17.5	12.7	11.3	32
D. diplesium, SNSB-BSPG 2014 XXXIV 69269	45	20.8	13	13	32
D. fischeri, SMNS 70658	24	12.5	(7)	7	37

Table 1. Measurements [mm] of *Diplesioceras* spp. D = diameter; Wh = whorl height; Ww = whorl width; Uw = umbilical width; R = secondary ribs per whorl.

L S I D : urn:lsid:zoobank.org:pub:042AA5E1-542E-4335-91A9-2EAFE748E3DE

E t y m o l o g y : Named after the finder, WOLFGANG FISCHER (Schriesheim).

H o l o t y p e : SMNS 70658, a macroconch, illustrated on Fig. 1b.

Type locality: Port-en-Bessin, Normandy, NW France.

Type horizon: Oolithe de Bayeux Formation, upper Bajocian, probably Parkinsoni Zone (for discussion of the age see below).

Material studied: Only holotype.

M e a s u r e m e n t s : See Table 1.

D i a g n o s i s: Fine-ribbed species of *Diplesioceras* with an accelerated ontogeny.

Description: The holotype is a laterally prepared specimen preserved with its calcitic replacement shell. At the end of the preserved part, the inner mould is discernible and clearly indicates that the specimen is a fully septate phragmocone, and hence, the body chamber is completely missing. The specimen has a diameter of 24 mm and an estimated maximum whorl width of 7 mm. The floored, ventrally well-rounded keel shows a prominent vertical undulation. Its lateral margins are slightly concave. Up to a diameter of c. 15 mm, the relatively broad whorl section and ribbing style resemble those of D. diplesium, but its ribs are finer than in the latter. Every fourth rib thickens ventrolaterally and ends in a node; these flared ribs are weaker than in D. diplesium. On the last half of the outer whorl, there is a marked habitual change. The whorl section becomes higher, the flares disappear, and the simple ribs touch laterally the vertically undulating keel. The rectiradiate ribbing in the juvenile stage is replaced by a slighty falcoid ribbing. The kinking point is less marked than in D. diplesium. In the latest part of the outer whorl, a few secondary ribs are intercalated.

D is c u s s i o n : In *Diplesioceras fischeri* n. sp. the change from the juvenile stage with flared ribs to the next stage with a discoidal whorl section occurs very early in ontogeny. Since at least the entire body chamber was lost prior to burial, we cannot reconstruct the final size of this specimen, but we assume that the maximum conch size was smaller than in *D. diplesium*. Of course, more material is needed to confirm this assumption. Superficially, *Diplesioceras fischeri* n. sp. resembles the holotype of *Phlycticeras aenigmaticum* (FERNÁNDEZ-LÓPEZ, 1985), with which it was originally confused (SCHWEIGERT & DIETZE 1998). The latter differs from *D. fischeri* n. sp. in the whorl section, the absence of flared ribs, a strictly falcoid ribbing style throughout ontogeny, and a sharpened keel. At Port-en-Bessin, most parts of the ferruginous Oolithe de Bayeux Formation belong to the Upper Bajocian Parkinsoni Zone. Only at its basalmost part, ammonites of the Niortense Zone sporadically occur (pers. comm. G. PAVIA 2022). Hence, this *Diplesioceras* specimen must originate either from the Niortense Zone or, much more likely, from the Parkinsoni Zone. Since the stage with flared ribs is significantly shorter than in all *Diplesioceras diplesium* specimens of the Garantiana Zone, we suspect that the unique specimen represents a younger chronospecies of the Parkinsoni Zone.

## 6. Comments on possible sexual dimorphism in *Diplesioceras*

FERNÁNDEZ-LÓPEZ (1985) considered Diplesioceras being the microconchiate counterpart of his newly introduced oxyconic genus Melendezia FERNÁNDEZ-LÓPEZ, 1985. Subsequently, however, Melendezia was interpreted as a younger subjective synonym of Phlycticeras HYATT, 1900 (Schweigert & Dietze 1998; Schweigert et al. 2000). Indeed, unequivocal Phlycticeras from the Garantiana Zone and slightly younger chronospecies of this lineage differ significantly from Diplesioceras in a strictly oxyconic whorl section with a sharp and narrow keel, complete absence of tubercles and, if the shell is well preserved, in the presence of a spiral strigation. Among the ammonites studied and illustrated by FERNÁNDEZ-LÓPEZ (1985), a relatively evolute specimen included by him in Melendezia (FERNÁNDEZ-LÓPEZ 1985, pl. 17, fig. 10) might well represent a microconchiate Diplesioceras, since it shares both the whorl section and subadult ribbing style of our new material presented here. The preserved body chamber with a length of about one whorl suggests it was nearly adult, although the aperture is not preserved. The corresponding microconchs of *Phlycticeras* spp. are represented by the small-sized Oecoptychius NEUMAYR, 1878 (Schweigert & Dietze 1998, 1999), which differs from this putative Diplesioceras microconch both in its ornamentation and eccentric coiling, a much smaller size and, most important, by the complete absence of a keel.

FERNÁNDEZ-LÓPEZ (1997, fig. 4B) illustrated a tiny globose ammonite from the Parkinsoni Zone of Ricla

(Zaragoza, Spain) as "*Diplesioceras* sp. nov." and interpreted the latter as the corresponding microconch of *Melendezia enigmatica* [recte: *aenigmatica*] FERNÁNDEZ-LÓPEZ, 1985. In our view, a generic assignment of this very strongly corroded internal mould is impossible due to its poor state of preservation.

The small and densely ribbed specimen illustrated by FERNÁNDEZ-LÓPEZ (1985, pl. 17, fig. 8) as "Diplesioceras sp. nov. 1" is not a microconchiate Diplesioceras either, but more likely a representative of Toxamblyites with a relatively broad section but lacking a keel. A fragmentary specimen which is well comparable in lateral view to the latter has been recorded from the Bajocian of Kenya (GALÁCZ 1997). The fine and truly biplicate ribbing of these specimens contradicts a determination as Diplesioceras.

In respect of their rather involute coiling and despite of their relatively small sizes, all herein newly recorded specimens of *Diplesioceras* are considered as juvenile macroconchs.

# 7. Discussion of the systematic relationships of *Diplesioceras*

The systematic placement of Diplesioceras was controversially discussed from the first mention until today. Originally, BUCKMAN (1920: 25) had placed it in Oppeliidae DOUVILLÉ, 1890. Subsequently, SPATH (1931) and later ARKELL (in ARKELL et al. 1957) tentatively included it in Sonniniidae BUCKMAN, 1892. The latter saw some morphological similarities with Poecilomorphus BUCKMAN, 1889, Sonninia BAYLE, 1878 or Zurcheria DOUVILLÉ, 1885. ARKELL's opinion, however, was strictly rejected by DONOVAN et al. (1981). These authors found little similarities with sonniniids and pointed to the relatively long stratigraphical gap between the latest sonniniids and the sudden appearance of Diplesioceras. Consequently, they kept the higher systematic placement as open. FERNÁNDEZ-LÓPEZ (1985) added Diplesioceras after the representatives of the subfamily Bradfordiinae CALLOMON in DONOVAN et al., 1981, thus suggesting a placement within Haploceratidae ZITTEL, 1884. More recently, HOWARTH (2013) concurred with CALLOMON et al.'s opinion and placed it again in "incertae sedis" within Hildoceratoidea. This was the state of the art until our new study.

The herein documented specimens add significantly to the knowledge of the morphological characters of *Diplesioceras*. The shell surface is absolutely smooth, lacking any remains of spiral strigation. Coarse, rectiradiate primaries occur from the earliest visible stage onwards. In the juvenile stage, parabolic nodes occur in a ventrolateral position and form flared ribs. In a slightly later ontogenetic stage, the conch becomes discoidal and below midflank some ribs are thickened. The floored keel is broadly rounded and undulates vertically. Last preserved (?adult) stage shows a weakening of ribbing. These additional characters must be safely taken into account when discussing the systematic affiliation of *Diplesioceras*.

An assignment of Diplesioceras to Oppeliidae as suggested by BUCKMAN (1920) and later repeated by DACQUÉ (1934) can be easily excluded because of the floored keel, which is not developed in typical oppeliid genera (Oppelia WAAGEN, 1869, Oxycerites Rollier, 1909, Oecotraustes WAAGEN, 1869, etc.). This important character also excludes the subfamily Bradfordiinae. FERNÁNDEZ-LÓPEZ (1985) suspected an origination of Diplesioceras from latest representatives of Poecilomorphus BUCKMAN, 1889, which itself is said to have evolved from Toxamblyites BUCKMAN, 1924 (STURANI 1971; DONOVAN et al. 1981). Striking differences in the ribbing style and the keel in Diplesioceras and Poecilomorphus, however, do not point to a closer phyletic relationship but only to a superficial homoeomorphy (in the juvenile stage) between these two genera.

In first view, the vertically undulating keel of Diplesioceras points to an assignment to Strigoceratidae, where an undulating keel occurred independently within two phyletic lineages. In Phlycticeratinae (Phlycticeras, Oxydiscites), the keel is very sharp and finely serrate, unlike that in Diplesioceras. In Strigoceratinae, the keel is smooth and rounded, closer to that of Diplesioceras. Within Strigoceratinae, only the genus Granulochetoceras GEYER, 1960 exhibits a floored undulating keel (SCHWEIGERT et al. 2007). Its earliest known representative, G. oppeliisculptum Schweigert et al., 2007, from the Garantiana Zone of Spain, is still closely reminiscent to a strigoceratid ancestor. In Diplesioceras diplesium occurring at the same time and even in the same area as G. oppeliisculptum, there is no spiral striation discernible despite of the perfect preservation of the shell. Additionally, Strigoceratinae (as well as Phlycticeratinae) never show a coarse ribbing already in the nucleus as in Diplesioceras. Thickened ribs at midflank occur in some Phlycticeratinae, but never in Strigoceratinae, and parabolic nodes as on the flared ribs of Diplesioceras are unknown yet in both subfamilies. If we have correctly interpreted the specimen illustrated by FERNÁNDEZ-LÓPEZ (1985, pl. 17, fig. 10) as a microconchiate Diplesioceras, this morphology is untypical of a member of Strigoceratinae, where the microconchs are much smaller and always show an eccentric coiling. Hence, this analysis of characters does not confirm an affiliation to Strigoceratidae (Strigoceratinae and Phlycticeratinae) except for the presence of an undulating keel.

Alternatively, a placement of *Diplesioceras* within Sonniniidae BUCKMAN, 1892 must be considered. Both the blunt ribbing and the undulating keel are unknown in sonniniids, whereas the presence of spines does not contradict such a placement. However, there is a significant stratigraphical gap between the last occurrence of sonniniids in the Humphriesianum Zone and the appearance of *Diplesioceras* in the Garantiana Zone.

Finally, we may suspect a placement of Diplesioceras in Hammatoceratidae. This is corroborated by the broad section of the nucleus, the rounded floored keel and the blunt ribbing style. Most important, the development of parabolic nodes on the ribs is a rather common character of hammatoceratids. But if Diplesioceras was a hammatoceratid, what are the ancestral forms? The voungest known hammatoceratids, sometimes included in Sonniniidae, are represented by Fissilobiceras BUCKMAN, 1919, a genus which ranges up to the Humphriesianum Zone (DIETZE et al. 2011). Morphologically, the poorly sculptured Fissilobiceras is no adequate candidate for an ancestral form leading to Diplesioceras. In contrast, hammatoceratids of the Eudmetoceras / Euaptetoceras group are morphologically much closer, especially in respect of their blunt ribbing style developed very early in ontogeny and their gradual ontogenetic change towards a discoidal whorl section (see e.g. LINARES 2002, KOVÁCS 2009). In any case, the undulation of the keel must be interpreted as a derived character. A placement of Diplesioceras within Hammatoceratidae, however, is questioned by the fact that both Eudmetoceras BUCKMAN, 1920 and Euaptetoceras BUCKMAN, 1922 are unknown from strata younger than the early Bajocian Discites Zone. There is a wide stratigraphical gap between the last occurrence of unequivocal hammatoceratids and the sudden appearance of Diplesioceras in the late Bajocian. Following its interpretation as a hammatoceratid, Diplesioceras must be considered as a "Lazarus taxon", the ancestors of which had survived in a still unknown area for millions of years. Similar arguments may be put forward for an alternative placement in Sonniniidae. As a conclusion, the systematic placement of Diplesioceras remains enigmatic.

#### Acknowledgements

We thank WOLFGANG FISCHER (Schriesheim) for donating the herein described unique specimen of *D. fischeri* illustrated in Fig. 1b as well as REINHARD SCHMODE (Rheurdt), HELMUT KEUPP (Berlin) and ALEXANDER NÜTZEL (Munich) for allowing to study the ammonite of Fig. 1c. DIETER BERGER (Wiesloch) provided access to a further *Diplesioceras* specimen from his collection and allowed the preparation of a plaster cast. ROGER FURZE (Überlingen) kindly donated the plaster cast illustrated in Fig. 1e. Finally, we acknowledge the constructive reviews by JOSÉ SANDOVAL (Granada) and ANDRÁS GALÁCZ (Budapest).

#### References

ARKELL, W. J., KUMMEL, B. & WRIGHT, C. W. (1957): Mesozoic Ammonoidea. In: MOORE, R. C. (ed.): Treatise on Invertebrate Paleontology, part L, L80–490; Lawrence (University of Kansas).

- BAYLE, E. (1878): Fossiles principaux des terrains jurassiques. Mémoire pour l'Explication de la Carte géologique de France, 4, Atlas.
- BUCKMAN, S. S. (1887–1907): A monograph of the ammonites of the "Inferior Oolite series". – Monographs of the Palaeontographical Society: cclxii + 456 pp.
- BUCKMAN, S. S. (1909–1930): Yorkshire Type Ammonites/ Type Ammonites, Vols. 1–7, 790 pls.; London (Wheldon & Wesley/Wesley).
- DACQUÉ, E. (1934): Wirbellose des Jura. In: GÜRICH, G. (ed.): Leitfossilien. 7(2): 273–582; Berlin (Borntraeger).
- DONOVAN, D. T., CALLOMON J. H. & HOWARTH, M. K. (1981): Classification of the Jurassic Ammonitina. In: HOUSE, M. R. & SENIOR, J. R. (eds.): The Ammonoidea: The Evolution, Classification, Mode of Life, and Geological Usefulness of a Major Fossil Group. – The Systematics Association, Special Volumes, 18: 101–155; London & New York (Academic Press).
- DIETL, G. (1974): Zur Stratigraphie und Ammonitenfauna des Dogger, insbesondere des Oberbajocium der westlichen Keltiberischen Ketten (Spanien). – Stuttgarter Beiträge zur Naturkunde, Serie B, 14: 1–21.
- DIETZE, V., SCHWEIGERT, G., FIDDER, U. & WANNENMACHER, N. (2011): The Giganteuston Member of Öschingen (Humphriesianum Zone, Lower Bajocian, Swabian Alb), with comments on the genera *Dorsetensia* Buckman, 1892 and *Nannina* Buckman, 1927. – Zitteliana, Reihe A, **51**: 209– 236.
- DOUVILLÉ, H. (1885): Sur quelques fossiles de la zone à Ammonites sowerbyi des environs de Toulon. – Bulletin de la Société géologique de France, (3), 13: 12–44.
- DOUVILLÉ, H. (1890): Sur la classification des Cératites de la Craie. Bulletin de la Société géologique de France, (3), 18: 275–292.
- FERNÁNDEZ-LÓPEZ, S. (1985): El Bajociense en la Cordillera Ibérica. 848 + 23 pp.; Madrid (Facultad de Ciencias Geologicas, Departamento de Paleontología, Universidad Complutense).
- FERNÁNDEZ-LÓPEZ, S. (1997): Factores determinantes de la conservación de los ammonites jurásicos. In: GÁMEZ VINTANED, J.A. & LIÑÁN, E. (eds.): Vida y ambientes del Jurásico: 137– 157; Zaragoza (Institución "Fernando El Católico").
- GALÁCZ, A. (1997): Bajocian (Middle Jurassic) ammonites of stratigraphical and palaeobiogeographical importance from Mombassa, Kenya, East Africa. – Geodiversitas, **39**(4): 717– 727.
- GEYER, O. F. (1960): Über Oxydiscites Dacqué. Ein Beitrag zur Kenntnis der Ochetoceratinae (Cephal. jurass.). – Neues Jahrbuch für Geologie und Paläontologie, Monatshefte, 1960: 417–425.
- HAAS, O. (1942): Recurrence of morphological types and evolutionary cycles in Mesozoic ammonites. – Journal of Paleontology, 16: 643–650.
- HOWARTH, M. K. (2013): Treatise Part L, Revised, Volume 3B, Chapter 4: Psiloceratoidea, Eodoceratoidea [sic], Hildoceratoidea. – Treatise Online, 57: 1–139.
- HUDLESTON, W. H. (1887): A monograph of the British Jurassic Gasteropoda. Part I. A monograph of the Inferior Oolite Gasteropoda. – Palaeontographical Society Monographs, 40: 1–56.
- HYATT, A. (1900): Cephalopoda. In: ZITTEL, K. A. VON & EASTMAN, C. R. (eds.): Textbook of Palaeontology: 502–604; London (Macmillan).

- Kovács, Z. (2009): Toarcian–Aalenian Hammatoceratinae (Ammonitina) from the Gerecse Mts (NE Transdanubian Range, Hungary). – Fragmenta Palaeontologica Hungarica, 27: 1–72.
- LAMARCK, J. B. (1819): Histoire naturelle des Animaux sans vertèbres. Classe onzième. Les Conchifères, 6: 1–258; Paris (Verdière).
- LINARES, A. (2002): Nuevos Hammatoceratinae (Ammonitina) del limite Aaleniense/Bajociense de la Zona Subbética (Cordillera Bética, Sur de España). – Revista Española de Paleontología, 17: 285–291.
- MELÉNDEZ, G., FERNÁNDEZ-LÓPEZ, S., SORIA LLOP, C., PÉREZ URRESTI, I., BELLO, J., DELVENE, G., COMAS-RENGIFO, M. J., GOY, A., CLEMENTE, E. & RODRÍGUEZ MORA, M. (2002): Los yacimientos paleontológicos del Jurásico de la Sierra de Albarracín (Rama Castellana de la Cordillera Ibérica): aspectos estratigráficos y patrimoniales. – In: MELÉNDEZ HEVIA, G. & PEÑALVER MOLLÁ, E. (coord.): El Patrimonio paleontológico de Teruel: 81–136; Teruel (Instituto de Estudios Turolenses).
- NEUMAYR, M. (1878): Über unvermittelt auftretende Cephalopoden-Typen im Jura Mitteleuropas. – Jahrbuch der kaiserlichköniglichen geologischen Reichsanstalt, 28: 37–80.
- PARSONS, C. (1974): The stratigraphy of the Stony Head Cutting. – Proceedings of the Dorset Natural History & Archaeological Society, 96: 8–13.
- PAVIA, G. (1994): Taphonomic remarks on d'Orbigny's type-Bajocian (Bayeux, West France). – Miscellanea Servizio Geologico Nazionale, 5: 93–111.
- PAVIA, G., DEFAVERI, A., MAERTEN, L., PAVIA, M. & ZUNINO, M. (2013): Ammonite taphonomy and stratigraphy of the Bajocian at Maizet, South of Caen (Calvados, NW France). – Comptes Rendus Palevol, 12: 137–148.
- RICHARDSON, L. (1928–1929): The Inferior Oolite and contiguous deposits of the Burton Bradstock – Broadwindsor districts, Dorset. – Proceedings of the Cotteswold Naturalists' Field Club, 22: 35–68 [1928], 149–185 [1929].

- ROLLIER, L. (1909): Phylogénie des principaux genres d'ammonoides de l'Oolithique (Dogger) et de l'Oxfordien. – Archives des Sciences physicales et naturelles, 28: 613–623.
- ROMAN, F. (1938): Les ammonites jurassiques et crétacées. Essai de génera. 554 pp.; Paris (Masson).
- SCHWEIGERT, G. & DIETZE, V. (1998): Revision der dimorphen Ammonitengattungen *Phlycticeras* Hyatt – *Oecoptychius* Neumayr (Strigoceratidae, Mitteljura). – Stuttgarter Beiträge zur Naturkunde, Serie B, **269**: 1–59
- SCHWEIGERT, G. & DIETZE, V. (1999): Oecoptychius des Rätsels Lösung. – Fossilien, 1999(1): 51–58.
- SCHWEIGERT, G., DIETZE, V. & DIETL, G. (2003): Neue Nachweise von *Phlycticeras* und *Oecoptychius* (Ammonitina: Strigoceratidae: Phlycticeratinae). – Stuttgarter Beiträge zur Naturkunde, Serie B, **335**: 1–21.
- SCHWEIGERT, G., DIETZE, V., CHANDLER, R. B. & MITTA, V. V. (2007): Revision of the Middle Jurassic dimorphic ammonite genera *Strigoceras/Cadomoceras* (Strigoceratidae) and related forms. – Stuttgarter Beiträge zur Naturkunde, Serie B, **373**: 1–74.
- SENIOR, J. R. (1977): The Jurassic ammonite *Bredyia* Buckman. Palaeontology, **20**: 675–693.
- SOWERBY, J. (1823): The Mineral Conchology of Great Britain, 4: 1–104; London (Arding).
- SPATH, L. F. (1923–1943): A monograph of the Ammonoidea of the Gault. – Palaeontographical Society Monographs: 1–787; London (The Palaeontographical Society).
- STURANI, C. (1971): Ammonites and stratigraphy of the "Posidonia alpina" beds of the Venetian Alps (Middle Jurassic, mainly Bajocian). – Memorie degli Istituti di Geologia e Mineralogia dell'Università di Padova, 28: 1–190.
- WAAGEN, W. (1869): Die Formenreihe des Ammonites subradiatus. Versuch einer paläontologischen Monographie. – Beneckes Geognostisch Paläontologische Beiträge, 2 (2): 181–256.
- ZITTEL, K. A. v. (1884): Handbuch der Palaeontologie, Abteilung 1 (2): 893 pp.; Leipzig (Oldenbourg).

Addresses of the authors:

VOLKER DIETZE, Meraner Str. 41, 86720 Nördlingen, Germany; e-mail: dietze.v@t-online.de

GÜNTER SCHWEIGERT, Staatliches Museum für Naturkunde, Rosenstein 1, 70191 Stuttgart, Germany; e-mail: guenter.schweigert@smns-bw.de

ROBERT B. CHANDLER, Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, United Kingdom; e-mail: baj1894@gmail.com

Manuscript received: 20 September 2022, revised version accepted: 11 October 2022.