



## **Errata for Yumi et al., Egg Transport in the Coelom of the Newt *Cynops pyrrhogaster*. I. The Ovulated Egg is Transported to the Ostium by the Ciliary Movement of Coelomic Epithelial Cells**

Source: Zoological Science, 19(9) : 1075

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/0289-0003-19.9.1075>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ERRATA

Volume 19, No. 8 (2002), In the article “Egg Transport in the Coelom of the Newt *Cynops pyrrhogaster*. I. The Ovulated Egg is Transported to the Ostium by the Ciliary Movement of Coelomic Epithelial Cells.” by Yumi Yamahama and Kazuo Onitake, page 899–906. for the reader’s convenience, the corrections appear as follows;

page 906, line 39–40

... compared with the VEs of coelomic eggs and uterine eggs.

(correct) ... compared with the VEs of coelomic eggs and uterine eggs.