



## **Time Scale for Cyclostome Evolution Inferred with a Phylogenetic Diagnosis of Hagfish and Lamprey cDNA Sequences**

Authors: Kuraku, Shigehiro, and Kuratani, Shigeru

Source: Zoological Science, 23(12) : 1053-1064

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.23.1053>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Time Scale for Cyclostome Evolution Inferred with a Phylogenetic Diagnosis of Hagfish and Lamprey cDNA Sequences

Shigehiro Kuraku\* and Shigeru Kuratani

Laboratory for Evolutionary Morphology, RIKEN Center for Developmental Biology,  
2-2-3 Minatojima-minami, Chuo-ku, Kobe 650-0047, Japan

The Cyclostomata consists of the two orders Myxiniiformes (hagfishes) and Petromyzoniiformes (lampreys), and its monophyly has been unequivocally supported by recent molecular phylogenetic studies. Under this updated vertebrate phylogeny, we performed *in silico* evolutionary analyses using currently available cDNA sequences of cyclostomes. We first calculated the GC-content at four-fold degenerate sites (GC<sub>4</sub>), which revealed that an extremely high GC-content is shared by all the lamprey species we surveyed, whereas no striking pattern in GC-content was observed in any of the hagfish species surveyed. We then estimated the timing of diversification in cyclostome evolution using nucleotide and amino acid sequences. We obtained divergence times of 470–390 million years ago (Mya) in the Ordovician–Silurian–Devonian Periods for the interordinal split between Myxiniiformes and Petromyzoniiformes; 90–60 Mya in the Cretaceous–Tertiary Periods for the split between the two hagfish subfamilies, Myxiniinae and Eptatretinae; 280–220 Mya in the Permian–Triassic Periods for the split between the two lamprey subfamilies, Geotriinae and Petromyzoninae; and 30–10 Mya in the Tertiary Period for the split between the two lamprey genera, *Petromyzon* and *Lethenteron*. This evolutionary configuration indicates that Myxiniiformes and Petromyzoniiformes diverged shortly after the common ancestor of cyclostomes split from the future gnathostome lineage. Our results also suggest that intra-subfamilial diversification in hagfish and lamprey lineages (especially those distributed in the northern hemisphere) occurred in the Cretaceous or Tertiary Periods.

**Key words:** cyclostome, hagfish, lamprey, GC-content, synonymous substitution, molecular clock

## INTRODUCTION

Extant agnathans, the cyclostomes, comprise the hagfishes (Hyperotreti; order Myxiniiformes) and lampreys (Hyperoartia; order Petromyzoniiformes [often misspelled as “Petromyzontiformes”]) (Hardisty and Potter, 1971; Forey and Janvier, 1993; Jørgensen, 1998; see also Ota and Kuratani, 2006) (Fig. 1). After a long-standing controversy on the phylogenetic positions of hagfishes and lampreys, the monophyly of cyclostomes has been unequivocally supported by molecular phylogenetics using a triad of molecules frequently used for reconstruction of species phylogeny, namely, mitochondrial genes (mtDNA), nuclear ribosomal RNA genes (rDNA), and nuclear protein-coding genes (nuDNA) (Fig. 1; Stock and Whitt, 1992; Mallatt and Sullivan, 1998; Kuraku *et al.*, 1999; Delarbre *et al.*, 2002; Furlong and Holland, 2002; Takezaki *et al.*, 2003; Blair and Hedges, 2005; Delsuc *et al.*, 2006). Therefore, our interest in cyclostome evolution has shifted to the topological and temporal aspects of divergence patterns within this animal group.

The order Myxiniiformes is thought to be monophyletic, based on molecular phylogenetic studies using mitochondrial 16S rDNA (Kuo *et al.*, 2003; Chen *et al.*, 2005). This order is divided into the two subfamilies Myxiniinae and Eptatretinae, based on morphological features (Fig. 1; Fernholm, 1998). The subfamily Myxiniinae consists of four genera, *Myxine* and three genera intrinsic to the southern hemisphere (*Neomyxine*, *Nemamyxine*, and *Notomyxine*). The other subfamily, Eptatretinae, consists of three genera, *Eptatretus*, *Paramyxine*, and *Rubicundus*. In contrast, the order Petromyzoniiformes is composed of three subfamilies, Mordaciinae, Geotriinae, and Petromyzoninae, in accordance with morphology such as dentition (Hubbs and Potter, 1971; Gill *et al.*, 2003). The subfamilies Mordaciinae and Geotriinae are endemic to the southern hemisphere, and each comprises a single genus, *Mordacia* and *Geotria*, respectively (Potter and Strahan, 1968). The subfamily Petromyzoninae is composed of at least six genera (Fig. 1; Hardisty and Potter, 1971; Hubbs and Potter, 1971; Potter and Gill, 2003). However, there are few detailed reports of molecular approaches to estimate divergence times in the cyclostome lineage. To address questions regarding the temporal pattern of cyclostome evolution, the accumulating nucleotide and amino acid sequences of hagfishes and lampreys will provide novel information.

\* Corresponding author. Phone: +81-78-306-3064;  
Fax : +81-78-306-3370;  
E-mail: venezia@cdb.riken.jp

doi:10.2108/zsj.23.1053

In this study, we analyzed the GC-content in cDNA sequences of hagfishes and lampreys and calculated the divergence times of several branching points in cyclostome phylogeny using nucleotide and amino acid sequences, based on an updated version of vertebrate phylogeny representing the monophyly of cyclostomes.

## MATERIALS AND METHODS


### GC<sub>4</sub> calculation

Currently available annotated nucleotide sequences (as of February 15, 2006) were retrieved for each cyclostome species from NCBI Entrez Nucleotide (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=Nucleotide>). Redundant sequences were manually removed. To avoid biased gene selection, in which a large proportion of the sequence population for one species is occupied by members of a limited number of gene families, cDNAs derived from variable leukocyte receptor genes in the sea lamprey, *Petromyzon marinus* (accession numbers CK988414-CK988652 in NCBI dbEST; Pancer *et al.*, 2004a), and the inshore hagfish, *Eptatretus*

*burgeri* (accession numbers AY964719-AY965612; Pancer *et al.*, 2005), were excluded from our sequence collection. The nucleotide sequences were used to calculate the GC-content at four-fold degenerate sites (GC<sub>4</sub>) with the Perl script, in which an open reading frame is automatically detected with an alignment generated by BLASTX (Altschul *et al.*, 1997). Sequences of mtDNA and nuclear rRNA genes were excluded from this GC calculation.

### Molecular phylogenetic tree inference

Sequences that showed significant similarity to a query in a BLASTP search (Altschul *et al.*, 1997) were retrieved from databases: GenBank (release 151), NCBI-refseq (release 06-02-16), SWISSPROT (release 49.0), and PIR (release 80.0). An optimal multiple alignment of these amino acid sequences was constructed using the alignment editor XCED implemented in the MAFFT program (Katoh *et al.*, 2002) in combination with manual inspection. Molecular phylogenetic trees were inferred by the neighbor-joining method (Saitou and Nei, 1987) with XCED and the maximum-likelihood method (Felsenstein, 1981; Kishino *et al.*, 1990) with PAML 3.1 (Yang, 1997), using amino acid sites at which the alignment was



**Fig. 1.** Taxonomy and molecular phylogeny of cyclostomes. Taxonomic information was based on NCBI Entrez Taxonomy, which is mainly consistent with the classification by Renaud (1997) and Fernholm (1998). Taxonomic ranks are indicated in the top row. References that suggest particular phylogenetic relationships are listed near the corresponding nodes in gray shades, with the type of molecule (nuDNA, rDNA, or mtDNA) employed in their analyses. Note that branch lengths do not correspond to evolutionary times. The numbers of nucleotide and amino acid sequences, respectively, found in NCBI Entrez Nucleotide and NCBI Entrez Protein (as of February 15, 2006) are shown in parentheses beside genus names. The hagfish genera *Eptatretus* and *Paramyxine* are not monophyletic to each other according to the molecular phylogenetic tree of the 16S rRNA gene (Kuo *et al.*, 2003; Chen *et al.*, 2005). The lamprey genera *Entosphenus*, *Lampetra*, and *Lethenteron* are not monophyletic to one another according to the molecular phylogenetic tree of the cytochrome *b* and NADH dehydrogenase 3 genes (Docker *et al.*, 1999; see DISCUSSION for details).

unambiguous with no gaps, with among-site rate heterogeneity taken into account (Yang, 1994).

#### Estimation of number of synonymous and nonsynonymous substitutions per site

Nucleotide sequences were prepared as described above in the procedures for GC<sub>4</sub> calculation. By inferring molecular phylogenetic trees, we selected genes with a homologue present as a single orthologue in a pair of species in question. Nucleotide sequences of the selected genes were aligned based on an alignment generated for their deduced amino acid sequences. The number of synonymous and nonsynonymous substitutions per site ( $K_s$  and  $K_a$ , respectively) was calculated with the codon-based maximum-likelihood method (Goldman and Yang, 1994). Computation was accomplished using PAML 3.1 (Yang, 1997).

#### Amino acid sequence-based divergence time estimation

Estimation of divergence times was processed without assuming a global molecular clock, using the MULTIDIVTIME program in which Markov-chain Monte-Carlo (MCMC) procedures for Bayesian analysis are implemented (Kishino *et al.*, 2001). The upper and lower limits of divergence times outside the cyclostomes were pre-set by referring to a set of fossil records (Young, 1962) used by Dickerson (1971), or by referring to molecular dating (Kumar and Hedges, 1998; Blair and Hedges, 2005). To estimate divergence times using mitochondrial genes, we used a modified version of the MULTIDIVTIME program as instructed on the developers' web page (<http://statgen.ncsu.edu/thorne/multidivtime.html>). Results were confirmed with the program R8S, which enables penalized rate smoothing (data not shown; Sanderson, 2002, 2003).

## RESULTS


#### GC-content in cyclostome cDNAs

GC<sub>4</sub> was calculated for cDNA sequences derived from nuclear protein-coding genes for each cyclostome species. The GC<sub>4</sub> of annotated cDNAs exhibited a unimodal distribution, with peaks at 40–60% in hagfish species (Figs. 2A–C) and at 70–90% in lamprey species (Figs. 2D–F). Non-annotated abundant cDNAs of *Eptatretus burgeri* (Suzuki *et al.*, 2004) and *Petromyzon marinus* (Pancer *et al.*, 2004b) showed similar GC<sub>4</sub> distributions to those of annotated cDNAs for hagfish and lamprey species, respectively (Fig. 2G). The results of the GC<sub>4</sub> calculation for genera or species with a small number of available cDNAs were as follows: *Paramyxine*, 41–61% (n=5); *Ichthyomyzon*, 73–87% (n=7); *Entosphenus*, 75–83% (n=2); *Mordacia mordax*, 72–82% (n=3); *Geotria australis*, 73–90% (n=9).

#### Estimated number of synonymous substitutions

We selected genes that existed as single orthologues in a pair of species in question with more than 600 bp of aligned nucleotide stretches. Estimation of  $K_s$  was processed using the maximum-likelihood method (Goldman and Yang, 1994). The average  $K_s$  between the two hagfish genera, *Myxine* and *Eptatretus*, was 0.24 (standard deviation (SD), 0.11; n=11; Table 1). In contrast, the average  $K_s$  for *Petromyzon*–*Lethenteron* and *Geotria*–*Lethenteron* pairs was 0.15 (SD, 0.09; n=19; Table 2A) and 1.03 (SD, 0.39; n=2; Table 2B), respectively. Sequence comparison between a hagfish species and a lamprey species always yielded an apparently saturated  $K_s$  (>3; data not shown).

We also estimated the  $K_s$  of mitochondrial protein-coding genes in the *Myxine*–*Eptatretus* and *Petromyzon*–*Lam-*



**Fig. 2.** Distribution of GC<sub>4</sub> for hagfish and lamprey cDNAs. Histograms representing the fraction of cDNAs for each GC<sub>4</sub> value are shown for three hagfish species, (A) *Myxine glutinosa*, (B) *Eptatretus stoutii*, and (C) *Eptatretus burgeri*, and three lamprey species, (D) *Petromyzon marinus*, (E) *Lethenteron japonicum*, and (F) *Lampetra fluviatilis*. (G) The distributions of GC<sub>4</sub> in assembled expressed sequence tags (ESTs) of *Eptatretus burgeri* and *Petromyzon marinus*, also shown as histograms. ESTs were retrieved from NCBI dbEST for *Eptatretus burgeri* (accession numbers, BJ644497–BJ668380; Suzuki *et al.*, 2004) and *Petromyzon marinus* (CO542795–CO553159; Pancer *et al.*, 2004b). These ESTs were automatically assembled with Phrap (<http://www.phrap.com>), so that one cDNA sequence represents one gene. The assembled sequences are available upon request. In analyses using the GC-content of the third positions (GC<sub>3</sub>), we obtained similar results for all the graphs in this Fig.

**Table 1.** Estimated numbers of synonymous and non-synonymous substitutions between Myxininae and Eptatretinae.

Gene Name	Gene symbols of potential human orthologues	Species		Aligned length (nt)	$K_s$	$K_a$
		Myxininae	Eptatretinae			
blood 5-aminolevulinate synthase	ALAS1, ALAS2	Mg	Eb	603	0.20	0.01
creatine kinase	CKM, CKB, CKMT1, CKMT2	Mg	Es	1137	0.30	0.05
eukaryotic translation elongation factor 1 gamma	EEF1G	Mg	Eb	630	0.14	0.05
guanine nucleotide-binding protein, beta polypeptide 2-like 1	GNB2L1	Mg	Eb	900	0.20	0.02
H <sup>+</sup> -transporting ATP synthase alpha subunit isoform 1	ATP5A1	Mg	Eb	1092	0.24	0.01
low molecular mass polypeptide	PSMB5, PSMB8	Mg	Eb	819	0.02	0.01
ribosomal protein S2	RPS2	Mg	Eb	603	0.31	0.00
ribosomal protein S3	RPS3	Mg	Eb	633	0.23	0.00
ribosomal protein S4	RPS4X, RPS4Y1	Mg	Eb	714	0.49	0.03
ribosomal protein L5	RPL5	Mg	Eb	888	0.23	0.04
ribosomal protein L7a	RPL7A	Mg	Eb	723	0.26	0.05

Species names are indicated as abbreviations: Mg, *Myxine glutinosa*; Eb, *Eptatretus burgeri*; Es, *Eptatretus stoutii*.

**Table 2.** Estimated numbers of synonymous and non-synonymous substitutions between lamprey species.A, *Petromyzon-Lethenteron*

Gene Name	Gene symbols of potential human orthologues	Species		Aligned length (nt)	$K_s$	$K_a$
		<i>Petromyzon</i>	<i>Lethenteron</i>			
bone morphogenetic protein (BMP) 2/4a	BMP2, BMP4	Pm	Lj	831	0.19	0.01
cytoplasmic actin (LjCA1)	ACTB, ACTG1	Pm	Lj	1128	0.22	0.00
Dlx1/6 (DlxD)	DLX1, DLX6	Pm	Lj	804	0.14	0.00
enolase-2	ENO1, ENO2, ENO3	Pm	Lr	1185	0.12	0.01
eukaryotic elongation factor-1 alpha	EEF1A1, EEF1A2	Pm	Lj	1389	0.07	0.00
eukaryotic translation elongation factor 1 gamma	EEF1G	Pm	Lj	1068	0.10	0.01
fructose-bisphosphate aldolase, muscle type (EJM8)	ALDOA, ALDOB, ALDOC	Pm	Lj	1065	0.22	0.02
guanine nucleotide binding protein, beta polypeptide 2-like 1	GNB2L1	Pm	Lj	951	0.12	0.00
HMG 1/2/3	HMGB1, HMGB2, HMGB3	Pm	Lf	624	0.10	0.00
Ikaros-like transcription factor IKLF1	ZNFN1A1, ZNFN1A2, ZNFN1A3, ZNFN1A4, ZNFN1A5	Pm	Lf	1197	0.05	0.02
intermediate filament protein type III	PRPH, DES, VIM	Pm	Lf	1419	0.26	0.01
low molecular mass polypeptide (LMPX)	PSMB5, PSMB8	Pm	Lj	615	0.01	0.00
myosin heavy chain (LjMyHC3)	MYH9, MYH10, MYH11, MYH14	Pm	Lj	786	0.09	0.01
Pax2/5/8	PAX2, PAX5, PAX8	Pm	Lj	834	0.24	0.00
phosphoglycerate kinase (PGK)	PGK1, PGK2	Pm	Lr	828	0.22	0.00
ribosomal protein S2	RPS2	Pm	Lj	603	0.11	0.00
ribosomal protein L7a	RPL7A	Pm	Lj	672	0.36	0.01
SH3-domain GRB2-like (endophilin SH3p4)	SH3GL1, SH3GL2	Pm	Lf	627	0.10	0.00
TATA-box binding protein	TBP, TBPL1, TBPL2	Pm	Lj	867	0.09	0.00

B, *Geotria-Lethenteron*


Gene Name	Gene symbols of potential human orthologues	Species		Aligned length (nt)	$K_s$	$K_a$
		<i>Petromyzon</i>	<i>Lethenteron</i>			
rhodopsin	RHO	Ga	Lj	1059	0.76	0.04
red opsin	OPN1LW	Ga	Lj	1044	1.31	0.06

Species names are indicated as abbreviations: Pm, *Petromyzon marinus*; Lj, *Lethenteron japonicum*; Lr, *Lethenteron reissneri*; Lf, *Lampetra fluviatilis*; Ga, *Geotria australis*. We treated the genus *Lampetra* as equally distant from *Lethenteron* in relation to *Petromyzon* based on the previous molecular phylogenetic study (Docker *et al.*, 1999).

**Table 3.** Divergence times and numbers of synonymous substitutions reported for closely related organism pairs.

Taxonomic group	Species pairs <sup>a</sup>	$K_s$				Divergence time	
		Average	Method	Gene number	Reference	Mya	Reference
Mammalia	1, human-mouse	0.56	ML	11084	RGSPC, 2004	87	Springer <i>et al.</i> , 2003
	2, mouse-rat	0.19	ML	11503	RGSPC, 2004	16	Springer <i>et al.</i> , 2003
	3, human-pig	0.31	ML	1120	Jørgensen <i>et al.</i> , 2005	94	Springer <i>et al.</i> , 2003
Amniota	4, human-chicken	1.66	ML	7529	ICGSC, 2004	310	Benton, 1993
Sauropsida	5, chicken-turkey	0.12	ML	155	Axelsson <i>et al.</i> , 2005	28	Dimcheff <i>et al.</i> , 2002
	6, chicken-turtle	0.96	ML	56	Kuraku <i>et al.</i> , 2006	222–276	Kumar and Hedges, 1998
Teleostei	7, <i>Tetraodon nigroviridis</i> - <i>Fugu rubripes</i>	0.42	PBL	5802	Jaillon <i>et al.</i> , 2003	18–30	Crnogorac-Jurcevic <i>et al.</i> , 1997
Cephalochordata	8, <i>Branchiostoma belcheri</i> - <i>B. floridae</i>	0.38	ML	14	this study <sup>b</sup>	112	Nohara <i>et al.</i> , 2005

ML, Calculated with the method of Goldman and Yang (1994); PBL, Calculated with the method by Pamilo and Bianchi (1993) and Li (1993). ICGSC, International Chicken Genome Sequencing Consortium. RGSPC, Rat Genome Sequencing Project Consortium. <sup>a</sup>Numbers 1–8 in the column of species pairs correspond to those in Fig. 3. <sup>b</sup>See Supplemental Table S2.



**Fig. 3.** Divergence times and number of synonymous substitutions per site. (A) The tentative clock calibrated by all of the divergences of species pairs (black circles) in Table 3. (B) The tentative clock calibrated by nodes 3, 5, 6, and 8 in Table 3. In this “Clock B”, nodes 1 and 2 were excluded because of an elevated evolutionary rate in the rodent lineage. Node 4 was excluded because of its unrelatively high  $K_s$  value ( $>1$ ). Node 7 was excluded because  $K_s$  for this species pair was estimated with a different method from the others. In both clocks, regression lines indicated as broken lines pass through the origin. For nodes 6 and 7, the intermediate values of estimated ranges of divergence times in Table 3 were used as divergence times in this Fig. Divergences in cyclostomata (e–g) that were dated with these clocks are indicated as open circles.

*petra* pairs for which the sequences were available in public databases (Lee and Kocher, 1995; Rasmussen *et al.*, 1998; Delarbre *et al.*, 2000; Delarbre *et al.*, 2001; Delarbre *et al.*, 2002). The average  $K_s$  for the *Petromyzon–Lampetra* pair was 1.19 (SD, 0.57;  $n=13$ ; Supplemental Table S1, <http://dx.doi.org/10.2108/zsj.23.1053>), whereas the *Myxine–Eptatretus* pair yielded an apparently saturated  $K_s$  ( $>2$ ; data not shown). The difference in  $K_s$  values between mitochondrial

and nuclear genes (7.9-fold for the *Petromyzon–Lampetra* pair) was roughly consistent with previous observations in mammals and amphibians (Miyata *et al.*, 1982; Crawford, 2003).

#### Rough estimation of divergence times based on number of synonymous substitutions

The number of synonymous substitutions per site has been reported for some pairs of chordate species (Table 3). In the present study, to supplement pre-existing data, we preliminarily estimated the  $K_s$  between two species of the cephalochordate genus *Branchiostoma* (Supplemental Table S2, <http://dx.doi.org/10.2108/zsj.23.1053>). The divergence times and  $K_s$  values in Table 3, including the data for *Branchiostoma*, were plotted two-dimensionally in Fig. 3. The overall rate of this putative clock was  $2.4 \times 10^{-9}$ /site/year (Clock A; Fig. 3A), whereas the rate of the clock for selected species pairs (for details, see legend for Fig. 3) was  $1.9 \times 10^{-9}$ /site/year (Clock B; Fig. 3B). By applying these clocks tentatively to cyclostome taxon pairs, we obtained divergence times of the inter-subfamilial split between Myxiniinae and Eptatretinae in the hagfish lineage at 93–28 Mya, the intergeneric split between *Petromyzon* and *Lethenteron* at 57–15 Mya, and the inter-subfamilial split between Geotriinae and Petromyzoninae in the lamprey lineage at 383–136 Mya (Table 4).

#### Estimation of divergence times using amino acid sequences

We selected 10 nuclear protein-coding genes with relatively long alignment lengths ( $>150$  amino acids; total length, 2947 amino acids; Table 5) in which no gene duplication was detected in major vertebrate lineages, as shown in Fig. 4 for the gene *GNB2L1*. The upper and lower limits of divergence times for branching points outside the cyclostomes were preset as shown in Table 6, and the tree topology shown in Fig. 5 was assumed. By executing the MULTIDIV-TIME program (Kishino *et al.*, 2001), we obtained divergence times for Myxiniiformes and Petromyzoniiformes at 671–391 Mya, Myxiniinae and Eptatretinae at 162–63 Mya, and *Petromyzon* and *Lethenteron* at 30–2 Mya (Table 7).

In addition, the timing of the above branching points was estimated using 12 mitochondrial protein-coding genes that had relatively long alignment lengths (total length, 3320 amino acids). ATP synthase F0 subunit 8 was excluded from this analysis because of its short alignment length. As a result, divergence times were estimated to be 728–459 Mya for the Myxiniiformes–Petromyzoniiformes split, 72–39 Mya for the Myxiniinae–Eptatretinae split, and 37–18 Mya for the *Petromyzon–Lampetra* split (Table 7).

**Table 4.** Divergence times estimated with tentative synonymous substitution clocks.


Node (Taxon pair)	e (Myxiniinae-Eptatretinae)		f ( <i>Petromyzon</i> - <i>Lethenteron</i> )		g (Geotriinae-Petromyzoninae)	
<i>K</i> <sub>s</sub>	0.24±0.11 (n=11)		0.14±0.07 (n=19)		1.03±0.39 (n=2)	
Clock	A	B	A	B	A	B
Estimated divergence time (Mya)	51±23	65±28	30±15	38±19	219±83	278±105

Clocks A and B correspond to those in Fig. 3. Nodes e–g correspond to those in Fig. 5.

**Table 5.** Genes on nuDNA used for amino acid-based estimation of divergence times.

Gene name	Gene symbols of a human ortholog	Species <sup>a</sup>		Aligned length (aa)
		Myxiniiformes	Petromyzoniformes	
DNA-directed RNA polymerase II largest subunit	<i>POLR2A</i>	<i>Eb</i>	<i>Lr</i>	595
DNA-directed RNA polymerase III largest subunit	<i>POLR3A</i>	<i>Eb</i>	<i>Lr</i>	493
eukaryotic translation elongation factor 1 gamma	<i>EEF1G</i>	<i>Mg, Eb</i>	<i>Pm, Lj</i>	275
guanine nucleotide binding protein, beta polypeptide 2-like 1	<i>GNB2L1</i>	<i>Mg, Eb</i>	<i>Pm, Lj</i>	299
H+-transporting ATP synthase alpha subunit isoform 1	<i>ATP5A1</i>	<i>Mg, Eb</i>	<i>Pm</i>	332
ribosomal protein S2	<i>RPS2</i>	<i>Mg, Eb</i>	<i>Pm, Lj</i>	201
ribosomal protein S3	<i>RPS3</i>	<i>Mg, Eb</i>	<i>Pm, Lj</i>	190
ribosomal protein L5	<i>RPL5</i>	<i>Mg, Eb</i>	<i>Pm, Lj</i>	165
ribosomal protein L7A	<i>RPL7A</i>	<i>Mg, Eb</i>	<i>Pm, Lj</i>	241
triosephosphate isomerase	<i>TPI1</i>	<i>Eb</i>	<i>Lr</i>	206

<sup>a</sup>Species names: *Mg*, *Myxine glutinosa*; *Eb*, *Eptatretus burgeri*; *Pm*, *Petromyzon marinus*; *Lj*, *Lethenteron japonicum*; *Lr*, *Lethenteron reissneri*.



**Fig. 4.** Molecular phylogenetic tree of a guanine nucleotide-binding protein, beta polypeptide 2-like 1 (*GNB2L1*) protein. The tree was inferred with the neighbor-joining method using 300 amino acid residues and assuming among-site rate heterogeneity (shape parameter  $\alpha=0.64$ ; see MATERIALS AND METHODS for details). The maximum-likelihood method yielded similar tree topologies, although they are not statistically significant. For amino acid-based divergence time estimation, we selected 10 genes with no gene duplication, as shown in this tree.

**Table 6.** Divergence times for constrained nodes.

Node	a		b		c		Reference
	Min	Max	Min	Max	Min	Max	
I	300	300	400	400	500	500	Dickerson, 1971
II	310	310	415	485	489	639	Kumar and Hedges, 1998
III	310	370	425	495	605	742	Blair and Hedges, 2005

Max, Upper limit of divergence time. Min, lower limit of divergence time. Letters indicating nodes (a–c) correspond to those in Fig. 5.


**DISCUSSION**

**GC<sub>4</sub> as a reflection of base composition in cyclostome genomes**

The GC-content at synonymous sites in a protein-coding gene is expected to positively correlate with the global GC-content of the genomic region where the gene is located (Clay *et al.*, 1996; Musto *et al.*, 1999; Kuraku *et al.*, 2006). Therefore, we focused on GC<sub>4</sub> in currently available cDNAs reported for cyclostomes (Fig. 2). Hagfish and lamprey cDNAs show similar levels of heterogeneity in GC-content (Fig. 2). However, there is a striking difference in the level of GC-content between hagfish and lamprey: every lamprey species we analyzed showed a high GC<sub>4</sub> (70–90%; Figs.

2A–C), whereas every hagfish species we analyzed showed a relatively moderate GC<sub>4</sub> (40–60%; Figs. 2D–F).

Cytogenetic studies have revealed that hagfishes have a relatively moderate number of relatively moderate-sized chromosomes compared to other vertebrates (2n=14–36 in somatic cells), whereas lampreys possess a much greater number of small, dot-like chromosomes (2n=76–178; Potter and Rothwell, 1970; Potter and Robinson, 1971; Robinson *et al.*, 1975; Nakai *et al.*, 1995; Animal Genome Size Database, <http://www.genomesize.com>). The contrast in chromosome size, chromosome number, and GC-content between hagfishes and lampreys is reminiscent of the intra-genomic difference between macrochromosomes and microchromosomes seen in sauropsids (Burt, 2002; Kuraku *et al.*, 2006).



**Fig. 5.** A tree topology preset for divergence time estimation. Letters at nodes correspond to those in Tables 6 and 7.

Further investigation will be required to understand the putative relationships among these genomic features.

#### **$K_s$ as a tool to standardize evolutionary distances**

The number of synonymous substitutions per site in a protein-coding region serves as an ideal standard for evolutionary distance when comparing closely related species, as long as the gene in question has evolved in a neutral manner (Miyata and Yasunaga, 1980; Perler *et al.*, 1980). In this study, we estimated  $K_s$  with the maximum-likelihood method (Goldman and Yang, 1994) because this method is expected to produce relatively appropriate estimates, even in species with highly biased base compositions, such as lampreys (Fig. 2). For all the genes analyzed in this study,  $K_s$  was larger than  $K_a$ , indicating that these genes have evolved neutrally without experiencing positive selection. In selecting pairs of cyclostome species for  $K_s$  estimation, we treated the lamprey genus *Lampetra* as having the same distance the genus *Lethenteron* has from *Petromyzon*. This is based on a previous phylogenetic study using the cytochrome *b* and NADH dehydrogenase subunit 3 (ND3) genes (Docker *et al.*, 1999), which is consistent with the classification by Potter (1980), who formerly considered that the three subgenera, *Entosphenus*, *Lethenteron*, and *Lampetra*, compose the single genus *Lampetra*. Despite the small number of genes sampled, similar  $K_s$  values were consistently obtained for *Petromyzon*–*Lampetra* and *Petromyzon*–*Lethenteron* pairs (Table 2). In contrast to inter-generic and inter-subfamilial comparisons in cyclostomes, our prelimi-

nary  $K_s$  estimates between two tunicate species in the same genus (*Ciona intestinalis* and *C. savignyi*) and two amphioxus species in the same genus (*Branchiostoma belcheri* and *B. floridae*) yielded relatively larger  $K_s$  values (see Supplemental Table S2, <http://dx.doi.org/10.2108/zsj.23.1053> for *Branchiostoma*;  $K_s > 3$  for *Ciona*; data not shown), despite their close taxonomic distances.

Theoretically, the  $K_s$  value is determined by the time that has elapsed since the divergence of two species in question, as long as neutrality holds. This feature of  $K_s$  is useful in judging orthology between genes (especially members of a gene family prone to gene duplications) of closely related species. If this idea is tentatively applied to previous cross-species comparisons in lamprey studies, for example, orthology between *L. japonicum Hox6w* and *L. fluviatilis HoxL6* is again confirmed with an extremely low  $K_s$  value ( $K_s = 0.039$ ), a reasonable estimate for intrasubfamilial comparison, as suggested previously (Takio *et al.*, 2004).

#### **Methodological aspects of molecular dating**

Proposing a constant rate of nucleotide and amino acid substitutions was one of the milestones for molecular dating of divergence times (“molecular clock”; Zuckerkandl and Pauling, 1962, 1965; also see Donoghue *et al.*, 2003; Bromham and Penny, 2003; and Kumar, 2005, for review). As shown in Fig. 3, the numbers of synonymous substitutions per site and divergence times behave in a clock-like manner, at least within the chordates, indicating that this may serve as a rough molecular clock for species pairs with an unsaturated  $K_s$ . However, this clock was calibrated by branching points in different lineages outside the cyclostomes (Table 3), because no branching point with a known divergence time was available within cyclostomes. In addition, variation of evolutionary rate among lineages, such as rate elevation in rodents (Kikuno *et al.*, 1985; Wu and Li, 1985; Rat Genome Sequencing Project Consortium, 2004), may confuse divergence time estimation. Therefore, our results need to be verified by refinement of this silent clock with reexamination of divergence times and estimation of  $K_s$  for more pairs of organisms.

We utilized amino acid sequences to obtain more robust estimates. Especially in our  $K_s$  analysis, synonymous substitutions between Myxiniformes and Petromyzoniformes were apparently saturated (data not shown), suggesting that this split would need to be dated with amino acid sequences rather than nucleotide sequences. However, as exemplified in Fig. 4, phylogenetic trees including cyclostome species often show a high degree of rate heterogeneity because of the accelerated evolutionary rate in hagfishes. To minimize the undesirable influence of this rate heterogeneity on divergence time estimation, we employed a non-parametric

**Table 7.** Divergence times estimated with amino acid sequences.

Molecule type	Genes on nuDNA (n=10)					Genes on mtDNA (n=12)			
Node	c	d	e	f	c	d	e	f	
Constraint set I	500± 0	432±41	88±25	11± 9	500± 0	478±19	44±5	21±3	
Constraint set II	629± 9	524±53	107±30	14±11	629± 8	602±24	55±6	27±4	
Constraint set III	730±11	612±59	127±35	17±13	725±15	698±30	65±7	32±5	

Letters indicating nodes (c–f) correspond to those in Fig. 5. For details of constraint sets (I–III), see Table 5. Divergence times are shown with standard errors representing 95% confidence intervals.



molecular dating method that does not assume rate constancy (Kishino *et al.*, 2001; Thorne and Kishino, 2002; see also Hasegawa *et al.*, 2003). Moreover, we paid close attention to the orthologous/paralogous relationships of multiple members of gene families between hagfishes, lampreys, and gnathostomes, because putative genome duplications in early vertebrate evolution (Ohno, 1970; McLysaght *et al.*, 2002) often confuse orthology identification; Kuraku *et al.* (1999) provided an example in the *enolase* gene family. In addition, inclusion of genes prone to gene duplications may also result in misleading estimates of divergence time, possibly because of an accelerated evolutionary rate caused by neofunctionalization or subfunctionalization of duplicates. For these reasons, we deliberately selected genes for which no gene duplication was detected in major vertebrate lineages (Table 5).

As calibration points outside the cyclostomes, we used three sets of divergence time constraints (Table 6). One of the three constraint sets was based on the fossil records used by Dickerson (1971) (constraint set I), whereas the other two were based on previous studies using molecular data (constraint set II, Kumar and Hedges, 1998; constraint set III, Blair and Hedges, 2005). However, these molecular studies included genes that underwent duplication events early in vertebrate evolution and do not represent a 1:1 relationship between a cyclostome gene and a gnathostome counterpart (e.g., *bone morphogenetic protein (BMP) 2/4*, *enolase-2*). Although fossil records inherently tend to yield more recent divergence times because of potential incomplete fossil sampling in more ancient eras, it is possible that inappropriate gene selection might have yielded much more ancient estimates in these studies (constraint sets II and III) compared with commonly accepted fossil records (constraint set I). This discrepancy emphasizes again that precision in gene selection cannot be sacrificed, even in the name of high-throughput analysis using large data sets. For this reason, we summarize our results below, along a temporal axis based on the fossil records.

### Temporal reconstruction of cyclostome phylogeny


The monophyly of cyclostomes has resulted in a dispute over when hagfishes and lampreys split from each other in the cyclostome lineage. To answer this question, our analyses using mitochondrial and nuclear genes consistently showed that Myxiniiformes and Petromyzoniiformes diverged from each other 30–110 million years after the cyclostome lineage split from the future gnathostome lineage (Table 7). When calibration by fossil records was applied, this Myxiniiformes–Petromyzoniiformes split dated back to 470–390 Mya in the Ordovician–Silurian–Devonian Periods (Fig. 6), when fossil agnathans are thought to have diversified (Forey and Janvier, 1993; Janvier, 1996). Although we still do not know the precise branching pattern among these agnathans, hagfishes and lampreys represent two distinct agnathan groups that diverged early in vertebrate evolution and have survived thereafter for more than 400 million years. Our results indicate that, although both groups are classified as cyclostomes, the distance between hagfishes and lampreys is similar to that between humans and cartilaginous fishes, in terms of the geological time that has elapsed since their divergence.

Later in the hagfish lineage, our molecular dating with amino acid-based relaxed clocks and synonymous substitution clocks indicated that there was no branching of extant taxa until the two hagfish subfamilies, Myxiniinae and Eptatretinae, split from each other 90–60 Mya in the Cretaceous–Tertiary Periods (Fig. 6). However, there is a paleontological report of a fossil species, *Myxiniakela siroka*, from the Carboniferous fauna (~300 Mya) that is regarded as a putative outgroup of extant hagfishes (Bardack, 1991). In our divergence time estimate using relaxed molecular clocks, even when we tentatively constrained the upper limit of the divergence time of the Myxiniinae–Eptatretinae split to 300 Mya, we obtained an identical result (data not shown), suggesting that this fossil species, *Myxiniakela siroka*, actually is an outgroup of extant hagfishes (Fig. 6).

All the three subfamilies (Mordaciinae, Geotriinae, and Petromyzoninae) in the lamprey lineage are thought to have diverged from one another in a considerably short period of time (Conlon *et al.*, 2001; Gill *et al.*, 2003), as indicated by ambiguous phylogenetic relationships between these three taxa in recent molecular studies (Baldwin *et al.*, 1988; Silver *et al.*, 2004; Takahashi *et al.*, 2006). Our synonymous substitution clock indicates that the Geotriinae–Petromyzoninae split occurred in the Permian–Triassic Periods (280–220 Mya; Fig. 6). Although this estimate needs to be reinforced with more robust analyses using amino acid sequences, this divergence time may coincide with the break-up of Gondwana (inhabited by species in Mordaciinae and Geotriinae) from Laurasia (inhabited by species in Petromyzoninae). Later, in the lineage of Petromyzoninae, an inter-generic split between *Petromyzon* and *Lethenteron/Lampetra* occurred 30–10 Mya in the Tertiary Period (Fig. 6). This result is consistent with the rough estimate by Docker *et al.* (1999), who simply assumed that a 2% divergence in mtDNA sequence corresponds to one million years (Brown *et al.*, 1979). To further confirm phylogenetic relationships in Petromyzoninae, reported previously based on morphological features (Gill *et al.*, 2003), molecular sequence data from other genera need to be included. Fossils of the lampreys *Hardistiella montanensis*, *Mayomyzon pieckoensis*, and *Pipiscius zangerli* were found in the Carboniferous fauna (~280 Mya) and have been treated as outgroups of all extant lampreys (Bardack and Zangerl, 1971; Bardack and Richardson, 1977; Janvier and Lund, 1983). In nuDNA-based and mtDNA-based analyses using relaxed molecular clocks, adding these paleontological data did not produce any substantial differences in results (data not shown), indicating that these fossil lampreys should still be regarded as outgroups of all extant lamprey species in Petromyzoniiformes (Fig. 6).

### Perspectives

Thanks to the efforts of researchers in various fields of biology (e.g., Kuratani *et al.*, 2002), nucleotide and amino acid sequences of hagfishes and lampreys are accumulating in public databases. However, information at the molecular level is still far from satisfactory for cyclostomes (Fig. 1), in two aspects. First, in terms of the coverage of species diversity, there is a paucity of molecular data for southern hemisphere species, and these data are crucial for inferences of phylogenetic relationships and divergence times. For exam-



**Fig. 6.** Hypothesized time scale of cyclostome evolution. The phylogeny and timing of cyclostome diversification are illustrated so that branch lengths correspond to evolutionary times (black solid lines). Numerical time scales corresponding to constraint sets I–III in Table 6 are indicated in the upper rows. The geological time scale based on the chart provided by the Geological Society of America (<http://www.geosociety.org/science/timescale/timescl.htm>) is added to the time scale based on constraint set I. Black circles indicate branching points whose divergence times were estimated in this study. The divergence time of the split between Myxiniinae and Eptatretinae and between *Petromyzon* and *Lethenteron/Lampetra* is based on both synonymous substitution clocks and amino acid-based relaxed clocks using mtDNA and nuDNA. The divergence time of the split between Myxiniformes and Petromyzoniformes is solely based on a relaxed clock analysis using amino acid sequences encoded by mtDNA and nuDNA. The divergence time of the split between Geotriinae and Petromyzoninae is estimated based on synonymous substitution clocks. Fossil records are indicated by solid grey lines, with the putative phylogenetic relationships proposed by Janvier (1997).

ple, in Myxiniformes, no molecular sequence data have been reported for *Notomyxine*, *Nemamyxine*, or *Neomyxine*, which are thought to belong in the subfamily Myxiniinae (Jørgensen, 1998). Similarly, in lampreys, the unavailability of appropriate nucleotide sequences for *Mordacia* hindered inclusion of Mordaciinae in our silent clock analysis, and the unavailability of appropriate amino acid sequences for *Mordacia* and *Geotria* did not allow us to include Mordaciinae and Geotriinae in our amino acid-based relaxed clock analysis.

Second, there are few reports of genomic DNA sequences for hagfishes and lampreys, and most of the reported nucleotide sequences are derived from mRNAs. In this study, based on the nucleotide sequences of protein-coding exons, we estimated accumulated levels of synonymous substitutions for inter-subfamilial and inter-generic species pairs, which highlighted a relatively low level of neutral nucleotide changes in the *Myxine*–*Eptatretus* and *Petromyzon*–*Lethenteron* pairs (Tables 1 and 2A). If the synonymous substitution rate in coding regions roughly cor-

responds to the neutral substitution rate in intergenic or intronic regions without regulatory functions, our estimates imply that these taxon pairs might be too phylogenetically close to discern potentially functional sequences, such as *cis*-regulatory elements or non-coding genes.

This principle is referred to as “phylogenetic footprinting” (Gumucio *et al.*, 1992; see also Zhang and Gerstein, 2003, for review), and selecting multiple species with appropriate levels of nucleotide substitution facilitates an efficient *in silico* detection of potentially functional genomic sequences (Uchikawa *et al.*, 2003; Johnson *et al.*, 2004; Kusakabe, 2005). In this context, using lampreys as an example, comparison of non-coding genomic sequences among multiple species in Petromyzoninae alone will not provide a sufficient level of resolution to highlight functional fractions. Instead, judging from the almost saturated  $K_s$  level between Geotriinae and Petromyzoninae (Table 2B), inclusion of southern hemisphere lampreys (*Mordacia* or *Geotria*) would be highly promising in comparisons with species in the northern hemisphere subfamily Petromyzoninae, such as *Petromyzon marinus*, whose genome sequencing project is now underway.

Despite the scarcity of sequence information, we attempted to overview the general features of base composition and evolutionary distance in cyclostomes, and obtained results that will serve as standards for future evolutionary and genomic studies of cyclostomes. We propose that, in any taxa, this sort of succinct evolutionary analysis should be a prerequisite for any biological studies involving multi-species comparisons. Even if the amount of available sequence information is limited, general trends embedded in sequence information can be extracted in light of the theories of molecular evolution.

## ACKNOWLEDGMENTS

We are very grateful to Kazutaka Katoh for computational advice and to Kinya Ota, David McCauley, and Yoichi Matsuda for insightful discussion. This work was supported by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

## REFERENCES

- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402
- Axelsson E, Webster MT, Smith NG, Burt DW, Ellegren H (2005) Comparison of the chicken and turkey genomes reveals a higher rate of nucleotide divergence on microchromosomes than macrochromosomes. *Genome Res* 15: 120–125
- Baldwin J, Mortimer K, Patak A (1988) Evolutionary relationships among lamprey families: amino acid composition analysis of lactate dehydrogenase. *Biochem Syst Ecol* 16: 351–353
- Bardack D (1991) First fossil hagfish (Myxinoidea): a record from the Pennsylvanian of Illinois. *Science* 254: 701–703
- Bardack D, Richardson ES (1977) New agnathous fishes from the Pennsylvanian of Illinois. *Fieldiana Geol* 33: 489–510
- Bardack D, Zangerl R (1971) Lampreys in the fossil record. In “The Biology of Lampreys” Ed by MW Hardisty, IC Potter, Academic Press, London & New York, pp 67–84
- Benton MJ (1993) The Fossil Record 2. 1st ed, Palaeontological Association, Royal Society (Great Britain), Linnean Society of London, Chapman and Hall, London
- Blair JE, Hedges SB (2005) Molecular phylogeny and divergence times of deuterostome animals. *Mol Biol Evol* 22: 2275–2284
- Bromham L, Penny D (2003) The modern molecular clock. *Nat Rev Genet* 4: 216–224
- Brown WM, George M Jr, Wilson AC (1979) Rapid evolution of animal mitochondrial DNA. *Proc Natl Acad Sci USA* 76: 1967–1971
- Burt DW (2002) Origin and evolution of avian microchromosomes. *Cytogenet Genome Res* 96: 97–112
- Chen Y, Chang H, Mok H (2005) Phylogenetic position of *Eptatretus chinensis* (Myxiniidae: Myxiniformes) inferred by 16S rRNA gene sequence and morphology. *Zool Stud* 44: 111–118
- Clay O, Caccio S, Zoubak S, Mouchiroud D, Bernardi G (1996) Human coding and noncoding DNA: compositional correlations. *Mol Phylogenet Evol* 5: 2–12
- Conlon JM, Wang Y, Potter IC (2001) The structure of *Mordacia mordax* insulin supports the monophyly of the Petromyzontiformes and an ancient divergence of Mordaciidae and Geotriidae. *Comp Biochem Physiol B Biochem Mol Biol* 129: 65–71
- Crawford AJ (2003) Relative rates of nucleotide substitution in frogs. *J Mol Evol* 57: 636–641
- Crnogorac-Jurcevic T, Brown JR, Lehrach H, Schalkwyk LC (1997) *Tetraodon fluviatilis*, a new puffer fish model for genome studies. *Genomics* 41: 177–184
- Delarbre C, Escriva H, Gallut C, Barriel V, Kourilsky P, Janvier P, Laudet V, Gachelin G (2000) The complete nucleotide sequence of the mitochondrial DNA of the agnathan *Lampetra fluviatilis*: bearings on the phylogeny of cyclostomes. *Mol Biol Evol* 17: 519–529
- Delarbre C, Rasmussen AS, Arnason U, Gachelin G (2001) The complete mitochondrial genome of the hagfish *Myxine glutinosa*: unique features of the control region. *J Mol Evol* 53: 634–641
- Delarbre C, Gallut C, Barriel V, Janvier P, Gachelin G (2002) Complete mitochondrial DNA of the hagfish, *Eptatretus burgeri*: the comparative analysis of mitochondrial DNA sequences strongly supports the cyclostome monophyly. *Mol Phylogenet Evol* 22: 184–192
- Delsuc F, Brinkmann H, Chourrout D, Philippe H (2006) Tunicates and not cephalochordates are the closest living relatives of vertebrates. *Nature* 439: 965–968
- Dickerson RE (1971) The structures of cytochrome c and the rates of molecular evolution. *J Mol Evol* 1: 26–45
- Dimcheff DE, Drovetski SV, Mindell DP (2002) Phylogeny of Tetraoninae and other galliform birds using mitochondrial 12S and ND2 genes. *Mol Phylogenet Evol* 24: 203–215
- Docker MF, Youson JH, Beamish RJ, Devlin RH (1999) Phylogeny of the lamprey genus *Lampetra* inferred from mitochondrial cytochrome *b* and ND3 gene sequences. *Can J Fish Aquat Sci* 56: 2340–2349
- Donoghue PCJ, Smith MP, Sansom IJ (2003) The origin and early evolution of chordates: molecular clocks and the fossil record. In “Telling the Evolutionary Time” Ed by PCJ Donoghue, MP Smith, Taylor & Francis, London, pp 190–223
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17: 368–376
- Fernholm B (1998) Hagfish systematics. In “The Biology of Hagfishes” Ed by JM Jørgensen, JP Lomholt, RE Weber, H Malte, Chapman & Hall, London, pp 33–44
- Forey P, Janvier P (1993) Agnathans and the origin of jawed vertebrates. *Nature* 361: 129–134
- Furlong RF, Holland PWH (2002) Bayesian phylogenetic analysis supports monophyly of ambulacraria and of cyclostomes. *Zool Sci* 19: 593–599
- Gill HS, Renaud CB, Chapleau F, Mayden RL, Potter IC (2003) Phylogeny of living parasitic lampreys (Petromyzontiformes) based on morphological data. *Copeia* 4: 687–703

- Goldman N, Yang Z (1994) A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Mol Biol Evol* 11: 725–736
- Gumucio DL, Heilstedt-Williamson H, Gray TA, Tarle SA, Shelton DA, Tagle DA, Slightom JL, Goodman M, Collins FS (1992) Phylogenetic footprinting reveals a nuclear protein which binds to silencer sequences in the human gamma and epsilon globin genes. *Mol Cell Biol* 12: 4919–4929
- Hardisty MW, Potter IC (1971) *The Biology of Lampreys*. Academic Press, London & New York
- Hasegawa M, Thorne JL, Kishino H (2003) Time scale of eutherian evolution estimated without assuming a constant rate of molecular evolution. *Genes Genet Syst* 78: 267–283
- Hubbs CL, Potter IC (1971) Distribution, phylogeny, and taxonomy. In “*The Biology of Lampreys*” Ed by MW Hardisty, IC Potter, Academic Press, London & New York, pp 1–66
- International Chicken Genome Sequencing Consortium (ICGSC) (2004) Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature* 432: 695–716
- Jaillon O, Aury JM, Brunet F, Petit JL, Stange-Thomann N, Mauceli E, Bouneau L, Fischer C, Ozouf-Costaz C, Bernot A *et al.* (2004) Genome duplication in the teleost fish *Tetraodon nigroviridis* reveals the early vertebrate proto-karyotype. *Nature* 431: 946–957
- Janke A, Arnason U (1997) The complete mitochondrial genome of *Alligator mississippiensis* and the separation between recent archosauria (birds and crocodiles). *Mol Biol Evol* 14: 1266–1272
- Janvier P, Lund R (1983) *Hardistiella montanensis* n. gen. et sp. (Petromyzontida) from the Lower Carboniferous of Montana, with remarks on the affinities of the lampreys. *J Vert Paleont* 2: 407–413
- Janvier P (1996) The dawn of the vertebrates: characters versus common ascent in the rise of current vertebrate phylogenies. *Paleontology* 39: 259–287
- Janvier P (1997) The Tree of Life Project (URL: <http://tolweb.org/Hyperoartia/14831>)
- Johnson DS, Davidson B, Brown CD, Smith WC, Sidow A (2004) Noncoding regulatory sequences of *Ciona* exhibit strong correspondence between evolutionary constraint and functional importance. *Genome Res* 14: 2448–2456
- Jørgensen JM (1998) *The Biology of Hagfishes*. 1st ed, Chapman and Hall, London
- Jørgensen FG, Hobolth A, Hornshøj H, Bendixen C, Fredholm M, Schierup MH (2005) Comparative analysis of protein coding sequences from human, mouse and the domesticated pig. *BMC Biol* 3: 2
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30: 3059–3066
- Kikuno R, Hayashida H, Miyata T (1985) Rapid rate of rodent evolution. *Proc Jpn Acad* 61: 153–156
- Kishino H, Miyata T, Hasegawa M (1990) Maximum likelihood inference of protein phylogeny and the origin of chloroplasts. *J Mol Evol* 30: 151–160
- Kishino H, Thorne JL, Bruno WJ (2001) Performance of a divergence-time estimation method under a probabilistic model of rate evolution. *Mol Biol Evol* 18: 352–361
- Kumar S (2005) Molecular clocks: four decades of evolution. *Nat Genet Rev* 6: 654–662
- Kumar S, Hedges SB (1998) A molecular timescale for vertebrate evolution. *Nature* 392: 917–920
- Kuo CH, Huang S, Lee SC (2003) Phylogeny of hagfish based on the mitochondrial 16S rRNA gene. *Mol Phylogenet Evol* 28: 448–457
- Kuraku S, Hoshiyama D, Katoh K, Suga H, Miyata T (1999) Monophyly of lampreys and hagfishes supported by nuclear DNA-coded genes. *J Mol Evol* 49: 729–735
- Kuraku S, Ishijima J, Nishida-Umehara C, Agata K, Kuratani S, Matsuda Y (2006) cDNA-based gene mapping and GC<sub>3</sub> profiling in soft-shelled turtle suggest chromosome size-dependent GC bias shared by sauropsids. *Chromosome Res* 14: 187–202
- Kuratani S, Kuraku S, Murakami Y (2002) Lamprey as an evo-devo model: lessons from comparative embryology and molecular phylogenetics. *Genesis* 34: 175–183
- Kusakabe T (2005) Decoding *cis*-regulatory systems in ascidians. *Zool Sci* 22: 129–146
- Lee WJ, Kocher TD (1995) Complete sequence of a sea lamprey (*Petromyzon marinus*) mitochondrial genome: early establishment of the vertebrate genome organization. *Genetics* 139: 873–887
- Mallatt J, Sullivan J (1998) 28S and 18S rDNA sequences support the monophyly of lampreys and hagfishes. *Mol Biol Evol* 15: 1706–1718
- McLysaght A, Hokamp K, Wolfe KH (2002) Extensive genomic duplication during early chordate evolution. *Nat Genet* 31: 200–204
- Miyata T, Yasunaga T (1980) Molecular evolution of mRNA: a method for estimating evolutionary rates of synonymous and amino acid substitutions from homologous nucleotide sequences and its application. *J Mol Evol* 16: 23–36
- Miyata T, Hayashida H, Kikuno R, Hasegawa M, Kobayashi M, Koike K (1982) Molecular clock of silent substitution: at least six-fold preponderance of silent changes in mitochondrial genes over those in nuclear genes. *J Mol Evol* 19: 28–35
- Musto H, Romero H, Zavala A, Bernardi G (1999) Compositional correlations in the chicken genome. *J Mol Evol* 49: 325–329
- Nakai Y, Kubota S, Goto Y, Ishibashi T, Davison W, Kohno S (1995) Chromosome elimination in three Baltic, south Pacific and north-east Pacific hagfish species. *Chromosome Res* 3: 321–330
- Nohara M, Nishida M, Manthacitra V, Nishikawa T (2004) Ancient phylogenetic separation between Pacific and Atlantic cephalochordates as revealed by mitochondrial genome analysis. *Zool Sci* 21: 203–210
- Ohno S (1970) *Evolution by Gene Duplication*. Springer-Verlag, Berlin
- Ota K, Kuratani S (2006) The history of scientific endeavors towards understanding hagfish embryology. *Zool Sci* 23: 403–418
- Pancer Z, Amemiya CT, Ehrhardt GR, Ceitlin J, Gartland GL, Cooper MD (2004a) Somatic diversification of variable lymphocyte receptors in the agnathan sea lamprey. *Nature* 430: 174–180
- Pancer Z, Mayer WE, Klein J, Cooper MD (2004b) Prototypic T cell receptor and CD4-like coreceptor are expressed by lymphocytes in the agnathan sea lamprey. *Proc Natl Acad Sci USA* 101: 13273–13278
- Pancer Z, Saha NR, Kasamatsu J, Suzuki T, Amemiya CT, Kasahara M, Cooper MD (2005) Variable lymphocyte receptors in hagfish. *Proc Natl Acad Sci USA* 102: 9224–9229
- Perler F, Efstratiadis A, Lomedico P, Gilbert W, Kolodner R, Dodgson J (1980) The evolution of genes: the chicken preproinsulin gene. *Cell* 20: 555–566
- Potter IC (1980) The Petromyzoniiformes with particular reference to paired species. *Can J Fish Aquat Sci* 37: 1595–1615
- Potter IC, Gill HS (2003) Adaptive radiation of lampreys. *J Great Lakes Res* 29: 95–112
- Potter IC, Robinson ES (1971) The chromosomes. In “*The Biology of Lampreys*” Ed by MW Hardisty, IC Potter, Academic Press, London, pp 279–293
- Potter IC, Rothwell B (1970) The mitotic chromosomes of the lamprey, *Petromyzon marinus* L. *Experientia* 26: 429–430
- Potter IC, Strahan R (1968) The taxonomy of the lampreys, *Geotria*

- and *Mordacia*, and their distribution in Australia. *Proc Linn Soc Lond* 179: 229–240
- Rasmussen AS, Janke A, Arnason U (1998) The mitochondrial DNA molecule of the hagfish (*Myxine glutinosa*) and vertebrate phylogeny. *J Mol Evol* 46: 382–388
- Rat Genome Sequencing Project Consortium (RGSP) (2004) Genome sequence of the brown Norway rat yields insights into mammalian evolution. *Nature* 428: 493–521
- Renaud CB (1997) Conservation status of northern hemisphere lampreys (Petromyzontidae). *J Appl Ichthyol* 13: 143–148
- Robinson ES, Potter IC, Atkin NB (1975) The nuclear DNA content of lampreys. *Experientia* 31: 912–913
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406–425
- Sanderson MJ (2002) Estimating absolute rates of molecular evolution and divergence times: a penalized likelihood approach. *Mol Biol Evol* 19: 101–109
- Sanderson MJ (2003) r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. *Bioinformatics* 19: 301–302
- Silver MR, Kawauchi H, Nozaki M, Sower SA (2004) Cloning and analysis of the lamprey GnRH-III cDNA from eight species of lamprey representing the three families of Petromyzoniformes. *Gen Comp Endocrinol* 139: 85–94
- Springer MS, Murphy WJ, Eizirik E, O'Brien SJ (2003) Placental mammal diversification and the Cretaceous-Tertiary boundary. *Proc Natl Acad Sci USA* 100: 1056–1061
- Stock DW, Whitt GS (1992) Evidence from 18S ribosomal RNA sequences that lampreys and hagfishes form a natural group. *Science* 257: 787–789
- Suzuki T, Shin IT, Kohara Y, Kasahara M (2004) Transcriptome analysis of hagfish leukocytes: a framework for understanding the immune system of jawless fishes. *Dev Comp Immunol* 28: 993–1003
- Takahashi A, Nakata O, Moriyama S, Nozaki M, Jos JMP, Sower SA, Kawauchi H (2006) Occurrence of two functionally distinct proopiomelanocortin genes in all modern lampreys. *Gen Comp Endocrinol* 148: 72–78
- Takezaki N, Figueroa F, Zaleska-Rutczynska Z, Klein J (2003) Molecular phylogeny of early vertebrates: monophyly of the agnathans as revealed by sequences of 35 genes. *Mol Biol Evol* 20: 287–292
- Takio Y, Pasqualetti M, Kuraku S, Hirano S, Rijli FM, Kuratani S (2004) Evolutionary biology: lamprey Hox genes and the evolution of jaws. *Nature* 429: 1 p following 262
- Thorne JL, Kishino H (2002) Divergence time and evolutionary rate estimation with multilocus data. *Syst Biol* 51: 689–702
- Uchikawa M, Ishida Y, Takemoto T, Kamachi Y, Kondoh H (2003) Functional analysis of chicken Sox2 enhancers highlights an array of diverse regulatory elements that are conserved in mammals. *Dev Cell* 4: 509–519
- Wu CI, Li WH (1985) Evidence for higher rates of nucleotide substitution in rodents than in man. *Proc Natl Acad Sci USA* 82: 1741–1745
- Yang Z (1994) Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *J Mol Evol* 39: 306–314
- Yang Z (1997) PAML: a program package for phylogenetic analysis by maximum likelihood. *Comput Appl Biosci* 13: 555–556
- Young JZ (1962) *The Life of Vertebrates*. 2nd ed, Oxford Univ Press, Oxford
- Zhang Z, Gerstein M (2003) Of mice and men: phylogenetic footprinting aids the discovery of regulatory elements. *J Biol* 2: 11
- Zuckerandl E, Pauling L (1962) Molecular disease, evolution and genetic heterogeneity. In "Horizons in Biochemistry" Ed by M Kasha, B Pullman, Academic Press, New York, pp 189–225
- Zuckerandl E, Pauling L (1965) Evolutionary divergence and convergence. In "Evolving Genes and Proteins" Ed by V Bryson, HJ Vogel, Academic Press, New York, pp 97–166

(Received May 1, 2006 / Accepted July 4, 2006)