

## **Madygella Humioi sp. nov. from the Upper Triassic Mine Group, Southwest Japan: The Oldest Record of a Sawfly (Hymenoptera: Symphyta) in East Asia**

Authors: Oyama, Nozomu, and Maeda, Haruyoshi

Source: Paleontological Research, 24(1) : 64-71

Published By: The Palaeontological Society of Japan

URL: <https://doi.org/10.2517/2019PR005>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# *Madygella humioi* sp. nov. from the Upper Triassic Mine Group, southwest Japan: the oldest record of a sawfly (Hymenoptera: Symphyta) in East Asia

NOZOMU OYAMA<sup>1</sup> AND HARUYOSHI MAEDA<sup>2</sup>

<sup>1</sup>Graduate School of Science, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka 819-0395, Japan (e-mail: 716o.nonn@gmail.com)

<sup>2</sup>The Kyushu University Museum, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

Received January 17, 2019; Revised manuscript accepted March 12, 2019

**Abstract.** A primitive sawfly, *Madygella humioi* sp. nov., belonging to the family Xyelidae (Hymenoptera: Symphyta), is newly described from the Upper Triassic Mine Group, Yamaguchi Prefecture, southwest Japan. The new species differs from the five previously known *Madygella* species in having a cell length of 1r plus 2r shorter than that of 3r+4r and a cell height of 3r+4r lower than 2r plus pterostigma in a forewing. To date, this is the oldest fossil record of sawflies in East Asia. Regarding genus *Madygella*, this is the first example found outside of the Kyrgyz Republic. This discovery provides an insight into the early evolution of the order Hymenoptera and suggests a widespread distribution of the pioneering genus *Madygella* during the Triassic period.

**Key words:** Hymenoptera, Japan, Mine Group, sawflies, Triassic, Xyelidae

## Introduction

The oldest representatives of the order Hymenoptera are xyelid sawflies, belonging to the family Xyelidae. Their fossil records include specimens from Middle to Upper Triassic deposits in the Kyrgyz Republic (Rasnitsyn, 1964, 1969; Kopylov, 2014) and from Upper Triassic deposits in Australia (Riek, 1955; Grimaldi and Engel, 2005), Argentina (Lara *et al.*, 2014), and South Africa (Schlüter, 2000). The Xyelidae family comprises four subfamilies: Xyelinae, Macroxyelinae, Archexyelinae, and Madygellinae. The extinct subfamily Madygellinae was a pioneering group of sawflies restricted to the Triassic period, comprising eight different species belonging to three genera occurring from the Middle to Upper Triassic in the Kyrgyz Republic (Rasnitsyn, 1969; Kopylov, 2014). Their existence in other areas is not known.


The Upper Triassic Mine Group, belonging to the Inner Belt of southwest Japan, yields abundant insect fossil assemblages (Fujiyama, 1973, 1991; Ueda, 1991; Figure 1). Fujiyama (1973) firstly described some parts of the orders Blattodea, Parapaleoptera, Hemiptera, and Coleoptera from this Group. Subsequently, examples of the orders Mecoptera (Ueda, 1991), Odonata (Fujiyama, 1991), Diptera, Ephemeroptera, Hymenoptera, Neuroptera, Orthoptera, Plecoptera, and Trichoptera (Takahashi *et*

*al.*, 1997) were reported. More than half of specimens described consists of wings of individuals of Coleoptera and Blattodea (Fujiyama, 1991), while the occurrence of Diptera is rare and important for the Triassic (Kopylov, personal comm.).

The insect fauna of the Mine Group is closely related to those of the Mount Crosby Formation in Australia, the Madygen Formation in the Kyrgyz Republic, the Djam Djam Formation in Vietnam, and the Yan-chen Formation in Inner Mongolia (Fujiyama, 1973). Particularly, the Madygen Formation yields one of the richest Triassic insect fauna comprising 20 orders (Shcherbakov, 2008), and among them, 13 orders are common to the Mine Group.

From the Mine Group, more than 6,000 specimens have been collected since the initial discovery. Most specimens are stored at the Mine City Museum of History and Folklore, Mine City, Yamaguchi Prefecture, Japan. However, these samples remain largely to be exploited in details.

This paper focuses on the first discovery of sawflies in Triassic deposits of East Asia and sheds light on the early evolution of sawflies in this region. It provides the first example of sawflies from the pioneering genus *Madygella* found outside of the Kyrgyz Republic.



**Figure 1.** Locality of the *Madygella humioi* sp. nov. holotype (MMHF3-00091) in Mine City, Yamaguchi Prefecture, Japan.

### Geological setting and fossil locality

The Triassic Mine Group, a thick clastic sequence, is lithologically divided into three units: the Hirabara, Momonoki, and Aso formations in ascending order in the Omine area, Mine City, Yamaguchi Prefecture, southwest Japan (Katayama, 1939; Kametaka, 1999; Figure 1). The lowermost Hirabara Formation is a marine deposit with a thickness of 250–1,000 m, which consists of conglomerate, sandstone, and mudstone intercalated with coal seams containing abundant marine bivalves (Hase, 1950; Tokuyama, 1960; Kametaka, 1999). The Momonoki Formation is a nonmarine deposit with a thickness of 800–1,200 m, which is composed of conglomerate, sandstone, and mudstone intercalated with coal seams (Takahashi

and Mikami, 1975; Kametaka, 1999). The uppermost Aso Formation is a marine deposit with a thickness more than 1,500 m and is composed of sandstone and mudstone with marine bivalves (Hase, 1950; Tokuyama, 1958; Kametaka, 1999). Some bivalves (Tokuyama, 1960), ammonites (Ishibashi *et al.*, 1990), plants (Katayama, 1939; Takahashi, 1973), and vertebrates (Hasegawa and Ota, 1975; Uyeno *et al.*, 1996) have been reported from the Mine Group. The geological age of the Momonoki Formation is assigned to the Carnian stage of the Upper Triassic (Fujiyama, 1973; Kametaka, 1999).

The material reported in this paper was collected from the Momonoki Formation exposed along Route 435 at Okubata, Mine City, Yamaguchi Prefecture (Figure 1). From the same locality, Fujiyama (1991) and Ueda

(1991) previously reported some fossil insects. The specimen described in the present study (MMHF3-00091) is stored at the Mine City Museum of History and Folklore.

Abbreviations of morphological terms utilized in this paper are as follows (according to Rasnitsyn, 1969; Lara *et al.*, 2014): C, costal vein; SC, subcostal vein; Pt, pterostigma; R, radial vein; RS (1-RS, 2-RS, 3-RS, 4-RS, 5-RS), radial sector and its sections; M (1-M, 2-M, 3-M, 4-M, 5-M), medial vein and its sections; Cu (1-Cu, 2-Cu, 3-Cu, 4-Cu), cubital vein and its sections; r (1r, 2r, 3r, 4r), radial cells; rm (1rm, 2rm, 3rm), radial-medal cells; mCu (1mCu, 2mCu, 3mCu), medio-cubital cells; cua (1cua, 2cua), cubito-anal cells; r-rs (1r-rs, 2r-rs), crossveins between R and RS; r-m (1r-m, 2r-m, 3r-m), radial-medal crossveins; m-cu (1m-cu, 2m-cu), medio-cubital crossveins; cu-a (1cu-a), cubito-anal crossveins.

### Systematic description

Order Hymenoptera Linnaeus, 1758  
 Suborder Symphyta Linnaeus, 1758  
 Superfamily Xyeloidea Newman, 1834  
 Family Xyelidae Newman, 1834  
 Subfamily Madygellinae Rasnitsyn, 1969 emend.  
 Kopylov, 2014

*Type genus.*—*Madygella* Rasnitsyn, 1969.

*Diagnosis of subfamily.*—Small sawfly. Forewings 2.8–5.8 mm long according to reconstructed proportions. Costal space narrow. C thickened toward apex, sometimes almost reaching R, but never touching it, separated from Pt. SC weak, joining only C, sometimes preserved only at wing base, transformed into fold, or completely lost. Pt usually not sclerotized, veins bordering it usually thickened. R straight up to base of RS, with slight bend near point of divergence of RS, usually straight within cell 1r, less often arcuate anteriorly, without gap at base of Pt. RS1 (anterior branch of RS) lost without traces. 1r-rs not longer than 2r-rs. Cell 1r along anterior margin subequal in length to 2r. 1-RS either shorter or longer than 1-M. M+Cu arcuate (very rarely slightly S-shaped). Cell 1mCu long and narrow, at least twice as long as wide (Rasnitsyn, 1969, p. 50–51; Kopylov, 2014, p. 611).

*Generic composition.*—The subfamily consists of three genera: *Madygella* Rasnitsyn, 1969; *Samarandakia* Kopylov, 2014; and *Chubakka* Kopylov, 2014 from the Middle-Upper Triassic of Kyrgyzstan.

*Comparison.*—The subfamily Madygellinae differs from other xyelid groups in the absence of RS1 (anterior branch) branches with rare exceptions: *Xyela alberta* (Curran, 1923) and *Lydoxyela excellens* (Rasnitsyn, 1966). The Madygellinae differs from subfamilies Xyelinae and Macroxyelinae in having the long cell 2r, and (for

the majority of representatives of these two subfamilies) in the narrow costal space. It also differs from subfamily Xyelinae in having the apically thickened C and from subfamily Macroxyelinae in the absence of characteristic sclerotization at the base of the Pt.

*Remarks.*—The original diagnosis (Rasnitsyn, 1969) was considerably expanded by Kopylov (2014). We add this study as a new locality to the distribution.

Genus *Madygella* Rasnitsyn, 1969  
*Madygella* Rasnitsyn, 1969, emend. Kopylov, 2014

*Type species.*—*Madygella analoga* Rasnitsyn, 1969.

*Revised generic diagnosis.*—1-RS shorter than RS+M and usually shorter than 1-M, rarely slightly longer. Length of cell 1r plus 2r longer than that of 3r+4r, sometimes length of cell 1r plus 2r shorter than that of 3r+4r; height of cell 3r+4r almost as high as that of 2r plus Pt, sometimes height of cell 3r+4r lower than that of 2r plus Pt. RS curved at point of 2r-m joining it.

*Species composition.*—*Madygella* genus consists of five species: *M. analoga* Rasnitsyn, 1969; *M. levivenosa* Kopylov, 2014; *M. kurochkini* Kopylov, 2014; *M. aristovi* Kopylov, 2014; *M. bashkuevi* Kopylov, 2014.

*Comparison.*—The present genus differs from *Samarandakia* in 1-RS shorter than 1-M, RS curved at point of 2r-m joining it and 3r-m longer (up to 1.3 times) than 2r-m. The genus also differs from *Chubakka* in 2r-rs diverging in apical one-third of Pt, cell 1mCu weakly dilating toward apex.

*Remarks.*—Most characters of *Madygella humioi* sp. nov. match with the previous generic diagnosis. Unlike other *Madygella* species, however, *M. humioi* sp. nov. shows cell length of 1r plus 2r shorter than that of 3r+4r and cell height of 3r+4r lower than 2r plus Pt as described below. Taking these features into consideration, the generic diagnosis is slightly revised and updated here.

*Madygella humioi* Oyama and Maeda, sp. nov.


Figure 2

*Etymology.*—In honor of Humio Takahashi, director of the Mine City Museum of History and Folklore.

*Holotype* (MMHF3-00091).—A specimen of well preserved forewing without anal area from the Carnian deposits of the Momonoki Formation at Okubata, Omine area, Mine City, Yamaguchi Prefecture, southwest Japan (Figure 1). It is stored at the Mine City Museum of History and Folklore. Impression of other insects is overprinted on the surface.

*Diagnosis.*—*Madygella* species having a cell length of 1r plus 2r shorter than that of 3r+4r, and a cell height of 3r+4r lower than 2r plus Pt in a forewing.





**Figure 2.** Forewing of *Madygella humioi* sp. nov., holotype MMHF3-00091 from the Upper Triassic Mine Group. **A**, photograph of the horizontal view; **B**, line drawing based on A with abbreviated terminology of vein and cell. Scale bars, 1 mm.

**Description.**—SC is developed over the entire length, joining C basal to RS base. Costal space is extremely wide. R at the base of RS has a weak curve, slightly dilates; within cell 1r, R is straight. The Pt is not sclerotized; the veins running along the periphery of the Pt are thickened. 2r-rs is inclinal, shorter than 4-RS, 1.3 times longer than 1r-rs; the base of 2r-rs is 2.6 times closer to the apex of the Pt than to 1r-rs. 1-RS is much shorter than 1-M; RS+M is about 2 times longer than 1-RS. 2-RS is straight (2-RS joins RS+M). 5-RS is 3.1 times longer than 4-RS. 2r-rm is strongly inclined, 1.9 times longer than 2r-rs. 3r-m is probably longer than 2r-m. 1m-cu is subequal in length to 3-Cu. The free end of M and Cu are lost; 3r-m gradually passes into 5-M; 2m-cu gradually passes into 4-Cu. 3-M has a weak curve at the base of 3-M.

**Measurements.**—Forewings impression length, 4.5 mm; width, 2.3 mm.

**Comparison.**—The newly described species differs from subfamily Madygellinae in its extremely wide cos-


tal space.

It is similar to the type species *Madygella analoga* and *M. levivenosa* in developing SC over the entire wing length. However, *M. humioi* sp. nov. differs from the type species in non-sclerotized Pt, strongly inclined 2r-m, same length of both 3-Cu and 1m-cu, absence of a free end of M with bend and slightly curved 3-M. *Madygella humioi* sp. nov. differs from *M. levivenosa* in inclinal 2r-rs, strongly inclined 2r-m, same length of both 3-Cu and 1m-cu and distinctly curved basal 3-M.

The newly described species is also similar to *Madygella bashkuevi* in strongly inclined 2r-m and curved 3-M. Strong 2r-m and curved 3-M are peculiar characteristic features to *M. bashkuevi* and *M. humioi* sp. nov. On the other hand, the new species is clearly distinct from *M. bashkuevi* in SC joining C basal to RS base and subequal 2r-rs and 1r-rs.

**Table 1.** Comparison of forewing characters in selected *Madygella* species. –, undescribed.

Characters/Taxa	<i>Madygella humioi</i> (this study)	<i>M. analoga</i> (Rasnitsyn, 1969; Kopylov, 2014)	<i>M. aristovi</i> (Kopylov, 2014)
Length (mm)	4.4	4.9	3.7
Width (mm)	2.3	1.4	–
Length/Width	1.9	3.5	–
Sc	joins C basal to RS base and is probably developed over the entire length	developed over the entire length, joining C slightly distal to RS base	transformed into a fold
R (at base of RS)	a weak curve slightly dilates	a rather small curve, slightly dilates	straight, slightly dilates
R (within 1r)	weakly arcuate	weakly arcuate	weakly arcuate
Pt	not sclerotized the vein is thickened	sclerotized	(sclerotized)
length of 1r plus 2r : 3r+4r	1r plus 2r < 3r+4r	1r plus 2r > 3r+4r	1r plus 2r > 3r+4r
height of 2r plus Pt : 3r+4r	2r plus Pt > 3r+4r	2r plus Pt < 3r+4r	(lost)
2r-rs	inclival, 2r-rs > 4-RS, 2r-rs = 1r-rs ( $\times 1.3$ )	inclival, 2r-rs > 4-RS, 2r-rs > 1r-rs ( $\times 1.8$ )	reclival, 2r-rs < 4-RS, (1r-rs lost)
2r-rs : 1r-rs (close to apex of Pt)	2.6 times	2.2 times	(1r-rs lost)
1-RS	1RS < 1-M (remarkably)	1-RS < 1-M (slightly)	1-RS < 1-M
RS+M	RS+M > 1-RS ( $\times$ about 2)	RS+M > 1-RS ( $\times 2.5$ )	(RS+M lost)
2-RS	straight (not joining)	straight	(lost)
5-RS	5RS > 4-RS ( $\times 3.1$ )	5RS > 4-RS ( $\times 3.5$ )	5RS > 4-RS ( $\times 4.5$ )
2r-m	incline (strongly), 2r-m > 2r-rs ( $\times 1.9$ )	(not incline), 2r-m = 2r-rs	(not incline), 2r-m > 2r-rs ( $\times 2.6$ )
3r-m	3r-m > 2r-m	3r-m > 2r-m ( $\times 1.2$ )	(lost)
3-Cu	3-Cu = 1m-cu (subequal)	3-Cu > 1m-cu ( $\times 1.5$ )	3-Cu ( $\times 1.6$ ) < 1m-cu
M (free end)	lost (3r-m passes into 5-M)	present	(lost)
Cu (free end)	lost (2m-cu passes into 4-Cu)	lost (2m-cu passes into 4-Cu)	present
3-M	slightly curved	straight	straight
Characters/Taxa	<i>M. bashkuevi</i> (Kopylov, 2014)	<i>M. kurochkini</i> (Kopylov, 2014)	<i>M. levivenosa</i> (Kopylov, 2014)
Length (mm)	4.2	3.2	3.4
Width (mm)	–	–	2.3
Length/Width	–	–	1.5
Sc	only at wing base	at wing base (apex of Sc is effaced in the impression)	over the entire length (costal space is crumpled and SC is tucked under R in the impression)
R (at base of RS)	a weak curve, slightly dilates	a visible curve, slightly dilates	a rather small curve, dilates
R (within 1r)	straight	arcuate	weakly arcuate, then straight
Pt	not sclerotized, the vein is not thickened to a greater extent than C or R	sclerotized	not sclerotized, the vein is thickened
length of 1r plus 2r : 3r+4r	1r plus 2r > 3r+4r	(lost)	1r plus 2r > 3r+4r
height of 2r plus Pt : 3r+4r	2r plus Pt < 3r+4r	2r plus Pt < 3r+4r	2r plus Pt < 3r+4r
2r-rs	inclival, 2r-rs < 4-RS, 2r-rs > 1r-rs ( $\times 1.9$ )	inclival, 2r-rs > 4-R, 2r-rs > 1r-rs ( $\times 1.3$ )	vertical, 2r-rs > 4-R, 2r-rs > 1r-rs ( $\times 2.0$ )
2r-rs : 1r-rs (close to apex of Pt)	1.8 times	2.3 times	1.6 times
1-RS	1-RS < 1-M (remarkably)	1-RS > 1-M	1-RS < 1-M (slightly)
RS+M	RS+M > 1-RS ( $\times 3$ )	RS+M > 1-RS ( $\times 1.6$ )	RS+M > 1-RS ( $\times 1.4$ )
2-RS	straight	arcuate	straight
5-RS	(5RS > 4-RS)	5RS > 4-RS ( $\times 3$ )	5RS > 4-RS ( $\times 3$ )
2r-m	incline (strongly), 2r-m > 2r-rs (distinctly)	(not incline), 2r-m > 2r-rs ( $\times 1.4$ )	(not incline), 2r-m > 2r-rs ( $\times 1.2$ )
3r-m	(lost)	(lost)	3r-m > 2r-m ( $\times 1.2$ )
3-Cu	(lost)	3-Cu = 1m-cu	3-Cu ( $\times 1.4$ ) < 1m-cu
M (free end)	(lost)	present	lost (3r-m passes into 5-M)
Cu (free end)	(lost)	lost (2m-cu joins 4-Cu with sharp bend)	lost (2m-cu passes into 4-Cu)
3-M	curved (strongly)	straight	straight



**Figure 3.** Insect fossils from the Mine Group not showing remarkable distortion. **A**, two partly overlapped wings of a sawfly (Wing P, Q, MMHF3-00095) from the same horizon of *Madygella humioi* sp. nov. **B**, outline of A with measurement axes and crossed-axes angle between the wings P and Q ( $67^\circ$ ). **C**, body parts of Coleoptera (MMHF3-00092) from the same horizon of *Madygella humioi* sp. nov. The body was vertically collapsed and flattened by compaction, but the planar geometry is not secondarily distorted and well preserved. **D**, measurements of B. Note aspect ratios of both wings are not disparate (3.00–3.35) even though orientations of the two wings are different. Scale bars, 5 mm.

## Discussion

The fossil records of Triassic Hymenoptera were previously found from four regions: Kyrgyz Republic (Rasnitsyn, 1964, 1969; Kopylov, 2014), Queensland (Riek, 1955; Grimaldi and Engel, 2005), South Africa (Schlüter, 2000), and Argentina (Lara *et al.*, 2014). Although East Asia has several contemporaneous fossil insect localities (Zheng *et al.*, 2018), there was no previous record of individuals belonging to the order Hymenoptera in the region. *Madygella humioi* sp. nov. is the oldest record of sawflies in East Asia, including Japan. This discovery reveals that primitive Hymenoptera (xyelid sawflies) were already widespread in five isolated realms, and further expands our knowledge regarding the early evolution and diversity of sawflies during the Triassic period.

The taxonomic study of Triassic sawflies has been

largely based on wing morphology as wings possess a higher potential for fossilization than other body parts (Grimaldi and Engel, 2005; Schlüter, 2000). However, the simple wing morphology is not always adequately diagnostic (Lara *et al.*, 2014), in part due to the potential post-burial deformation of morphological characters (Kopylov, 2014). For example, *Madygella bashkuevi* from the Middle–Upper Triassic in the Kyrgyz Republic has inclined 2r-m and curved 3-M, but these could have resulted from secondary distortion by shear stress of tectonism (Kopylov, 2014). Thus, Kopylov (2014) adopted a standardizing proportion formula to reconstruct the original wing geometry.

*Madygella humioi* sp. nov. from the Mine Group also shows inclined 2r-m and curved 3-M (see Table 1). Taking Kopylov's results into consideration, we examined the mode of secondary distortion in our samples. Many insect wings from the Mine Group are variously scat-



tered on a bedding plane without showing any preferred orientation. Therefore, the geometric disparity would be expected depending on their orientation if secondary distortion occurred. However, a close taphonomic observation reveals that most insect fossils in the Mine Group are not remarkably distorted, unlike Kyrgyz Republic samples. For example, two partly overlapped wings of a sawfly are preserved in a bedding plane of the shale (Figure 3A). Their crossed-axes angle is quite large, i.e., 67° (Figure 3B), while remarkable geometric disparity is not recognized between the two wings. Their aspect ratios are not disparate (3.00–3.35) even though orientations are different (Figure 3D). The minimum secondary distortion is also suggested by the preservational state of an intact fossil beetle (Coleoptera) found from the same horizon as *M. humioi* sp. nov. (Takahashi *et al.*, 1997, pl. 8, fig. 3C). The body of the beetle was vertically collapsed and flattened by compaction, while the planar geometry was still intact and not remarkably distorted (Figure 3C). The line of evidence suggests that the inclined 2r-m and curved 3-M shown in *M. humioi* sp. nov. can be interpreted as an essential morphological character, rather than a secondary distortion.

### Acknowledgements

The authors are highly indebted to Humio Takahashi (Mine City Museum of History and Folklore) and Kenji Shinoda (Mine City) who allowed to borrow the material for studying, and to Koji Wakita (Yamaguchi University) for his valuable comments and help in the field work. Constructive critical comments by Aleksandr P. Rasnitsyn and Dmitry Kopylov were very helpful for improvement of the draft. The authors would like to express their sincere gratitude toward Yusuke Okazaki (Kyushu University), Munetoshi Maruyama (The Kyushu University Museum), and Toshiharu Mita (Kyushu University) for their valuable suggestions and comments. We also thank Hirokazu Yukawa (Fukui Prefectural Dinosaur Museum) for allowing us to study fossil insects from the Mine Group. In addition, we would like to thank members of the Mine Fossil Museum and Mine City Museum of History and Folklore; and Syougo Yoshinaga, Yukiko Arakawa, Fumiko Inakura, and Yuko Kazitani for their support during the field work. This study was conducted with the support of The Late Professor Tatsuro Matsumoto Scholarship, and was partly supported by a JSPS KAKENHI Grant (number JP18K03827).

### References

Curran, C. H., 1923: A new genus and species of Xyelidae (Tenthredinoidea, Hymenoptera) from western Canada. *Canadian Entomol-*

*ogist*, vol. 55, p. 20.

- Fujiyama, I., 1973: Mesozoic insect fauna of East Asia, Part 1. Introduction and Upper Triassic faunas. *Bulletin of the National Science Museum*, vol. 16, p. 331–386.
- Fujiyama, I., 1991: Late Triassic insect from Mine, Yamaguchi, Japan, Part 1. Odonata. *Bulletin of the National Science Museum, Series C (Geology and Paleontology)*, vol. 17, p. 49–56.
- Grimaldi, D. and Engel, M. S., 2005: A new sawfly from the Triassic of Queensland (Hymenoptera: Xylidae). *Memoirs of the Queensland Museum*, vol. 51, p. 558.
- Hase, A., 1950: The stratigraphy of the Triassic system in the southwestern part of Yamaguchi Prefecture. *Science Reports, Department of Geology, Kyushu University*, vol. 2, p. 101–128. (in Japanese)
- Hasegawa, Y. and Ota, M., 1975: The Triassic reptilian tooth from the Hirabara Formation, Mine Group, Yamaguchi Prefecture, Japan. *Bulletin of the Akiyoshi-dai Museum of Natural History*, no. 11, p. 25–28. (in Japanese with English abstract)
- Ishibashi, T., Hashimoto, K., Nohara, M. and Yoshida, K., 1990: Triassic ammonites newly discovered in the western district of Yamaguchi Prefecture, Japan. *Journal of the Geological Society of Japan*, vol. 96, p. 771–774. (in Japanese)
- Kametaka, M., 1999: Provenance of the Upper Triassic Mine Group, Southwest Japan. *Journal of the Geological Society of Japan*, vol. 105, p. 651–667. (in Japanese)
- Katayama, M., 1939: Stratigraphical study on the Mine Series. *Journal of the Geological Society of Japan*, vol. 46, p. 127–141. (in Japanese)
- Kopylov, D. S., 2014: New sawflies of the Subfamily Madygellinae (Hymenoptera, Xyelidae) from the Middle–Upper Triassic of Kyrgyzstan. *Paleontologicheskij Zhurnal*, vol. 48, p. 610–620.
- Lara, M. B., Rasnitsyn, A. P. and Zavattieri, A. M., 2014: *Potereilloxyela menendezii* gen. et sp. nov. from the Late Triassic of Argentina: The oldest representative of Xyelidae (Hymenoptera: Symphyta) for Americas. *Paleontologicheskij Zhurnal*, vol. 48, p. 182–190.
- Linnaeus, C., 1758: Hymenoptera. In, *Systema Naturae, per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Editio decima, reformata. Tomus I*, p. 553–583. Laurentii Salvii, Holmiae (Stockholm).
- Newman, E., 1834: Attempted division of British insects into natural orders. *Entomological Magazine*, vol. 11, p. 379–430.
- Rasnitsyn, A. P., 1964: New Triassic Hymenoptera from Central Asia. *Paleontologicheskij Zhurnal*, no. 1, p. 88–96. (in Russian; original title translated)
- Rasnitsyn, A. P., 1966: New Xyelidae (Hymenoptera) from the Mesozoic of Asia. *Paleontologicheskij Zhurnal*, no. 4, p. 69–85. (in Russian; original title translated)
- Rasnitsyn, A. P., 1969: The origin and evolution of the lower Hymenoptera. *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR*, vol. 123, p. 1–196. (in Russian; original title translated)
- Riek, E. F., 1955: Fossil insects from the Triassic beds at Mt. Crosby, Queensland. *Australian Journal of Zoology*, vol. 3, p. 654–691.
- Schlüter, T., 2000: *Moltenia riei* n. gen., n. sp. (Hymenoptera: Xyelidae?), a tentative sawfly from the Molteno Formation (Upper Triassic), South Africa. *Paläontologische Zeitschrift*, vol. 74, p. 75–78.
- Shcherbakov, D. E., 2008: Madygen, Triassic Lagerstätte number one, before and after Sharov. *Alavesia*, vol. 2, p. 113–124.
- Takahashi, E., 1973: Stratigraphic distribution of the Mine-type and the Toyara-type floras. *Science Report of Yamaguchi University*, vol. 20, p. 15–28. (in Japanese with English abstract)
- Takahashi, E. and Mikami, T., 1975: Triassic. *Geology of Yamaguchi*



- Prefecture*, no. 3, p. 93–108. Yamaguchi Museum Press, Japan. (in Japanese)
- Takahashi, H., Ishida, H., Nohara, M., Doi, E. and Taniguchi, S., 1997: Occurrence of insect fossils from the Late Triassic Mine Group. *Bulletin of the Mine City Museum, Yamaguchi Prefecture, Japan*, no. 13, p. 1–27. (in Japanese)
- Tokuyama, A., 1958: Die obertriadische Molasse im Mine-Gebiet Westjapans 1. Teil Stratigraphie und Werdegang. *Journal of the Geological Society of Japan*, vol. 64, p. 454–643. (in Japanese with German abstract)
- Tokuyama, A., 1960: On the pelecypod fauna of the Late Triassic Hirabara Formation in West Japan. *Japanese Journal of Geology and Geography*, vol. 31, p. 201–217.
- Ueda, K., 1991: A Triassic fossil of scorpion fly from Mine, Japan. *Bulletin of the Kitakyushu Museum of National History*, vol. 10, p. 99–103.
- Uyeno, T., Takahashi, H., Doi, E. and Naito, G., 1996: Triassic fishes collected in Mine City and Sanyocho in Yamaguchi Prefecture. *Bulletin of the Mine City Museum, Yamaguchi Prefecture, Japan*, no. 12, p. 1–27. (in Japanese)
- Zheng, D., Chang, S. C., Wang, H., Fang, Y., Wang, J., Feng, C., Xie, G., Jarzembowski, E. A., Zhang, H. and Wang, B., 2018: Middle–Late Triassic insect radiation revealed by diverse fossils and isotopic ages from China. *Science Advance*, vol. 4, p. 1–7.

### Authors contribution

Nozomu Oyama initiated the study and was primarily responsible for taxonomic aspects.

Haruyoshi Maeda revised the manuscript critically for important intellectual content. All authors contributed to the writing of the paper.