

## **A New Genus and Species of Microphysidae (Hemiptera: Heteroptera) with Long Labium in Late Cretaceous Iwaki Amber from Futaba Group of Iwaki City, Fukushima Prefecture, Japan**

Authors: Aiba, Hiroaki, Souma, Jun, and Inose, Hiroaki

Source: Paleontological Research, 29(1) : 44-53

Published By: The Palaeontological Society of Japan

URL: <https://doi.org/10.2517/prpsj.240017>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A new genus and species of Microphysidae (Hemiptera: Heteroptera) with long labium in Late Cretaceous Iwaki amber from Futaba Group of Iwaki City, Fukushima Prefecture, Japan

Hiroaki AIBA<sup>1\*</sup>, Jun SOUMA<sup>2</sup>, Hiroaki INOSE<sup>3</sup>
<sup>1</sup>Institute for Educational Practice Studies, 1-12-16, Kichijoji-kitamachi, Musashino-shi, Tokyo 180-0001, Japan

<sup>2</sup>Shirakami Research Center for Environmental Sciences, Faculty of Agriculture and Life Science, Hirosaki University, 3 Bunkyo-cho, Hirosaki-shi, Aomori 036-8152, Japan


<sup>3</sup>Fukushima Museum, 1-25, Jyoto Machi, Aizuwakamatsu City, Fukushima 965-0807, Japan

[Received August 22, 2024; Revised manuscript accepted December 25, 2024; Published online March 17, 2025]

## ABSTRACT

The Iwaki insect-bearing amber from Fukushima Prefecture, northeastern Honshu, Japan, is considered from the Late Cretaceous Coniacian age. The insect-bearing amber localities from that period are few worldwide. Here, we report *Iwakia longilabiata* gen. et sp. nov., a new fossil genus and species of the infraorder Cimicomorpha Leston, Pendergrast and Southwood, 1954 (Hemiptera: Heteroptera) from the Futaba Group of Iwaki City. Although the preservation is incomplete, this fossil is provisionally placed in the family Microphysidae Dohrn, 1859 based on the general consistency of the diagnostic characters of the family. The fossil is distinguishable from any genus of Cimicomorpha including Microphysidae because of its long labium and brachypterous female.

Additionally, the fossil shares the female diagnostic characteristics of the Palearctic and Nearctic taxa, suggesting that it is an early stage in the evolution of sexual dimorphism of the family Microphysidae. The evolution of sexual dimorphism in Microphysidae may have already occurred in East Asia during the Late Cretaceous Coniacian age. This fossil may be the first discovery from Asia and the oldest record of the family Microphysidae.



**ZooBank registration:** urn:lsid:zoobank.org:pub:CC4DAF02-7182-4D9E-871F-A3B81BFFBDFD

**Keywords:** fossil insect, Iwaki amber, labium, Late Cretaceous, Microphysidae

## Introduction

The Iwaki insect-bearing amber (hereinafter Iwaki amber) which was excavated from Fukushima Prefecture in northeastern Honshu, Japan, is considered to have originated from the Late Cretaceous Coniacian age (Saegusa and Tomida, 2011), and few fossils from this

period are available worldwide (Rasnitsyn and Quicke, 2002). In addition, little is known about the occurrence of insect-bearing amber in East Asia, except for Burmese amber (Cenomanian) (Seyfullah *et al.*, 2018; Ross, 2023). Therefore, the study of insect fossils from the Iwaki amber is important from both paleobiogeographic and geological perspectives and will provide valuable data for unraveling the evolutionary history of insects. However, little paleontological research has been conducted on Iwaki amber. Aiba and Inose (2024) described a new species, *Archaeromma chisatoi*, which is a small group of microscopic parasitic wasps from Iwaki amber. This

\* Corresponding author: Hiroaki AIBA (erimomisa@gmail.com)



This article is licensed under a Creative Commons [Attribution 4.0 International] license.  
© 2025 The Authors.

discovery represents the first description of a new species from Iwaki amber.

The true bug family Microphysidae Dohrn, 1859 (Hemiptera: Heteroptera: Cimicomorpha: Microphysoidea), little pirate bugs or minute bugs, is one of the smallest heteropteran families, approximately 30 species in five extant genera that have not been well studied because of their small size (1.5 to 3.0 mm), cryptic habitats, and limited distribution (Schuh and Slater, 1995; Péricart, 1996; Yasunaga, 2001; Schuh and Weirauch, 2020). This taxon was first treated as a separate family by Fieber (1861) but classified by Reuter (1884) as a subfamily Microphysina in Anthocoridae. However, Microphysidae has been treated as an independent family belonging to the superfamily Microphysoidea of the infraorder Cimicomorpha by molecular phylogenetic analyses in recent studies (Schuh *et al.*, 2009; Weirauch *et al.*, 2019; Ye *et al.*, 2022). This family comprises approximately 30 species belonging to the following five extant genera: *Chinaola* Blatchley, 1928; *Ciorulla* Péricart, 1974; *Loricula* Curtis, 1833; *Mallochiola* Bergroth, 1925; and *Nabidomorpha* Poppius, 1910 (Yasunaga and Yamada, 2017). The genera *Ciorulla* and *Loricula* were originally distributed in the Palearctic region. They had sexual dimorphism, in which males resemble the Anthocoridae and possess rather long, delicate forewings with well-developed membranes. However, females lack membrane of forewing and possess coleopteroid, macropterous, or staphylinoid forewings and sometimes rounded and posteriorly widened abdomen (Yasunaga and Yamada, 2017). However, six species are distributed in the Nearctic Region. Among these, only two monotypic genera, *Chinaola* and *Mallochiola*, are considered native. The other four species are immigrants from the Palearctic Region (Blinn, 2011). Genera *Chinaola* and *Mallochiola* do not exhibit sexual dimorphism. Females do not differ noticeably from males (Wheeler, 1992).

To date, eleven fossil species of Microphysidae have been known, ten of which are from the Eocene Baltic and Ukraine Rovno amber (Putshkov and Popov, 2003; Popov, 2004; Popov, 2006; Popov *et al.*, 2008; Popov and Herczek, 2009). Only one from the Mesozoic is from the Late Cretaceous (Campanian) Canadian amber (Mckellar and Engel, 2011). All these fossils are from Europe and North America, and none have yet been found in Asia.

Recently, we examined true bug fossils from Iwaki amber. Thus, we identified it as a new genus and species of the infraorder Cimicomorpha Leston, Pendergrast and Southwood, 1954 and provisionally assigned it to the family Microphysidae. This fossil may represent the first discovery from Asia and may be the oldest record of the family Microphysidae. This study is the second paleontological description of a new species from Iwaki amber.

## Geological setting

The Futaba Group is mainly distributed along the Pacific coast of northeastern Japan, from Naraha-machi, Futaba-gun, Iwaki City in Fukushima Prefecture. It is subdivided into the Ashizawa, Kasamatsu, and Tamayama formations (Figure 1). The Tamayama Formation is subdivided into the Kohisagawa and Irimazawa members (Ando *et al.*, 1995; Kubo *et al.*, 2002). A fossil insect dealt with in this study was discovered in amber from the Kohisagawa Member of the Tamayama Formation (Aiba *et al.*, 2023; Aiba and Inose, 2024).

The Irimazawa Member is a shallow marine deposit above the Kohisagawa Member that yielded marine invertebrate and vertebrate fossils dated from the late Coniacian to early Santonian (Obata and Suzuki, 1969). The lower Kohisagawa Member of the Kasamatsu Formation has not yielded any age-determining fossils. However, Matsumoto *et al.* (1982) considered it to be upper Coniacian based on marine faunal fossils from the overlying stratum, the Irimazawa Member, and the underlying stratum, the Ashizawa Formation. Recently, Hasegawa *et al.* (2020) suggested that the uppermost part of the Tamayama Formation extends into the early Campanian based on U-Pb ages. Therefore, it is reasonable to consider the age of the Kohisagawa Member of the Tamayama Formation, which bears fossil amber, as late Coniacian (86–87 Ma) (Gradstein *et al.*, 2004; Saegusa and Tomida, 2011).


## Material and methods

Chisato Suzuki discovered this fossil on August 23, 1986. The studied specimen was deposited at the Fukushima Museum, Fukushima Prefecture, Japan, under repository number FM-N202200012.

The amber was thinly sliced, and the surfaces were polished using an abrasive. Observations were performed using a Leica M205C stereomicroscope (Leica Corporation, Wetzlar, Germany). Thin, light-transparent materials were observed at low magnification using an Olympus CX43 optical microscope (Olympus Corporation, Tokyo, Japan). Photographs and measurements were obtained using a Leica MC170HD Macroscopic with Leica Application Suite Version 4.1.3 (Leica Corporation) and an Olympus TG4 optical microscope (Olympus Corporation). The morphological terminology was according to Yasunaga and Yamada (2017), Yasunaga *et al.* (2020), and Rédei (2024).

## Systematic paleontology

Order Hemiptera Linnaeus, 1758  
Suborder Heteroptera Latreille, 1810



**Figure 1.** Geographic and stratigraphic location of the Iwaki amber from Futaba Group. Fossil location denoted with a red star. Stratigraphic position denoted with a blue star. Stratigraphic chart modified from Ando *et al.* (1995).

Infraorder Cimicomorpha Leston, Pendergrast and Southwood, 1954

Superfamily Microphysoidea Dohrn, 1859

Family Microphysidae? Dohrn, 1859

***Iwakia* gen. nov.**

[New Japanese name: Kuchinaga-futagatakamemushizoku]

ZooBank lsid: urn:lsid:zoobank.org:act:C26BB7DA-C44D-4F18-862E-0C6FB82808E6

**Type species.**—*Iwakia longilabiata* sp. nov.

**Etymology.**—The genus epithet is Iwaki City, referring to the fossil locality; the gender is feminine.

**Diagnosis.**—Female brachypterous morph. Head prognathous, elongated, triangular, flattened, 1.5 times as long as maximum width across compound eyes. Ocelli absent in female. Labium extremely long, reaching beyond apex of abdomen, 1.3 times as long as body length. Forewing covering anterior half of abdomen, without membrane, reaching posterior margin of abdominal tergite V, with claval suture. Tarsi 2-segmented.

***Iwakia longilabiata* sp. nov.**

[New Japanese name: Kuchinaga-futagatakamemushi]

Figures 2–4

ZooBank lsid: urn:lsid:zoobank.org:act:E55E8C93-2E67-409C-AFFA-4B8DDF67DEF3

**Diagnosis.**—As for the genus.

**Material.**—Holotype: FM-N202200012; Incompletely preserved dorsal (Figure 2A), ventral (Figure 2B), and lateral views (Figure 2C).


**Type locality and horizon.**—The specimen was collected from amber in the muddy sandstone of the Upper Cretaceous (Coniacian) Kohisagawa Member of the Tamayama Formation of the Futaba Group, exposed at an outcrop quarry (37°10'51"N, 140°57'54"E) located in the upstream Suetsugi River in Suetsugi, Hisanohama-machi, Iwaki City, Fukushima Prefecture, Northeast Japan.

**Etymology.**—The species epithet is the Latin adjective “longus” (meaning long) and “labiatus” (meaning labiate), referring to the long labium.

**Description of holotype.**—*Brachypterous Female*: Body oblong, flattened, weakly sclerotized. Lateral margins parallel to each other (Figures 2A, 2B, 2C, and 4A).

**Head:** Prognathous, elongated, triangular, flattened, 1.5 times as long as maximum width across compound eyes, not punctate. Clypeus darker, reaching beyond apex of mandibular plate. Buccula visible in dorsal view, widened laterally, pale, bearing fine setae (Figures 3A, B). Compound eyes conspicuously protruding laterally (Figure 2A). Ocelli absent. Labrum very short, visible in






**Figure 2.** *Iwakia longilabiata* gen. et sp. nov., holotype (FM-N202200012). **A**, habitus, dorsal view; **B**, habitus, ventral view; **C**, habitus, lateral view. Abbreviations: lb II – IV, labial segment II – IV; st, stylets.

lateral and ventral views (Figure 3B). Antenna (Figure 3C) filiform, relatively short, 1.7 times as long as head length; segment I somewhat swollen, not bearing setae, not reaching beyond apex of head; segment II longest, 1.7 times as long as segment I, bearing short setae; segment III as long as segment II; segment IV as long as segment I. Labium extremely long, semitransparent (Figures 2B–F, 4), reaching beyond apex of abdomen, 1.3 times as long as body length; segment I very short (boundary between segments I and II not visible but presumed

behind labrum); segment II 2.3 times as long as segment I; segment III 2.3 times as long as segment II, gradually widened posteriorly, constricted in posterior quarter, bearing short setae (Figure 4B); segment IV longest among labial segments, 2.9 times as long as segment III, gradually narrowed posteriorly slightly swollen at apex, serrate in lateral margin, bearing short setae (Figures 3E, F, 4C). Stylet darker, completely visible.

Thorax: Pronotum campanulate, 3.7 times as wide as long, uniformly covered with minute punctures; anterior






**Figure 3.** *Iwakia longilabiata* gen. et sp. nov., holotype (FM-N202200012). **A**, details of the head (dorsal view); **B**, details of the head (lateral view); **C**, details of the antennae; **D**, details of the labium and stylets; **E**, enlargement of the boundary between labial segment III and IV; **F**, enlargement of the apex of labial segment IV; **G**, details of the fore legs; **H**, details of the right mid leg; **I**, details of the right hind leg; **J**, details of the left hind leg; **K**, details of the forewing. Abbreviations: a I – a IV, antennal segment I – IV; bu, buccula; cc, claval commissure; cl, claw; cs, claval suture; cv, costal vein; cy, clypeus; hf, hind leg femur; ht, hind leg tibia; lb II – IV, labial segment II – IV; mf, medial fracture; mp, mandibular plate; st, stylets; t I – t II, tarsus segments I – II.

margin slightly concave; posterior margin deeply concave. Humeral angles preserved on left side and deformed on right side; apex acute on left side and shrunk on right side (Figures 2A, 4A). Calli absent. Scutellum triangular, coarsely punctate in posterior half. Forewing cover-

ing anterior half of abdomen, semi-transparent, sparsely punctate (Figures 3K, 4A); puncture with short setae, denser towards apex; claval suture and medial fracture slightly visible (Figure 3K), costal fracture and membranous portion absent. Cuneus unclear. Forewing mem-



**Figure 4.** Line drawing of *Iwakia longilabiata* gen. et sp. nov. (FM-N202200012). **A**, habitus, dorsal view; **B**, labial segment I – III; **C**, labial segment IV, XY line is continuous. Abbreviations: a I – a IV, antennal segment I – IV; lb I – IV, labial segment I – IV.

brane absent or invisible. Fore and middle legs same in length (Figure 3G, H); hind leg twice as long as fore leg (Figure 3I, J); femora slightly swollen; tibiae with short

spines in fore and middle legs, and long spines in hind leg; tarsi 2-segmented; claws simple, without pulvillus.

Abdomen: Abdomen oblong-oval, semitransparent

coarsely punctate; puncture bearing setae. Lateral margin straight, parallel to each other. Abdominal tergites III–VI subequal in length; tergite VII longest among abdominal tergite; tergite VIII and IX 0.5 times as long as tergite VII; connexivum undeveloped. Small spiracles visible on dorsal and ventral sides because of semitransparent abdomen (Figure 2A, B). Detailed structures of terminalia not clearly visible, making presence or absence of ovipositor uncertain.

Measurements (in mm): Body length 3.38; length of head 0.83; maximum width across compound eyes of head 0.54; diameter of compound eye 0.12; lengths of antennal segments I to IV 0.22, 0.38, 0.32, 0.26, respectively; length of labrum 0.12; total length of labium 4.34; length of labial segments I to IV 0.19?, 0.41?, 0.94, and 2.80, respectively; maximum width of labial segment IV 0.05; length of pronotum 0.51; anterior pronotal width 0.45; maximum width of pronotum across humeri 1.14; median length of scutellum 0.37; maximum width of scutellum 0.57; maximum length of forewing 1.37; maximum width of forewing 0.55; length of profemur 0.61; length of protibia 0.55; length of protarsus 0.18; length of mesofemur 0.45; length of mesotibia 0.37; length of mesotarsus 0.16; length of metafemur 1.10; length of metatibia 0.37; length of metatarsus 0.16; lengths of abdominal segments III to IX 0.23, 0.22, 0.24, 0.26, 0.33, 0.16, and 0.11, respectively; maximum width of abdominal segments III to IX 1.10, 1.11, 1.09, 1.06, 1.02, 0.66, and 0.24, respectively.

## Discussion

### Morphological comparison

The fossil species described above resembles the family Aradidae (Pentatomomorpha: Aradoidea), and its long mandibular and maxillary stylets and flattened body are consistent with diagnostic characteristics of Aradidae. However, this fossil is clearly distinguishable from Aradidae in the following characteristics (e.g. Larivière and Laroche, 2006; Schuh and Weirauch, 2020): dorsal structure semitransparent (granular or rugose in Aradidae); labium extremely long, reaching beyond apex of abdomen (usually short and stout, not reaching abdominal sternite in Aradidae); connexivum not developed (broadly developed in Aradidae); claw without pulvilli (with pulvilli in most members of Aradidae); and abdomen with microtrichia (without microtrichia in Aradidae).

Additionally, the fossil is distinguishable from other species of the infraorder Pentatomomorpha based on the following features (Schuh and Weirauch, 2020): antennae fragelliform (never fragelliform in Pentatomomorpha); labial segment I short and not developed (well-developed in Pentatomomorpha). We consider the fossil to belong to the infraorder Cimicomorpha because of its small,

wispy body, prognathous head, labium anteriorly inserted on head, and flagellated antennae (Schuh and Weirauch, 2020).

Moreover, the fossil is identified as belonging to the Superfamily Microphysoidea based on its tiny body, short labial segment I, 2-segmented tarsi, and simple claw without pulvilli (Schuh and Weirauch, 2020). The Superfamily Microphysoidea includes two families, Microphysidae and Joppeicidae Reuter, 1910 (Weirauch *et al.*, 2019), and the Joppeicidae consists of only one species, *Joppeicus paradoxus* Puton, 1881 (Schuh and Weirauch, 2020). The fossil is similar to *J. paradoxus* in that the head morphology is long and triangular, and the buccula is visible in the dorsal view and widened laterally (Schuh and Štys, 1991). However, the fossil is clearly distinguishable from *J. paradoxus* based on many features, such as the absence of hind wing, ocelli, and membranes. Schuh and Weirauch (2020) summarized the characteristics of the Family Microphysidae as follows: body length 1.5–3.0 mm; many species similar to small cimicoids; head weakly prognathous; labium 4-segmented; labial segment I shortest among labial segments; and tarsi 2-segmented. Furthermore, they mentioned that the genera of Microphysidae from the Palaearctic Region exhibit sexual dimorphism, and females lack membrane of forewing. All these characteristics are consistent with those of the fossil, except the slightly larger size. Yasunaga and Yamada (2017) also characterized the genus *Loricula*, the type genus of Microphysidae, as follows: females lack ocelli; antennal segment II usually longer than segment IV; female pronotum trapezoid or campanulate; and posterior margin of pronotum concave. These characteristics are also consistent with fossil features. Based on the general consistency of the diagnostic characters, this fossil is considered to belong to Microphysidae or a closely related new family. However, because this fossil is incompletely preserved, we could not examine all the detailed morphological characteristics and avoided establishing a new family here. Therefore, we provisionally identify the fossil as a member of Microphysidae.

Microphysidae comprises five extant genera (*Ciorulla*, *Loricula*, *Myrmedobia*, *Chinaola*, and *Mallochiola*) and two fossil genera (*Tytthophysa* Popov and Herczek, 2009 and *Popovophysa* McKellar and Engel, 2011) in the Holarctic Region (Schuh and Weirauch, 2020). This fossil is easily distinguished from any of those genera in the extremely long labium 1.3 times as long as body. In addition, females of the genera *Ciorulla* and *Loricula* from the Palaearctic region, which are sexually dimorphic, are clearly distinguishable from the studied fossil in that they possess smaller bodies (approximately 1.5 mm) and a broad and thickened abdomen (Yasunaga and Yamada, 2017). Furthermore, this fossil specimen is distinguished



from both genera in the brachypterous female. The genera *Chinaola* and *Mallochiola* from the Nearctic Region do not exhibit sexual dimorphism; therefore, females possess the same oblong body with well-developed membrane of forewing as males. The ground plan of the Nearctic species is the same as that of the other known fossil species (Popov, 2006; Schuh and Weirauch, 2020). Moreover, the studied fossil is distinctly differentiated from the other fossils by the absence of ocelli and forewing membranes.

In conclusion, this fossil specimen is not consistent with any of the five extant and two fossil genera of Microphysidae, and *Iwakia* gen. nov. was described to accommodate the fossil specimen in the above section.

### Considerations from fossil morphology

The fossil characteristics of the absence of ocelli and membranes in females are consistent with the diagnostic characteristics of females of the Palaearctic genera (Yasunaga and Yamada, 2017). However, the oblong body and attributes of the forewing (presence of clavus) in females are consistent with the diagnostic characteristics of the Nearctic genera (Wheeler, 1992; Blinn, 2011).

The morphological characteristics of the studied fossil, which share the female diagnostic characters of the Palaearctic and Nearctic taxa, probably represent an early stage in the evolution of sexual dimorphism in the family Microphysidae and suggest that the evolution of sexual dimorphism in this family had already occurred in East Asia during the Late Cretaceous Coniacian. On the other hand, the following four fossil microphysid species, which are macropterous females and unlikely to possess sexual dimorphism, have been reported from the Middle Eocene Baltic amber of the western Palaearctic Region: *Loricula (Eocenophysa) damzeni* Popov, 2004, *L. (Myrmericula) ocellata* Popov, 2006, *L. (M.) samlandi* Popov, 2006, and *Tytthophysa sylwiae* Popov and Herczek, 2009, suggesting that microphysid species possessing sexual dimorphism may not have been distributed in the western Palaearctic Region, at least during the Middle Eocene period.

The studied fossil has an extremely long labium compared to the other species of Microphysidae. The length of the labium is insufficient to judge the feeding habits of the fossil species, but the fact that the labium is longer than the body length may indicate that the food of this fossil is different from that of the other species of the family.

According to the previous studies (Slater, 1975; Henry, 2012), heteropterous wing forms are divided into aptery, microptery, staphylinoidy, coleoptery, brachyptery, submacroptery, and macroptery. The studied fossil resembles macroptery or submacroptery in that it has the claval suture, making corium and clavus are differentiated.

However, this fossil can be clearly distinguished from macroptery and submacroptery by the absence of membranes. Consequently, the fossil is inconsistent with any of the seven heteropterous wing forms but is provisionally treated as brachyptery in the present study based on the absence of a forewing membrane and the fully exposed abdominal tergite VI.

In this study, we provisionally identified the fossil as a member of Microphysidae. However, the studied fossil has unique morphological features, such as an extremely long labium and a forewing with a claval suture and without a membrane, which are partly inconsistent with the diagnostic characters of the family. Therefore, the identification of the family of this fossil may be changed by further observation of the morphological characteristics with new methods or by the discovery of additional specimens.

### Acknowledgments

We thank Chisato Suzuki for discovering the specimen. Additionally, we are grateful to the editor and the two anonymous reviewers of Paleontological Research for their critical comments to the manuscript.

### References

- Aiba, H. and Inose, H., 2024: A new false fairy wasp (Hymenoptera: Mymarommatoidea: Mymaromatidae) in Late Cretaceous Iwaki amber from Futaba Group of Iwaki City, Fukushima Prefecture, Japan. *Paleontological Research*, vol. 28, p. 337–344.
- Aiba, H., Suzuki, C. and Inose, H., 2023: Amber fossils from the Tamayama Formation, Upper Cretaceous, Futaba Group, Iwaki City, Fukushima Prefecture, Japan, and their significance. *Bulletin Fukushima Museum*, vol. 37, p. 19–28. (in Japanese with English abstract)
- Ando, H., Seishi, M., Oshima, M. and Matsumaru, T., 1995: Fluvial-shallow marine depositional systems of the Futaba Group (Upper Cretaceous)—Depositional facies and sequences—. *Journal of Geography*, vol. 104, p. 284–303. (in Japanese with English abstract)
- Bergroth, E., 1925: On the “annectant bugs” of Messrs. McAtee and Malloch. *Bulletin of the Brooklyn Entomological Society*, vol. 20, p. 159–164.
- Blatchley, W. S., 1928: Two new Anthocoridae and a new microphysid from Florida (Heteroptera). *Entomological News*, vol. 39, p. 85–88.
- Blinn, R. L., 2011: New state records for *Mallochiola gagates* (McAtee & Malloch, 1924), a little-known microphysid in eastern North America (Hemiptera: Heteroptera: Microphysidae). *Zootaxa*, vol. 3009, p. 67–68.
- Curtis, J., 1833: Characters of some undescribed genera and species indicated in the guide to an arrangement of British insects. *Entomologist's Monthly Magazine*, vol. 1, p. 186–199.
- Dohrn, A., 1859: *Catalogus Hemipterorum*, 112 p. Herrcke & Lebeling, Stettin.
- Fieber, F. X., 1861: *Die europäischen Hemiptera. Halbflügler (Rhynchocha Heteroptera)*, 112 p. Gerold's Sohn, Wien.
- Gradstein, F. M., Ogg, J. G. and Smith, A. G., 2004: *A Geologic Time*

- Scale 2004, 589 p. Cambridge University Press, Cambridge.
- Hasegawa, R., Isozaki, Y. and Tsutsumi, Y., 2020: Fragmented ancient fore-arc basin: Cretaceous–Paleogene sandstones sporadically found in Kanto and southern Tohoku, Japan, as eastern extensions of the Izumi Group. *Journal of Geography*, vol. 129, p. 49–70. (in Japanese with English abstract)
- Henry, T. J., 2012: Revision of the plant bug genus *Tytthus* (Hemiptera, Heteroptera, Miridae, Phyllinae). *ZooKeys*, vol. 220, p. 1–114.
- Kubo, K., Yanagisawa, Y., Toshimitsu, S., Banno, Y., Kaneko, N., Yoshioka, T. *et al.*, 2002: *Geology of the Kawamae and Ide District*, 1:50,000, 136 p. Geological Survey of Japan, Tokyo. (in Japanese)
- Larivière, M. C. and Laroche, A., 2006: An overview of flat bug genera (Hemiptera, Aradidae) from New Zealand, with considerations of faunal diversification and affinities. In: Rabitsch, W. ed., *Hug the Bug: For Love of True Bugs: Festschrift zum 70. Geburtstag von Ernst Heiss. Denisia*, vol. 9, p. 181–214. Plöchl-Druck, Freistadt.
- Latreille, P. A., 1810: *Considérations Générales sur l'Orde Naturel des Animaux Composant les Classes des Crustacés, des Arachnides, et des Insectes*, 444 p. Frédéric Schoell, Paris.
- Leston, D., Pendergrast, J. G. and Southwood, T. R. E., 1954: Classification of the terrestrial Heteroptera (Geocorisae). *Nature*, vol. 174, p. 91–92.
- Linnaeus, C., 1758: *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, locis. Editio decima, reformata, (1758–1759)*, 823 p. Impensis Direct, Laurentii Salvii, Holmiae (Stockholm).
- Matsumoto, T., Obata, I., Tashiro, M., Ohta, Y., Tamura, M., Matsukawa, M. *et al.*, 1982: Correlation of marine and non-marine formations in the Cretaceous of Japan. *Fossils (Palaeontological Society of Japan)*, no. 31, p. 1–26. (in Japanese with English abstract)
- McKellar, R. C. and Engel, M. S., 2011: First Mesozoic Microphysidae (Hemiptera): a new genus and species in Late Cretaceous amber from Canada. *Canadian Entomologist*, vol. 143, p. 349–357.
- Obata, I. and Suzuki, T., 1969: Additional note on the upper limit of the Cretaceous Futaba Group. *Journal of the Geological Society of Japan*, vol. 75, p. 443–445. (in Japanese)
- Péricart, J., 1974: Une espèce nouvelle de Microphysidae représentant un genre nouveau, et une espèce nouvelle d'Anthocoridae (Rhynchota, Hemiptera). *Bulletin de la Société Entomologique de France*, vol. 79, p. 253–257.
- Péricart, J., 1996: Family Microphysidae Dohrn, 1859 — flower bugs, little pirate bugs, minute bugs. In: Aukema, B. and Rieger, C. H. eds., *Catalogue of the Heteroptera of the Palaearctic Region. vol. 2. Cimicomorpha I*, p. 108–140, The Netherlands Entomological Society, Amsterdam.
- Popov, Y. A., 2004: New microphysids (Heteroptera: Cimicomorpha, Microphysidae) from Baltic amber and taxonomy of this family. *Prace Museum Ziemi*, vol. 47, p. 97–107.
- Popov, Y. A., 2006: New microphysids (Heteroptera, Cimicomorpha, Microphysidae) from Baltic Eocene amber from the collection of Ernst Heiss. *Denisia*, vol. 19, p. 571–579.
- Popov, Y. A. and Herczek, A., 2009: A new peculiar minute bug (Hemiptera, Heteroptera, Cimicomorpha, Microphysidae) from the Eocene Baltic amber. *Denisia*, vol. 26, p. 151–154.
- Popov, Y. A., Herczek, A. and Kanja, I., 2008: One more microphysid (Heteroptera: Cimicomorpha, Microphysidae) from the Eocene Baltic amber. *Genus*, vol. 19, p. 611–617.
- Poppus, B., 1910: 12. Hemiptera, 4. Miridae, Anthocoridae, Teratophyllidae, Microphyidae and Nabidae. In: Sjöstedt, Y. ed., *Wissenschaftliche Ergebnisse der Swedischen Zoologischen Expedition nach dem Kilimandjaro, dem Meru und den Umgebenden Masaisteppen Deutsch-Ostafrikas 1905–1906, 2 Band*, p. 57–58. Palmquist AB., Stockholm.
- Puton, J. B. A., 1881: Enumeration des Hemipteres recoltés en Syrie par M. Abeille de Perrin avec la description des especes nouvelles. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, vol. 6, p. 119–129.
- Putshkov, P. V. and Popov, Y. A., 2003: The first find Microphysidae from Ukrainian (Rovno) amber (Heteroptera, Cimicomorpha). *Annals of the Upper Silesian Museum in Bytom, Entomology*, vol. 12, p. 81–85.
- Rasnitsyn, A. P. and Quicke, D. L. J., 2002: *History of Insects*, 517 p. Kluwer Academic Publisher, Dordrecht.
- Rédei, D., 2024: A re-assessment of *Palaeotanyrhina* (Hemiptera: Heteroptera) elucidates the phylogeny of Leptopodoidea. *Insect Systematics and Diversity*, vol. 8, doi:10.1093/isd/ixae030.
- Reuter, O. M., 1884: *Monographia Anthocoridarum Orbis terrestris*, 204 p. Forgotten Book, London.
- Reuter, O. M., 1910: Neue beiträge zur Phylogenie und systematik der Miriden, nebst einleitenden bemerkungen über die phylogenie der Heteropteren-Familien. *Acta Societatis Scientiarum Fennicae*, vol. 37, p. 1–172.
- Ross, A. J., 2023: *Burmese (Myanmar) amber taxa, on-line supplement v.2023.1*. [online]. [Cited 22 July 2024]. Available from: [https://www.nms.ac.uk/media/1164144/burmese-amber-taxa-v2021\\_1.pdf](https://www.nms.ac.uk/media/1164144/burmese-amber-taxa-v2021_1.pdf).
- Saegusa, H. and Tomida, Y., 2011: Titanosauriform teeth from the Cretaceous of Japan. In: Kellner, A. W. and Tomida, Y. eds., *Proceedings of the Third Gondwanan Dinosaur Symposium, Anais da Academia Brasileira de Ciencias Anais de Academia Brasileira de Ciencias*, vol. 83, p. 247–265. Academia Brasileira de Ciências Rio de Janeiro, Brasil.
- Schuh, R. T. and Slater, J. A., 1995: *True Bugs of the World (Hemiptera: Heteroptera): Classification and Natural History*, 416 p. Cornell University Press, Ithaca.
- Schuh, R. T. and Štys, P., 1991: Phylogenetic analysis of cimicomorphan family relationships (Heteroptera). *Journal of the New York Entomological Society*, vol. 99, p. 298–350.
- Schuh, R. T. and Weirauch, C., 2020: *True Bugs of the World (Hemiptera: Heteroptera): Classification and Natural History (Second Edition)*, 768 p. Siri Scientific Press, Manchester.
- Schuh, R. T., Weirauch, C. and Wheeler, W. C., 2009: Phylogenetic relationships within the Cimicomorpha (Hemiptera: Heteroptera): a total-evidence analysis. *Systematic Entomology*, vol. 34, p. 15–48.
- Seyfullah, L. J., Beimforde, C., Dal Corso, J., Perrichot, V., Rikkinen, J. and Schmidt, A. R., 2018: Production and preservation of resins — past and present. *Biological Reviews*, vol. 93, p. 1684–1714.
- Slater, J. A., 1975: On the biology and zoogeography of Australian Lygaeidae (Hemiptera: Heteroptera) with special reference to the southwest fauna. *Journal of the Australian Entomological Society*, vol. 14, p. 47–64.
- Weirauch, C., Schuh, R. T., Cassis, G. and Wheeler, W. C., 2019: Revisiting habitat and lifestyle transitions in Heteroptera (Insecta: Hemiptera): insights from a combined morphological and molecular phylogeny. *Cladistics*, vol. 35, p. 67–105.
- Yasunaga, T., 2001: Family Microphysidae Dohrn, microphysids. In: Yasunaga, T., Takai, M. and Kawasawa, T. eds., *A Field Guide to Japanese Bugs II—Terrestrial Heteropterans*, 350 p. Zenkoku Noson Kyoiku Kyokai, Tokyo. (in Japanese)
- Yasunaga, T. and Yamada, K., 2017: Review of the microphysid genus *Loricula* Curtis in Japan (Hemiptera: Heteroptera: Microphysidae), with descriptions of three new species. *Journal of Natural History*, vol. 51, p. 1209–1227.
- Yasunaga, T., Yamada, K. and Ohno, T., 2020: A new species of the

genus *Loricula* Curtis from central Honshu, Japan (Heteroptera: Microphysidae). *Zootaxa*, vol. 4759, p. 98–106.

Ye, F., Kment, P., Rédei, D., Luo, J. Y., Wang, Y. H., Kuechler, S. M. *et al.*, 2022: Diversification of the phytophagous lineages of true bugs (Insecta: Hemiptera: Heteroptera) shortly after that of the flowering plants. *Cladistics*, vol. 38, p. 403–428.

### Author contributions

H. A. initiated the study, drafted the manuscript, and compiled all the figures. J. S. revised the manuscript and provided taxonomic input. H. I. revised the manuscript and provided geological input. All authors contributed to the writing of the manuscript.