

A New Species of Roldana (Asteraceae) from Southern Oaxaca

Authors: Quedensley, Taylor Sultan, and Villaseñor, José Luis

Source: Lundellia, 2009(12) : 28-30

Published By: The Plant Resources Center, The University of Texas at Austin

URL: https://doi.org/10.25224/1097-993X-12.1.28

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A NEW SPECIES OF *ROLDANA* (ASTERACEAE) FROM SOUTHERN OAXACA

Taylor Sultan Quedensley¹ and José Luis Villaseñor²

¹Plant Biology Graduate Program, The University of Texas, 1 University Station A6720, Austin, Texas 78712 ²Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado Postal 70-367, 04510 México, D.F., México

Abstract: Roldana uxordecora is described from Oaxaca, Mexico (Municipio de San Mateo Río Hondo). The genus *Roldana* is common in the montane regions of the state of Oaxaca (22 species), and this addition to *Roldana* adds to a list of endemic species, now five, already collected from this montane region.

Resumen: Se describen **Roldana uxordecora** de Oaxaca, México (Municipio de *San* Mateo Río Hondo). El genero *Roldana* esta frecuente en los regiones montañosas del estado de Oaxaca (22 especies), y ese adición a *Roldana* agrega a una lista de especies endémicas, ahora cinco, ya colectado de ese región montañosa.

Keywords: Asteraceae, Senecioneae, Mexico, montane forests, endemism.

During field work in 2009, the authors discovered an interesting species of *Roldana* La Llave (Asteraceae: Senecioneae) that did not appear to match any of the currently described species in the genus. After close examination of the material collected, it was apparent that it represents a new species of *Roldana* and it is described below.

Roldana uxordecora T.S. Quedensley & J.L. Villaseñor, sp. nov. (Fig. 1).

TYPE: **MEXICO. OAXACA**: Sierra Madre del Sur, Mpio. San Mateo Río Hondo, along Highway 175 to San Pachutla from Miahuatlán, 1 km northeast of San Jose del Pacifico; rocky slopes and secondary pineoak forest with *Pinus* sp., *Baccharis heterophylla, Holodiscus* sp., and *Chiranthodendron pentadactylon*; 2482 m; 16°10'32.2"N, 96°30'24.0"W, 14 Jan 2009, *T. Sultan Quedensley, J. L. Villaseñor, T. Buxton, & O. Hinojosa 7050* (HOLOTYPE: MEXU!; ISO-TYPES: BIGU!, CAS!, F!, MICH!, MO!, NY!, TEX!, UC!, US!)

Roldana schaffneri (Sch. Bip. ex Klatt) H. Rob. & Brettell similis sed differt foliis subter floccoso-tomentosis, petiolis ad basim late expansis, caulibus juvenibus floccoso-tomentosis, et flosculis radii semper tres flosculis disci semper quinque.

SHRUBS erect, 1.5-2.5 m tall; stems woody, striate; young stems floccose-tomentose, old growth glabrous. LEAVES simple, pinnately-veined, petiolate; petioles 7-12 cm long, floccose, greatly expanded at base; blades ovate to oblanceolate, 11-15 cm long, 7-8 cm wide, weakly crenate with callose denticles; acute to acuminate at apex, cordate to truncate at base, dark green and glabrous above, densely floccose-tomentose below. HEADS numerous, arranged in terminal rounded cymes 5-8 cm wide, ultimate peduncles 7-11 mm, floccose. PHYLLARIES uniseriate, five, 8-10 mm long, 2-3 mm wide; brown in the center with scarious margins, apex floccose. RAY FLORETS three per head, 10-11 mm long, ligules yellow, 6-7 mm long. DISC FLORETS five per head, yellow, 10-12 mm long; corolla tube 7-8 mm long. CYPSELAE glabrous, brown, 1-2 mm long; pappus of numerous white bristles; 7–8 mm long.

Roldana is distributed throughout the montane regions of Mexico and Central America. It is a medium-sized Neotropical genus of suffrutescent herbs, shrubs, and small trees currently including forty-eight species (Funston, 2008) and a typical taxon in montane forests of southern Mexico, becoming most prevalent at elevations

Downloaded From: https://complete.bioone.org/journals/Lundellia on 16 Jul 2025 Terms of Use: https://complete.bioone.org/terms-of-use

LUNDELLIA 12:28-30. 2009



FIG. 1. A. Flowering branch. B. Abaxial leaf surface and expanded petiole base. C. Head. D. Ray floret. E. Disc floret. All images were drawn from isotypes and photos of living material in the field (T. *S. Quedensley et al. 7050*) by Cindy Quedensley.

between 2000 and 3000 meters. Currently, in the state of Oaxaca there are twenty-two recorded species of this genus, four of which are endemic to that state.

Although *Roldana uxordecora* is morphologically distinct in several features within the genus, it keys out to *R. schaffneri* (Sch. Bip. ex Klatt) H. Rob. & Brettell in all published treatments (Robinson & Brettell, 1974; Turner, 2005; Funston, 2008). Of the three previously-described species of *Roldana* with pinnately-veined leaves that occur in Oaxaca, *R. uxordecora* differs from *R. schaffneri* in having a dense floccose-tomentum on the underside of the leaves, on the young stems, and on the flowering branches. It differs from *R. barba-johannis* (DC.) H. Rob. & Brettell in having five phyllaries instead of ten to thirteen. *Roldana lineolata* (DC.) H. Rob. & Brettell has lobed, sinuate leaves with petioles that are not expanded at the base and with blades that are only lightly tomentose abaxially.

KEY TO THE PINNATELY-VEINED SPECIES OF ROLDANA THAT OCCUR IN OAXACA

| 1. | Leaves abaxially pubescent only along veins R. schaffneri |
|----|---|
| 1. | Leaves abaxially pubescent on entire leaf surface |
| | 2. Phyllaries 5; petioles expanded at base of leaves R. uxordecora |
| | 2. Phyllaries 7-13; petioles not expanded at base of leaves |
| | 3. Leaves deeply lobed, sinus depth to one-third of way to midrib R. lineolata |
| | 3. Leaves sinuate to weakly lobed, sinus depth never approaching midrib R. barba-johannis |

Presently, Roldana uxordecora is only known from the type locality and more collections along the Sierra Madre del Sur, from southern Oaxaca would more accurately illustrate the species' distribution. Prior to this description, 22 species of Roldana have been collected in the montane regions of Oaxaca, representing 45% of the known species in the genus. Of these 22 species, four are strictly endemic to Oaxaca, and three species are found additionally to only one neighboring state. Endemism in the Asteraceae has been used to establish conservation priorities for Mexico, and Oaxaca is one of the most endemic-rich states in regards to this diverse flowering plant family (Villaseñor et al., 1998).

The Latin epithet means "beautiful wife," and the authors take pride in naming the species after their wives whom have supported their research with unwavering devotion.

ACKNOWLEDGMENTS

Funding for field work from The University of Texas at Austin was provided by the Plant Biology Graduate Program. Extramural funding was provided by Sterling Keeley of the University of Hawaii-Manoa. We thank Timothy Buxton (Cabrillo College) and Oscar Hinojosa (UNAM) for assistance in the field; Tom Wendt and the University of Texas at Austin Plant Resources Center for technical support in the herbarium; and Guy Nesom for the Latin diagnosis. We thank Beryl Simpson and Michael Gruenstaeudl (University of Texas at Austin), Michele Funston (Missouri Botanical Garden), and Robert Kowal (University of Wisconsin-Madison) for providing comments on the manuscript. Lastly, we are extremely grateful to Cindy Quedensley for the fine illustrations provided.

LITERATURE CITED

- Funston, A. M. 2008. Taxonomic Revision of *Roldana* (Asteraceae: Senecioneae), a genus of the Southwestern U.S.A., Mexico, and Central America. Ann. Missouri Bot. Gard. 95: 282–337.
- Robinson, H. and R. Brettell. 1974. Studies in the Senecioneae (Asteraceae). Phytologia 27: 402–439.
- Turner, B. L. 2005. A recension of the Mexican species of *Roldana* (Asteraceae: Senecioneae). Phytologia 87: 204–263.
- Villaseñor, J. L., G. Ibarra, and D. Ocaña. 1998. Strategies for the conservation of Asteraceae in Mexico. Conservation Biol. 12: 1066–1075.