

Slate-crowned Antpitta *Grallaricula nana* in the western Andes of Ecuador

Authors: Freile, Juan F., Loaiza, José M., and Molina, Pablo

Source: Bulletin of the British Ornithologists' Club, 138(2) : 140-142

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v138i2.2018.a9>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Slate-crowned Antpitta *Grallaricula nana* in the western Andes of Ecuador

by Juan F. Freile, José M. Loaiza & Pablo Molina

Received 6 March 2018; revised 26 April 2018; published 22 June 2018

<http://zoobank.org/urn:lsid:zoobank.org:pub:6B045FD6-E8C7-4366-8815-940EA6802478>


Slate-crowned Antpitta *Grallaricula nana* has a widespread Andean distribution, with up to eight subspecies recognised (Krabbe & Schulenberg 2003, Donegan 2008). It ranges from the coastal mountains of Venezuela, west and south along the three Andean ranges in Colombia, and in the eastern Andes of Ecuador to northernmost Peru, with an isolated population in the tepuis of south-east Venezuela and adjacent Guyana (Greeney 2018). The taxonomy of *G. nana* is not fully resolved, as indicated by ongoing discussion as to the validity of *G. nana cumanensis* (with *G. n. pariae*) and *G. n. kukenamensis* as separate species (Donegan 2008, Remsen *et al.* 2017, Greeney 2018).

Between southern Colombia and northern Peru, *G. nana* has been recorded to date only on the Amazonian slopes of the eastern Andes (Krabbe *et al.* 2017). These populations have been assigned to *G. n. occidentalis* by some authorities (Donegan 2008, Krabbe *et al.* 2017, Greeney 2018) but to the nominate subspecies by Krabbe & Schulenberg (2003) who synonymised *G. n. occidentalis* with it (see also Ridgely & Greenfield 2001).

During field work on 3–7 February 2017 at El Corte (00°49'13.28"N, 78°05'40.63"W; 2,500–2,620 m elevation), Carchi province, north-west Ecuador, we discovered *G. nana* in natural secondary forests on very steep slopes adjacent to small streams. We located five territories in very dense and tangled understorey, with patchy bamboo and a broken canopy. Despite intensive efforts, we were unable to mist-net or photograph adults, but we recorded the vocalisations of two different individuals (Fig. 1; archived on xeno-canto.org; XC404700, 404702, 404706, 404707).

The geographically closest population to north-west Carchi is in the eastern Andes of Ecuador, which corresponds to *G. n. occidentalis* (*fide* Donegan 2008, Krabbe *et al.* 2017, Greeney 2018). The *nudo de Pasto* of southern Colombia—where the three Andean ranges of Colombia merge into a single plateau—reaches c.4,000 m at high passes, and isolates the eastern and western slopes of the Andes. However, several taxa (i.e. Golden-faced Tyrannulet *Zimmerius chrysops*, Sulphur-bellied Tyrannulet *Mecocerculus minor*, Rufous-tailed Tyrant *Knipolegus poecilurus*, Bicoloured Antvireo *Dysithamnus occidentalis*) that occur on the eastern Andean slopes of southern Colombia and northern Ecuador 'spill over' the Andes into north-west Ecuador, north of the dry Mira Valley (Krabbe *et al.* 1998, Krabbe 2008), in some cases ranging into south-west Colombia as well. This pattern is probably shared by *G. nana* given the lack of records from the western Andes of Imbabura and ornithologically well-known Pichincha province, south of the Mira Valley. To date, there is a single, undocumented record of *G. nana* from the West Andes of southern Colombia, at Laguna de Cumbal (00°57'02.5"N, 77°49'05.7"W), dpto. Nariño, just 30 km north of the El Corte region, in 2015 (J. Zuleta; <https://ebird.org/view/checklist/S25166185>). This record, however, requires confirmation as the locality appears unsuitable by elevation and habitat for *G. nana*. Other localities on the Pacific slope of the West Andes of Colombia are further north—beyond the Patía Valley, from dpto. Cauca northwards (Donegan 2008, Greeney 2018).

A



B




Figure 1. Sonogram of four phrases of song (A); sonogram and oscillogram of a single phrase (B) of Slate-crowned Antpitta *Grallaricula nana*, El Corte, Carchi province, north-west Ecuador, February 2017. Recorded by JFF using a Zoom H4N Pro (Zoom North America, NY) with built-in microphone; sonogram and oscillograms prepared by PM using Raven Pro (Cornell Lab of Ornithology, Ithaca, NY).

Donegan (2010) suggested that the population in the West Andes of north-central Colombia and the headwaters of the Magdalena Valley, where the West and Central Andes meet, might represent an unnamed subspecies, due to its vocal characters. Curiously, Donegan (2008) included two localities from eastern dpto. Nariño, close to the Ecuadorian border on the east-facing slope of the Andes (Llorente and La Victoria; Paynter 1997), as being occupied by this putative subspecies. Our recordings suggest that the vocalisations of *G. nana* in north-west Carchi are similar to those of widespread *G. n. occidentalis*. Number of notes, song duration and pace are within or close to the range of variation in this subspecies (Donegan 2008), suggesting a closer relationship with cis-Andean populations of *G. n. occidentalis* and not to the West Andes population in northern and central Colombia (Donegan 2008). It remains to be determined if *G. n. occidentalis* ranges north to the Patía Valley onto the Pacific slopes of dpto. Cauca and northwards, and whether it meets the population in West Andes of Colombia (*vide* Donegan 2008).

We tentatively assign the population in north-west Carchi to *G. n. occidentalis*, but suggest the need for additional exploration of the area—including adjacent dpto. Nariño of southern Colombia—to obtain further audio-recordings and to collect specimens, considering the apparent geographic isolation from both *G. n. occidentalis* and the West Andes ‘vocal type’ of Donegan (2008, 2010). A better understanding of the distribution of *G. nana* subspecies in south-west Colombia will also aid in understanding the geographic variation in song postulated by Donegan (2008).

Acknowledgements

Field work formed part of a project searching for Turquoise-throated Puffleg *Eriocnemis godini*, funded by the American Bird Conservancy. Thanks to Wendy Willis and Daniel Lebbin for their assistance; to Jordan Karubian for his permanent support and advice; and to Lenin Quintanchala and Rubén Quintanchala for hospitality and company in the field. Harold Greeney provided an insightful revision of the submitted manuscript.

References:

- Donegan, T. M. 2008. Geographical variation in Slate-crowned Antpitta *Grallaricula nana*, with two new subspecies from Colombia and Venezuela. *Bull. Brit. Orn. Cl.* 128: 150–178.
- Donegan, T. M. 2010. *Grallaricula nana*: subspecies and species limits. www.xeno-canto.org/article/34 (accessed 10 January 2018).
- Greeney, H. F. 2018. *Antpittas and gnateaters*. Christopher Helm, London.
- Krabbe, N. 2008. Arid valleys as dispersal barriers to high-Andean forest birds in Ecuador. *Cotinga* 29: 28–30.
- Krabbe, N. & Schulenberg, T. S. 2003. Family Formicariidae (ground antbirds). Pp. 682–731 in del Hoyo, J., Elliott, A. & Christie, D. A. (eds.) *Handbook of the birds of the world*, vol. 8. Lynx Edicions, Barcelona.
- Krabbe, N., Skov, F., Fjeldså, J. & Petersen, I. K. 1998. *Avian diversity in the Ecuadorian Andes. An atlas of distribution of Andean forest birds and conservation priorities*. DIVA Tech. Rep. 4. Centre for Research on the Cultural and Biological Diversity of Andean Rainforests (DIVA), Rønde.
- Krabbe, N. K., Schulenberg, T. S., de Juana, E. & Kirwan, G. M. 2017. Slate-crowned Antpitta (*Grallaricula nana*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from www.hbw.com/node/56919 on 22 December 2017).
- Paynter, R. A. 1997. *Ornithological gazetteer of Colombia*. Second edn. Mus. Comp. Zool., Cambridge, MA.
- Remsen, J. V., Areta, J. I., Cadena, C. D., Jaramillo, A., Nores, M., Pacheco, J. F., Pérez-Emán, J., Robbins, M. B., Stiles, F. G., Stotz, D. F. & Zimmer, K. J. 2017. A classification of the bird species of South America. www.museum.lsu.edu/~Remsen/SACCBaseline.html (accessed 22 December 2017).
- Ridgely, R. S. & Greenfield, P. J. 2001. *The birds of Ecuador*, vol. 1. Cornell Univ. Press, Ithaca, NY.
- Addresses:* Juan F. Freile, Comité Ecuatoriano de Registros Ornitológicos, Pasaje El Moro E4-216 y Norberto Salazar, Quito, Ecuador, e-mail: jfreileo@yahoo.com. José M. Loaiza, Fundación ECOMINGA Red de Bosques Protectores Amenazados, 12 de Noviembre 270 y Luis A. Martínez, Baños, Ecuador. Pablo Molina, San Antonio de Ibarra, Calle Ramón Teanga, Ecuador.