

## **First record of Lesson's Seedeater *Sporophila bouvronides* in Costa Rica**

Authors: Gatgens-García, Johan, Chaves-Sánchez, Alexander, and Sandoval, Luis

Source: Bulletin of the British Ornithologists' Club, 142(4) : 526-528

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v142i4.2022.a11>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First record of Lesson's Seedeater *Sporophila bouvronides* in Costa Rica

by Johan Gatgens-García, Alexander Chaves-Sánchez & Luis Sandoval

Received 27 July 2022; revised 5 October 2022; published 7 December 2022

<http://zoobank.org/urn:lsid:zoobank.org:pub:4BB7BB88-8A9A-4D91-8EBD-A98E7638A6DC>

**SUMMARY.**—We report the first record of Lesson's Seedeater *Sporophila bouvronides*, a South American species, in Costa Rica. It was seen at Tortuguero, in the lowlands of north-east Costa Rica in June 2022. The record is perhaps explained by the passage of a tropical storm that moved north-west from off northern South America.

The genus *Sporophila* comprises c.40 species, distributed from Texas in the USA to Argentina (Machado & Silveira 2010, Rising 2011). Most species live in grasslands, but a few inhabit forest interiors (Ridgely & Tudor 1989, Areta *et al.* 2013). Due to urban development and the expansion of agricultural monocultures (e.g., pineapple, banana or soybean), the areas inhabited by these species (especially grasslands) have been declining, which may explain changes in their distributions (Silva 1999).

Although the majority of *Sporophila* species are sedentary, approximately 25% perform migrations between the breeding and non-breeding ranges (Rising 2011). One species with long-distance movements is Lesson's Seedeater *Sporophila bouvronides*, which breeds in north-east Colombia, northern Venezuela, Trinidad & Tobago and the Guianas, and migrates to western Amazonia, reaching northern Bolivia, western Brazil and eastern Peru during the non-breeding season (Ridgely & Tudor 1989, Rising 2011). It inhabits shrubby savannas, forest edges, cultivated areas and grasslands near water (Rising 2011). Clearly, the species is capable of moving long distances, although the only previous evidence of vagrancy is a few records in eastern Panama (Angehr & Dean 2010). Here we present the first report in Costa Rica.

## First record in Costa Rica

The observation was made at Tortuguero (10°32'23.9"N, 83°30'06.1"W), Pococí, Limón province, in the lowlands of north-east Costa Rica (Fig. 1) on 25 June 2022, on a private




Figure 1. Male Lesson's Seedeater *Sporophila bouvronides*, Tortuguero, Limón province, Costa Rica, June, with a male Variable Seedeater *Sporophila corvina* (\*) (Johan Gatgens-García and Alexander Chaves-Sánchez)




Figure 2. Distribution of Lesson's Seedeater *Sporophila bouvronides* in South America, showing the closest report to Costa Rica (blue dot) and the new record in Tortuguero, Limón, Costa Rica (red dot). Breeding range in green and non-breeding range in yellow.

property with an open area and lawn, and involved an adult male (Fig. 1) with a flock of Variable Seedeaters *Sporophila corvina*, near the beach and foraging under a tropical almond *Terminalia catappa*. It immediately drew our attention, because the two species' plumages are very different, and the bird remained alert and wary. The observation lasted c.7 minutes, during which time it stayed with the Variable Seedeaters, then flew c.10 m from the other seedeaters, before flying further and disappearing.

We separated the Lesson's Seedeater from other *Sporophila* in Costa Rica as follows. Male Morelet's Seedeater *S. moreletti* has a white collar and white wingbars, a white spot below the eye, and the lower underparts are off-white or cream. Males of the Caribbean subspecies of Variable Seedeater are all black, whilst the Pacific subspecies (not recorded or expected in the area) has a white collar. Male Lined Seedeater *S. lineola*, an accidental species in Costa Rica (Garrigues & Dean 2014), has a white median crown-stripe (Rising 2011).

## Discussion

There are no previous reports of Lesson's Seedeater in Costa Rica. The closest records are in Darién, Panama, where an eBird (<https://ebird.org/checklist/S56064460>) report in May 2019 also involved a bird in the company of Variable Seedeaters. Our record is c.615 km north-west of this and 760 km from the closest area where the species breeds (Fig. 2). The date of our sighting coincides loosely with the species' return to the breeding range (from April but mainly May or June; Ridgely & Tudor 1989, Rising 2011). With the passage of tropical storm Bonnie that began on 25 June in the southern Caribbean, then moved east to

west, and on 1 July impacted Barra del Colorado north of Costa Rica with gusts of wind of 75 km/h (<https://cap-sources.s3.amazonaws.com/cr-imm-es/2022-07-01-17-56-34.xml>; <https://www.nhc.noaa.gov/archive/2022/al02/al022022.discus.018.shtml>), the bird could have been blown off course to eastern Costa Rica. Potentially similar vagrancy has occurred in a closely related species, with a male Lined Seedeater found on the Pacific slope of Costa Rica in October 2013 (Obando-Calderón *et al.* 2013).

#### References:

- Angehr, G. R. & Dean, R. 2010. *The birds of Panama a field guide*. Zona Tropical Publications, Ithaca, NY.
- Areta, J. I., Bodrati, A., Thom, G., Rupp, A. E., Velazquez, M., Holzmann, I., Carrano, I. & Zimmermann, C. E. 2013. Natural history, distribution, and conservation of two nomadic *Sporophila* seedeaters specializing on bamboo in the Atlantic Forest. *Condor* 115: 237–252.
- Garrigues, R. & Dean, R. 2014. *The birds of Costa Rica a field guide*. Second edn. Zona Tropical, San José.
- Machado, É. & Silveira, L. F. 2010. Geographical and seasonal distributions of the seedeaters *Sporophila bouvreuil* and *Sporophila pileata* (Aves: Emberizidae). *Pap. Avuls. Zool.* 50: 517–533.
- Obando-Calderón, G., Chaves-Campos, J., Garrigues, R., Montoya, M. & Ramírez, O. 2013. Listado oficial de las aves de Costa Rica. *Zeledonia* 17(2): 44–59.
- Ridgely, R. S. & Tudor, G. 1989. *The birds of South America*, vol. 1. Univ. of Texas Press, Austin.
- Rising, J. D. 2011. Family Emberizidae (buntings and New World sparrows). Pp. 428–683 in del Hoyo, J., Elliott, A. & Christie, D. A. (eds.) *Handbook of the birds of the world*, vol. 16. Lynx Edicions, Barcelona.
- Silva, J. M. C. 1999. Seasonal movements and conservation of seedeaters of the genus *Sporophila* in South America. *Stud. Avian Biol.* 19: 272–280.

*Addresses:* Johan Gatgens-García, Programa de Posgrado en Biología, Sistema de Estudios de Posgrado, Universidad de Costa Rica, 11501-2060, San José, Costa Rica, e-mail:jgatgensgarcia12@gmail.com. Alexander Chaves-Sánchez, Universidad de Costa Rica, Sede del Caribe, Limón, Costa Rica. Luis Sandoval, Laboratorio de Ecología Urbana y Comunicación Animal, Escuela de Biología, Centro de Investigación en Biodiversidad y Ecología Tropical (CIBET), Universidad de Costa Rica, San José, Costa Rica.