

Factors affecting changes in waterfowl populations in eutrophic wetlands in the Finnish lake district

Authors: Kauppinen, Jukka, and Väänänen, Veli-Matti

Source: Wildlife Biology, 5(2) : 73-81

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/wlb.1999.011>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Factors affecting changes in waterfowl populations in eutrophic wetlands in the Finnish lake district

Jukka Kauppinen & Veli-Matti Väänänen

Kauppinen, J. & Väänänen, V-M. 1999: Factors affecting changes in waterfowl populations in eutrophic wetlands in the Finnish lake district. - *Wildl. Biol.* 5: 73-81.

From the 1970s to the 1990s, changes in breeding waterfowl populations were monitored at 26 well-vegetated lakes in southern Finland. Extreme total population levels were found to differ by 40%, but between successive years fluctuations did not exceed 20%, the most unstable populations being garganey *Anas querquedula*, tufted duck *Aythya fuligula*, coot *Fulica atra* and pochard *Aythya ferina*. Multiple regression analysis was used to determine the effects of weather factors during wintering, spring and brood seasons on the variation in breeding waterfowl populations. The following weather factors were statistically significant in explaining fluctuations observed in four species; in garganey: spring temperature; in pintail *Anas acuta*: brood season temperature, winter severity in France and spring temperature; in goldeneye *Bucephala clangula*: winter severity in the Baltic and spring temperature; and in mallard *Anas platyrhynchos*: brood season temperature of the previous year. Mallard, teal *Anas crecca*, wigeon *Anas penelope* and goldeneye were shot in larger numbers than would be expected considering their respective proportions of the breeding waterfowl community. Hunting pressure on mallard, tufted duck and goldeneye was clearly higher in our study area than in other parts of Finland. Our results indicate that the level of hunting pressure in our study area may affect population densities of mallard.

Key words: hunting, waterfowl populations, weather factors, wetlands

Jukka Kauppinen, Kuopio Natural History Museum, Myhkyrinkatu 22, FIN-70100 Kuopio, Finland

Veli-Matti Väänänen, Department of Applied Zoology, P.O. Box 27 (Latokartanonkaari 5), FIN-00014 University of Helsinki, Finland - e-mail: veli-matti.vaananen@helsinki.fi

Received 14 December 1998, accepted 23 February 1999

Associate Editor: Hannu Pöysä

During the 20th century, many waterfowl species of eutrophic lakes with a southern distribution have expanded into northern Europe (e.g. Kalela 1946, Onno 1965, Yarker & Atkinson-Willes 1971, von Haartman 1973). The expansion has been attributed to both climatic and environmental changes (e.g. Kalela 1949, von Haartman 1973). Although eutrophication of lakes has produced good habitats for many waterfowl species, the process may also have unfavourable

effects on waterfowl habitats (e.g. von Haartman 1975, Nilsson 1985).

In North America, many researchers have investigated the relationship between waterfowl and environmental factors in pothole areas of northern prairies and parklands. Habitats in these areas have been shown to be unstable primarily due to climatic conditions: water level fluctuates widely both between years and during the breeding season, as do the

amount of open water and mosaics of vegetation. The major reason for population and community changes has been found to be alteration and destruction of breeding habitats (e.g. Stoudt 1971, Trauger & Stoudt 1978, Boyd 1981, Leitch & Kaminski 1985, Bethke 1993). In contrast, waterfowl habitats in northern Europe may be fairly stable between years (Nummi & Pöysä 1993), and only slow changes in waterfowl communities occur (Pöysä 1989, Kauppinen 1995). Although it is evident that habitat factors can greatly affect the structure and long-term changes in waterfowl communities, other factors may play a more crucial role in the short-term dynamics of these communities (Pöysä 1984, Kauppinen 1995).

It has been suggested that weather factors may regulate short-term fluctuations of some waterfowl populations in northern Europe (e.g. von Haartman 1945, 1957, Grenquist 1965, Hildén 1966, Nilsson 1979, 1984), and these changes may also affect populations in the long term. For example, it is known that harsh weather in wintering areas can influence populations either directly by causing winter mortality or indirectly by lowering breeding success in the following season (Boyd 1964, Nilsson 1979, 1984, Andersen-Harild 1981). Spring temperatures may also prolong or shorten the spring migration period. Moreover, weather conditions in spring and summer may influence breeding success and, hence, also the breeding population in the following year.

Apart from weather factors, hunting may be the most important mortality factor affecting fledged waterfowl populations (e.g. Stoudt & Cornwell 1976, Rogers, Nichols, Martin, Kimball & Posphala 1979). Caithness, Williams & Nichols (1991) reported age-specific mortality in ducks; more young than old individuals are shot. Other studies have shown that the survival rates of ducks vary between years, and in theory, the increase in hunting mortality appears to be compensated for by a decrease in non-hunting mortality (Anderson & Burnham 1976, Rogers et al. 1979). However, few studies have focused on the role of hunting mortality in regulating waterfowl populations, especially waterfowl populations in Europe.

In this study, we investigate recent changes in the breeding waterfowl populations in the wetlands of the Finnish lake district, and examine the role of weather factors and hunting in explaining changes in waterfowl populations.

Study area, material and methods

Data were gathered from richly vegetated lakes in northern Savo, (63°N, 27°E) at the northern border of the Finnish lake district. Breeding waterfowl populations were censused in 26 lakes covering a total area of 22.4 km², once at the beginning of the 1970s (1972-1975), and then again at the beginning of the 1980s (1983-1985). Eleven of these lakes, in total covering 9.46 km², were monitored for waterfowl yearly during 1984-1995.

The area of the 11 lakes ranged within 18-226 ha, with most lakes comprising <100 ha (mean = 94.6 ha). Emergent vegetation covered 25-90% (mean = 54.1%) of the total lake area. Stands of sedge *Carex* sp. and horsetail *Equisetum fluviatile* were dominant, though reed *Phragmites australis* and bullrush *Schoenoplectus lacustris* were sometimes also found. Maximum water depth varied within 1-8 m, with depths of 1-2 m predominating (mean = 2.8 m). Transparency of the water usually varied between 0.5 and 1.5 m, and the pH-value of the water ranged within 6.0-7.0. Both the characteristics of the lake types examined and the methods used to measure the environmental parameters of the lakes are described in more detail in Kauppinen & Väisänen (1993).

Censuses of breeding waterfowl were conducted using the 'round count' method (Kauppinen 1980, 1983, Kauppinen, Koskimies & Väisänen 1991). Lakes were censused twice during the breeding season, i.e. 10-15 and 25-30 May. The criteria used to determine whether pairs were breeding or not are described in Kauppinen & Väisänen (1993). All censuses were performed by the authors, with the same lake always being monitored by the same person. When calculating coefficient of variation there may be more inherent stochastic variation in small samples than in large samples and therefore the results may only reflect natural noise (Helle & Mönkkönen 1986, see also Wiens 1981). To minimise this problem, coefficient of variation of densities (CV%) was not calculated for less abundant species, i.e. for species with mean densities of <1.5 pairs/km².

Wing and ringing data were used to indicate the hunting pressure on waterfowl in our study area. Species, age and sex composition of the annual hunting bag were determined from wing data using criteria described by Salminen (1983). Wing data accounting for ca 30% of the annual bag were obtained annually from hunters in a restricted area in the municipality of Maaninka in 1985-1989. In the area

Table 1. Indices of winter severity expressed as maximum ice cover in the Baltic Sea (%), mean temperatures in December - January in northern France, and mean temperatures (°C) in northern Savo in spring (pair season) and at the beginning of summer (brood season) during 1983-1995.

Winter severity	1983/84	84/85	85/86	86/87	87/88	88/89	89/90	90/91	91/92	92/93	93/94	94/95	
Baltic sea ice cover	45	85	80	96	35	12	16	29	16	17	49	16	
Mean temperature in northern France	4.1	0.9	4.6	1.5	5.5	5.3	4.9	3.5	2.8	5.4	5.9	5.5	
Mean temperatures in northern Savo	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
15 April - 24 May	8.0	7.9	3.9	6.8	5.3	5.2	8.7	8.4	4.3	4.9	9.8	5.9	4.8
1 June - 10 July	15.1	14.4	14.0	17.1	14.1	17.2	17.1	14.5	14.7	15.3	11.8	14.7	

where the wings were collected, the breeding population of tufted duck was smaller than that observed for the whole study area, whereas populations of shoveler *Anas clypeata* and pintail were somewhat larger. Waterfowl ringing data covering 1970-1991 were obtained from the Finnish Museum of Natural History. Ringing data which were used to compare the recovery rate of young ducks (see hunting mortality, Table 6) in northern Savo with those in other parts of Finland were available for the years 1980-1991.

Maximum ice cover in the Baltic Sea, which is an important wintering area for mallard, tufted duck, goldeneye and coot, and the mean temperatures for December and January in northern France, which is an important wintering area for all other species, were used as indices of winter severity in the wintering areas of the waterfowl studied (Finnish Marine Research Institute, Météo-France). Mean temperatures in spring, i.e. 15 April - 24 May, were used as indices for determining the timing of spring, and mean temperatures in the beginning of the summer,

Table 2. Numbers of pairs of 12 waterfowl species occurring in 26 lakes covering a total area of 22.5 km² in southern Finland in the beginning of 1970s and 1980s. Differences in pair numbers was tested using G-test; level of significance: *** P < 0.001, ** = P < 0.01, * = P < 0.05.

Species	1970s	1980s	Level of significance
Great crested grebe	36	44	
Slavonian grebe	15	15	
Wigeon	59	108	**
Teal	125	193	***
Mallard	94	78	
Pintail	41	51	
Shoveler	24	48	*
Garganey	32	27	
Pochard	26	24	
Tufted duck	132	197	***
Goldeneye	39	61	*
Coot	11	42	***
Total	634	893	***

i.e. between 1 June and 10 July, were used to indicate the conditions of the brood season (Kuopio Meteorological Station). Indices describing weather conditions are presented in Table 1. Winter severity indices and spring and summer temperature indices were uncorrelated (in all cases $R_s < 0.463$, $P > 0.05$), as were the indices indicating winter severity in the Baltic sea area and in France ($R_s = -0.366$, $P > 0.05$).

Multiple regression analysis was used to study if and how weather factors affect waterfowl populations. Indices of spring and summer temperatures of the previous year and winter severity were used as independent variables. The statistical significance of the models was tested using an analysis of variance. Goldeneye data were analysed without brood season temperature; as the goldeneye starts breeding at the age of 2-4 years (Dow & Fredga 1984), offspring




Figure 1. Total number of breeding pairs in the waterfowl population in 11 eutrophic wetlands of northern Savo, Finland, during 1984-1995. The solid line indicates the winter severity index for the Baltic Sea during the winters of 1983/84-1994/95.

Table 3. Densities of 14 waterfowl species expressed as pairs/km² (range, mean and standard deviation), coefficients of variations in densities (CV%) and Spearman rank correlations (R_s) of densities and year in 11 wetlands covering a total area of 9.46 km² during 1984-1995. Level of significance: ***P < 0.001, **P < 0.01, *P < 0.05.

Species	Range	Mean	SD	CV%	R _s	Level of significance
Great crested grebe	3.9 - 5.6	4.55	0.52	11.4	-0.413	
Slavonian grebe	0.1 - 0.5	0.26	0.12	-	0.116	
Red-necked grebe	0.0 - 0.1	-	-	-	-	
Whooper swan	0.1 - 0.5	0.24	0.15	-	0.921	**
Wigeon	3.2 - 8.0	5.87	1.19	20.3	0.031	
Teal	5.9 - 13.0	8.63	2.36	27.4	-0.529	
Mallard	3.5 - 5.9	4.52	0.91	20.1	-0.172	
Pintail	2.0 - 5.2	3.63	1.01	27.8	0.350	
Shoveler	2.7 - 5.3	3.79	0.67	17.7	0.305	
Garganey	1.0 - 2.6	1.62	0.57	35.2	-0.299	
Pochard	0.8 - 2.8	1.81	0.59	32.6	-0.902	***
Tufted duck	2.7 - 10.5	6.52	2.24	34.4	-0.867	**
Goldeneye	2.3 - 4.4	3.43	0.70	20.4	0.712	*
Coot	3.6 - 9.2	5.86	1.97	33.6	0.322	

production does not affect breeding numbers until after two or more years, therefore, the results for goldeneye consist of breeding females only.

Results

Changes in waterfowl populations

In the beginning of the 1980s, the total waterfowl population was ca 40% higher than at the beginning of the 1970s (Table 2). Populations of most species increased during this decade; the increase was significant for six species (see Table 2). Differences between extreme population levels during the 1980s and 1990s was 30%, and between successive years not above 20%.

We chose data from 12 successive years (1984-1995) for a more detailed analysis. During this period the wintering conditions of waterfowl clearly varied, but most of the winters in the Baltic Sea area were milder than average (Finnish Marine Research Institute). After the severe winter of 1984/85 followed by a cool spring, the total population decreased by 16%, and by 26% from the highest level recorded in 1984 to the lowest level recorded in 1986. The winter of 1986/87 was also severe, thus causing the total population to remain at a low level within our study area. The situation did not improve until spring 1989; the mean brood size of the previous summer was high (V-M. Väänänen, unpubl. data), and the previous winter was mild (Fig. 1, see Table 1).

Table 4. Results of multiple regression analysis used to explain the importance of weather factors (wintering, spring and brood seasons) affecting variation in breeding mallard, pintail, garganey and goldeneye populations during 1985-1995. The final model has been calculated including only significant independent variables. Explanations: Winter B = winter severity index in the Baltic Sea, Winter Fr = winter severity index in northern France, Spring x = spring temperature in year x, Spring x-1 = Spring temperature in year x-1, Brood s. x-1 = brood season temperature in year x-1 (see Methods for details).

Species	Variable	DF	Parameter estimate	Standard error	T	P	Final Model			
							Analysis of variance			
							DF	F-value	P	Adjusted R ²
Mallard	Winter B	1	-0.075	0.065	-1.244	0.254	1	11.384	0.007	0.486
	Spring x	1	0.290	1.102	0.263	0.800				
	Spring x-1	1	1.414	0.963	1.468	0.185				
	Brood s. x-1	1	4.012	1.300	3.089	0.018				
Pintail	Winter Fr	1	5.565	1.667	3.338	0.012	3	3.278	0.080	0.383
	Spring x	1	-4.136	1.519	-2.723	0.030				
	Spring x-1	1	1.955	1.039	1.881	0.102				
	Brood s. x-1	1	5.787	1.686	3.432	0.011				
Garganey	Winter Fr	1	-1.250	0.627	-1.994	0.086	1	20.178	0.001	0.636
	Spring x	1	2.616	0.571	4.580	0.003				
	Spring x-1	1	-0.673	0.391	-1.723	0.129				
	Brood s. x-1	1	0.297	0.634	0.469	0.654				
Goldeneye	Winter B	1	-0.190	0.044	-4.320	0.002	2	10.169	0.005	0.625
	Spring x	1	-1.919	0.685	-2.812	0.020				

Table 5. Proportions (%) of 10 waterfowl species of the total breeding population, and their respective proportions (%) of the hunting bag in August, September-November and in total during 1985-1989. The proportion of juveniles in the bag is given separately and the total number of observations is given in parenthesis.

	Mallard	Teal	Wigeon	Shoveler	Pintail	Garganey	Tufted duck	Pochard	Goldeneye	Coot
Breeding population	10.3 (232)	20.2 (454)	12.5 (280)	7.8 (174)	7.1 (159)	3.5 (79)	19.0 (427)	4.6 (103)	6.4 (144)	8.5 (191)
Hunting bag										
August	23.6 (393)	30.8 (504)	13.4 (223)	4.9 (82)	5.0 (84)	2.8 (47)	4.3 (72)	2.0 (33)	10.3 (172)	2.2 (36)
September-November	41.2 (361)	14.7 (129)	19.5 (171)	2.2 (19)	2.0 (17)	0.1 (1)	5.5 (48)	2.1 (18)	6.8 (60)	5.9 (52)
Total	29.8 (754)	25.0 (633)	15.5 (394)	4.0 (101)	4.0 (101)	1.9 (48)	4.7 (120)	2.0 (51)	9.2 (232)	3.5 (88)
Juveniles (%)	86.7	80.9	87.5	95.0	89.0	91.7	90.0	100.0	98.3	-

Populations of whooper swan *Cygnus cygnus* and goldeneye showed a statistically significant increasing trend, whereas populations of pochard and tufted duck showed decreasing trends (Table 3). Populations of garganey, tufted duck, coot and pochard were found to be the most unstable (see coefficients of variation in Table 3).

Weather factors affecting species populations

Our multiple regression model examining the relationship between the number of pairs and weather factors during wintering, spring and brood seasons was statistically significant for garganey, mallard and goldeneye and almost significant for pintail (Table 4), but not significant for all other species.

In garganey, spring temperatures explained most of the fluctuation: in warm springs numbers of garganey were highest. In mallard, cold brood seasons negatively affected the population and explained most of the fluctuations (see Table 4). In goldeneye, both severe winters in the Baltic and warm springs had negative effects on breeding numbers. Both winter severity in France and cold brood seasons in our study area had negative effects on the numbers of pintail. In cold springs pintails were found to be most abundant.

Hunting

Mallard, teal, wigeon and goldeneye were shot in

higher numbers than would be expected considering their respective proportion of the breeding communities (Table 5). All in all, a statistically significant difference was observed between proportions in the breeding duck community and in the hunting bag (G-test, $G = 706.9$, $P < 0.001$).

For the most hunted duck species, i.e. mallard, teal and wigeon, 81-87% of the hunting bag consisted of young individuals (see Table 5). The corresponding figures for other duck species were 90-100 %. The proportion of bagged adults was larger in dabbling ducks (14 %) than in diving ducks (<4%) ($G = 35.8$, $P < 0.001$). Only a small proportion (<2%) of the hunting bag consisted of adult goldeneyes.

The hunting pressure on mallard, tufted duck and goldeneye was clearly higher in our study area than elsewhere in Finland (Table 6). According to Finnish ringing data (young ringed), 77.5% of the mallards shot during 1970-1991 originated from Finland.

Discussion

Weather factors affecting populations

Fluctuations in total population levels may be related to weather factors, especially through the wintering success of the species wintering in the North. In Europe, there was a period of mild winters in the beginning of the 1970s; at that time, the wintering

Table 6. Recovery rates of young wigeons, mallards, pintails, shovelers, tufted ducks and goldeneyes banded in northern Savo and 'Other parts of Finland' during 1980-1991, and subsequently shot in Finland. Differences in recovery rates between northern Savo and 'Other parts of Finland' (mallard, tufted duck, goldeneye and total) were tested using G-test; *** $P < 0.001$.

Species	Northern Savo		Other parts of Finland		G-test
	Nos ringed	Nos recovered	Nos ringed	Nos recovered	
Wigeon	55	7 (12.7%)	110	5 (4.6%)	$G = 20.8$ ***
Mallard	79	21 (26.6%)	528	32 (6.1%)	
Pintail	71	7 (9.9%)	41	3 (7.3%)	
Shoveler	63	8 (12.7%)	47	2 (4.3%)	
Tufted duck	184	36 (19.8%)	954	17 (1.8%)	$G = 62.0$ ***
Goldeneye	3364	489 (14.5%)	3148	115 (3.7%)	$G = 207.5$ ***
Total	3816	568 (14.9%)	4828	174 (3.6%)	$G = 298.3$ ***

populations of many waterfowl species increased (e.g. Rüger, Prentice & Owen 1986, Monval & Piro 1989).

In the beginning of the 1980s, the total breeding waterfowl population in our study area was ca 40% higher than in the beginning of the 1970s. Our census results for the 1970s were obtained before the effects of mild winters could be seen in populations. The second census was performed mainly during the population peak of the 1980s. After the severe winters of the 1980s, the total population decreased by ca 30%.

In our study, we analysed population fluctuations of different waterfowl species in the 1980s and 1990s in more detail. With the exception of several winters in the 1980s, winters were generally mild in Europe during this period. Nevertheless, winter temperature explained population fluctuations in the goldeneye.

Spring temperature is known to cause population fluctuations in southern/northern species at the limit of their distribution area (in waterfowl, see e.g. Siira & Eskelinen 1983). In our study area, the annual breeding populations of garganey and pintail are primarily affected by spring temperatures, with southern garganey being found to be most abundant in warm springs and northern pintail in cold springs.

Early onset of spring may be important to waterfowl populations. After late springs, local hunting of early-arriving autumn migrants such as shoveler, pintail, garganey and widgeon may be more intense and larger proportions of ducklings may be less developed when the hunting season begins. In our study area, only late broods of pintail and shoveler were observed to be in their natal area at the beginning of the hunting season on 20 August (Väänänen 1996). These results indicate that after late springs ducks may be more vulnerable to local hunting.

In North America, the most important waterfowl production areas are very unstable (Evans & Black 1956, Eiserlohr 1969, Pospkala, Anders & Henny 1974). In these areas, brood production is significantly correlated with the size of the following year's breeding population (Reynolds & Sauer 1991). Variation in the quality, and especially in the number of August wetlands of brood production areas are known to bring about a strong annual fluctuation in the brood production (Leitch & Kaminski 1985, Sutherland 1991). In contrast, the quality of wetlands in northern Europe does not change dramatically, neither within nor between successive years (e.g. Nummi & Pöysä 1993), and thus are not very likely to cause considerable variation in brood production.

Brood production may also occasionally play an important role in the fluctuations of populations in our study area but, unfortunately, we do not have the data to examine this in detail.

Hunting

The total effect of hunting (compensatory or additive mortality) on breeding populations is difficult to evaluate, because waterfowl hunting occurs in the breeding, gathering, migrating and wintering areas. Mortality due to hunting also fluctuates between years (Rogers et al. 1979); for example, in severe winters, the hunting pressure may be high, because the ice cover forces ducks to move to areas with limited open water, thus making them more vulnerable to hunting (e.g. Grenquist 1965, Ogilvie 1981).

In our study area, the hunting pressure was most heavy on mallards (see also Pirkola & Lindén 1972, V-M. Väänänen, unpubl. data) and might have had adverse effects on breeding numbers. For example, in 1986-1988, the densities of the mallard populations in our study area were clearly below those observed in other parts of Finland, such as Satakunta, Päijät-Häme, southeastern Finland and southern Lapland (E. Lammi & R.A. Väisänen, unpubl. data). However, densities of the southern duck species, garganey and shoveler, were higher in our study area than in other parts of Finland (E. Lammi & R.A. Väisänen, unpubl. data). Thus, the quality of the habitats does not explain the low mallard densities in our study area.

Furthermore, in our study area, unlike in other areas of Finland, the rapid increase in the waterfowl populations during 1987-1991 was not observed for mallard (Spearman $R_s = 0.100$, $P > 0.1$) even though an overall increase in total waterfowl numbers was evident (see Fig. 1). Conversely, during the same period in the Päijät-Häme area of southern Finland (E. Lammi, unpubl. data), the mallard population showed a steady increase (Spearman $R_s = 1.000$, $P < 0.001$). Similar increasing trends for mallards were also observed by H. Pöysä (unpubl. data) in southeastern Finland (Spearman $R_s = 0.670$, $P < 0.1$), and are further corroborated by the findings of the nationwide monitoring program of breeding waterfowl in Finland showing an increasing trend in mallard numbers during the same period (Pöysä, Lammi, Väisänen & Wikman 1993). During 1987-1991, the annual bag of mallard in our study area increased (Spearman $R_s = 0.900$, $P = 0.072$); this result, however, was only marginally significant possibly due to a study

period of only five years. These findings seem to indicate that local hunting could have had a remarkably adverse effect on the densities of mallard breeding populations in our study area.

The proportion of adult females bagged in our study area is, except for teal, smaller than that reported in Denmark (Clausager 1987, 1988, 1989). In Denmark, most hunting of adult diving ducks is concentrated in wintering areas, where the proportion of, for example, adult goldeneye in the bag may exceed 50% (Clausager 1987, 1988, 1989), whereas in our study area adult goldeneye accounted for less than 2% of the bag. The survival of breeding females is very important because experienced breeders have considerably higher breeding success than first-time breeders (e.g. Owen & Black 1990).

Even though large numbers of goldeneye were shot, the breeding population of this duck species seemed to increase. During our study, the bag of goldeneye mainly consisted of young ducks (see Table 5). This is due to the fact that adult females migrate to the Finnish sea area in order to moult prior to the opening of the Finnish hunting season (Runko & Väänänen 1989), and that males typically leave the breeding areas before the middle of June.

Goldeneye start breeding at the age of 3-5 years (Fredga & Dow 1984). Because of this late breeding age, survival of old females is crucial for the stability of the population. In Sweden, an annual survival rate of 71.8% has been reported for adult goldeneye females (Fredga & Dow 1984); in our study area approximately 81% of breeding females have been observed to breed again (Runko & Väänänen 1989). Thus, the exceptionally high survival rate of breeding females evidently contributed to the increase in the goldeneye population.

Our ringing data indicate that the hunting pressure on mallard, tufted duck and goldeneye is stronger in our study area than in other parts of Finland. In many cases, hunting is likely to have an adverse effect only in years when weather conditions are unfavourable (e.g. late springs and severe winters).

Garganey, shoveler and pintail are early autumn migrants, as demonstrated by the banding and bag data (Mihelsons, Mednis & Blums 1986, Clausager 1987, 1988, 1989, Väänänen 1996, V-M. Väänänen, unpubl. data). In contrast to the findings of Siira & Eskelinen (1983) from the Liminka Bay wetland in western Finland, our data suggest that local hunting hardly affects the breeding populations of these early migrating species in our study area.

Concluding remarks

In summary, fluctuations of waterfowl populations appear to reflect many interacting factors. Weather is an important factor affecting breeding numbers. In pothole areas of Northern America, habitat alteration and destruction are crucial factors influencing fluctuations. In northern Europe, the conditions of habitats do not generally change so dramatically.

Expansion of tall vegetation, such as *Salix* and *Phragmites*, into open sedge stands and shore meadows affects populations of certain wetland species (e.g. von Haartman 1975, Soikkeli & Salo 1979). Most dabbling ducks have been shown to favour open habitats in our study area (Kauppinen 1993). These species have not shown parallel population trends that can be easily explained by such habitat alterations. In our study area the shores of many eutrophic lakes are still pastured, with cultivated fields in the near vicinity, so the features of landscape are still open. In contrast, the shores of many mixotrophic lakes have never been pastured, and their habitat structure has remained fairly stable for a long time.

Predation, especially nest predation, is an important factor affecting the production of offspring in birds (e.g. Lack 1968, Wiens 1989). In our study area, nest predation of *Aythya* species is high outside gull colonies, whereas inside colonies gulls successfully protect all nests (Väänänen in press). Decreasing numbers of breeding black-headed gulls *Larus ridibundus* may play an important role also in the decreasing population trends of tufted duck and pochard (J. Kauppinen & V-M. Väänänen, unpubl. data).

In our study area, the total population of all waterfowl species was in a phase of increase during the 1970s, and reached its peak level at the beginning of the 1990s. Nevertheless, the mallard population level was lower in the 1990s. We suggest that hunting has had a negative effect on the densities of the breeding mallard population.

Acknowledgements - we are grateful to Esa Lammi, Hannu Pöysä and Risto A. Väisänen for providing comparative data from the Finnish waterfowl monitoring project and to the Finnish Marine Research Institute, Kuopio Meteorological Station and Météo-France for providing meteorological data. We also wish to thank associate editor Hannu Pöysä, Pekka Helle, Petri Nummi and two anonymous referees for useful discussions and valuable comments on the manuscript. This study was financially supported by the Kuopio Naturalist Society and the Finnish Game Foundation.

References

- Andersen-Harild, P. 1981: Population dynamics of *Cygnus* olor in Denmark. - Proceedings of the Second International Swan Symposium, Sapporo: 176-190.
- Anderson, D.R. & Burnham, K.P. 1976: Population ecology of the mallard VI. The effect of exploitation on survival. - U.S. Fish & Wildlife Service Resources Publications 128: 1-66.
- Boyd, H. 1964: Wildfowl and other waterfowl found dead in England and Wales in January-March 1963. - Wildfowl Trust Annual Reports 15: 20-23.
- Boyd, H. 1981: Prairie dabbling ducks, 1941-1990. - Canadian Wildlife Service (Progress Notes) 119: 1-9.
- Bethke, R.W. 1993: Geographical patterns of persistence in duck guilds. - *Oecologia* 93: 102-128.
- Caithness, T., Williams, M. & Nichols, J.D. 1991: Survival and band recovery rates of sympatric grey ducks and mallard in the New Zealand. - *Journal of Wildlife Management* 55: 111-118.
- Clausager, I. 1987: Vingeindsamling fra jagtsæsonen 1986/87 i Danmark. - Rapport fra Vildtbiologisk Station, Landbrugsministeriets Vildtforvaltning, Juli 1987, 31 pp. (In Danish with English summary).
- Clausager, I. 1988: Vingeindsamling fra jagtsæsonen 1987/88 i Danmark. - Rapport fra Vildtbiologisk Station, Landbrugsministeriets Vildtforvaltning, Juli 1988, 32 pp. (In Danish with English summary).
- Clausager, I. 1989: Vingeindsamling fra jagtsæsonen 1988/89 i Danmark. - Rapport fra Vildtbiologisk Station, Landbrugsministeriets Vildtforvaltning, Juli 1989, 38 pp. (In Danish with English summary).
- Danell, K. & Sjöberg, K. 1979: Abundance and productivity of ducks on boreal lakes in northern Sweden. - *Annales Zoologica Fennici* 16: 123-128.
- Dow, H. & Fredga, S. 1984: Factors affecting reproductive output of the goldeneye duck *Bucephala clangula*. - *Journal of Animal Ecology* 53: 679-692.
- Eisenlohr, W.S. 1969: Hydrology of small water areas in the prairie pothole region. - Canadian Wildlife Services, Report Series 6: 35-39.
- Evans, C.D. & Black, K.E. 1956: Duck production studies on the prairie potholes of South Dakota. - U.S. Fish & Wildlife Service, Special Scientific Report 32: 1-59.
- Fredga, S. & Dow, H. 1984: Factors affecting the size of a local population of goldeneye, *Bucephala clangula* (L.). - *Viltrevy* 13: 225-252.
- Grenquist, P. 1965: Changes in abundance of some duck and seabird populations off the coast of Finland 1949-1963. - *Finnish Game Research* 27: 1-114.
- Helle, P. & Mönkkönen, M. 1986: Annual fluctuations of land bird communities in different successional stages of boreal forests. - *Annales Zoologica Fennici* 23: 269-280.
- Hildén, O. 1966: Changes in the bird fauna of Valassaaret, Gulf of Bothnia, during recent decades. - *Annales Zoologica Fennici* 3: 245-269.
- Kalela, O. 1946: Zur Ausbreitungsgeschichte der Vögel vegetationsreicher Seen. - *Annales Academiae Scientiarum Fennicae Series A, IV. Biologica* 12: 1-81. (In German).
- Kalela, O. 1949: Changes in geographical ranges in the avifauna of northern and central Europe in relation to recent changes in climate. - *Bird Banding* 20: 77-103.
- Kauppinen, J. 1980: Sorsalintujen pesivän kannan laskentameteodeista ja niiden virhelähteistä. (In Finnish with English summary: Sources of error in estimation methods of breeding waterfowl populations. - *Lintumies* 15: 74-82.
- Kauppinen, J. 1983: Methods used in the census of breeding ducks in Northern Savo (Finland) at the beginning of the breeding season. - *Finnish Game Research* 40: 49-81.
- Kauppinen, J. 1993: Densities and habitat distribution of breeding waterfowl in boreal lakes in Finland. - *Finnish Game Research* 48: 24-45.
- Kauppinen, J. 1995: Temporal variation in the wetland waterfowl communities of the Finnish lake district. - *Ornis Fennica* 72: 145-153.
- Kauppinen, J., Koskimies, P. & Väisänen, R.A. 1991: Waterfowl round count. - In: Koskimies, P. & Väisänen, R.A. (Eds.); *Monitoring bird populations*. - Zoological Museum, Finnish Museum of Natural History, Helsinki, pp. 45-53.
- Kauppinen, J. & Väisänen, R.A. 1993: Ordination and classification of waterfowl communities in south boreal lakes. - *Finnish Game Research* 48: 3-23.
- Lack, D. 1968: Ecological adaptations for breeding in birds. - Methuen, London, 409 pp.
- Leitch, W.G. & Kaminski, R.M. 1985: Long-term wetland-waterfowl trends in Saskatchewan grassland. - *Journal of Wildlife Management* 49: 212-222.
- Linkola, P. 1962: Notes on the breeding success of ducks in Central Häme. - *Suomen Riista* 15: 157-174. (In Finnish with English summary).
- Mihelsons, H., Mednis, A. & Blums, P. 1986: Population ecology of migratory ducks in Latvia. - "Zinatne", Riga, 111 pp.
- Monval, J-Y. & Piro, J-Y. 1989: Results of the IWRB International Waterfowl Census 1967-1986. - IWRB Special Publication 8: 1-145.
- Nilsson, L. 1979: Variation in the production of young of swans wintering in Sweden. - *Wildfowl* 30: 129-134.
- Nilsson, L. 1984: The impact of hard winters on waterfowl populations of south Sweden. - *Wildfowl* 35: 71-80.
- Nilsson, L. 1985: Bestandsdichte und Vergesellschaftung brütender Wasservögel Südschweden in Beziehung zur Produktivität der Seen. - *Journal of Ornithology* 126: 85-92. (In German).
- Nummi, P. & Pöysä, H. 1993: Habitat association of ducks during different phases of the breeding season. - *Ecography* 16: 319-328.
- Nummi, P. & Pöysä, H. 1995: Breeding success of ducks in relation to different habitat factors. - *Ibis* 137: 145-150.

- Ogilvie, M.A. 1981: Winter 1978/79 hard weather movements and mortality of ducks ringed in the United Kingdom. - Proceedings of the second Technical Meeting, Western Palearctic Migratory Bird Management, Paris: 174-180.
- Onno, S. 1965: The numbers of waterfowl in Estonia. - Wildlife Trust Annual Reports 16: 110-114.
- Owen, M. & Black, J.M. 1990: Waterfowl ecology. - Chapman & Hall, New York, 194 pp.
- Pirkola, M.K. & Lindén, H. 1972: Sorsansiipikeräyksen tuloksia vuosilta 1969 ja 1970. (In Finnish with English summary: Results of duck wing collection surveys in Finland 1969 and 1970. - Suomen Riista 24: 97-106.
- Posphala, R.S., Anders, D.R. & Henny, C.J. 1974: Breeding habitat conditions, size of the breeding populations, and production indices. - U.S. Fish & Wildlife Service Resources Publications 115, 73 pp.
- Pöysä, H. 1984: Temporal and spatial dynamics of waterfowl populations in a wetland area - a community ecological approach. - Ornis Fennica 61: 99-108.
- Pöysä, H. 1989: Vesilintuyhteisöjen vakauden alueellisista eroista. (In Finnish with English summary: Geographical gradients in the stability of waterfowl communities in Finland. - Suomen Riista 35: 7-16.
- Pöysä, H., Lammi, E., Väisänen, R.A. & Wikman, M. 1993: Monitoring of waterbirds in the breeding season: the programme used in Finland in 1986-92. - In: Moser, M., Prentice, R.C. & van Vessem, J. (Eds.); Waterfowl and wetland conservation in 1990s - a global perspective. Proceedings of an IWRB Symposium, St. Petersburg Beach, Florida, USA, IWRB Special Publication 26, Slimbridge, pp. 7-12.
- Reynolds, R.E. & Sauer, J.R. 1991: Changes in mallard breeding populations in relation to production and harvest rates. - Journal of Wildlife Management 55: 483-487.
- Rogers, J.P., Nichols, J.D., Martin, F.W., Kimball, C.F. & Posphala, R.S. 1979: An examination of harvest and survival rates of ducks in relation to hunting. - Transactions North American Wildlife Natural Resources Conference 44: 114-126.
- Rüger, A., Prentice, C. & Owen, M. 1986: Results of the IWRB international waterfowl census 1967-1983. - IWRB Special Publications 6: 1-118.
- Runko, P. & Väänänen, V.-M. 1989: Telkän (*Bucephala clangula*) rengastusten tuloksia Maaningalla. - Linnut 1989, II Suomen Lintutieteen kongressi (Abstract in Finnish), p. 35.
- Salminen, A. 1983: Suomen sorsalinnut. - SLY:n Lintutieto Oy, Helsinki, 206 pp. (In Finnish).
- Soikkeli, M. & Salo, J. 1979: The bird fauna of abandoned shore pastures. - Ornis Fennica 56: 124-132.
- Siira, J. & Eskelinen, O. 1983: Changes in the abundance of breeding waterfowl in the Liminka bay in 1954-81. - Finnish Game Research 40: 106-121.
- Stoudt, J.H. 1971: Ecological factors affecting waterfowl production in the Saskatchewan parklands. - U.S. Fish & Wildlife Service Resources Publications 99: 1-58.
- Stoudt, J.J. & Cornwell, G.W. 1976: Non-hunting mortality of fledged North American waterfowl. - Journal of Wildlife Management 40: 681-691.
- Sutherland, J.M. 1991: Effects of drought on American coot (*Fulica americana*) reproduction in Saskatchewan parklands. - Canadian Field Naturalist 105: 267-273.
- Trauger, D.L. & Stoudt, J.H. 1978: Trends in waterfowl populations and habitats on study areas in Canadian parklands. - Transactions North American Wildlife Natural Resources Conference 43: 187-205.
- Väänänen, V.-M. 1996: Vesilintujen metsästysverotus rengastusaineistojen valossa. (In Finnish with English summary: Hunting pressure of ducks in Finland according to ringing data. - Suomen Riista 42: 40-46.
- Väänänen, V.-M. in press: Predation risk associated with nesting in gull colonies by two *Aythya* species: observations and an experimental test. - Journal of Avian Biology 30.
- von Haartman, L. 1945: Zur Biologie der Wasser- und Ufervögel im Schärenmeer Südwestfinnlands. - Acta Zoologica Fennica 44: 1-120.
- von Haartman, L. 1957: Population changes in the tufted duck, *Aythya fuligula* (L.). - Societas Scientiarum Fennica Commentationes Biologicae 16, 5: 1-11.
- von Haartman, L. 1973: Changes in the breeding bird fauna of north Europe. - In: Farner, D.S. (Ed.); Breeding Biology of Birds. National Academy of Sciences, Washington, pp. 448-481.
- von Haartman, L. 1975: Changes in the breeding bird fauna of coastal bays in southwestern Finland. - Ornis Fennica 52: 57-67.
- Wiens, J.A. 1981: Scale problems in avian censusing. - Studies in Avian Biology 6: 513-521.
- Wiens, J.A. 1989: The ecology of birds communities. Vol 2. Processes and variations. - Cambridge University Press, Cambridge, 316 pp.
- Yarker, B. & Atkinson-Willes, G.L. 1971: The numerical distribution of some British breeding ducks. - Wildfowl 22: 63-70.