

Echeveria lyonsii new species from Tamaulipas, Mexico

Author: Kimnach, Myron

Source: Cactus and Succulent Journal, 79(5) : 215-219

Published By: Cactus and Succulent Society of America

URL: [https://doi.org/10.2985/0007-9367\(2007\)79\[215:ELNSFT\]2.0.CO;2](https://doi.org/10.2985/0007-9367(2007)79[215:ELNSFT]2.0.CO;2)

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Echeveria lyonsii

a new species from
Tamaulipas, Mexico

Summary. A new species, *Echeveria lyonsii*, is described from Tamaulipas, Mexico. It appears to be related to *E. shaviana* and *E. bifida*, both with 13 chromosomes. From these, *E. lyonsii* differs in having 12 chromosomes and smaller flowers. Also discussed is a natural hybrid apparently allied with *E. lyonsii*.

Echeveria lyonsii

In 1971 Gary Lyons and I were exploring along the road from Ciudad Victoria to Jaumave, Tamaulipas, Mexico. About 19 miles from Ciudad Victoria we found a number of interesting plants growing on an oak-strewn slope much disturbed by livestock: *Mammillaria candida*, *M. picta* (?), *M. macrocentra* (?), *Opuntia microdasys*, *O. stenopetala*, a coryphantha, *Dolichothele baumii*, two agave species, *Villadia cucullata* ssp *apiculata*, a dasyliion, and *Hechtia* aff *capituligera*. We dug up a plant of the latter and, while I was carrying it back to the truck, Gary pointed out that below it an echeveria was dangling by its roots.

This species—so serendipitously discovered—is named for Gary Lyons, Curator of the Desert Garden at the Huntington. Besides being apparently the first to have seen the species, Gary is noted for having designed the Ethel Mars Cactus Garden in Las Vegas, Nevada, authored a book (*Desert plants, a curator's introduction to the Huntington Desert Garden*, Huntington Library, 2007) and co-authored another (*Desert gardens*, by Levick and Lyons, 2000). In 1966 and 1971 he participated in two Huntington botanical expeditions to Mexico.

The purplish leaves of *Echeveria lyonsii* closely resemble those of *E. bifida*, though they are somewhat more spatulate; the leaves of *E. lyonsii* also lack the undulation found in *E. shaviana*. Florally, *E.*

lyonsii is distinct in series *Angulatae* in having very small flowers—only 9 mm long. An apparently allied natural hybrid found by Jack Davies in 1965 near Jaumave is discussed below.


In the ensuing 40 years or so since these two collections were made, both have been propagated, photographed, and studied. However, their final classification has been delayed because of uncertainty as to the relationship between them and allied species.


In the course of his work with the Mexican Crassulaceae, Charles Uhl made eight hybrids of *E. lyonsii* with other *Echeveria* species and one with a *graptopetalum*¹.

Inflorescence of
E. lyonsii type
plant.



▼ The natural hybrid allied with *E. lyonsii*. 1. Flowering plant, ($\times 0.4$). 2, 3. Leaf, ($\times 0.5$). 4, 5. Flower, lateral and apical views, ($\times 2$). 6. Petals and stamens, ($\times 2$). 7. Gynoecium, ($\times 3$). 8. Nectary, greatly enlarged. Drawing by Diana Jacobs, 1985, financed by a grant from the Cactus and Succulent Society of America.





J. Traeger

▲ **Rosette of *Echeveria lyonsii*.**

Echeveria lyonsii is easily grown, offsets prolifically and can produce young plants from detached leaves or bracts. Its flowering stems are multiple and vigorous, sometimes weakening the plant during hot summers.

An allied natural hybrid

Six years before the discovery of *E. lyonsii*, Jack Davies of Monrovia, California, was looking for plants along the same road. About 10 km southwest of Jaumave (some 25 miles southwest of the type locality of *E. lyonsii*) he stopped at a

▼ **Gary Lyons at the Huntington.**



J. Traeger

»» GLOSSARY

androecium the male part of a flower, that is, the stamens

aristate with a bristle at the apex

bracteoles small bracts on the pedicels or upper part of an inflorescence

cincinnus (pl. cincinnati) a curving branch of an inflorescence with the flowers alternating in orientation

epipetalous/antisepalous stamens opposite the petals / opposite the sepals

glaucous with a gray or white, waxy coating

gynoecium the female part of a flower, that is, the follicles and stigma

lenticular lens- or lentil-shaped

pedicel the stalk immediately below the flower

pyriform pear-shaped

rachis the central stem of an inflorescence or compound leaf

spathulate spatula-shaped; rounded and wider at the apex, tapering below



➤➤ DIAGNOSIS & DESCRIPTION

***Echeveria lyonsii* Kimnach sp. nov.** *Caulis minus quam 3 cm altus, rosula 11–14 cm lata, folia spathulata acuta 6–7 cm longa 1–18 mm lata 2–2.5 mm crassa purpureo-grisea. Caules florentes ad 35 cm alti, cincinni 2 vel plerumque, pedicelli minus quam 2 mm longi, sepala ascendens 2–5 mm longa, corolla 9 mm longa rubello-lutea.*

Plant proliferous, **stem** to 6 cm tall or more, 8–10 mm thick, with brown, horizontal, lenticular leaf-scars. Rosette 11–14 cm wide, **leaves** ascending to horizontal, spathulate, acute, 6–7 cm long, 17–19 mm wide at widest part 2–2.5 cm from apex, 17 mm wide at middle, ca. 7 mm wide near base, mostly 2–2.5 mm thick, ca. 4 mm thick near base, convex and bluntly keeled on lower surface, concave along upper half of upper surface, purplish gray, lightly glaucous.

Flowering stems 2–5-parted, sometimes with up to three additional secondary branches, peduncle ca. 25–45 cm long, ca. 5 mm thick near base, ca. 4 mm thick just below first cincinnus, pinkish gray, lightly glaucous; **bracts** 22–24, attached ca. 1.5–3 cm apart, ascending, the lower ones slightly recurving on apical half, narrowly elliptical-linear, acute, 22–35 mm long, 5–7 mm wide, 3–4 mm thick; **cincinni** 2–8 per flowering stem, 4–16 cm long, unbranched, 3–17-flowered, rachis 1.2–2 mm thick, flowers ca. 5–10 mm apart; **bracteoles** 1 between each pair of flowers, oblong-elliptical, acute, 6–10 mm long, 3–4 mm wide, 1.5–2 mm thick, colored like the leaves and bracts; **pedicels** less than 2 mm long, ca. 1.5 mm thick; **sepals** ascending, ovate-elliptical, acute, 2–5 mm long, 1–2 mm wide, 0.5–1.75 mm thick; **corolla** ± straight-sided, strongly angled; **petals** linear-lanceolate, acute, ca. 9 mm long, ca. 1.5 mm wide, pinkish orange, darker near apex, the apex expanding to 45°; epipetalous **stamens** 2 mm long, the antisepalous 3–3.5 mm long, anthers oblong, ca. 1.25 mm long, pale yellow; **gynoecium** pyriform, acute, ca. 6 mm long and 4 mm thick, cream below, stigmas light green; **nectaries** shaped like incisor teeth, 0.25 mm wide, light yellowish.

Chromosomes: $n = 12$.

Mexico: Tamaulipas: 19 miles SW of Ciudad Victoria, 4400', May 24, 1971, *M. Kimnach and G. Lyons* 1445, Huntington Botanical Gardens 28005, *Uhl U2428* (HNT, holotype [sheet number 3561]; MEXU, US, isotypes).

➤➤ DESCRIPTION

Echeveria hybrid

Plant short-stemmed, proliferous, entirely glaucous except for androecium and gynoecium. **Stems** to 4–5 cm high or more, 1–2 cm thick. **Leaves** densely arranged in rosettes 12–18 cm wide, each with 15–35 leaves, at first strongly ascending, oblanceolate to linear-oblanceolate, subobtuse to acute, with a nearly aristate cusp 2–3 mm long, leaf blade 6.5–8.5 cm long, 7–10 mm wide at base, narrowing to 5–8 mm ca. 5 mm above base, 15–24 mm wide at widest part ca. 2 cm from apex, 3–5 mm thick, thickest ca. 2 cm from base, upper side convex along basal half, deeply concave along apical half, lower side strongly convex, obtusely to indistinctly keeled, apex upcurving, margin obtuse below, upper portion subacute, somewhat undulate on apical half but usually only on one side, somewhat glaucous, pinkish gray, apical portion darker pink.

Flowering stems 1 or 2 to a rosette, erect, 30–40 cm tall, with 2–4 cincinni, some of these often bifurcate; rachis terete, 20–28 cm long and 2–7 mm thick up to first cincinnus; **bracts** 18–27, lacking for basal 1–3 cm, then attached 1–2 cm apart, ascending, similar in shape to rosette leaves, lower ones more linear-oblanceolate, the upper subelliptical, 1.5–6 cm long, 2–10 mm wide, 2–5 mm thick, at base with 1–3 acute, whitish spurs to 2 mm long, lower bracts incurving near apex, upper ones slightly recurving near apex; **inflorescence** ca. 3–8 cm tall and wide when young, later to about 12 cm tall and wide when cincinni are fully expanded; cincinni usually 1 or bifurcate, often with another 1 or 2 lower down on opposite sides of rachis, arching-pendent when young, later nearly erect, 6–14 cm long; bracteoles 7–20 mm long, with (8–)15–21 flowers; **pedicels** less than 2 mm long, ca. 2 mm thick; **sepals** ascending-incurving, very unequal, 1.5–4 mm wide, 2.5–14 mm long; **corolla** conical-pentagonal, exterior prominently keeled, 10–12 mm long, ca. 6 mm thick, orange-pink

Mexico: Tamaulipas: 10 km SW of Jaumave on road from Ciudad Victoria to Tula, 1965, *Jack Davies s.n.*, Huntington Botanical Gardens 41537, *Uhl U2845*, *ISI 1848* (BH, HNT, MEXU, SD, US).

rocky, lightly forested slope. While returning to his car he found a solitary echeveria growing underneath a shrub. He collected the plant, lacking time to explore the higher, steeper slopes where more of the plants probably grew. In 1988 International Succulent Introductions distributed his collection² as ISI 1848, “*E. shaviana* × *E. bifida*.” It is not known if any of the plants still exist in collections.

Uhl¹ provided a thorough and rather technical discussion of this plant that explains that the plant is, indeed, a natural hybrid: “In 1965, 10 km SW of Jaumave... on the road to Palmillas and ca. 40 km S of known localities of *E. shaviana*, Jack Davies found a single attractive plant (HNT 41537 = U2845) with pinkish leaves having more or less undulate margins, a bit like those of *E. shaviana* but thicker. Thought at first to be a new species, it is highly irregular at meiosis and clearly a hybrid. At metaphase I it forms 7–11 paired elements, some of them large multivalents, and 1–5 unpaired chromosomes... At anaphase it has chromosome bridges, at metaphase II the number of chromosomes on each plate ranges from 11 to 14, and only about 1% of its pollen was stainable. The International Succulent Institute offered this in 1988... as an apparent natural hybrid between *E. shaviana* ($n = 13$) and *E. bifida* ($n = 12$), and the cytology is compatible with this.”

Although this parentage of the hybrid is possible, it may be that *E. lyonsii* is in its ancestry. One of the striking characters of the hybrid is its long, pendulous cincinni, especially pronounced under good cultivation. Aside from this and its cytology, the hybrid differs mostly in several somewhat larger dimensions:

	<i>E. LYONSII</i>	HYBRID
Leaves	2–4 mm thick	3–5 mm thick
Bracts	22–35 mm long	15–60 mm long
Sepals	2–5 mm long	2.5–15 mm long
Corolla	9 mm long	10–12 mm long

It is obvious that publishing a new species and discussing an allied natural hybrid on the basis of a single collection of each is not desirable. However, Mexican regulations now greatly hinder foreign botanists who wish to collect and export Mexican plants for research. It is hoped that this article will encourage Mexican botanists to study these two taxa in the field—where each was found only once nearly 40 years ago—so their interrelationships may be clarified. ❖

ACKNOWLEDGEMENTS

Thanks are due to Charles Uhl for his critique of this paper and to Jack Davies for photos and plant material.



▲ Inflorescence of the hybrid as cultivated by Frank Reinelt, Santa Cruz, California.

▼ The probable natural hybrid in cultivation.



REFERENCES

- 1 UHL CH. 1999 (“1998”). Chromosomes and hybrids of *Echeveria* (Crassulaceae) VI. Series Angulatae Walther and Pruinosaes Walther. *Haseltonia* 6: 70, 85, 88–89; Figs 13, 42. 2 DAVIS L, AND OTHERS. 1988. The 1988 offering of International Succulent Introductions. *Cact Succ J* (US) 60: 93.



1



2



3



4



5



6



7



8

*More
plants
described
by Myron*

Photos by
John Trager.

1 *Graptopetalum pentandrum* ssp. *pentandrum* 2 *Peperomia kinnachii* 3 \times *Tacipetalum* 'Stardust' 4 *Graptopetalum paraguayense* ssp. *paraguayense* 5 *Graptopetalum paraguayense* ssp. *bernalense* 6 *Graptopetalum paraguayense* (left) and ssp. *bernalense* (right) 7 *Peperomia dolabella* 8 *Graptopetalum saxifragoides* variety *saxifragoides*.