

## **Assessing Potential Habitat for Freshwater Mussels by Transferring a Habitat Suitability Model within the Ozark Ecoregion, Missouri**

Authors: Hartman, Jordan H., Rosenberger, Amanda E., Key, Kayla N., and Lindner, Garth

Source: Freshwater Mollusk Biology and Conservation, 26(1) : 32-44

Published By: Freshwater Mollusk Conservation Society

URL: <https://doi.org/10.31931/fmbc-d-21-00005>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REGULAR ARTICLE

# ASSESSING POTENTIAL HABITAT FOR FRESHWATER MUSSELS BY TRANSFERRING A HABITAT SUITABILITY MODEL WITHIN THE OZARK ECOREGION, MISSOURI

Jordan H. Hartman<sup>\*†1</sup>, Amanda E. Rosenberger<sup>2</sup>, Kayla N. Key<sup>‡1</sup>, and Garth Lindner<sup>§3</sup>

<sup>1</sup> Tennessee Cooperative Fisheries Research Unit, Department of Biology, Tennessee Technological University, Cookeville, TN 38501 USA

<sup>2</sup> U.S. Geological Survey, Tennessee Cooperative Fisheries Research Unit, Department of Biology, Cookeville, TN 38501 USA

<sup>3</sup> University of Missouri, Missouri Cooperative Fisheries Research Unit, School of Natural Resources, Columbia, MO 65211 USA

---

## ABSTRACT

Habitat suitability models for freshwater mussels can inform conservation of these imperiled animals. Riverscape-scale hydrogeomorphic variables were previously used to predict suitable mussel habitat in the Meramec River basin, Missouri. We evaluated transferability of the Meramec River habitat suitability model to the Gasconade and Little Black rivers, in the Ozark Highlands ecoregion, Missouri. The best-fit models relied on transferring and adapting the original modeling framework to better represent the unique habitat characteristics of each river. Mussel bed occurrence in both rivers was associated with reaches that were classified as pools. Mussel beds in the Gasconade River were also associated with laterally stable reaches adjacent to small bluffs, distant from gravel bars, and with higher stream power indices. Mussel beds in the Little Black River were associated with reaches with higher surface water availability during low-flow conditions, lower stream power indices, and bluffs located downstream. Our results show that existing habitat models can be transferred to other streams with similar environmental conditions, but differences in watershed characteristics can affect transferability.

**KEY WORDS:** freshwater mussels, habitat suitability modeling, hydrogeomorphology, MaxEnt, riverscape scale, transferability

---

## INTRODUCTION

Understanding habitat and environmental associations of freshwater mussels is essential for the conservation of these highly imperiled animals (FMCS 2016). The occurrence of

large, multispecies mussel aggregations, or mussel beds, suggests that common habitat preferences influence or limit mussel establishment and persistence across multiple species (Vaughn 1997). Reach-scale factors such as microhabitat characteristics and host-fish distributions typically have little explanatory power for predicting mussel distribution and abundance (Strayer and Ralley 1993; Johnson and Brown 2000; Vaughn 2012; Pandolfo et al. 2016; Randklev et al. 2019). Mussel occurrence can be predicted at watershed scales based on geology, soils, land use, and topography (Strayer 1993; Arbuttle and Downing 2002; Daniel and Brown 2014; Walters et al. 2017), but these factors are not tractable for

---

\*Corresponding Author: Jnhnd@illinois.edu

†Current Address: University of Illinois, Department of Natural Resources and Environmental Science, Urbana-Champaign, Illinois 61829 USA

‡Current Address: West Tennessee River Basin Authority, Tennessee Department of Environment and Conservation, Humboldt, Tennessee 38343 USA

§Current Address: Indiana Department of Natural Resources, Indianapolis, Indiana 46204 USA

management actions (Fausch et al. 2002). More recently, hydrogeomorphic variables corresponding to in-channel stability (e.g., shear stress, hydraulic stability, and presence of refugia during high- and low-flow events) show promise for predicting mussel occurrence at the reach scale (Allen and Vaughn 2010; Drew et al. 2018). However, understanding mussel habitat associations at the riverscape scale may be most useful for prioritizing management efforts (Bouska et al. 2018).

The riverscape scale represents the continuous, longitudinal river gradient as intermediate between reach-scale microhabitat characteristics and watershed-scale factors. Therefore, the riverscape scale is small enough to be influenced by management actions but large enough to encompass the continuous, hierarchical, and heterogeneous river system in its entirety (Schlosser and Angermeier 1995; Fausch et al. 2002; Bouska et al. 2018). However, at a riverscape scale, it is time-consuming and expensive to generate hydrogeomorphic data needed to predict mussel occurrence while providing inferences relevant to management (Bouska et al. 2018).

Key et al. (2021) developed a habitat suitability model using open-source, remotely sensed data to predict mussel bed occurrence at the riverscape scale in the Meramec River basin, Missouri. The Meramec River habitat suitability model (MRHSM) assessed the association of mussel beds with hydrogeomorphic variables reflecting water availability, channel stability, and the presence of stable gravel substrate. Habitat suitability models, such as the MRHSM, may be transferred to other areas with similar environmental conditions by obtaining remotely sensed data for those areas (Randin et al. 2006; Barbosa et al. 2009; Werkowska et al. 2016).

We investigated the transferability of the MRHSM (Key et al. 2021) to the Gasconade and Little Black rivers, two other Ozark rivers in Missouri. Transferring the MRHSM could inform mussel conservation throughout the Ozark region and provide more information about mussel habitat associations in general. Our objectives were to derive a dataset of spatial layers for our study streams that represent hydrogeomorphic variables used in the MRHSM and determine the best method for transferring the MRHSM to the Gasconade and Little Black rivers. We discuss how well the MRHSM can be transferred to the Gasconade and Little Black rivers and how the hydrogeomorphology of those watersheds affects transferability.

## METHODS

### Study Areas

The Meramec, Gasconade, and Little Black river basins are within the Ozark Highlands ecoregion of Missouri (Fig. 1) and share similar physiographic and watershed features. In the interior of the region, dolomite and sandstone comprise the dominant bedrock, while the western outer regions are

dominated by Mississippian limestone (Ozark Ecoregional Assessment Team 2003). All three watersheds have steep bluffs along streams, narrow valleys, and karst features, and many of their streams are spring-fed. Seasonal patterns of discharge are similar among all three streams (Fig. 2), but discharge in the Little Black River is much lower than the Gasconade and Meramec rivers because of its smaller watershed (990 km<sup>2</sup> and 7,268 km<sup>2</sup> for the Little Black and Gasconade rivers, respectively). We describe additional features of the Gasconade and Little Black rivers below; a description of the Meramec River basin can be found in Key et al. (2021).

*Gasconade River.*—The mainstem Gasconade River flows north for 436 river-km (rkm) before joining the Missouri River (Blanc 2001). Our habitat suitability models included about 800 rkm including the mainstem Gasconade River and three of its tributaries, Osage Fork, Big Piney River, and Roubidoux Creek (Fig. 1). These streams are not channelized or impounded, but in-channel gravel mining has altered and destabilized some segments (Blanc 2001), and decreased riparian vegetation has also contributed to channel instability and erosion (Jacobson and Primm 1997). Forty-six mussel species are reported from the Gasconade River basin (Blanc 2001).

*Little Black River.*—The mainstem Little Black River flows south 137 rkm into Arkansas before joining the Current River. Most of the Little Black River and its tributaries are within the Ozark Highlands, but the downstream portion of the mainstem flows through the Mississippi Alluvial Plains (Fig. 1; Wilkerson 2003). Because of the physiographic differences between the Ozark Highlands and Mississippi Alluvial Plains, we did not include that portion of the stream in our habitat suitability models. Our habitat suitability models included 120 rkm comprising the mainstem Little Black River and three of its tributaries, North and South prongs and Beaverdam Creek (Fig. 1). The Little Black River is highly altered with 13 impoundments and about 98 rkm of channelized streams (Wilkerson 2003). Thirty-nine mussel species are reported from the Little Black River basin (Wilkerson 2003).

### Mussel Survey Dataset

We determined mussel bed locations in the Gasconade and Little Black rivers from the Missouri Department of Conservation mussel database (data available upon request to and subject to the approval of the Missouri Department of Conservation, 3500 East Gans Road, Columbia, MO 65201). This database includes mussel survey information for specific locations across Missouri, including GPS points, survey methods, lists of species found, and numbers of individuals found. We used mussel survey data from 1994 to 2013, following the MRHSM (Key et al. 2021). We filtered the data to include only timed-search samples; incidental collections, collections using a groping technique, or entries with missing sampling method were excluded. We considered sites within 180 m of each other to represent the same mussel bed




Figure 1. Map of Missouri, USA, showing the Gasconade (pink), Little Black (dark pink), and Meramec (blue) river watersheds and major streams included in habitat suitability models. Other text and boundaries within the state boundary are Level IV ecoregions.

(Lueckenhoff 2015; Schrum 2017; Key et al. 2021). Our resulting dataset included 130 unique mussel bed locations. We selected a subset of 85 mussel beds that had the highest species richness (hereafter species-rich mussel beds; Key et al. 2021) to develop our habitat suitability models. The remaining 45 mussel beds (hereafter validation mussel beds) were used to validate our models by determining how many of these beds fell within habitat deemed suitable by our model.

### Generation of Hydrogeomorphic Variables

We derived 12 hydrogeomorphic variables for the Gasconade and Little Black rivers, including all 10 variables used in the MRHSM and two additional variables that we created (see below; Table 1). These variables represent habitat characteristics thought to correspond to suitable habitat for mussels at a riverscape scale, including bluff adjacency, presence of and proximity to gravel bars, lateral channel stability, low-flow surface water availability, and stream power index (Table 1; Key et al. 2021). Mussel beds in Ozark rivers are often found in the vicinity of bluffs, possibly because bluffs exert channel control and stabilization that is amenable to mussel establishment and persistence (Vannote and Minshall 1982; Key et al. 2021). Mussel beds often are

associated with gravel bars, and the presence of persistent gravel bars after high-flow events can indicate channel stability (Bates 1962; Peck 2005; Zigler et al. 2008; Key et al. 2021). Lateral channel movement is indicative of bank erosion and sediment deposition, which can destabilize substrate and limit mussel occurrence (Strayer 1999; Strayer et al. 2004). Low-flow surface water availability is intended to represent a proxy for the existence of permanently watered areas that serve as refugia during drought periods (Table 1; Golladay et al. 2004; Key et al. 2021). Stream power is an index of potential energy in the channel and influences channel erosion and stability.

We derived estimates for the hydrogeomorphic variables from high-resolution, open-source datasets of aerial imagery and topography following Key et al. (2021) and summarized as follows. We began our workflow by defining the stream dimensions and location of the river channel and subsequently creating a stream centerline. We then generated points on the stream centerline at 10-m cross sections to create a spatially continuous dataset. After we defined our stream dimensions and stream centerline, we derived our 12 hydrogeomorphic variables and assigned the data to each point on the stream centerline. The 10-m points were then interpolated using natural neighbors into continuous grids representing our final




Figure 2. (A) Mean monthly unit discharge ( $\text{ft}^3/\text{s}/\text{mi}^2$ ) and (B) mean monthly discharge ( $\text{ft}^3/\text{s}$ ) for the Meramec, Gasconade, and Little Black rivers. Flow data are from <https://waterdata.usgs.gov/mo/nwis/rt> (accessed January 23, 2023).

hydrogeomorphic variables. All spatial analyses were performed in ArcGIS and projected to NAD 1983 UTM zone 15N (ESRI 2011; Key et al. 2021).

We used a combination of 1-m light detection and ranging (LiDAR) and 10-m digital elevation model (DEM) coverage of both study areas (MSDIS 2011) to generate six hydrogeomorphic variables related to bluff adjacency and stream power. Two bluff adjacency variables represented whether a bluff is present within one channel width of each bank from a mussel bed (binary variable) and, if so, the total bluff area

(continuous variable). In addition to the bluff adjacency variables from the MRHSM, we generated two new variables representing the total bluff area within 500 m upstream or downstream of a mussel bed. We added these variables to explore whether bluffs located upstream or downstream, versus directly adjacent to the stream channel, are associated with mussel bed occurrence. Because of the limited availability of LiDAR for the Gasconade River, we used 10-m DEMs to extend the remotely sensed data across the entire drainage area. The 1-m horizontal resolution LiDAR tiles and 10-m

Table 1. Justification and description of hydrogeomorphic variables evaluated in habitat suitability models for the Meramec, Gasconade, and Little Black rivers. Variables that were included in the final, best-fit models differed among streams and transferability levels (see text). “Type” refers to whether the variable was continuous or binary.

Habitat Characteristic: Type	Justification	Description
Bluff adjacency area: continuous	Mussel beds are usually found in the vicinity of bluffs adjacent to the stream channel.	Total bluff area (m <sup>2</sup> ) within one channel width of each bank
Bluff adjacency: binary		Whether there is a bluff within one channel width of each bank
Longitudinal bluff adjacency area upstream: continuous	Stream power influences oxygen, food supply, successful host infestation, and offspring dispersal.	Total bluff area (m <sup>2</sup> ) within 500 m upstream
Longitudinal bluff adjacency area downstream: continuous		Total bluff area (m <sup>2</sup> ) within 500 m downstream
Stream power index: continuous		Index of potential energy of water in the channel, using $SPI = \ln(A_d) * S_{500}$
Stream power class: binary		Potential energy of water in the stream channel, classed as either high or low using the mean
Lateral channel stability: binary	Lateral channel movement can disrupt habitat condition.	Lateral channel movement of > 10 m between 1990–95 and 2015, classed as unstable, all else classed as stable
Gravel/pool class: binary	Reaches with persistent gravel bars can indicate in-stream stability after high-flow events. In smaller streams, however, they can also indicate reaches that dry during low-flow events.	Reaches dominated by gravel are classed at gravel, all else classed as pool reaches
Gravel bar proximity: binary		All locations within 100 m of a gravel bar are classed as adjacent to a gravel bar
Distance to gravel bar: continuous	Refuge during drought periods is necessary for mussel survival.	Euclidean distance (m) to nearest gravel bar
Low-flow surface water availability index: continuous		The number of water pixels surrounding each cell
Low-flow surface water availability class: binary		The number of water pixels surrounding each cell, classed as high or low using the mean

DEMs were mosaicked into a single DEM and resampled to 10-m resolution for analysis (Key et al. 2021). We then used a slope and range criteria to define bluffs in the watersheds and performed a zonal search at each point on the stream centerline.

For the stream power variables, we estimated the watershed area at each stream centerline point and estimated the stream power index (SPI) using the bankfull elevation (see Key et al. 2021 for methods to determine bankfull elevation) as  $SPI = S \times \ln(A_d)$ , where, for any location along the stream centerline, SPI is the stream power index,  $S$  is the channel slope, and  $A_d$  is the watershed area (Moore et al. 1991). Slope was averaged over a 500-m interval, 250 m upstream and 250 m downstream from each point on the stream centerline, and then smoothed using a 50-m moving average. The binary stream power variable was derived by classifying each pixel as either high or low stream power using the mean value as the break between the two classes.

We used National Agriculture Imagery Program leaf-off aerial imagery to generate the six remaining hydrogeomorphic variables related to gravel bars, lateral channel stability, and low-flow surface water availability. We derived three variables reflecting the presence of or proximity to gravel bars for each

mussel bed (Table 1). We classified mussel bed locations as “gravel” if the stream reach was dominated by persistent gravel before and after a high-flow year, and as “pool” if the reach was dominated by water (binary variable). We then determined whether each mussel bed was located within 100 m of a gravel bar (binary variable), and we created a continuous variable representing the Euclidean distance from a mussel bed to the nearest gravel bar. We derived these variables with a differencing technique between two sets of aerial imagery. In the Little Black River, 2007 and 2015 were low-water years, while 2013 was a high-water year. In the Gasconade River, 2012 and 2014 were low-water years, and 2013 was a high-water year. Pixels that changed state (water or gravel) between the two images were classified as a pool, and pixels that did not change state were classified as either gravel or a pool. The gravel/pool class therefore does not represent the underlying sediment (gravel versus depositional sediments) but rather areas that had persistently exposed gravel bar versus areas that were predominantly water during low-flow conditions. Without ground-truthing, we cannot differentiate whether the areas classified as pools had gravel or depositional sediments.

For the lateral channel stability variable, we created two polygons representing the stream banks based on visual cues

such as shadow, vegetation, and scour lines in the leaf-off imagery from 1990–95 and 2015. We defined each point on the stream centerline as unstable if the channel moved  $> 10$  m between the two time periods or stable if the channel moved  $\leq 10$  m (binary variable; Key et al. 2021).

We derived two variables for low-flow surface water availability index using imagery taken at the time of lowest discharge available (2007 for the Little Black River and 2012 for the Gasconade River). We performed a focal search to estimate the number of pixels classified as water that were adjacent to the focal pixel (continuous variable). We then used the median value to categorize high- and low-surface water availability in a low-flow period to create the binary variable. We acknowledge that these variables do not directly represent vulnerability to drying because water depth is not accounted for (see Key et al. 2021). However, we used these variables as proxies for drought refugia because bathymetric data were not available from our imagery.

### Habitat Suitability Models

We used maximum entropy modeling (MaxEnt; Phillips and Dudík 2008) to generate habitat suitability models for mussels in the Meramec, Gasconade, and Little Black rivers. This method uses presence-only occurrence data in combination with environmental data layers to produce a model of habitat suitability spanning a specified geographic area (Phillips and Dudík 2008). For models that included the Meramec River, we used the presence-only occurrence data and environmental layers created by Key et al. (2021). We spatially constricted our habitat suitability models to each drainage and used the location of species-rich mussel beds in each river system in combination with the hydrogeomorphic variables determined for those locations. We used the same settings in MaxEnt as used for the MRHSM (Key et al. 2021). Specifically, we set the run type to bootstrap to generate training and test occurrence data (80% and 20% of the species-rich mussel bed locations, respectively), and we ran models with 10,000 background points and 5,000 iterations. None of the hydrogeomorphic variables included in each model were correlated with each other (correlation coefficient  $< 0.40$ ).

We converted the raw model results to a binary map of suitable and unsuitable reaches based on the equal test sensitivity and specificity logistic threshold of each model (Key et al. 2021). The equal test sensitivity and specificity logistic threshold is a commonly used threshold that sets the sensitivity equal to the specificity (Cao et al. 2013; Phillips 2017; Key et al. 2021). After suitable and unsuitable reaches were delineated, we used a buffer of 40 m to separate the suitable and unsuitable habitats to account for areas of transition (following Key et al. 2021). We then used jackknife analysis and the test gain values to assess the relative contribution of each hydrogeomorphic variable and to determine which variables were most important for model fit (Phillips 2017; Key et al. 2021). We used a stepwise model selection approach of our hydrogeomorphic variables to select

the best-fit model. The area under the receiver operating curve (AUC) values from MaxEnt provided relative values for comparing the performance of models that were built with the same data (Phillips et al. 2006). Therefore, we selected variables that led to higher AUC values and contained hydrogeomorphic variables with sizeable individual effects on model results when others were removed (following Elith 2002 and Key et al. 2021). We also created response curves to investigate the relationships between suitable and unsuitable habitats (y-axis) and our hydrogeomorphic variables (x-axis). The results for our continuous hydrogeomorphic variables were presented as curves spanning the range of values for that layer, whereas the binary hydrogeomorphic variables were presented as two bars representing the binary. The range in values of the continuous—or bars of the binary—hydrogeomorphic variables were classified as suitable if they were equal to or above the equal test sensitivity and specificity logistic threshold on the response curves. For low-flow surface water availability and stream power, higher values in the response curves represented more contiguous surface water availability during low-flow conditions and higher stream power, respectively. Although AUC values provided comparisons of model performance, they did not provide a measure of the accuracy of habitat suitability (Jiménez-Valverde and Lobo 2006). Therefore, we used the location of the validation mussel beds (not used in model development) to assess the accuracy of our best-fit models. We calculated the percentage of the validation mussel beds that fell within a reach predicted to be suitable by the best-fit models across the entire spatial extent. For models that included the Meramec and Gasconade rivers or the Meramec and Little Black rivers, we also calculated validation per drainage as the proportion of validation mussel bed locations that fell within a reach predicted to be suitable for each drainage, separately.

### Transferability

We separated our methods of transferability into three categories representing different levels of dependence on the original MRHSM: Level 1, transferring the original model; Level 2, transferring the modeling framework; and Level 3, adapting the modeling framework (Fig. 3).

*Level 1: Transferring the original model.*—To transfer the original model from the Meramec River to the Gasconade and Little Black rivers, we utilized the species-rich mussel bed locations and hydrogeomorphic variables used in the MRHSM (Key et al. 2021) and species-rich mussel bed locations and hydrogeomorphic variables that we derived for the Gasconade and Little Black rivers. Specifically, we combined the species-rich mussel bed locations and hydrogeomorphic variables to include the spatial extent of both the Meramec and Gasconade rivers or the Meramec and Little Black rivers. We started this level of transferability with all 10 hydrogeomorphic variables from the MRHSM (Fig. 3). We then used the jackknife analysis and stepwise model selection approach as described previously to find the best-fit model. The results from the best-




Figure 3. General framework used to test the transferability of the Meramec River habitat suitability model to the Gasconade and Little Black rivers, including three levels of transferability and the spatial extent, required data, and evaluation criteria for each level.



Table 2. Results for the Meramec River habitat suitability model (Key et al. 2021) and evaluation of transferability of that model to the Gasconade and Little Black rivers. All models are best-fit models for each river and level of transferability. We considered model transfer successful if the best-fit model had a test AUC  $\geq$  0.70 and total validation  $\geq$  0.70. Equal test sensitivity and specificity logistic threshold is the value used to delineate suitable and unsuitable habitats for all variables in each model. Total validation is the proportion of validation mussel bed locations that were identified by the model as suitable habitat across all rivers in the model. Validation/drainage is the proportion of validation mussel-bed locations identified as suitable habitat within each river.

River(s)	Level of Transferability	Test AUC	Equal Test Sensitivity and Specificity Logistic Threshold	Total Validation	Validation/ Drainage
Meramec River	Original model	0.62	0.45	0.90	—
Meramec and Gasconade rivers	Level 1	0.69	0.41	0.68	Meramec: 1.00 Gasconade: 0.24
Gasconade River	Level 2	0.70	0.34	0.82	—
Meramec and Little Black rivers	Level 1	0.64	0.42	0.64	Meramec: 0.65 Little Black: 0.60
Little Black River	Level 2	0.74	0.48	0.60	—
Little Black River	Level 3	0.72	0.44	0.80	—

fit model were converted to a binary map and validated using the validation mussel bed locations. We considered model transfer to be successful if the best-fit model had a test AUC  $\geq$  0.70 and total validation  $\geq$  0.70 (Fig. 3). If either the test AUC or model validation was  $<$  0.70, we progressed to Level 2 (Fig. 3).

*Level 2: Transferring the modeling framework.*—To reduce our dependence on the MRHSM, we transferred only the modeling framework by building our MaxEnt models without data from the Meramec River. Specifically, we used only species-rich mussel bed locations and 10 hydrogeomorphic variables in either the Gasconade River or Little Black River (Fig. 3). Following the same methodology as Level 1, we determined the best-fit model and validated those results with the validation mussel bed locations. Again, if the best-fit model had a test AUC  $\geq$  0.70 and total validation  $\geq$  0.70, it was considered a successful transfer of the modeling framework. If either the test AUC or model validation was  $<$  0.70, we progressed to Level 3 (Fig. 3).

*Level 3: Adapting the model.*—If neither of the previous transferability methods produced an adequate model, we adapted the modeling framework used in the MRHSM by including two additional hydrogeomorphic variables, longitudinal bluff adjacency upstream and downstream. Similar to Level 2, models were built only with the species-rich mussel bed locations and hydrogeomorphic variables from the Gasconade or Little Black rivers. We started this level with the 10 original hydrogeomorphic variables plus the two additional bluff adjacency variables that we created. We followed the same stepwise model selection approach as Levels 1 and 2 to find the best-fit model and then created the binary suitability map and validated the results with the locations of the validation mussel beds. If the best-fit model had a test AUC  $\geq$  0.70 and total validation  $\geq$  0.70, we considered this a successful adaptation to the modeling framework (Fig. 3). If either the test AUC or total validation was  $<$  0.70, we considered model transfer unsuccessful (Fig. 3).

## RESULTS

### Gasconade River

*Level 1: Transferring the original model.*—The best-fit habitat suitability model for Level 1 had a test AUC of 0.69 (Table 2). The best-fit model included six hydrogeomorphic variables: lateral channel stability, distance to gravel bar, gravel/pool class, stream power index, bluff adjacency area, and low-flow surface water availability index. Jackknife analysis indicated that bluff adjacency area, distance to gravel bar, gravel/pool class, and lateral channel stability contributed significantly to the final model (Table 3).

An equal test sensitivity and specificity logistic threshold of 0.41 separated habitats into suitable and unsuitable. Response curves indicated that suitable habitat was represented by reaches with small bluffs, 0–700 m or  $>$  1,500 m from gravel bars, areas with a low-flow surface water availability index  $>$  0 but  $<$  10, and intermediate stream power. While 68% of our validation mussel bed locations were found in areas identified as suitable in both the Meramec and Gasconade rivers, only 24% of the validation mussel beds were found in areas identified as suitable in the Gasconade River alone. Because the test AUC and total validation were  $<$  0.70, we considered model transfer to the Gasconade River unsuccessful at this level and continued to Level 2 (Table 2).

*Level 2: Transferring the modeling framework.*—The best-fit habitat suitability model for Level 2 had a test AUC of 0.82 (Table 2). The best-fit model included six hydrogeomorphic variables: lateral channel stability, distance to gravel bar, gravel/pool class, stream power index, bluff adjacency area, and low-flow surface water availability index. Jackknife analysis indicated that bluff adjacency area, distance to gravel bars, gravel/pool class, and low-flow surface water availability contributed significantly to the model (Table 3).

An equal test sensitivity and specificity logistic threshold of 0.34 separated habitats into suitable and unsuitable. Response curves indicated that suitable habitat was represent-

Table 3. Results of the jackknife analyses for the final, best-fit models of the original Meramec River habitat suitability model and transfer of that model to the Gasconade River and Little Black rivers. An asterisk (\*) indicates hydrogeomorphic variables that contributed significantly to the model based on the jackknife analyses.

Variable	Gasconade River	Meramec River	Little Black River
Bluff adjacency area	Near small bluffs*	Near small bluffs	—
Distance to gravel bar	Farther than 1,250 m*	Less than 400 m*	Any distance outside the reach*
Gravel/pool class	Pool*	—	Pool*
Low-flow surface water availability index	Greater than 0*	Greater than 3*	Greater than 7*
Stream power index	Greater than 0	Greater than 0*	Less than 0.05*
Lateral channel stability	Laterally stable	Laterally stable*	—
Bluff adjacency area downstream	—	—	Any amount of bluff area downstream*

ed by reaches classified as pools, near small bluffs, 100–300 m or > 1,250 m from persistent gravel bars, with low-flow surface water availability and stream power indices > 0, and laterally stable channels (Fig. 4 and Table 3). Because the test AUC was > 0.70 and total validation was 0.82, we concluded that transfer of the modeling framework to the Gasconade River was successful, and we did not evaluate Level 3 transferability (Table 2).

### Little Black River

*Level 1: Transferring the original model.*—The best-fit habitat suitability model for Level 1 had a test AUC of 0.64 (Table 2). The best-fit model included the same six hydrogeomorphic variables as for the Gasconade River: lateral channel stability, distance to gravel bar, gravel/pool class, stream power index, bluff adjacency area, and low-flow




Figure 4. Response curves for hydrogeomorphic variables that contributed significantly to transferring the Meramec River modeling framework to the Gasconade River at Level 2. The dashed line represents the equal sensitivity and specificity logistic threshold used to delineate suitable and unsuitable habitats.

surface water availability index. Jackknife analysis indicated that distance to gravel bar, gravel/pool class, low-flow surface water availability index, stream power index, and lateral channel stability contributed significantly to the final model.

An equal test sensitivity and specificity logistic threshold of 0.42 separated habitats into suitable and unsuitable. Response curves indicated that suitable habitat was represented by reaches near small bluffs, 0–500 m or > 2,000 m from gravel bars, with higher low-flow surface water availability, low-intermediate stream power indices, and laterally unstable channels. Sixty-four percent of validation mussel bed locations were found in areas identified as suitable by the model, and validation in the Little Black River was 60% (Table 2). Because the test AUC and total validation were < 0.70, we considered model transfer to the Little Black River unsuccessful at this level and continued to Level 2.

*Level 2: Transferring the modeling framework.*—The best-fit habitat suitability model for Level 2 had a test AUC of 0.74 (Table 2). The best-fit model included the same six hydrogeomorphic variables as Level 1. Jackknife analysis indicated that distance to gravel bar, gravel/pool class, stream power index, and lateral channel stability contributed significantly to the final model.

An equal test sensitivity and specificity logistic threshold of 0.48 separated habitats into suitable and unsuitable. Response curves indicated that suitable habitat was represented by reaches near small bluffs, with high water availability, lower stream power indices, and areas classified as pools. The test AUC was > 0.70, but because total validation was only 60%, we concluded that transfer of the modeling framework to the Little Black River was unsuccessful and continued to Level 3 (Table 2).

*Level 3: Adapting the model.*—The best-fit model for Level 3 had a test AUC of 0.72 (Table 2). The best-fit model included downstream bluff adjacency area, distance to gravel bar, gravel/pool class, low-flow surface water availability, and stream power index. Jackknife analysis indicated that downstream bluff adjacency area, gravel/pool class, low-flow surface water availability, and stream power index contributed significantly to the final model (Table 3).

An equal test sensitivity and specificity logistic threshold of 0.44 separated habitats into suitable and unsuitable. Response curves indicated that suitable habitat was represented by reaches classified as pools, with higher surface water availability and lower stream power indices (Fig. 5 and Table 3). Suitable habitat also was represented by reaches with any amount of downstream bluff area and persistent gravel bars at any distance. The best-fit model at this level had a total validation of 0.80 (Table 2). Because the test AUC was > 0.70 and total validation was 0.80, we concluded that transfer of the model to the Little Black River at this level was successful.

## DISCUSSION

Our study successfully identified suitable habitat for freshwater mussels in the Gasconade and Little Black rivers.

Mussel beds in both rivers were associated with reaches classified as pools based on the absence of exposed gravel bars. In the Gasconade River, laterally stable reaches near small bluffs, with gravel bars farther than 1,250 m away and higher stream power indices, were considered more suitable. In the Little Black River, suitable habitat was related to reaches with higher surface water availability during low-flow conditions, lower stream power indices, and bluffs located downstream.

In the MRHSM, distance to gravel bars, low-flow surface water availability index, stream power index, and lateral channel stability contributed significantly to the final model based on jackknife analysis (Table 3; Key et al. 2021). An equal test sensitivity and specificity logistic threshold of 0.45 separated habitats into suitable and unsuitable. Based on the response curves, locations identified as suitable were in reaches close to small bluffs, near persistent gravel bars, with higher stream power indices, laterally stable channels, and in reaches with greater low-flow surface water availability (Key et al. 2021).

The similarity of some features of our models to the original MRHSM suggests that mussel beds in the Gasconade and Little Black rivers are associated with some of the same habitat characteristics as those in the Meramec River. This is not surprising because the Meramec, Gasconade, and Little Black rivers all are in the Ozark Highlands ecoregion and share similar physiographic and watershed features. Most conspicuously, the best-fit habitat suitability model for all three rivers included a hydrogeomorphic variable representing bluff adjacency. Response curves show similar trends in increased habitat suitability associated with smaller bluffs (in relation to the amount of bluff area in the system) and decreased habitat suitability associated with larger bluffs. While bluffs can exert lateral channel control (Vannote and Minshall 1982), larger bluffs could reduce flow and sediment transport causing areas of unstable gravel deposition (Jacobson and Gran 1999; Owen et al. 2011). The best-fit model for the Little Black River included the bluff area downstream of mussel beds, but we do not necessarily know how downstream bluffs may influence channel stability or other habitat features.

Differences in other aspects of our models between all three rivers suggest that factors associated with mussel bed location differ according to watershed characteristics specific to each system. We were unable to transfer the MRHSM to either river at Level 1, which shows that direct transfer of the MRHSM was not possible. We were able to transfer the MRHSM to the Gasconade River at Level 2 and able to adapt it to the Little Black River at Level 3. Our results suggest that the unique features of each watershed affect model transferability, and additional variables (e.g., downstream bluff adjacency) may be needed to predict mussel occurrence in some streams.

Although the Meramec, Gasconade, and Little Black river watersheds share features characteristic of the Ozark Highlands ecoregion, each stream has unique features that may influence mussel bed habitat associations. Stream drying is an important factor in the disturbance regime of many rivers in




Figure 5. Response curves for hydrogeomorphic variables that contributed significantly to adapting the Meramec River modeling framework to the Little Black River at Level 3. The dashed line represents the equal sensitivity and specificity logistic threshold used to delineate suitable and unsuitable habitats.

the Ozark Highlands (Lynch et al. 2018). We observed numerous large, dry stretches of streambed during low-flow conditions in the aerial imagery for both the Gasconade and Little Black rivers. In contrast, we did not observe streambed drying throughout the Meramec River, even during low-flow conditions. The association of mussel beds in the Gasconade and Little Black rivers with reaches classified as pools, at greater distances from gravel bars, and with higher surface water availability may indicate mussel occurrence in reaches that are less prone to drying during drought (Gagnon et al. 2004; Haag and Warren 2008; Atkinson et al. 2014).

Our use of remotely sensed, large-scale hydrogeomorphic data instead of direct measurements of stream habitat characteristics affects the interpretation of our habitat suitability models. We showed an association of mussel beds with areas classified as pools. However, our aerial imagery did not provide bathymetric or flow data necessary to differentiate between low-flow, depositional pools and gravel-bottomed runs with no exposed gravel. Typically, mussel beds do not occur in depositional pools, but gravel-bottomed runs can be optimal mussel habitat (Vannote and Minshall 1982; Vaughn

and Taylor 1999). Similarly, without bathymetric data we cannot fully evaluate the extent to which low-flow surface water availability represents vulnerability to emersion during drought. Nevertheless, these variables provide useful information with which to broadly characterize reaches that support mussel beds in our study streams.

Many other factors that influence mussel presence were not included in our model, including species-specific differences in habitat requirements, anthropogenic factors, and fish-host relationships. However, at the riverscape scale, our hydrogeomorphic variables can identify broad habitat characteristics necessary to support mussels. By providing longitudinally continuous characterization of habitat suitability at the riverscape scale, our models provide a baseline that can allow evaluation of the effects of other factors on mussel occurrence (Bouska et al. 2018; Key et al. 2021).

#### ACKNOWLEDGMENTS

Data on freshwater mussel distributions were provided by the Missouri Department of Conservation (MDC) and

Matthew Schrum, who is working on a related project in the Little Black River with support from the MDC. Funding for this research was provided by the MDC, Tennessee Technological University Water Center, and Tennessee Technological University Department of Biology. As data were acquired from existing sources, no federal, state, or Animal Care permits were required for completion of this study. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. We also would like to thank Kristin Womble for editing the manuscript prior to submission, as well as an anonymous reviewer and Wendell Haag for constructive reviews of the manuscript.

### LITERATURE CITED

- Allen, D. C., and C. C. Vaughn. 2010. Complex hydraulic and substrate variables limit freshwater mussel species richness and abundance. *Journal of the North American Benthological Society* 29:383–394.
- Arbuckle, K. E., and J. A. Downing. 2002. Freshwater mussel abundance and species richness; GIS relationships with watershed land use and geology. *Canadian Journal of Fisheries and Aquatic Sciences* 59:310–316.
- Atkinson, C. L., J. P. Julian, and C. C. Vaughn. 2014. Species and function lost: Role of drought in structuring stream communities. *Biological Conservation* 176:30–38.
- Barbosa, A. M., R. Real, and J. Mario Vargas. 2009. Transferability of environmental favourability models in geographic space: The case of the Iberian desman (*Galemys pyrenaicus*) in Portugal and Spain. *Ecological Modelling* 220:747–754.
- Bates, J. M. 1962. The impact of impoundment on the mussel fauna of Kentucky Reservoir, Tennessee River. *American Midland Naturalist* 68:232–236.
- Blanc, T. J. 2001. Gasconade River watershed inventory and assessment. Missouri Department of Conservation Fisheries Division, Jefferson City. Available at [https://mdc.mo.gov/sites/default/files/2021-12/130\\_2021\\_GasconadeRiver.pdf](https://mdc.mo.gov/sites/default/files/2021-12/130_2021_GasconadeRiver.pdf) (accessed January 31, 2023).
- Bouska, K. L., A. Rosenberger, S. E. McMurray, G. A. Lindner, and K. N. Key. 2018. State-level freshwater mussel programs: Current status and a research framework to aid in mussel management and conservation. *Fisheries* 43:345–360.
- Cao, Y., E. R. DeWalt, J. L. Robinson, T. Tweddale, L. Hinz, and M. Pessino. 2013. Using Maxent to model the historic distributions of stonefly species in Illinois streams: The effects of regularization and threshold selections. *Ecological Modelling* 259:30–39.
- Daniel, W., and K. M. Brown. 2014. The role of life history and behavior in explaining unionid mussel distributions. *Hydrobiologia* 734:57–68.
- Drew, C. A., M. Eddy, T. J. Kwak, W. G. Cope, and T. Augspurger. 2018. Hydrologic characteristics of freshwater mussel habitat: novel insights from modeled flows. *Freshwater Science* 37:343–356.
- Elith, J. 2002. Quantitative methods for modeling species habitat: comparative performance and an application to Australian plants. Pages 39–59 in S. Ferson and M. Burgman, editors. *Quantitative Methods for Conservation Biology*. Springer, New York.
- ESRI (Environmental Systems Research Institute). 2011. ArcGIS Desktop: Release 10.8. Environmental Systems Research Institute, Redlands, CA.
- Fausch, K. D., C. E. Torgersen, C. V. Baxter, and H. W. Li. 2002. Landscapes to riverscapes: Bridging the gap between research and conservation of stream fishes. *BioScience* 52:483–498.
- FMCS (Freshwater Mollusk Conservation Society). 2016. A national strategy for the conservation of native freshwater mollusks. *Freshwater Mollusk Biology and Conservation* 19:1–21.
- Gagnon, P. M., S. W. Golladay, W. K. Michener, and M. C. Freeman. 2004. Drought responses of freshwater mussels (Unionidae) in Coastal Plain tributaries of the Flint River Basin, Georgia. *Journal of Freshwater Ecology* 19:667–679.
- Golladay, S. W., P. Gagnon, M. Kerans, J. M. Battle, and D. W. Hicks. 2004. Response of freshwater assemblages (Bivalvia: Unionidae) to a record drought in the Gulf Coastal Plain of southwestern Georgia. *Journal of the North American Benthological Society* 23:494–506.
- Haag, W. R., and M. L. Warren. 2008. Effects of severe drought on freshwater mussel assemblages. *Transactions of the American Fisheries Society* 137:1165–1178.
- Jacobson, R. B., and K. B. Gran. 1999. Gravel sediment routing from widespread, low-intensity landscape disturbance, Current River Basin, Missouri. *Earth Surface Processes and Landforms* 24:897–917.
- Jacobson, R. B., and A. T. Primm. 1997. Historical land-use changes and potential effects on stream disturbance in the Ozark Plateaus, Missouri. US Geological Survey. Available at <https://pubs.usgs.gov/wsp/2484/report.pdf> (accessed January 31, 2023).
- Jiménez-Valverde, A., and J. M. Lobo. 2006. The ghost of unbalanced species distribution data in geographical model predictions. *Diversity and Distributions* 12:521–524.
- Johnson, P. D., and K. M. Brown. 2000. The importance of microhabitat factors and habitat stability to the threatened Louisiana pearl shell, *Margaritifera hembeli* (Conrad). *Canadian Journal of Zoology* 78:271–277.
- Key, K. N., A. E. Rosenberger, G. A. Lindner, K. Bouska, and S. E. McMurray. 2021. Riverscape-scale modeling of fundamentally suitable habitat for mussel assemblages in an Ozark River System, Missouri. *Freshwater Mollusk Biology and Conservation* 24:43–58.
- Lueckenhoff, L. 2015. Development of standardized visual sampling methods for assessing community metrics of unionoid mussel species and tribal groups in Missouri. Master's thesis, University of Missouri, Columbia.
- Lynch, D. T., D. R. Leasure, and D. D. Magoulick. 2018. The influence of drought on flow-ecology relationships in Ozark Highland streams. *Freshwater Biology* 63:946–968.
- Moore, I. D., R. B. Grayson, and A. R. Ladson. 1991. Digital terrain modelling: A review of hydrological, geomorphological, and biological applications. *Hydrological Processes* 5:3–30.
- MSDIS (Missouri Spatial Data Information Service). 2011. Curators of the University of Missouri-Columbia. Available at <http://msdis.missouri.edu> (accessed January 31, 2023).
- Owen, M. R., R. T. Pavlowsky, and P. J. Womble. 2011. Historical disturbance and contemporary floodplain development along an Ozark river, southwest Missouri. *Physical Geography* 32:423–444.
- Ozark Ecoregional Assessment Team. 2003. Ozarks ecoregional conservation assessment. The Nature Conservancy, Midwestern Resource Office, Minneapolis, Minnesota.
- Pandolfo, T. J., T. J. Kwak, and W. G. Cope. 2016. Microhabitat suitability and niche breadth of common and imperiled Atlantic Slope freshwater mussels. *Freshwater Mollusk Biology and Conservation* 19:27–50.
- Peck, A. J. 2005. A reach scale comparison of fluvial geomorphological conditions between current and historic freshwater mussel beds in the White River, Arkansas. Master's thesis, Arkansas State University, Jonesboro.
- Phillips, S. J. 2017. A brief tutorial on MaxEnt. Available at [http://biodiversityinformatics.amnh.org/open\\_source/maxent/](http://biodiversityinformatics.amnh.org/open_source/maxent/) (accessed January 31, 2023).
- Phillips, S. J., R. P. Anderson, and R. E. Schapire. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190:231–259.
- Phillips, S. J., and M. Dudik. 2008. Modeling of species distributions with MaxEnt: New extensions and a comprehensive evaluation. *Ecography* 31:161–175.
- Randin, C. F., T. Dirnböck, S. Dullinger, N. E. Zimmermann, M. Zappa, and

- A. Guisan. 2006. Are niche-based species distribution models transferable in space? *Journal of Biogeography* 33:1689–1703.
- Randklev, C. R., M. A. Hart, J. M. Khan, E. T. Tsakiris, and C. R. Robertson. 2019. Hydraulic requirements of freshwater mussels (Unionidae) and a conceptual framework for how they respond to high flows. *Ecosphere* 10:e02975. doi: 10.1002/ecs2.297
- Schlosser, I. J., and P. L. Angermeier. 1995. Spatial variation in demographic processes of lotic fishes: Conceptual models, empirical evidence, and implications for conservation. Pages 392–401 in J. L. Nielsen, editor. *Evolution and the Aquatic Ecosystem symposia proceedings, Evolution and the Aquatic Ecosystem: Defining Unique Units in Population Conservation*. American Fisheries Society, New York.
- Schrum, M. C. 2017. Development and validation of standardized sampling protocols for assessing freshwater mussel populations in Missouri. Master's thesis. University of Missouri, Columbia.
- Strayer, D. L. 1993. Macrohabitats of freshwater mussels (Bivalvia:Unionacea) in streams of the Northern Atlantic Slope. *Journal of the North American Benthological Society* 12:236–246.
- Strayer, D. L. 1999. Use of flow refuges by Unionid mussels in rivers. *Journal of the North American Benthological Society* 18:468–476.
- Strayer, D. L., J. A. Downing, W. R. Haag, T. L. King, J. B. Layzer, T. J. Newton, and S. Jerrine Nichols. 2004. Changing perspectives on pearly mussels, North America's most imperiled animals. *Bioscience* 54:429–439.
- Strayer, D. L., and J. Ralley. 1993. Microhabitat use by an assemblage of stream-dwelling Unionaceans (Bivalvia), including two rare species of Alasmidonta. *Journal of the North American Benthological Society* 12:247–258.
- Vannote, R. L., and G. W. Minshall. 1982. Fluvial processes and local lithology controlling abundance, structure, and composition of mussel beds. *Proceedings of the National Academy of Sciences* 79:4103–4107.
- Vaughn, C. C. 1997. Regional patterns of mussel species distributions in North American rivers. *Ecography* 20:107–115.
- Vaughn, C. C. 2012. Life history traits and abundance can predict local colonisation and extinction rates of freshwater mussels. *Freshwater Biology* 57:982–992.
- Vaughn, C. C., and C. M. Taylor. 1999. Impoundments and the decline of freshwater mussels: A case study of an extinction gradient. *Conservation Biology* 13:912–920.
- Walters, A. D., D. Ford, E. R. Chong, M. G. Williams, N. B. Ford, L. R. Williams, and J. A. Banta. 2017. High-resolution ecological niche modeling of threatened freshwater mussels in east Texas, USA. *Aquatic Conservation: Marine and Freshwater Ecosystems* 27:1251–1260.
- Werkowska, W., A. L. Márquez, R. Real, and P. Acevedo. 2016. A practical overview of transferability in species distribution modeling. *Environmental Reviews* 25:127–133.
- Wilkerson, T. F. 2003. Current River watershed inventory and assessment. Missouri Department of Conservation Fisheries Division, Jefferson City. Available at <https://mdc.mo.gov/sites/default/files/mdcd7/downloads/page/080Current%20River.pdf> (accessed January 31, 2023).
- Zigler, S. J., T. J. Newton, J. J. Steuer, M. R. Bartsch, and J. S. Sauer. 2008. Importance of physical and hydraulic characteristics to unionid mussels: A retrospective analysis in a reach of large river. *Hydrobiologia* 598:343–360.