

Genetic Structure of Faucet Snail, *Bithynia tentaculata* Populations in North America, Based on Microsatellite Markers

Authors: Perez, Kathryn E., Werren, Rebecca L., Lynum, Christopher A., Hartman, Levi A., Majores, Gabor, et al.

Source: Freshwater Mollusk Biology and Conservation, 19(2) : 56-68

Published By: Freshwater Mollusk Conservation Society

URL: <https://doi.org/10.31931/fmbc.v19i2.2016.56-68>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REGULAR ARTICLE

GENETIC STRUCTURE OF FAUCET SNAIL, *BITHYNIA TENTACULATA* POPULATIONS IN NORTH AMERICA, BASED ON MICROSATELLITE MARKERS

Kathryn E. Perez^{1*}, Rebecca L. Werren², Christopher A. Lynum³, Levi A. Hartman², Gabor Majoros⁴, and Rebecca A. Cole⁵

¹ Department of Biology, University of Texas Rio Grande Valley, 1201 W. University Dr., Edinburg, TX 78539 USA

² Department of Biology, University of Wisconsin-La Crosse, 1725 State St., La Crosse, WI 54601 USA

³ Department of Biology, University of Massachusetts, Boston, 100 Morrissey Blvd, Boston, MA 02125 USA

⁴ Department of Parasitology and Zoology, Faculty of Veterinary Science, Szent István University Budapest, P.O.B. 2. H-1400, Hungary

⁵ United States Geological Survey, National Wildlife Health Center, 6006 Schroeder Rd, Madison, WI 53711 USA

ABSTRACT

Bithynia tentaculata is believed to have been extirpated from North America during the last glacial maximum. It was reintroduced into North America via the Great Lakes basin in the 1800's and has recently been expanding its geographic range. This snail serves as intermediate host for three trematodes that cause extensive recurring morbidity and mortality events in migratory water birds along the Mississippi River. Using twelve microsatellite loci for ~200 individual snails from 11 populations in North America and Europe, we examined one of the three major geographic regions from which founding populations into the Great Lakes typically originate. Our data supports a single recolonization of North America into the Great Lakes Basin followed by subsequent introduction events from the Great Lakes to other large watersheds in North America. However, additional watersheds in Europe require sampling to confirm this result. No populations with genetic signatures indicative of North American glacial relics were found. The initial invasion of North America was likely not from the Ponto-Caspian basin, the usual source of freshwater invasive species to the Laurentian Great Lakes.

KEYWORDS - faucet snail, phylogeography, invasive species, Mississippi River

INTRODUCTION

The Laurentian Great Lakes of North America have been a hotspot for invasion by exotic species. Many ecologically damaging aquatic invasive species have been introduced into the United States (U.S.) via this route (Mills et al. 1993). Molecular data have been used to determine the source of invasion of various aquatic invaders. For example, using the

mitochondrial cytochrome oxidase I gene, Gelembiuk et al. (2006) concluded that the source of invasion of zebra (*Dreissena polymorpha*) and quagga mussels (*D. bugensis*) into the Great Lakes was the Ponto-Caspian Sea basin (the Black, Caspian, and Azov Seas and their surrounding watersheds). This is congruent with other studies that have shown that the Ponto-Caspian Sea basin has been an important source of many aquatic invaders into the Great Lakes (Lee & Bell 1999, Ricciardi & MacIsaac 2000). Up to 70% of recent invaders in the Great Lakes (1985-2000) trace their source

*Corresponding Author: perezke@gmail.com

population to this region (Ricciardi & MacIsaac 2000, Brown & Stepien 2009, Keller et al. 2010). However, this is not the only possible source of aquatic invasive species. Shipping routes from European waters to the Great Lakes commonly originate in three locations: (1) the Black/Mediterranean Seas, (2) the North Sea, (3) or the Baltic Sea (Ricciardi & MacIsaac 2000, Grigorovich et al. 2003, Brown & Stepien 2009). These usual source regions for invaders into the Great Lakes provide an excellent starting point for comparing the genetic structure of invasive and potential source populations.

Bithynia tentaculata (L., 1758) had a Holarctic distribution prior to the last glacial maximum with shells found in Pleistocene fossil deposits from Lake Michigan, Illinois, U.S.A (Baker 1928). It is believed to have been extirpated from North America by glaciation events with subsequent recolonization through human-mediated introduction. Following the last glacial maximum, the first North American record of *B. tentaculata* was in Lake Michigan (presumably by passage through the Great Lakes Waterway, via the Hudson River) in 1871 (Baker 1928, Mills et al. 1993). It was speculated at that time that the snail was carried into Lake Michigan through ballast of timber ships arriving from Europe (Baker 1928). The species then spread throughout the Great Lakes region and into other U.S. waterways. It is now widespread in the Great Lakes, Northern Atlantic Coast drainages, isolated lakes in Montana, and most recently, in the Upper Mississippi River and Wolf River drainages, WI (Sauer et al. 2007)

Since the snail's introduction into the Mississippi River, first recorded in 2002 (National Wildlife Health Center, Madison, WI, unpublished data), parasites carried by *B. tentaculata* have caused recurring morbidity and mortality events in water bird populations during spring and fall migrations. The intestinal trematodes *Cyathocotyle bushiensis*, *Sphaeridiotrema globulus*, *S. pseudoglobulus* and *Leyogoni-mus polyoon* (Sauer et al. 2007, Mitchell & Cole 2008) cause intestinal hemorrhage and extensive mucosal damage. One snail can be infected with hundreds of infectious larval trematodes (Cole, unpublished data) and thus, by eating a small number of snails, a bird can receive a lethal infection in a short period of time (Sauer et al. 2007). From 2002-14 over 135,000 water birds consisting of 17 species have died in mortality events in Wisconsin, Minnesota and Illinois. These mortalities have been attributed to the four trematodes transmitted by *B. tentaculata*. The majority of these events have occurred in navigation pools (long stretches of river between dams) 7-11 of the Mississippi River and were predominately lesser scaup (*Aythya affinis*) and American coot (*Fulica americana*) (National Wildlife Health Center, Madison, WI, unpublished data).

Negative interactions between invasive species and native species are a leading cause of animal extinctions (Claver & Garcia-Berthou 2005) and freshwater gastropods are a highly threatened freshwater fauna with 74% of species categorized as imperiled or extinct (Johnson et al. 2013). In eutrophic lakes in upstate New York, *B. tentaculata* contributed to the decline

of populations of the pleurocerids *Elimia vivipara*, *E. virginica*, and *Pleurocera acuta* (Harman 1968, Harman 1968, Harman & Forney 1970, Jokinen 1992). The proposal that the ability of *B. tentaculata* to both graze and filter-feed contributed to their competitive ability was supported by a finding that *B. tentaculata* adds biomass approximately 10 times faster than pleurocerids (Harman & Forney 1970) due to the higher efficiency of carbon and nitrogen assimilation associated with filter-feeding (Tashiro & Colman 1982). While these pleurocerids are relatively common – this is indicative that this introduced species is a potential competitor with other native pleurocerids. Furthermore, an initial study indicated native snails could suffer negative consequences from *B. tentaculata* invasion, largely due to increased exposure to trematode parasite larvae transmitted at high densities of *B. tentaculata* (Sandland et al. 2013). A further study in an experimental setting found several native snail species and *B. tentaculata* were equally infected with the larval stage (metacercariae) of an echinostome parasite suggesting a potentially positive effect of the invasive snail on natives may occur by diluting the parasite load of the entire snail community (Gladosky & Sandland 2014); however, this does not consider the ability of *B. tentaculata* to form very high population densities, which would serve to enhance overall parasite abundance and pose a threat to native snails.

Understanding the history of invasion and parent populations of *B. tentaculata* may lead to precautionary steps to be implemented to limit the spread of this species. Use of microsatellite data can be helpful in understanding the routes of introduction and pinpointing parent populations (Stepien et al. 2005, Brown & Stepien 2009). In this study, we used microsatellite data to determine the colonization route of the invasive populations of *B. tentaculata* into and throughout North America. We distinguish between four alternative hypotheses of potential colonization routes: (1) a single population of *B. tentaculata* was introduced into the Great Lakes from a single source population and has since dispersed; (2) *Bithynia tentaculata* were introduced multiple times into the Great Lakes from multiple sources and have since dispersed; (3) there were multiple introductions of *B. tentaculata* from Europe into geographically distinct locations within North America; (4) while some invasive populations may have been introduced from Europe, some populations may be glacial relics that persisted in North America.

METHODS

Bithynia tentaculata samples (Figure 1) were stored at -20°C in 70-100% ethanol after collection and were deposited in the Field Museum of Natural History (F numbers 344681-344697). This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. As an invertebrate, this species is exempt from the approval process of the Institutional Animal Care and Use Committees at UWL



Figure 1. Collection sites of *Bithynia tentaculata* are indicated with filled black squares. Stars represent potential origination shipping routes for colonization from Europe. Locality information presented in Table 1.

and University of Texas Rio Grande Valley. Total cellular DNA was extracted from a snip of foot tissue using the CTAB (Cetyltrimethylammonium bromide) method (Saghai-Marooft et al. 1984). Primers that target twelve loci were used to determine allele frequencies as described by Henningsen et al. (2010). DNA was amplified via PCR with a CAG tagged primer along with the associated primers, with and without a tag. PCR was performed in a 20 μ L reaction volume with the amplification mixture at concentrations as follows: *Taq* DNA Polymerase Thermopol Buff-2 0.05 U/ μ L, 0.15 mM dNTP, 1.5 mM MgCl₂, 25 μ g/mL BSA, and 1X *Taq* Thermopol buff-2 buffer. Tagged primers were included at a concentration of 0.025 μ M, untagged primers at a concentration of 0.25 μ M, and 2 μ L of DNA template was added to each reaction. Cycling conditions consisted of 4 min at 94°C; followed by 32 cycles of: 94°C for 30 s, 55°C for 30 s, 65°C for one min followed by a final extension at 65°C for 3 min. Samples were then diluted with water 1:10 and genotyped at the University of Wisconsin Biotechnology center, on an ABI 3730xl Genetic Analyzer. Output files were analyzed using the auto run setting in GeneMarker® (Holland et al. 2008) with a GS500 size standard and ABI template, to determine the size of alleles present at each locus.

A Bayesian analysis in STRUCTURE v 2.2 (Pritchard et al. 2000, version 2. from http://pritch.bsd.uchicago.edu/software/structure2_1.html.) was used to infer the number of populations (K) using Markov Chain Monte Carlo (MCMC), with five independent runs of 100,000 steps following a 100,000 step burn-in for each K from K=1 to K=12 (these

represent the maximum and minimum possible values for K based on the number of populations sampled plus one). A test run of 200,000 steps and a 200,000 burn in was conducted to see if this level of iterations were required. The model assumed correlated allele frequencies among populations, sampling locations were informative about ancestry (LOCPRIOR), and followed an admixture model with a single value of lambda ($\lambda=1.0$) inferred for all populations. K was estimated based on the log likelihood score and posterior probability of K, Ln P(D) also known as $L(K)$ (Falush et al. 2007) as well as the rate of change in the log likelihood score (ΔK) (Evanno et al. 2005). The log likelihood score was calculated following Evanno et al. (2005): $\Delta K = m|L(K+1) - 2L(K) + L(K-1)| / s[L(K)]$ for each K. Three values were used to estimate K, $L(K)$, ΔK , and α . The best estimate of K is identified as the maximal value of $L(K)$, yet as the true K is reached, $L(K)$ at larger K values will plateau or even increase slightly (Evanno et al. 2005). The rate of change in the log likelihood score, ΔK , will be the highest at the true K. Finally, the lowest value for α indicates that most individuals are essentially from one population or another. The posterior probability of K, $L(K)$, Ln(K), and α were output directly from the program, and ΔK was calculated using the equation above for K=1 to K=12. Once K was estimated, 5 runs of STRUCTURE was used to calculate F_{ST} , H_E , and H_O for each of the K populations. Average expected and observed heterozygosity (H_O , N_A) for each of the three populations clusters determined by STRUCTURE were calculated in Microsoft Excel. Finally, the North American populations

Table 1. Eleven populations examined for 12 microsatellite loci with statistics summarizing genetic variation within populations. Data presented are latitude, longitude, number of individuals (n), number of microsatellite alleles (N_A), observed heterozygosity (H_O) and expected heterozygosity (H_E) averaged across loci. Dunaremete, Baracska, and Botanical Garden are all collection sites in the Danube River drainage. The order of populations matches the population number in Figure 2.

	Population	Latitude	Longitude	N	N_A	H_O	H_E
1	River Zala, Hungary	46.871633	16.787572	17	22	0.344	0.511
2	Lake Balaton, Hungary	46.763097	17.266496	9	8	0.381	0.403
3	Dunaremete, Hungary	47.884563	17.436472	6	18	0.339	0.515
4	Baracska, Hungary	47.287274	18.757078	1	1	—	—
5	Botanical Garden, Budapest, Hungary	47.485031	19.085412	21	43	0.589	0.582
6	Lake Winnibigoshish, Minnesota, U.S.	47.431292	−94.196227	10	17	0.466	0.620
7	Georgetown Lake, Montana, U.S.	46.181239	−113.286868	3	7	0.750	0.750
8	Rattlesnake Reservoir, Montana, U.S.	45.90345	−108.426982	18	17	0.463	0.629
9	Upper Mississippi River (Pool 7), Wisconsin, U.S.	43.8669095	−91.3070842	65	39	0.409	0.557
10	Lake Winnebago, Wisconsin, U.S.	43.806288	−88.402219	6	12	0.521	0.533
11	Ottawa River, Canada	45.793924	−76.99684	7	9	0.389	0.589

were run separately to determine if the greater European diversity masked internal structure in North American populations.

Genepop v 4.0.10 (Raymond & Rousset 1995), was used to estimate number of alleles (N_A) for each population. GenAlEx 6.501 (Peakall & P.E. 2006, Peakall & Smouse 2012) was used to perform an analysis of molecular variance to calculate F_{ST} and F'_{ST} of the three population clusters (regions) determined by STRUCTURE. We used these combinational, regional groups rather than individual populations to increase the sample size of each group. This conforms with the findings of Hale et al. (2012) that 25–30 individuals are needed per “group” for accuracy of microsatellite data. When grouping by the regions Danube ($n=27$), Lake Balaton ($n=26$), and North America ($n=109$) we have sufficient sampling for comparison among regions, although not comprehensive sampling for any region. GenAlEx v. 6.501 was also used to perform a genetic distance based Principal Coordinates Analysis (PCoA) on alleles from all populations.

As a final examination of patterns of population genetic structure we used Bottleneck 1.2.02 (Cornuet & Luikart 1996) to examine each of the 3 regions for signs of a recent genetic bottleneck using a Wilcoxon sign-rank test, to accommodate our limited loci and sampling under the two-phase-model of evolution as recommended for microsatellite loci (Luikart & Cornuet 1997). In a genetic bottleneck, reduced population size results in loss of alleles and a decline in heterozygosity, recent bottlenecks should appear to have higher than expected genetic diversity compared to expectations from Hardy-Weinberg equilibrium.

RESULTS

Eleven populations of *Bithynia tentaculata* were sampled for 12 microsatellite loci from the native European range and from North America (Table 1). Structure runs with 200,000 burn-in and iterations were not different from 100,000 burn-in

and 100,000 iterations all further analyses were carried out using the latter settings.

All three *ad hoc* estimates for K from our analysis using STRUCTURE, considered together, suggest a best estimate of $K=3$ (data not figured). All eleven populations grouped into one of these 3 clusters and provided the rationale for combining populations in further analyses. The three clusters include 1) the Danube population (all populations in the Danube River Basin, Hungary), 2) Lake Balaton population (includes Lake Balaton and River Zala—which flows into Lake Balaton, Hungary), and 3) North America. The F_{ST} value for the combined Danube populations is much lower than that for North America or Lake Balaton (Table 2). The North American populations analyzed without the European data resulted in $K=1$.

In Figure 2 which illustrates all eleven populations as well as the three combined populations, Lake Balaton stands out as being the least intermixed (also Table 2 with the highest among population F_{ST} value, 0.3379). The Danube has some contribution from Lake Balaton and from the population that is the source of the North American populations. The North American populations in Canada and Montana are the most heterogeneous, although lacking unique alleles. Examination for genetic signatures of a recent population bottleneck found no signature of this event in the Danube (Wilcoxon sign-rank test for two-phase-model, $P=0.57$), and Lake Balaton regions ($P=0.688$), however the North American region shows the signature of a recent bottleneck ($P=0.012$).

Table 2. Results of AMOVA on populations grouped by regions. Genetic differentiation among populations, F_{ST} (+SD) and expected heterozygosity, H_E (+SD) for $K=3$.

Cluster	n	F_{ST}	H_E
Danube	28	0.1614+0.0010	0.6261+0.0001
North America	109	0.3133+0.0015	0.5238+0.0003
Lake Balaton	26	0.3379+0.0025	0.5613+0.0001




Figure 2. STRUCTURE output of Q (or proportion of each individual attributed to each cluster). Clusters indicated by color, population by number: (Lake Balaton Cluster: 1-River Zala HUN, 2-Lake Balaton HUN; Danube River Cluster: 3-Dunaremete HUN, 4-Baracska HUN, 5-Botanic Garden HUN; North American Cluster: 6-Lake Winnibigoshish MN, 7-Georgetown Lake MT, 8-Rattlesnake Reservoir MT, 9-Upper Mississippi River, 10-Lake Winnebago WI, 11-Ottawa River, Canada).

The number of alleles at 12 microsatellite loci ranged from 1–39. In most populations, observed heterozygosity was lower than expected under Hardy-Weinberg equilibrium (Table 1). Allele-frequency divergence among these three populations is shown in Table 3. The two European populations are more similar to each other (0.1160) than to the North American population, which is roughly equally different from the two European populations (0.1484, 0.1547). The Danube River population had the largest number of private alleles (21), followed by North America (11) and the fewest in Lake Balaton (8).

Most loci not only had different frequencies, but usually unique alleles in each population. In general, allelic diversity is highest in the Danube populations, followed by North America, then Lake Balaton (Table 1). For example, a single well-sampled locus, Bt03, displays very different allele frequencies across all populations, each population also has unique alleles at this locus (not figured). This pattern is repeated in most other loci. However, some population structure was observed. Allele frequencies are more similar in populations such as the Montana Lakes and Ottawa River (i.e. those outside the areas adjacent to Lake Michigan) and have fewer alleles, none unique to those populations (not figured).

A principal coordinates analysis (PCoA) of the microsatellite allele data across all populations (excluding Baracska, a Hungarian population represented by a single individual) resulted in three significant PCoA axes, axis one explains 17% of the variation present, axis two 13.19 %, and axis three 9.28 %. A scatterplot showing all individuals grouped by population on axis one and two is shown in Figure 3. On both axis one and two (Figure 3) the Upper Mississippi River population and Lake Winnibigoshish populations have the widest variation in allelic diversity. The other populations are restricted to the lower left quadrant of the graph, encompassed within the diversity of those two populations. Axis three (not figured) distinguishes the European populations from Lake Winnebago, WI and the Montana populations.

Table 3. Allele frequency divergence among regions (net nucleotide distance) calculated using STRUCTURE.

	Danube	North America
Danube	—	—
North America	0.1483	—
Lake Balaton	0.1160	0.1547

DISCUSSION

The goal of this study was to distinguish among four alternative hypotheses for introduction of *B. tentaculata* into North America. Hypothetical scenarios were proposed considering what is known of the possible invasion route and history of the species. The expected genetic consequences of each scenario are proposed based on the patterns observed in a review of genetic consequences of invasion in 80 species of animals, plants, and fungi by Dlugosch and Parker (2008) and glacial refugia by Maggs et al. (2009).

Hypothesis One incorporated a single introduction scenario: *B. tentaculata* was introduced once into the Great Lakes and dispersed into other North American watersheds. If this hypothesis were supported we would expect a low F_{ST} in North American populations compared to European populations, similar allele frequencies to the source, and very few private alleles in North America compared to European source. Hypothesis Two was developed around a multiple introduction scenario into the Great Lakes; after which *B. tentaculata* dispersed. If this hypothesis were correct, we would expect a higher F_{ST} , and greater heterogeneity in North America and many alleles from all across the founding European populations (albeit at lesser frequencies) and many private alleles if comparing North American population against a single source population. Hypothesis Three described a scenario with introductions from more than one European source population into the different North American watersheds in which *B. tentaculata* has now been found. If this hypothesis were correct, we would expect a high F_{ST} , very different allele frequencies both within North American populations and between North American and European populations, as well as private alleles unique to different North American populations. Hypothesis Four incorporated a European source of introduction for some of the North American populations, while other North American populations were assumed glacial relics. If this hypothesis were correct, we would expect a signal similar to Hypothesis Three but with private alleles in the glacial relic populations that are different from other North American and European populations.

Populations in Hungary compared to populations in North America

The *B. tentaculata* samples collected in Hungary are all part of the Ponto-Caspian Basin which contributes to the Black sea colonization route, and has contributed >70% of Great




Figure 3. Genetic comparisons between populations using a Principal coordinates analysis of the genetic distance matrix from the microsatellite allele data. Similar shapes represents populations from the same region or drainage basin. Axis 1 represents 17% and axis 2 represents 13.19% of the variation present.

Lakes invaders (Ricciardi 2001, Gelembiuk et al. 2006). The concordance of the *ad hoc* estimates given in STRUCTURE (Figure 2) grouped all 11 populations into one of three clusters (K). However, Evanno et al. (2005) warn that these estimates must also coincide with scenarios that are biologically significant. Considering the *a priori* knowledge of sampling sites, K=3 corresponds with what would be predicted based on sampling localities from North America and two different Hungarian watersheds, the Danube River (Dunaremete, Baracska, and the Botanic Garden in Budapest sites) and Lake Balaton (Lake Balaton and River Zala sites) watersheds.

Examination of allele frequencies and variability (Figure 2, Figure 3, Table 3) from the Danube River and Lake Balaton supports the idea that the populations could be genetically distinct with the North American samples grouped as a third distinct population. There appears to be some shared alleles with alleles found in the Hungarian populations also present in the North American populations (Figure 2). This may indicate some contribution to ancestry of or a shared common ancestry with the North American populations, however additional European populations must be sampled for a robust comparison to be made. Some preliminary estimates can be made from the allele-frequency divergences of each population (Table 3). Even though the two Hungarian populations are distinct (Figure 3), they are more similar to each other than either is to the North American population (Table 2). This may indicate that the Hungarian populations are not the source of

the North American populations. This inference is also supported by the number of private alleles, as there are more private alleles in samples from Hungary than in samples from North America, supporting the idea that Hungary is not the source of the North American populations. However, this is not definitive; the source could be from further downstream on the Danube, which would explain some of the same alleles present seen in the STRUCTURE figure (Figure 2) and as overlap in the PCoA (Figure 3).

These data do not support the Black sea colonization route as the source of the North American invasion, however this route cannot be definitely ruled out considering its large range and our limited sampling. Samples from further downstream on the Danube River, or within the Black Sea, will be necessary before this entire watershed can be definitively excluded as a source for the North American invasion. Sampling of the other two likely colonization routes through the North Sea and the Baltic Sea is also needed to determine if those routes may be the source population of the North American invasion.

Populations within North America

The North American populations were all combined into a single population by the *ad hoc* estimates in STRUCTURE which points toward a recent shared ancestry of all the populations within North America. Further when analyzed

separately to determine if the European diversity masked variation within North American, STRUCTURE found the most support for a single North American population. This evidence supports Hypotheses one or two that *B. tentaculata* was introduced into the Great Lakes and dispersed from there. This is supported by few private alleles in North America (compared to the European populations) and that the North American F_{ST} is relatively low compared to the native European populations. Another source of data offers some support for this scenario. In a relatively recent M.S. thesis Whalen (2011) used 11 microsatellite loci of which 4 overlap with this study and several of the same populations but also including 2 population not included in this study which are from Eastern Wisconsin near the Great Lakes. They found the populations near the Great Lakes (e.g. Lake Winnebago, WI) were probably “parent” populations to the Lake Onalaska and Lake Winnibigoshish populations included in both studies.

The STRUCTURE analysis does not appear to support Hypothesis Three, that there are multiple European source populations for the North American populations, unless Europe has very homogeneous populations, which is unlikely with the amount of divergence we observed in just the two Hungarian watersheds. Hypothesis Four also appears unlikely given our data set, though it is hard to distinguish private alleles from potential glacial relics without more extensive sampling of specimens from Europe for comparison. It is difficult at this point to confirm whether Hypothesis One or Two is more likely, as there were not enough data from European specimens to compare source populations. However, by comparing allele frequencies among populations, we can get a hint of which hypothesis (One or Two) is more likely.

With the exception of two loci, Bt22 and Bt40, the allele frequencies across the three populations differ, once again supporting the hypothesis of a different source population for invasion than those sampled in Hungary (data not figured). There were alleles present in snails from North America that were absent in snails collected in Hungary. While these may be attributed to new alleles arising in the population, such a scenario is unlikely across so many loci. There were also alleles present in the Hungarian populations that are not in snails from North America. Founder effects could account for this; however, if this were the case, the other alleles present would most likely be at similar frequencies, which they were not. It is possible that some snails from Hungary were mixed with other European source populations that then gave rise to the North American populations. If this were the case, there would likely be more heterogeneity in the North American populations (Figure 2). Most of the North American loci are dominated by one allele indicating a significant recent genetic bottleneck occurred, this is also supported by the Bottleneck analysis finding the signature of a recent bottleneck only in the North American population. This signature could be due to a relatively small initial invasive population from a single source population and subsequent bottlenecks with colonization of additional watersheds. This coincides with other invasions into the Great Lakes that have been shown to be from a single

source population (Ricciardi & MacIsaac 2000, Brown & Stepien 2009). At this point it is still speculative, but Hypothesis One is more likely, and the higher F_{ST} (Table 3) may be an artifact of founder effects which can elevate F_{ST} levels (Weir & Cockerham 1984).

Our data suggest *B. tentaculata* has dispersed across the U.S. from a single initial colonization, not from multiple invasions from different sources. It also appears that all sampled North American populations are recent recolonizations, not glacial relic populations. Given the few populations sampled in the European range of *B. tentaculata*, the European source of the introduction into North America is still unknown, and will require further study of the European range. However, it does appear, based on the data available, that the most common route for invasion into the Great Lakes, from the Ponto-Caspian Sea Basin through the Black sea, is not the likely source of introduced *Bithynia tentaculata* in North America.

ACKNOWLEDGEMENTS

The authors would like to thank individuals that contributed their insight, time, and specimens to the project: Russell Minton, Anindo Choudbury, Jenny Sauer, Roger Haro, Mark Sandheinrich, Ben Walker, Greg Sandland, Rachelle Amundson, Adam Prise, Kirk Gallant, and Josh Laurila. This project was funded by the UWL River Studies Program, United States Geological Survey, Upper Mississippi Environmental Research Center, National Great Rivers Research and Education Center, University of Wisconsin Institute for Race and Ethnicity, the UWL faculty research grant program, and the UWL College of Science and Health Dean's Fellowship program. Several anonymous reviewers contributed greatly to the improvement of the manuscript, we appreciate their critical insights. Use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the US Government.

LITERATURE CITED

- Baker, F.C. 1928. The american *Bithynia* not wholly an introduced species. *Illinois State Academy of Science* 46: 56-62, 51 pl.
- Brown, J.E. & Stepien, C.A. 2009. Invasion genetics of the Eurasian round goby in North America: tracing sources and spread patterns. *Molecular Ecology* 18: 64-79.
- Claver, M. & Garcia-Berthou, E. 2005. Invasive species are a leading cause of animal extinctions. *Trends in Ecology and Evolution* 20: 110.
- Cornuet, J.M. & Luikart, G. 1996. Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* 144: 2001-2014.
- Dlugosch, K.M. & Parker, I.M. 2008. Founding events in species invasions: genetic variation, adaptive evolution, and the role of multiple introductions. *Molecular Ecology* 17: 431-449.
- Evanno, G., Regnaut, S. & Goudet, J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* 14: 2611-2620.
- Gelembiuk, G.W., May, G.E. & Lee, C.E. 2006. Phylogeography and

- systematics of zebra mussels and related species. *Molecular Ecology* 15: 1033-1050.
- Gladosky, C. & Sandland, G. 2014. Assessing host competency between native and invasive snail species exposed to the native parasite *Echinostoma revolutum*. *Aquatic Invasions* 9: 87-93.
- Grigorovich, I., Colautti, R., Mills, E., Holeck, K. & MacIsaac, H. 2003. Ballast mediated animal introductions in the Laurentian Great Lakes: retrospective and prospective analyses. *Canadian Journal of Fisheries and Aquatic Sciences* 60: 740-756.
- Hale, M.L., Burg, T.M. & Steeves, T.E. 2012. Sampling for microsatellite-based population genetic studies: 25 to 30 individuals per population is enough to accurately estimate allele frequencies. *PLoS ONE* 7: doi:10.1371/journal.pone.0045170.
- Harman, W.N. 1968. Interspecific competition between *Bithynia* and pleurocerids. *Nautilus* 82: 72-73.
- Harman, W.N. 1968. Replacement of pleurocerids by *Bithynia* in polluted waters of central New York. *Nautilus* 81: 77-83.
- Harman, W.N. & Fomey, J.L. 1970. Fifty years of change in the molluscan fauna of Oneida Lake, New York. *Limnology and Oceanography* 15: 454-460.
- Henningsen, J.P., Lance, S.L., Jones, K.L., Hagen, C., Laurila, J., Cole, R.A. & Perez, K.E. 2010. Development and characterization of 17 polymorphic microsatellite loci in the faucet snail, *Bithynia tentaculata* (Gastropoda: Caenogastropoda: Bithyniidae). *Conservation Genetics Resources* 2: 247-250.
- Holland, B., Clarke, A. & Meudt, H. 2008. Optimizing automated AFLP scoring parameters to improve phylogenetic resolution. *Systematic Biology* 57: 347-366.
- Johnson, P.D., Bogan, A.E., Brown, K.M., Burkhead, N.M., Cordeiro, J.R., Garner, J.T., Hartfield, P.D., Lepitzki, D.A.W., Mackie, G.L., Pip, E., Tarpley, T.A., Tiemann, J.S., Whelan, N.V. & Strong, E.E. 2013. Conservation Status of Freshwater Gastropods of Canada and the United States. *Fisheries* 38: 247-282.
- Jokinen, E. 1992. *The Freshwater Snails (Mollusca: Gastropoda) of New York State*. The University of the State of New York, The State Education Department, The New York State Museum, Albany, New York.
- Keller, R.P., Drake, J.M., Drew, M.B. & Lodge, D.M. 2010. Linking environmental conditions and ship movements to estimate invasive species transport across the global shipping network. *Diversity & Distributions* 17: 93-102.
- Lee, C. & Bell, M. 1999. Causes and consequences of recent freshwater invasions by saltwater animals. *Trends in Ecology and Evolution* 14: 284-288.
- Luikart, G. & Cornuet, J.M. 1997. Empirical evaluation of a test for identifying recently bottlenecked populations from allele frequency data. *Conservation Biology* 12: 228-237.
- Maggs, C.A., Castilho, R., Foltz, D., Hensler, C., Jolly, T., Olsen, J., Perez, K.E., Stam, W., Vainola, R., Viard, F. & Wares, J. 2009. Evaluating signatures of glacial refugia for North Atlantic marine organisms. *Ecology* 89: s108-122.
- Mills, E.L., Leach, J.H., Carlton, J.T. & Secor, C.L. 1993. Exotic species in the Great Lakes - a history of biotic crises and anthropogenic introductions. *Journal of Great Lakes Research* 19: 1-54.
- Mitchell, A.J. & Cole, R.A. 2008. Survival of the faucet snail after chemical disinfection, pH extremes, and heated water bath treatments. *North American Journal of Fisheries Management* 28: 1597-1600.
- Peakall, R. & P.E., S. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288-295.
- Peakall, R. & Smouse, P. 2012. GenAIEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research-an update. *Bioinformatics* 28: 2537-2539.
- Pritchard, J.K., Stephens, M. & Donnelly, P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155: 945-959.
- Raymond, M. & Rousset, F. 1995. GenePop (version 3.4): population genetics software for exact test and ecumenicism. *Journal of Heredity* 86: 248-249.
- Ricciardi, A. 2001. Facilitative interactions among aquatic invaders: is an 'invasional meltdown' occurring in the Great Lakes? *Canadian Journal of Fisheries and Aquatic Sciences* 58: 13p.
- Ricciardi, A. & MacIsaac, H. 2000. Recent mass invasion of the North American Great Lakes by Ponto-Caspian species. *Trends in Ecology and Evolution* 15: 62-65.
- Saghai-Marouf, M.A., Soliman, K.M., Jorgensen, R.A. & Allard, R.W. 1984. Ribosomal DNA spacer length variation in barley: Mendelian inheritance, chromosomal location and population dynamics. *Proceedings of the National Academy of Sciences of the United States of America* 81: 8018-8021.
- Sandland, G.J., Houk, S., Walker, B., Haro, R.J. & Gillis, R. 2013. Differential patterns of infection and life-history expression in native and invasive hosts exposed to a trematode parasite. *Hydrobiologia* 701: 89-98.
- Sauer, J.S., Cole, R.A., Nissen, J.M., Upper Midwest Environmental Sciences Center (Geological Survey) & Geological Survey (U.S.) (2007). Finding the exotic faucet snail (*Bithynia tentaculata*): investigation of waterbird die-offs on the Upper Mississippi River National Wildlife and Fish Refuge. Reston, Va., U.S. Geological Survey: 3 p.
- Stepien, C.A., Brown, J.E., Neilson, M.E. & Tumeo, M.A. 2005. Genetic diversity of invasive species in the Great Lakes versus their Eurasian source populations: Insights for risk analysis. *Risk Analysis* 25: 1043-1060.
- Tashiro, J.S. & Colman, S.D. 1982. Filter-feeding in the freshwater prosobranch snail *Bithynia tentaculata*: bioenergetic partitioning of ingested carbon and nitrogen. *American Midland Naturalist* 107: 114-132.
- Weir, B.S. & Cockerham, C.C. 1984. Estimating *F*-statistics for the analysis of population structure. *Evolution* 38: 1358-1370.
- Whalen, S.J. (2011). Analysis of the genetic structure of *Bithynia tentaculata* snail populations in Wisconsin and Minnesota. Master's thesis, University of Minnesota, Mankato.

Appendix 1. List of all specimens genotyped. Localities, latitude, and longitude are listed as well as specimen identification number and Field Museum accession number (F-number).

F Number	DNA Numbers	Locality	latitude	longitude
344681.1	1669	Ottawa River, Canada	45.7939	–76.9968
344681.2	1671	Ottawa River, Canada	45.7939	–76.9968
344681.3	1672	Ottawa River, Canada	45.7939	–76.9968
344681.4	1673	Ottawa River, Canada	45.7939	–76.9968
344681.5	1675	Ottawa River, Canada	45.7939	–76.9968
344681.6	1874	Ottawa River, Canada	45.7939	–76.9968
344681.7	1875	Ottawa River, Canada	45.7939	–76.9968
344681.8	1877	Ottawa River, Canada	45.7939	–76.9968
344681.9	1893	Ottawa River, Canada	45.7939	–76.9968
344681.10	1894	Ottawa River, Canada	45.7939	–76.9968
344682.1	1868	Baracska, Hungary	47.2873	18.7571
344683.1	1820	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.2	1896	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.3	1897	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.4	1899	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.5	1900	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.6	1902	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.7	1905	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.8	1909	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.9	1915	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.10	1923	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.11	1927	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.12	1928	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.13	1930	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.14	1931	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.15	1934	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.16	1937	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.17	1941	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.18	1944	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.19	1945	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.20	1947	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.21	1949	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.22	1950	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344683.23	1951	Botanic Garden near Eotvos Lorant Univ., Budapest, Hungary	47.4850	19.0854
344684.1	1854	Dunaremete, Hungary	47.8846	17.4365
344684.2	1855	Dunaremete, Hungary	47.8846	17.4365
344684.3	1856	Dunaremete, Hungary	47.8846	17.4365
344684.4	1857	Dunaremete, Hungary	47.8846	17.4365
344684.5	1858	Dunaremete, Hungary	47.8846	17.4365
344684.6	1859	Dunaremete, Hungary	47.8846	17.4365
344685.1	1788	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.2	1792	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.3	1793	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.4	1794	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.5	1795	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.6	1797	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.7	1801	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.8	1802	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.9	1803	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.10	1807	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665

Appendix 1, continued.

F Number	DNA Numbers	Locality	latitude	longitude
344685.11	1809	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.12	1810	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.13	1811	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.14	1812	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.15	1813	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.16	1814	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.17	1816	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.18	1817	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.19	1818	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.20	1826	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.21	1827	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344685.22	1828	Lake Balaton, near Keszthely, Hungary	46.7631	17.2665
344686.1	1865	Lipót, Hungary	47.8661	17.4860
344687.1	1849	Northern Part of Budapest, Hungary	47.5128	19.0427
344687.2	1850	Northern Part of Budapest, Hungary	47.5128	19.0427
344687.3	1851	Northern Part of Budapest, Hungary	47.5128	19.0427
344687.4	1852	Northern Part of Budapest, Hungary	47.5128	19.0427
344687.5	1853	Northern Part of Budapest, Hungary	47.5128	19.0427
344688.1	1765	River Zala, Hungary	46.8716	16.7876
344688.2	1766	River Zala, Hungary	46.8716	16.7876
344688.3	1767	River Zala, Hungary	46.8716	16.7876
344688.4	1768	River Zala, Hungary	46.8716	16.7876
344688.5	1769	River Zala, Hungary	46.8716	16.7876
344688.6	1770	River Zala, Hungary	46.8716	16.7876
344688.7	1771	River Zala, Hungary	46.8716	16.7876
344688.8	1772	River Zala, Hungary	46.8716	16.7876
344688.9	1773	River Zala, Hungary	46.8716	16.7876
344688.10	1774	River Zala, Hungary	46.8716	16.7876
344688.11	1775	River Zala, Hungary	46.8716	16.7876
344688.12	1776	River Zala, Hungary	46.8716	16.7876
344688.13	1777	River Zala, Hungary	46.8716	16.7876
344688.14	1778	River Zala, Hungary	46.8716	16.7876
344688.15	1779	River Zala, Hungary	46.8716	16.7876
344688.16	1780	River Zala, Hungary	46.8716	16.7876
344688.17	1781	River Zala, Hungary	46.8716	16.7876
344688.18	1782	River Zala, Hungary	46.8716	16.7876
344688.19	1783	River Zala, Hungary	46.8716	16.7876
344688.20	1784	River Zala, Hungary	46.8716	16.7876
344688.21	1785	River Zala, Hungary	46.8716	16.7876
344689.1	1703	Lake Winnebago, MN	47.4313	-94.1962
344689.2	1704	Lake Winnebago, MN	47.4313	-94.1962
344689.3	1705	Lake Winnebago, MN	47.4313	-94.1962
344689.4	1706	Lake Winnebago, MN	47.4313	-94.1962
344689.5	1707	Lake Winnebago, MN	47.4313	-94.1962
344689.6	1708	Lake Winnebago, MN	47.4313	-94.1962
344689.7	1709	Lake Winnebago, MN	47.4313	-94.1962
344689.8	1710	Lake Winnebago, MN	47.4313	-94.1962
344689.9	1713	Lake Winnebago, MN	47.4313	-94.1962
344689.10	1714	Lake Winnebago, MN	47.4313	-94.1962
344689.11	1715	Lake Winnebago, MN	47.4313	-94.1962
344689.12	1717	Lake Winnebago, MN	47.4313	-94.1962

Appendix 1, continued.

F Number	DNA Numbers	Locality	latitude	longitude
344689.13	1720	Lake Winnebagoishish, MN	47.4313	-94.1962
344689.14	1722	Lake Winnebagoishish, MN	47.4313	-94.1962
344689.15	1724	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.1	1878	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.2	1879	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.3	1880	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.4	1881	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.5	1882	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.6	1883	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.7	1884	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.8	1885	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.9	1886	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.10	1887	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.11	1888	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.12	1889	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.13	1890	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.14	1891	Lake Winnebagoishish, MN	47.4313	-94.1962
344697.15	1892	Lake Winnebagoishish, MN	47.4313	-94.1962
344690.1	1655	Georgetown Lake, Montana	46.1812	-113.2869
344690.2	1656	Georgetown Lake, Montana	46.1812	-113.2869
344690.3	1657	Georgetown Lake, Montana	46.1812	-113.2869
344690.4	1658	Georgetown Lake, Montana	46.1812	-113.2869
344691.1	1659	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.2	1660	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.3	1661	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.4	1663	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.5	1666	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.6	1727	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.7	1728	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.8	1729	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.9	1730	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.10	1732	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.11	1734	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.12	1735	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.13	1742	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.14	1746	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.15	1755	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.16	1756	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.17	1757	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.18	1760	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.19	1761	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.20	1763	Rattlesnake Reservoir, MT	45.9035	-108.4270
344691.21	1764	Rattlesnake Reservoir, MT	45.9035	-108.4270
344692.1	1456	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.2	1457	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.3	1458	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.4	1460	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.5	1462	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.6	1463	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.7	1464	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.8	1465	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985

Appendix 1, continued.

F Number	DNA Numbers	Locality	latitude	longitude
344693.1	1466	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.2	1467	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.3	1469	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.4	1470	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.5	1471	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.6	1472	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.7	1473	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.8	1474	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.9	1475	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344692.9	1478	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.10	1479	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344692.11	1480	Pool 7, Cormorant Is. La Crosse, WI	43.9028	-91.2985
344693.10	1484	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.11	1485	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.12	1488	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.13	1489	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344693.14	1491	Pool 7, Broken Gun Is. E. La Crosse, WI	43.9133	-91.2889
344694.1	1493	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.2	1494	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.3	1496	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.4	1497	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.5	1498	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.6	1500	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.7	1501	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.8	1520	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.9	1521	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.10	1523	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.11	1524	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.12	1525	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.13	1526	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.14	1527	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.15	1528	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344694.16	1529	Pool 7, Arrow Head Is. E., La Crosse, WI	43.8996	-91.2863
344695.1	1624	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.2	1625	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.3	1628	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.4	1629	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.5	1631	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.6	1632	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.7	1634	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.8	1636	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.9	1637	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.10	1638	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.11	1639	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.12	1641	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.13	1644	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.14	1645	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.15	1646	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.16	1647	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.17	1650	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863
344695.18	1653	Pool 7 of Mississippi River, La Crosse, WI	43.8996	-91.2863

Appendix 1, continued.

F Number	DNA Numbers	Locality	latitude	longitude
344695.19	1679	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.20	1680	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.21	1681	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.22	1682	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.23	1683	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.24	1684	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.25	1685	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.26	1687	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.27	1689	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.28	1691	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.29	1693	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.30	1694	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.31	1695	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.32	1696	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.33	1697	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.34	1698	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.35	1699	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.36	1701	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344695.37	1702	Pool 7 of Mississippi River, La Crosse, WI	43.8996	−91.2863
344696.1	2056	Lake Winnebago, WI	43.9670	−88.5228
344696.2	2058	Lake Winnebago, WI	43.9670	−88.5228
344696.3	2059	Lake Winnebago, WI	43.9670	−88.5228
344696.4	2063	Lake Winnebago, WI	43.9670	−88.5228
344696.5	2065	Lake Winnebago, WI	43.9670	−88.5228
344696.6	2066	Lake Winnebago, WI	43.9670	−88.5228
344696.7	2068	Lake Winnebago, WI	43.9670	−88.5228
344696.8	2069	Lake Winnebago, WI	43.9670	−88.5228
344696.9	2071	Lake Winnebago, WI	43.9670	−88.5228
344696.10	2072	Lake Winnebago, WI	43.9670	−88.5228
344696.11	2073	Lake Winnebago, WI	43.9670	−88.5228
344696.12	2074	Lake Winnebago, WI	43.9670	−88.5228
344696.13	2076	Lake Winnebago, WI	43.9670	−88.5228