

A new species of *Bufonia* (Caryophyllaceae) from Israel: *B. ramonensis*

Author: Danin, Avinoam

Source: Willdenowia, 31(1) : 95-100

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.31.31108>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

AVINOAM DANIN

A new species of *Bufonia* (*Caryophyllaceae*) from Israel: *B. ramonensis*

Abstract

Danin, A.: A new species of *Bufonia* (*Caryophyllaceae*) from Israel: *B. ramonensis*. – Willdenowia 31: 95-100. 2001. – ISSN 0511-9618.

Bufonia ramonensis is a small chamaephyte with lignified branches, described from one mountain slope at the Negev Highlands, where it is confined to crevices and soil pockets of smooth-faced limestone. The new species is closely related to *B. multiceps*, which is a hemicryptophyte with herbaceous stems endemic to similar microhabitats in magmatic rocks of southern Sinai.

Following previous investigations during the last 35 years (Danin 1983, 1999, Danin & Solomeshch 1999) I kept telling my colleagues that smooth-faced rock outcrops in desert areas are to be expected to support unexpected findings. In the last decades I passed many times half a kilometer north, east or west of the particular slope that I visited at the beginning of February 2001. It was thus an almost expected surprise to find a population of several hundred small shrubs of a so far undescribed species in rock crevices and soil pockets in an area of a shrub-steppe of *Moricandia nitentis*-*Artemisietum sieberi*.

Bufonia ramonensis Danin, *sp. nova*

Holotype: Israel, Negev Highlands, 2 km SW of Har Ramon, 30°30'N, 34°37'E, 900 m, in crevices and soil pockets of smooth-faced limestone outcrops, a gentle north-facing slope, 9.2.2001, Danin (HUI); isotypes: B, K, E) – Fig. 1, 4.

A *Bufonia multiceps* Decne. ramis prostrato-ascendentibus lignosis et ramulis floriferis induratis perennantibus (nec ramis ramulisque herbaceis), sepalis 3(-6)-nervis (nec 6-7-nervis), margine ciliatis (nec glabris), ovarium subaequantibus (nec tertia parte longioribus) et seminibus omnino tuberculatis (nec dorso tenuiter tuberculatis facie planiusculis cellulis stellatis) differt.

A low *chamaephyte*, stems up to 30 cm long, 1-20 mm diam., much branched, densely appressed-pubescent with recurved hairs; the previous year's and older twigs greyish, giving rise to new branches from the lower 2/3 or 4/5 of previous year's stems; flowering branches 5-10 cm long. Leaves 5-12 × 0.2-0.5 mm, narrowly linear-lanceolate with ciliate margins, connate at the base and appressed to the stem, base with scarious margin. *Inflorescence* a dense terminal, short-



Fig. 1. *Bufonia ramonensis* – branched stems. – Scale 1 cm; from the holotype.

pedicelled dichasial cyme of 4-12 flowers, 5-7 mm in diam. resembling a capitulum, in addition with 3-6 single lateral flowers or 2-3 flowers in monochasia at the axiles of bracts below it; often only the lateral flowers occur. *Pedicels* up to 1 mm, appressed hairy, in dense cymes with longitudinal depression. *Flowers* 3-4 mm long; sepals 3 mm, subequal, ovate-lanceolate, 3(-6)-nerved, margins scarious, ciliate; petals white, shorter than sepals; stamens 8. *Capsule* ovate, 2.5×1.5 mm, as long as, slightly longer, or slightly shorter than calyx. *Seeds* 2, brown, reniform, laterally compressed, 1.5-1.8 mm long, 0.8-1.5 mm wide, tuberculate throughout (Fig. 4).




Fig. 2. Distribution of *Bufonia ramonensis* (■) and *B. multiceps* (▲) as documented in HUI.

Distribution and ecology

So far known only from the type locality (Fig. 2), at an area of less than half a hectare, where the number of individuals of *Bufonia ramonensis* is estimated as 200-300. Until discovered in additional places it may be regarded as one of the rarest endemic plants of Israel.

The companions of *B. ramonensis* are shrub-steppe species of the plant community *Moricandia nitentis-Artemisietum sieberi* Danin & Solomeshch 1999 and of the *Pistacio atlanticae-Chiliadenetum iphionoidis* Danin, Orshan & Zohary ex Danin & Solomeshch 1999 (cf. Danin & Solomeshch 1999). Those of the *Artemisietum* are: *Artemisia sieberi* Besser, *Astragalus sanctus* Boiss., *Diplotaxis harra* (Forssk.) Boiss., *Erodium crassifolium* L'Her., *Gymnocarpus decander* Forssk., *Helianthemum vesicarium* Boiss., *Moricandia nitens* (Viv.) E. A. Durand & Barratte, *Noaea mucronata* (Forssk.) Asch. & Schweinf. and *Salvia lanigera* Poir. Those of the *Chiliadenetum* are: *Astragalus amalecitanus* Boiss., *A. bethlehemiticus* Boiss., *Ankyropetalum gypsophiloides* Fenzl, *Chiliadenus iphionoides* (Boiss. & Blanche) Brullo, *Delphinium ithaburense* Boiss., *Dianthus sinaicus* Boiss., *Eryngium glomeratum* Lam., *Fumana thymifolia* (L.) Webb, *Haplophyllum poorei* C. C. Towns., *Micromeria sinaica* Benth., *Phagnalon rupestre* (L.) DC., *Pteroccephalus pulverulentus* Boiss. & Balansa, *Rhamnus disperma* Boiss., *Tanacetum sinaicum* (Fresen.) Decne. ex K. Bremer & Humphries, *Urospermum picroides* (L.) F. W. Schmidt and *Umbilicus intermedius* Boiss.




Fig. 3. Seed of *Bufonia multiceps* – a: whole seed; b: lateral faces with stellate epidermal cells; c: anti-micropylar pole with 70-80 μm long, distinct and spaced appendages; d: stellate epidermal cells. – Scale bars: a-c = 100 μm , d = 10 μm ; specimen from S Sinai, 17.7.1968, leg. Tadmor (HUJ).




Fig. 4. Seed of *Bufonia ramonensis* – a: whole seed; b: lateral faces with tubercles; c: anti-micropylar pole with 50-60 μm long densely packed appendages; d: densely packed tubercular epidermal cells. – Scale bars: a-c = 100 μm , d = 10 μm ; from the type collection.

A live plant collected together with the type collection, planted in a pot and raised in Jerusalem started to bloom on 20.5.2001.

Relationship

There are one or two additional species of *Bufonia* in the Flora Palaestina area. *B. virgata* Boiss. is an annual. *B. ephedrina* Sam., recorded by Mouterde (1966) and by Chrtek & Křisa (1999) from Jordan, is described by Mouterde (1966) as pubescent at base and glabrous above, with 7-nerved sepals and smooth seeds. These properties are not shared with *B. ramonensis*. Our new species neither agrees with any of the *Bufonia* species treated by Rechinger (1988) nor with any of those recognized by Chrtek & Křisa (1999) in their revision of the genus in Asia. Geographically and morphologically the closest species is *B. multiceps* (see distribution map, Fig. 2) as described by Decaisne (1835), Boissier (1867) and Boulos (1999). Its differentiating characters are presented in the Latin diagnosis and in the English description of *B. ramonensis*, above.

The description of the seed of *B. multiceps* by Boissier (1867: “seminibus dorso tenuiter tuberculatis facie planiusculis”) agrees with specimens of this species deposited in HUJ better than the original description by Decaisne (1835: “semina reniformia, compressa, tuberculata”) and by Boulos (1999: “seeds ... muricate-tuberculate”). In fact there is a part of the seed with muricate-tuberculate surface; this is the pole opposite the micropyle (Fig. 3a, c). The seed of *B. ramonensis*, in contrast, is covered by tubercles throughout (Fig. 4). The stellate cells of the lateral seed surface of *B. multiceps* (Fig. 3d) markedly differ from the tubercular surface of *B. ramonensis* (Fig. 4d).

Specimens of *Bufonia multiceps* seen: S Sinai, Gebel Katherina, 2 km S of top, 2400 m, 17.7. 1968, *Tadmor* (HUJ); Gebel Abu Hisheib (N of wadi Sulaf), 9.9.1968, *Orshan* (HUJ).

Acknowledgements

I thank Mr M. Dvorachek, The Geological Survey of Israel, for the scanning electron micrographs, Dr N. Kilian for the critical reading of a draft of the manuscript and checking the Latin diagnosis, Dr J. McNeill and Dr J. Chrtek, Senior for peer-reviewing the manuscript and Dr U. Ravid and Dr N. Dudai, Ministry of Agriculture, Israel, for their help in the fieldwork.

References

- Boissier, E. 1867: *Flora orientalis* **1**. – Genève.
- Boulos, L. 1999: *Flora of Egypt* **1**. – Cairo.
- Chrtek, J. & Křisa, B. 1999: A revision of Asian species of the genus *Bufonia* L. – *Acta Univ. Carol., Biol.* **43**: 77-118.
- Danin, A. 1983: *Desert vegetation of Israel and Sinai*. – Jerusalem.
- 1999: Desert rocks as plant refugia in the Near East. – *Bot. Rev.* **65**(2): 93-170.
- & Solomeshch, I. A. 1999: Synopsis of the vegetation and enumeration of the associations. – Pp. 33-316 in: Danin, A. & Orshan, G. (ed.), *Vegetation of Israel I. Desert and coastal vegetation*. – Leiden.
- Decaisne, M. J. 1835: *Florula sinaica* 2. – *Ann. Sci. Nat. Bot.*, ser. 2, **3**: 257-291.
- Mouterde, P. S. J. 1966: *Nouvelle flore du Liban et de la Syrie* **1**. – Beyrouth.
- Rechinger, K. H. 1988: *Buffonia*. – Pp. 114-124 in: Rechinger, K. H. (ed.), *Flora iranica* **163**. – Graz.

Address of the author:

Avinoam Danin, Department of Evolution, Systematics, and Ecology, The Hebrew University of Jerusalem, Jerusalem, Israel 91904; e-mail: danin@vms.huji.ac.il