



## The problem of *Citharexylum spinosum* (Verbenaceae) in the Antilles

Author: Méndez, Isidro

Source: Willdenowia, 31(2) : 419-424

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.31.31209>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISIDRO MÉNDEZ

## The problem of *Citharexylum spinosum* (*Verbenaceae*) in the Antilles

### Abstract

Méndez, I.: The problem of *Citharexylum spinosum* (*Verbenaceae*) in the Antilles. – Willdenowia 31: 419-424. 2001. – ISSN 0511-9618.

As part of the revision of the genus *Citharexylum* for the projects “Flora of the Cuban Republic” and “Flora of the Greater Antilles”, conspecificity of *C. spinosum* and plants assigned by many authors to a different species, misnamed “*C. fruticosum*” (an illegitimate name), is demonstrated. Intraspecific classification of *C. spinosum* is critically reconsidered, resulting in five new combinations, three of them at a new rank.

### Introduction

Numerous authors have approached the taxonomic problem surrounding *Citharexylum spinosum* L., especially in relation to *C. fruticosum* L. (Schulz 1909, Alain 1957, Moldenke 1958a-b, 1959, Görts-van Rijn 1988, Howard 1989, Liogier 1994, 1995, Acevedo 1996, among others). Méndez & Cafferty (2001) demonstrated that *C. fruticosum* is an illegitimate name used by Linnaeus in his work “Systema Naturae ed. 10” (1759), in replacement of *C. spinosum*. The present article summarizes the taxonomic and nomenclatural approaches that the author has assumed for the purposes of the projects “Flora of the Cuban Republic” and “Flora of the Greater Antilles.”

Linnaeus established *Citharexylum* in 1753 as a monotypic genus for the species *C. spinosum*. In 1759, although he continued recognizing the existence of a single species, he substituted the previous name with *C. fruticosum*. During the first three quarters of the twentieth century, all authors who studied this group in the Caribbean region accepted the existence of two different species and made some effort to justify their distinction (Schulz 1909, Moldenke 1939, Alain 1957, Moldenke 1958a-b, 1959, Liogier 1994, 1995). More recently, the existence of two independent taxa has been questioned, among others by: (1) Görts-van Rijn (1988), who considered that in Guyana only *C. spinosum* exists, and included *C. fruticosum* L. as a synonym; (2) Howard (1989), who suggested that, in the Lesser Antilles, *C. fruticosum* is only an ecological variant of *C. spinosum*; (3) Acevedo (1996), who predicted that exhaustive studies could demonstrate that both binomials are synonymous, with *C. spinosum* having priority; (4) Méndez & Cafferty (2001), who demonstrated the illegitimacy of *C. fruticosum*.

Nevertheless, the taxonomic and nomenclatural problems are not completely clarified, because numerous heterotypic synonyms exist. An infraspecific system under *Citharexylum fruti-*

*cosum* auct. (non L.) has been established and hybrids have been pointed out between the two supposed species and between each of them and *C. caudatum* L., whose situation should be clarified.

The author who with the most precision established the distinction between *Citharexylum spinosum* L. and *C. fruticosum* auct. (non L.) was Schulz (1909). It was based on the consistency of the foliar lamina, the prominence of the veins on both surfaces of the leaves, the length of the floral pedicels, the configuration of the calyx border and the form of the fruit. Moldenke (1939) accepted this approach without major modifications and, based on it, he distinguished hybrids between the two species and between each of them and *C. caudatum* (Moldenke 1959).

Field observations (carried out by the author during 17 years of work in Cuba and Puerto Rico) and the study of more than 300 herbarium specimens (from B, BM, BR, C, F, FL, GH, HAC, HAJB, HIPC, HIVC, HPPR, JBSD, JE, M, NY, PR, S and US), including six types, have confirmed that the characters presented by Schulz (1909) are expressed in a continuous way and that the differential details cited are extreme manifestations that rarely appear jointly in a single individual, as had been described. The existence of a single species (previously outlined by Görts-van Rijn (1988) and suggested by Howard (1989) and Acevedo (1996)) was verified, in spite of the variability that sustains the infraspecific system established under *C. fruticosum* auct. (non L.)

A critical revision of this system is summarised below, with the new combinations that are necessary in the light of our current knowledge.

- Citharexylum spinosum* L., Sp. Pl.: 625. 1753  $\equiv$  *Citharexylum cinereum* L., Sp. Pl., ed. 2: 872. 1763, nom. illeg.  $\equiv$  *Citharexylum fruticosum* L., Syst. Nat., ed. 10: 1115. 1759, nom. illeg. – Lectotype (Howard in Bot. J. Linn. Soc. 79: 82. 1979): [icon] “*Citharexylum americanum* alterum, foliis ad marginem dentatis” in Plukenet, Phytographia: t. 161, f. 5. 1692. Epitype (Méndez & Cafferty in Taxon 50, in press. 2001): “Barbados, St. Andrew,” 9.1940, *Gooding* 228 (BM [photo!]).
- $\equiv$  *Citharexylum quadrangulare* Jacq., Enum. Syst. Pl. Carib.: 26. 1760. – Type not designated.
- $\equiv$  *Citharexylum teres* Jacq., Select. Stirp. Amer. Hist.: 185, t. 118. 1763. – Lectotype (designated here): [icon] Jacquin, Select. Stirp. Amer. Hist.: t. 118. 1763!, based on material from Martinique.
- $\equiv$  *Citharexylum cinereum* Jacq. ex J. F. Gmel., Syst. Nat., ed. 13, 2: 492. 1789, nom. illeg. – Holotype [icon] Gmelin, Syst. Nat., ed. 13, 2: t. 178. 1789 [n.v.], based on material from Martinique.
- $\equiv$  *Citharexylum pentandrum* Vent., Descr. Pl. Jard. Cels: 47. 1800. – Type not designated.
- $\equiv$  *Citharexylum surrectum* Griseb., Fl. Brit. W. I.: 497. 1861. – Type not designated.
- $\equiv$  *Citharexylum hibridum* Mold. in Lilloa 4: 313. 1939 (pro hybr. *C. fruticosum* L.  $\times$  *C. spinosum* L.). – Holotype: Tobago, Rockkley Vale, 21.7.1910, *Brodway* 4064 (NY [n.v.]).
- *Citharexylum fruticosum* auct. (non L., Syst. Nat., ed. 10: 1115. 1759).

*Shrub or tree*, 4-18 m tall; trunk slender. *Branches* generally 4-angled, sometimes cylindrical and grooved; glabrous except at the nodes. *Leaves* opposite-decussate; petioles 5-30 mm, glabrous, with a gland at the apex; lamina orbicular, elliptic, oblong, lanceolate-elliptic, lanceolate, or linear-lanceolate, 4-20 cm, generally chartaceous, rarely membranaceous, with 1-3 pairs of glands near the base; acute, obtuse or emarginate at the apex; cuneate at the base; glabrous above, glabrous or with variable pubescence below; margin somewhat revolute, usually entire, some juvenile leaves sometimes subserrate toward the apex, with teeth up to 4 mm; veins prominent on both surfaces. *Inflorescences* mostly terminal or, more rarely, axillary, simple or branched with two or more basal branches, 9-30 cm long, laxly many-flowered; axis glabrous; bracteoles filiform to lanceolate, sometimes with hairs. *Pedicels* 1-1.5 mm long. *Flowers* fragrant. *Calyx* pale green, 2-4 mm, glabrous outside; margin ciliate, irregular or 5-toothed. *Corolla* white, pale yellowish or pale reddish; tube 4-10 mm, 2-3 times as long as the calyx; limb 5-parted, 6-7 mm in diameter. *Drupe* oblong, 6-10 mm in diameter, initially yellow-orange, black when mature; fruiting calyx hardened, cup-shaped.

*Distribution.* – Bermuda, Antilles, Venezuela and Guyana.

The species is extremely variable as to indumentum, form consistency and border of the leaf lamina. A thorough study of various populations revealed that the phenotypic expression of these characters is continuous and that the morphological segregation is unrelated to phytogeographical or ecological patterns. For such variation, the form is the most appropriate rank at the infra-specific level.

### Key to the formae

1. Leaves mostly glabrous below, if pubescent then only in the axils of the secondary veins . . . 2
- Leaves pubescent below, at least along the midvein (not only in the axils of the secondary veins) . . . . . 4
2. Leaves pubescent below in the axils of the secondary veins, all with an entire margin . . . . . 1. *C. spinosum* f. *spinosum*
- Leaves totally glabrous, some subserrate at the apex . . . . . 3
3. Leaves linear-lanceolate, up to 6 times as long as wide. . . . . 2. *C. spinosum* f. *smallii*
- Leaves elliptic, lanceolate-elliptic or rarely lanceolate, but in such cases only some leaves up to 4 times as long as wide . . . . . 3. *C. spinosum* f. *subserratum*
4. Hairs present throughout the lower surface of the leaf, margin entire or subserrate toward the apex . . . . . 4. *C. fruticosum* f. *villosum*
- Hairs dispersed, mostly limited to the veins, margin entire . . . . . 5
5. Leaves membranaceous, veins scarcely prominent . . . . . 5. *C. spinosum* f. *brittonii*
- Leaves chartaceous or subcoriaceous, veins prominent . . . . . 6. *C. spinosum* f. *subvillosum*

#### 1. *Citharexylum spinosum* L. f. *spinosum*

*Leaves* elliptic, lanceolate-elliptic or rarely lanceolate (but in such cases only some leaves up to 4 times as long as wide), margin entire, pubescent below only in the axils of the secondary veins.

*Distribution.* – Bermuda, Antilles, Venezuela and Guyana. For additional specimens examined see the electronic supplement at <http://www.bgbm.fu-berlin.de/bgbm/library/publikat/willd31/mendez.htm>.

The lectotype selected by Howard (1979) offers very limited information about the varying morphology to which the binomial was originally assigned, but the most evident character relates to the leaf margin, a detail that he repeats in three of the lines of variation within the species. The typical form is assigned to this phenotype, based on the epitype selected by Méndez & Cafferty (2001).

2. *Citharexylum spinosum* f. *smallii* (Mold.) I. E. Méndez, **comb. & stat. nov.** ≡ *Citharexylum fruticosum* var. *smallii* Mold. in Repert. Spec. Nov. Regni Veg. 37: 223. 1934. – Holotype: Bahamas, Andros, coppice near Deep Oreek, Long Bay Cay Sections, 20.-22.1.1910, *Small & Carter 8583* (NY!).

*Leaves* linear-lanceolate, up to 6 times as long as wide; margin entire or 2-4-toothed toward the apex (on some leaves of young branches of certain specimens).

*Distribution.* – Bahamas, Jamaica, Puerto Rico. Alain (1957) and Moldenke (1958, 1980) reported this form also from Cuba. For additional specimens examined see the electronic supplement at <http://www.bgbm.fu-berlin.de/bgbm/library/publikat/willd31/mendez.htm>.

3. *Citharexylum spinosum* f. *subserratum* (Sw.) I. E. Méndez, **comb. nova** ≡ *Citharexylum subserratum* Sw., Prodr.: 91. 1788 ≡ *Citharexylum fruticosum* var. *subserratum* (Sw.) Mold. in

Phytologia 1: 17. 1933 ≡ *Citharexylum fruticosum* f. *subserratum* (Sw.) Mold. in Phytologia 36: 164. 1977. – Lectotype (Moldenke in Phytologia 6: 384. 1958): Española, Swartz (S [n.v.]).

*Leaves* elliptic, lanceolate-elliptic, or rarely lanceolate (in such cases only some up to 4 times as long as wide), totally glabrous below; margin entire or sometimes subserrate toward the apex (teeth in extreme cases up to 4 mm).

*Distribution.* – Hispaniola (Alain 1957, Moldenke 1958, 1980); Liogier (1994) reported this form also from Cuba. For additional specimens examined see the electronic supplement at <http://www.bgbm.fu-berlin.de/bgbm/library/publikat/willd31/mendez.htm>.

**4. *Citharexylum spinosum* f. *villosum*** (Jacq.) I. E. Méndez, **comb. & stat. nov.** ≡ *Citharexylum villosum* Jacq., Icon. Pl. Rar. 1: 12. 1786 ≡ *Citharexylum fruticosum* var. *villosum* (Jacq.) O. E. Schulz in Urban, Symb. Antill. 6: 63. 1900. – Holotype: [icon] Jacquin, Icon. Pl. Rar. 1: 12. 1786 [n.v.], based on a cultivated plant in the Botanic Garden of Vienna, Austria, apparently grown from seeds obtained from Santo Domingo.

= *Citharexylum polystachyum* Turcz. in Bull. Soc. Imp. Naturalistes Moscou 36: 209. 1863 – Holotype: “Cuba, from Santiago to Río Guaso, Oriente”, 1837, *Linden 1975* (KWU [n.v.]).

*Leaves* elliptic or oblong, rarely orbicular; pubescent, with dense hairs on all veins and on leaf tissue between the veins; margin entire or subdentate toward the apex.

*Distribution.* – Cuba, Hispaniola, Puerto Rico, Bahamas. For additional specimens examined see the electronic supplements at <http://www.bgbm.fu-berlin.de/bgbm/library/publikat/willd31/mendez.htm>.

**5. *Citharexylum spinosum* f. *brittonii*** (Mold.) I. E. Méndez, **comb. & stat. nov.** ≡ *Citharexylum fruticosum* var. *brittonii* Mold. in Lilloa 4: 311. 1939 ≡ *Citharexylum broadwayi* O. E. Schulz in Urban, Symb. Antill. 7: 354. 1912. – Holotype: Cedros Islands, 15.1.1908, *Broadway 3198* (B†).

*Leaves* elliptic, lanceolate-elliptic, or rarely lanceolate (in such cases only some up to 4 times as long as wide), membranaceous, veins hardly prominent, pubescent below, hairs scattered, mostly limited to the nerves, margin entire.

*Distribution.* – Trinidad.

*Note.* – For this project it was not possible to study either the holotype (destroyed in B) or additional specimens from Trinidad. The description upon which the new combination and status are based, is taken from Moldenke (1958a: 366).

**6. *Citharexylum spinosum* f. *subvillosum*** (Mold.) I. E. Méndez, **comb. nova** ≡ *Citharexylum fruticosum* f. *subvillosum* (Mold.) Mold. in Phytologia 36: 164. 1977 ≡ *Citharexylum fruticosum* var. *subvillosum* Mold. in Repert. Spec. Nov. Regni Veg. 37: 223. 1934. – Holotype: Cuba, Guantánamo Bay, 17.-30.3.1909, *Britton 2101* (NY!).

= *Citharexylum bahamense* Millsp. in Bull. New York Bot. Gard. 3: 450. 1905 ≡ *Citharexylum fruticosum* var. *bahamense* (Millsp.) Mold. in Phytologia 13: 242. 1966. – Holotype: Bahamas, New Providence, coppice Farrington Road, 24.7.1904, *Britton & Brace 233a* (NY!; isotype US).

*Leaves* lanceolate, chartaceous or subcoriaceous, pubescent below, hairs scattered, on the nerves and leaf tissue, margin entire.

*Citharexylum fruticosum* f. *subvillosum* (Mold.) Mold. was established originally to differentiate individuals with prominent veins. This is actually unfounded in practice, because the conspicuous veins are inherent to the whole species.

*Distribution.* – Cuba, Hispaniola, Bahamas, Virgin Islands. For additional specimens examined see the electronic supplement at <http://www.bgbm.fu-berlin.de/bgbm/library/publikat/willd31/mendez.htm>.

### Supposed hybrids

Harold Moldenke has also described two hybrids involving the foregoing species.

*Citharexylum* × *jamaicense* Mold. in Phytologia 6: 416. 1958 (*C. caudatum* L. × *C. fruticosum* L.). – Holotype: Jamaica, *Perkins 419* (B<sup>†</sup>).

*Citharexylum* × *perkinsii* Mold. in Phytologia 6: 496. 1959 (*C. caudatum* L. × *C. spinosum* L.). – Holotype: Jamaica, near Troy, in pasture, 1500 m, *Perkins 1320* (B<sup>†</sup>; isotypes G, N?, Z [n.v.]). – Moldenke (1980) reported this hybrid for Jamaica, Hispaniola and Puerto Rico.

It was not possible to study the holotypes (destroyed in B), but numerous specimens coming from Jamaica were evaluated, identified by Moldenke with these names. All fall within the range of variability of *C. caudatum* L., and the intermediate characters between the supposed parents are not evident in any of them.

### Acknowledgements

I thank the Alexander von Humboldt Foundation for the George Foster fellowship to undertake a postdoctoral study at the Botanic Garden and Botanical Museum Berlin-Dahlem, thanks to which it was possible to conclude these results. I am also grateful to the directors and curators of many herbaria (BM, BR, C, F, FL, GH, JBSD, JE, M, NY, PR, S and US) for sending material of *Verbenaceae* s.str. on loan for the purpose of this study. Thanks are due to Steve Cafferty for helping me with the clarification of the Linnaean nomenclature and for sending photos of herbarium specimens from BM. Thanks to the Atlantea Project of the University of Puerto Rico it was possible to do field and herbarium studies on that island. Dr Werner Greuter and Dr Duane Kolterman I thank for the revision of the text.

### References

- Acevedo-Rodríguez, P. 1996: Flora of St. John, U. S. Virgin Islands. – Mem. New York Bot. Gard. **78**.
- Alain, 1957: Flora de Cuba 4. – Contr. Ocas. Mus. Hist. Nat. Colegio “De La Salle” La Habana **16**.
- Gooding, E. G. B., Loveless, A. R. & Proctor, G. R. 1965: Flora of Barbados. – London.
- Görts-van Rijn, A. R. A. 1988: *Verbenaceae*. – In: Jansen-Jacobs, M. J. (ed.), Flora of the Guianas **4**. – Königstein.
- Howard, R. A. 1979: Early botanical records from the West Indies, particularly Barbados: Ligon (1657) to Lord Seaforth (1806). – Bot. J. Linn. Soc. **79**: 65-96.
- 1989: Flora of the Lesser Antilles **6**. – Cambridge, MA.
- Liogier, A. H. 1994: Flora de La Española VI. – Universidad Central del Este, San Pedro de Macorís, República Dominicana. Ser. Ci. **27**.
- 1995: Descriptive flora of Puerto Rico and adjacent islands **4**. – Río Piedras, PR.
- Méndez, I. & Cafferty, S. 2001: Typification of Linnaean names of taxa of *Verbenaceae* s.s. described from the Greater Antilles. – Taxon **50** (in press).
- Moldenke, H. N. 1939: The “*Verbenaceae*” and “*Avicenniaceae*” of Trinidad and Tobago. – Lilloa **4**: 283-336.
- 1958a: Materials toward a monograph of the genus *Citharexylum* III. – Phytologia **6**: 332-368.
- 1958b: Materials toward a monograph of the genus *Citharexylum* IV. – Phytologia **6**: 383-432.
- 1959: Materials towards a monograph of the genus *Citharexylum* VII. – Phytologia **7**: 49-77.

- 1980: A sixth summary of the *Verbenaceae*, *Avicenniaceae*, *Stilbaceae*, *Chloanthaceae*, *Symphoremaceae*, *Nyctanthaceae*, and *Eriocaulaceae* of the world as to valid taxa, geographic distribution, and synonymy. – *Phytologia Mem.* **2**.
- Schulz, O. E. 1909: *Citharexylum* Linn. – In: Urban, I. (ed.), *Symbolae antillanae* **6**: 56-69. – Leipzig, etc.

Address of the author:

Isidro E. Méndez Santos, Pedagogical College “José Martí”, Camagüey-6, CP-74670, Cuba;  
e-mail: mendez@ispcmw.rimed.cu.