

## **Euro Med Notulae, 5**

Authors: Greuter, Werner, and Raab-Straube, Eckhard Von

Source: Willdenowia, 41(1) : 129-138

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.41.41117>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

WERNER GREUTER<sup>1</sup> & ECKHARD VON RAAB-STRAUBE<sup>1\*</sup> (ed.)**Euro+Med Notulae, 5****Abstract**

Greuter W. & Raab-Straube E. von (ed.): Euro+Med Notulae, 5 [Notulae ad floram euro-mediterraneam pertinentes 27]. – Willdenowia 41: 129–138. – Online ISSN 1868-6397; © 2011 BGBM Berlin-Dahlem.  
doi: 10.3372/wi.41.41117 (available via <http://dx.doi.org/>)

This is the fifth of a series of miscellaneous contributions, by various authors, where hitherto unpublished data relevant to the Euro+Med (or Sisyphus) Project are presented. This instalment deals with the families *Betulaceae*, *Chenopodiaceae*, *Leguminosae*, *Plumbaginaceae*, *Portulacaceae*, *Ranunculaceae*, *Salicaceae* and *Gramineae*, including new country and area records for taxa of *Avena*, *Chamaecytisus* (*Cytisus*), *Dichanthium*, *Eragrostis*, *Festuca*, *Holcus*, *Poa* and *Portulaca*, and the validation of names in the genera *Alnus*, *Anemonastrum*, *Dactylis*, *Homalotrichon*, *Limonium*, *Salix* and *Salsola*.

Additional key words: Europe, Mediterranean area, vascular plants, taxonomy, distribution

**Notice**

A succinct description of the Euro+Med Project, with a list of recognised territories and their abbreviations, and the conventions used to indicate the status and presence of taxa, can be found in the introduction to the first instalment (Greuter & Raab-Straube 2005: 223–226) and on the Euro+Med Plantbase website (Euro+Med 2006+). As of 30.4.2011, Euro+Med Plantbase provides access to 135 families, corresponding to approximately 85 % of the Euro-Mediterranean flora of vascular plants. For the previous instalment of the Euro+Med Notulae, see Greuter & Raab-Straube (2009).

The following have contributed entries to the present instalment: Z. Barina, K. Baumann, M. Breitfeld, A. Danin, G. Domina, H. Freitag, D. Pifkó, T. Raus, M. Ristow, H. Scholz, A. Sennikov, P. Uotila and B. Valdés.

*Betulaceae*

*Alnus alnobetula* subsp. *crispa* (Aiton) Raus, **comb. nov.** ≡ *Betula crispa* Aiton, Hort. Kew. 3: 339. 1789 ≡ *Alnus viridis* subsp. *crispa* (Aiton) Turrill in Bot. Mag. 173:

382. 1962 ≡ *Duschekia crispa* (Aiton) Pouzar in Preslia 36: 339. 1964.

*Alnus alnobetula* subsp. *fruticosa* (Rupr.) Raus, **comb. nov.** ≡ *Alnus fruticosa* Rupr. in Beitr. Pflanzenk. Russ. Reiches 2: 53. 1845 ≡ *Alnus viridis* subsp. *fruticosa* (Rupr.) Nyman, Consp. Fl. Eur.: 672. 1881 ≡ *Duschekia fruticosa* (Rupr.) Pouzar in Preslia 36: 339. 1964.

*Alnus alnobetula* subsp. *sinuata* (Regel) Raus, **comb. nov.** ≡ *Alnus viridis* var. *sinuata* Regel in Bull. Soc. Imp. Naturalistes Moscou 38: 422. 1865 ≡ *Alnus sinuata* (Regel) Rydb. in Bull. Torrey Bot. Club 24: 190. 1897 ≡ *Alnus viridis* subsp. *sinuata* (Regel) Á. Löve & D. Löve in Univ. Colorado Stud., Ser. Biol. 17: 20. 1965 ≡ *Duschekia sinuata* (Regel) Pouzar in Preslia 36: 339. 1964.

According to Pouzar (1982) and Holub (1986), *Alnus viridis* (Chaix) DC. is based on *Betula viridis* Chaix (1785, in Villars 1786; for bibliographical details see Perret & Burdet 1981), a name preceded by *B. alnobetula* Ehrh. (Ehrhart 1783; for bibliographical details see

<sup>1</sup> Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin-Luise-Str. 6–8, 14195 Berlin, Germany; \*e-mail: [e.raab-straube@bgbm.org](mailto:e.raab-straube@bgbm.org) (author for correspondence), [w.greuter@bgbm.org](mailto:w.greuter@bgbm.org)

Manitz 1975). That is why the taxon's correct name in *Alnus* is *A. alnobetula* (Ehrh.) K. Koch (first invalidly published as a synonym of *A. ovata* (Schränk) Lodd. in Hartig 1843: 372, later validated by Koch 1872). From a nomenclatural point of view, this affects all infraspecific taxa of the species accepted and keyed out on subspecific level in circumpolar basic floras (Tutin & al. 1993; Morin & al. 1997), necessitating the above combinations in addition to the Corsican endemic, *A. alnobetula* subsp. *suaveolens* (Req.) Lambinon & Kerguélen (Lambinon & Kerguélen 1988). *A. alnobetula* subsp. *alnobetula* (syn. *A. viridis* subsp. *viridis*) is endemic to the mountains of Central Europe and the Balkan Peninsula. The N Eurasian-NW North American *A. alnobetula* subsp. *fruticosa* is, in Europe, confined to Arctic Russia and the Urals E of 45°W and N of 60°30'N (Komarov 1936; see also map V-47 in Tolmachev & al. 2000: 138, under *Alnaster fruticosa* (Rupr.) Ledeb.). *A. alnobetula* subsp. *crispa* from NE North America and Greenland and *A. alnobetula* subsp. *sinuata* from NW North America are both reported to be planted for shelter in Iceland, the Faroe Islands and Scandinavia, locally frequently so (Jonsell 2000); they may escape from cultivation.

Th. Raus

### Chenopodiaceae

The heterogeneity of the traditional genus *Salsola* has been known for some time (e.g., Freitag & Rilke 1997, Kadereit & al. 2003). Akhiani & al. (2007) dismembered *Salsola* on the basis of molecular results. However, for the time being, a broad generic concept is followed in Euro+Med Plantbase, because no consensus on a subdivision of the genus has yet been reached, and because of missing information on the appropriate placement in the new segregates for some of the taxa involved. Likewise, the genus *Bassia* has been treated in a wide sense. For a molecular-based new classification see Kadereit & Freitag (2011). We thank Alexander Sennikov for nomenclatural advice.

P. Uotila & H. Freitag

***Salsola boissieri* subsp. *serpentinicola*** (Freitag & Özhatay) Freitag & Uotila, **comb. nov.** = *Salsola canescens* subsp. *serpentinicola* Freitag & Özhatay in *Willdenowia* 27: 185. 1997. – A new combination is needed because *Salsola canescens* (Moq.) Boiss., Fl. Orient. 4: 963. 1879, is a later homonym of *Salsola canescens* Pers., Syn. Pl. 1: 296. 1805. The new name *S. boissieri* was introduced by Botschantzev (1968: 1442), who believed that the earlier name *S. canescens* Desf., Tabl. École Bot.: 41. 1804 was validly published, whereas in fact *S. canescens* Desf. appears in an enumeration of garden plants without any associated descriptive matter. Therefore, *S. boissieri* was rejected as nomen superfluum by Freitag (in Freitag & Rilke 1997: 231). However, both Botschantzev and Freitag overlooked *S. canescens* Pers. (Syn. Pl. 1: 296. 1805), which was neither included in Index Kewensis nor in IPNI, but had already been detected by Greuter & al.

(1984: 308). Although *S. canescens* Pers. is a superfluous substitute for *Chenopodium sinense* Willem. (today's *Chenolea diffusa* Thunb.) and therefore illegitimate, it is validly published and thus an earlier homonym of *S. canescens* (Moq.) Boiss. Consequently *S. boissieri* must be considered the correct name.

P. Uotila & H. Freitag

### Leguminosae

***Chamaecytisus leiocarpus*** (A. Kern.) Rothm. (*Cytisus leiocarpus* A. Kern.) – Flowers of *Chamaecytisus leiocarpus* are all arranged in leafy racemes, the shoots are covered with appressed hairs and the legume is glabrous. Based on this combination of features, the taxon is clearly distinct from all other *Chamaecytisus* taxa with flowers exclusively or partly arranged in leafy racemes. The flowers of two subspecies of the related *C. triflorus* (Lam.) Skalická (*Chamaecytisus hirsutus* auct. non (L.) Link), subsp. *ciliatus* (Wahlenb.) Holub and subsp. *falcatus* (Waldst. & Kit.) Pifkó, are arranged in racemes and their legumes are glabrous, but the shoots are either covered with patent hairs or glabrescent. The legume is also glabrous in *C. supinus* subsp. *alpestris* (Schur) Pifkó, which has patent hairs on the shoots (Pifkó 2009). The related *C. ratisbonensis* (Schaeff.) Rothm. and *C. elongatus* (Waldst. & Kit.) Link also have flowers in leafy racemes and shoots covered with appressed hairs, but their legumes are densely hairy.

+ **Mk:** Former Yugoslav Republic of Macedonia: W slope of Jakupica Mts, above village Belica, in rocky pine forest, 689 m, 41.6867°N, 21.3143°E, 7.2.2010, Z. Barina & D. Pifkó 17701 (BP). – New to the country. Micevski (2001) does not discuss this name nor any of its synonyms. *Chamaecytisus leiocarpus* occurs in the NE part of the Balkan Peninsula and in the adjacent part of the Carpathian Basin: in Bulgaria (Kuzmanov 1976), in Bosnia and Herzegovina (Beck 1887; as *Cytisus ciliatus* var. *bosniacus* Beck), in Serbia (Diklić 1972) and in the southwestern part of Romania (Oprea 2005).

? **Al:** A record of *Chamaecytisus leiocarpus* from Albania is included in Flora Europaea (Heywood & Frodin 1968); however, its basis is unknown. Later mentions from Albania (Greuter & al. 1989; Cristofolini 1991; Qosja & al. 1992) are presumably based on that work and also lack mention of localities. The line drawings in the Albanian Floras (Demiri 1983; Qosja & al. 1992) lack essential details and, anyway, are just copies of those in the Bulgarian Flora (Kuzmanov 1976: 83). In the herbaria revised by us (BEO, BEOU, BP, KRA, KRAM, SO, SOM) and those revised by Cristofolini (1991), no specimens of *C. leiocarpus* (or originally determined as *C. leiocarpus*) have been found, and we could not confirm the presence of the species in Albania during our field trips to the country from 2004 until today.

- ? **Tu(E)**: The only record from Turkey (Cristofolini 1991) should also be treated as questionable because of doubt on the precise origin of the only herbarium specimen from “Turkey (Trakya!)”, because the territory of formerly Turkish Thrace extends to today’s Bulgaria, Turkey, Greece and the Former Yugoslav Republic of Macedonia. Gibbs (1969) does not mention the taxon from Turkey.
- **Gr**: The report from Greece is erroneous. The single Greek record (Cristofolini 1991) is based on a reference to Flora Europaea (Heywood & Frodin 1968) where, however, the species is not mentioned from Greece.
- **Uk(U)**: There is a single record of this taxon from Ukraine (Fodor 1974), repeated later by Yakovlev & al. (1996) and Kagalo (1999). However, the only specimen (collected in 1966, at Onokivci, Zakarpattia Oblast) determined as *C. leiocarpus* in the herbarium of István Fodor (at UU) proved to be *C. ratisbonensis* (Schaeff.) Rothm. (rev. Pifkó 2007).  
D. Pifkó & Z. Barina

### Plumbaginaceae

Pollen-stigma dimorphism and self-incompatibility are common within *Statioideae* and widespread in *Limonium*. Facultative apomixis is common in *Limonium*, whereas truly sexual species are infrequent. Hybridisation, apomixis and polyploidy are putative key factors in the radiation of *Limonium* in the Mediterranean basin (Artelari & Georgiou 2003; Cowan & al. 1998; Erben 1979; Lledó & al. 2005). As a consequence, more than 400 microspecies and a large number of hybrids have been described from the Mediterranean area. According to the current trend in *Limonium* classification, it is appropriate to consider the taxa below at specific rather than subspecific rank.

G. Domina

***Limonium afrum*** (Pignatti) Domina ≡ *Limonium delicatulum* subsp. *afrum* Pignatti in Collect. Bot. (Barcelona) 6: 308. 1962.

***Limonium ammophilon*** (Papatsou & Phitos) Domina ≡ *Limonium graecum* subsp. *ammophilon* Papatsou & Phitos in Notes Roy. Bot. Gard. Edinburgh 34: 203. 1975.

***Limonium gallicum*** (Pignatti) Domina ≡ *Limonium delicatulum* subsp. *gallicum* Pignatti in Collect. Bot. (Barcelona) 6: 316. 1962. – Type gathering: Soc. Dauphinoise No. 1817; lectotype (designated here): “*Statice ovalifolia* Poir., Ile d’Aix (Char.-Inf.)”, 26.7.1875. A. Guillon in Société Dauphinoise 1817 (FI [fototeca 2635/A!]; isolecotype: FI [fototeca 2635!]).

***Limonium maroccanum*** (Batt. & Trab.) Domina ≡ *Statice tubiflora* var. *maroccana* Batt. & Trab. in Bull. Soc. Hist. Nat. Afrique N. 9: 15. 1918. ≡ *Limonium tubiflorum*

subsp. *maroccanum* (Batt. & Trab.) Maire & Weiller in Bull. Soc. Hist. Nat. Afrique N. 27: 244. 1936.

***Limonium romanum*** (Täckh. & Boulos) Domina ≡ *Limonium sinuatum* subsp. *romanum* Täckh. & Boulos in Publ. Cairo Univ. Herb. 5: 90. 1974.

***Limonium zanonii*** (Pamp.) Domina ≡ *Statice tubiflora* var. *zanonii* Pamp. in Nuovo Giorn. Bot. Ital., ser. 2, 24: 148. 1917 ≡ *Limonium tubiflorum* subsp. *zanonii* (Pamp.) Brullo in Webbia 33: 142. 1978.

### Portulacaceae

Since my first publication on the *Portulaca oleracea* complex (Danin & al. 1978), I have tried to determine every *Portulaca* specimen I came across, using the keys in Danin & Reyes-Betancort (2006), Danin & al. (2008), and Danin & al. (2011). Many of my determinations remain unpublished, most of which, when relevant to the Euro+Med area, are assembled here. An in-depth description of my research on *P. oleracea* can be found on the website: <http://flora.huji.ac.il/browse.asp?action=specie&specie=POROLE>.

From the random sample of Euro-Mediterranean collections of *Portulaca oleracea* aggr. studied by me, derived from several hundred populations, one may draw the following preliminary conclusions. The most common microspecies is *P. granulatostellulata*, recorded from 28 territories. Second is *P. trituberculata* (24 territories). The status of occurrence is still poorly understood. In the following list, the taxa have been considered as native when there is no evidence to the contrary. Native (including archaeophytic) status is supported by early findings of *P. trituberculata* in Germany in the 4th century AD. (Danin & al. 1978). New findings from the Po Valley near Ferrara (Bosi & al. 2009; and unpublished data) support such early presence. The introduction and extinction of microspecies of the *P. oleracea* aggr. is a dynamic processes, far from being fully understood.

I acknowledge the advice, and help in obtaining specimens, of P. Uotila, T. Raus, D. Blanár, C. Dobes, V. Ferák-ová, M. Hassan, P. Pils, G. Schneeweiss, W. Starmühler, A. Sukhorukov, J. Walter, A. Zeddám; and the herbarium keepers of B, E, G, H HUJ and MW.  
A. Danin

***Portulaca canariensis*** Danin & Reyes-Betancort

+ **Ca(C)**: Canary Islands (Spain): Gran Canaria, Agüimes, Barranco de Agüimes, 24.3.1978, I. Kukkonen 9733 (H).

+ **Ca(G)**: Canary Islands (Spain): La Gomera, Chejelpes, 4.4.2007, A. Kurtto & L. Helynranta (H).

***Portulaca cypria*** Danin

+ **Ag**: Algeria: City of Algiers, centre, 13.8.2006, A. Zeddám (B); Algiers, Institut Agronomique, 28.8.2006, A. Zeddám (B).

- + **Co**: Corsica: Entre Ajaccio et la Cove de Povata, 19.7.1900, *E. Burnat, J. Briquet & F. Cavillier* (G).
- + **Eg**: Egypt: Cairo, 5.11.1982, *M. A. Monier* 4278 (E).
- + **Ir**: Israel: Philistean Plain, Rehovot, 28.1.2011, *A. Danin* (HUI); Judean Mts, Jerusalem, 10.10.2005, *A. Danin* (HUI).
- + **Sy**: Syria: Palmyra, 3.8.1944, *Mouterde* 8268 (G).

***Portulaca granulatostellulata*** (Poelln.) Ricceri & Arrigoni

- + **Ag**: Algeria: Algiers, Institut National Agronomique, 28.8.2006, *A. Zeddami* (B); Algiers, Hydra, 28.8.2006, *A. Zeddami* J77 (B).
- + **Au(A)**: Austria: Niederösterreich, Dep. Breitenau, 8.9.1992, *J. Walter* (WU); Wien, 28.7.1975, *J. Walter* (WU).
- + **Az(J)**: Azores (Portugal): São Jorge, Calheta, 28.6.1938, *C. Cedercantz* (H).
- + **Co**: Corsica (France): Aléria, 7.2004, *J.-M. Tison* (HUI); 12.7.2000, *C. Dobes* (herb. C. Dobes).
- + **Ct**: Croatia: Korčula, Lumbarda, 22.6.1971, *K. Kaleva* 1241 (H); Istria, Cres / Nerezio, 9.8.1997, *J. Walter* (herb. J. Walter).
- + **Eg**: Egypt: Cairo, American University, 2.7.2009, *M. Hassan EG-04* (B); southern Egypt, Safaga, 29.10.2009, *M. Hassan EG-25* (B); Giza, on the Nile bank, 2.6.1965, *M. El Mahdi* (G).
- A Fe**: Finland: Ta. Janakkala, Turenki, 7.8.1959, *H. Niininen* (H); Obu. Kemi, Laitakari, 19.7.1914, *V. Räsänen* (H).
- + **Ga(F)**: France: Alpes-Maritimes, Mandelieu, 9.2000, *J.-M. Tison* (HUI); Var, Hyères, 9.2000, *J.-M. Tison* (HUI).
- + **Gg**: Georgia: Tbilisi, 27.8.2000, *G. Schneeweiß* (herb. G. Schneeweiß).
- + **Gr**: Greece: Macedonia, Grevena – Kozani, 18.8.1976, *W. Greuter* 14477 (G); Athens airport, 7.10.2004, *A. Danin* Gr2004Aa (HUI).
- + **He**: Switzerland: Alpes Lemaniennes, 7.9.1904, *J. Briquet* (G); Genève, 10.9.1970, *Terretaz* 2021 (G).
- + **Hu**: Hungary: Zala, Karmacs, Keszthely, 27.9.2010, *P. Uotila* 48917 (H).
- + **Ir**: Israel: Esdraelon Plain, Afula, 7.11.2005, *A. Danin* (HUI); Philistean Plain, Gan Yavne, 29.7.2005, *A. Danin* (HUI).
- + **Md(M)**: Madeira (Portugal): Ponta de San Lorenzo, cono volcanico, 14.9.2004, *A. Reyes-Betancort & A. Santos* (HUI).
- + **Po**: Poland: Kornik, 8.2007, *Boratyński* (HUI).
- + **Rf(C)**: Russia: Moscow region, Orekhovo-Zuevo distr., Krutoe, 10.9.1983, *I. V. Ivanova* (MW).
- + **Sk**: Slovakia: Muránska planina plateau, Tisovec, 18.8.2009, *D. Blánár* (herb. Blánár); Slovenské rudohorie, Lubeník, railway station, 18.8.2009, *D. Blánár* (herb. Blánár).
- + **Sl**: Slovenia: Portorož, 16.9.1983, *A. Leskinen* (H).

- + **Sr**: Serbia: Beograd, Botanical Garden, 10.9.2004, *A. Danin* (HUI); Deliblato, 8.9.2004, *A. Danin* (HUI); Pessara, 8.9.2004, *A. Danin* (HUI); Kladovo, 12.9.2009, *P. Uotila* 48316 (H); Đerdap gorge near Golubac, 12.9.2009, *P. Uotila* 48316 (H); Maljen, Divičbare, 9.9.2009, *P. Uotila* 48278a (H); Šumadija, Lajkovac, 9.9.2009, *P. Uotila* 48280 (H); Niš, 14.9.2009, *P. Uotila* 48423 (H).
- A Su**: Sweden: Blekinge, Sälvesborg, 1.8.1936, *B. Holmgren* (H).
- + **Tu(A)**: Turkey: B3 Eskişehir, Eskişehir, Anadolu University campus, 29.6.2007, *A. Danin* (HUI); A2(A) Istanbul, Istanbul, Bostancı, 1.7.2007, *A. Danin* (HUI); C2 Muğla, c. 10 km SE of Köyceğiz, Beyobasi, 19.5.1990, *P. Uotila* 37854 (H).
- + **Tu(E)**: Turkey-in-Europe: A1(E) Kırklareli, Babae-ski, 1.7.2007, *A. Danin* (HUI); A1(E) Çanakkale, Gelibolu, 3.7.2007, *A. Danin* (HUI).

***Portulaca nicaraguensis*** (Danin & H. G. Baker) Danin

- D Az(J)**: Azores (Portugal): São Jorge, Calheta, 3.7.1938, *C. Cedercantz* (H).
- D Ca(G)**: Canary Islands (Spain): La Gomera, Valle Gran Rey, 5.4.2007, *A. Kurtto & L. Helynranta* (H), *ibid.*, 7.4.2007, *A. Kurtto & L. Helynranta* (H). – For the Macaronesian islands, this predominantly North and Central American taxon was hitherto only known from Tenerife (Danin & Reyes-Betancort 2006).

***Portulaca nitida*** (Danin & H. G. Baker) Ricceri & Arrigoni

- + **Ag**: Algeria: Algiers, Brossette, 14.9.2006, *A. Zeddami* (B); Algiers, Hydra, 29.8.2006, *A. Zeddami* (B).
- + **Au(A)**: Austria: Wien XII, 25.6.2003, *J. Walter* (WU); Salzburg, 15.9.2002, *P. Pilsl* (herb. Pilsl).
- + **Be(B)**: Belgium: Liège, 9.10.1989, *J. Rousselle* (H).
- + **Cg**: Montenegro: Miločer, 1.7.1973, *J. Jalas* 2428 (H).
- + **Ct**: Croatia: Istria, Inner-Istria, SSE Lupoglav, 4.9.2003, *W. Starmühler* (herb. Starmühler); Istria, 15.10.1998, *W. Starmühler* (herb. Starmühler).
- + **Eg**: Egypt: Cairo, American University, 2.7.2009, *M. Hassan EG-08* (B); Giza, on the Nile bank, 2.6.1965, *M. El Mahdi* (G).
- A Fe**: Finland: Ab. Sauvo, Karuna, Voilahti, 17.9.1999, *L. Hörkö* (H); N. Pernaja, Isnäs, Labby, 20.9.2003, *P. Alanko* 119799 (H); N. Helsinki, Kaivopuisto, 20.9.1995, *C. af Forselles* (H).
- + **Ga(F)**: France: Allier, Chassenard, 22.9.1961, *A. Charpin* (G).
- + **Ge**: Germany: Bayern, Passau, Platz bei der Nibelungenhalle, 27.8.2003, *M. Hohla* (WU).
- + **Gr**: Greece: Athens airport, 7.10.2004, *A. Danin* (HUI); Mesogia, 10 km N of Athens Airport, 7.10.2004, *A. Danin* (HUI); Grammatiko, 7.10.2004, *A. Danin* (HUI).
- + **He**: Switzerland: Genève, 8.1877, *E. Ayasse* (G).

- + **Ir:** Israel: Hula Plain, Kibbutz Dan, 25.7.2005, *A. Danin* (HUI); Philistean Plain, Gan Yavne, 29.7.2005, *A. Danin* (HUI).
- + **Md(M):** Madeira (Portugal), 25.8.2004, *A. Reyes-Betancort* (HUI).
- + **Po:** Poland: Kornik, 8.2007, *Boratyński* (HUI).
- + **Rf(C):** Russia: Tambov province, Kirsanov, railroad bed near Kirsanov station, 3.7.1997, *A. Sukhorukov* (MW).
- + **Rf(S):** Russia: Kalmykia, Chernoziemsky, 24.7.1964, *V. Pavlov* (MW).
- + **Tu(A):** A2(A) Istanbul, Istanbul, Bostancı, 1.7.2007, *A. Danin* (HUI).
- + **Tu(E):** Turkey-in-Europe: A1(E) Kırklareli, Babae-ski, 1.7.2007, *A. Danin* (HUI); A1(E) Çanakkale, Gelibolu, 3.7.2007, *A. Danin* (HUI).

*Portulaca oleracea* L. s.str. [= *P. stellata* (Danin & H. G. Baker) Ricceri & Arrigoni]

- A **Fe:** Finland: Sa. Imatra, Neitsytniemi, 10.1999, *K. Rutanen* (H).
- + **Ga(F):** France: Bouches-du-Rhône, Fontvieille, 10.2001, *J.-M. Tison* (HUI).
- + **Ge:** Germany: Rheinland-Pfalz, Rheinhessen, Mainz-Mombach, 20.8.1978, *H. Kahlheber* (H).
- + **He:** Switzerland: Genève, 8.1877, *E. Ayasse* (G); Plan-tae Helvetiae et Sabaudiae, 1864, *H. Bernet* (G).
- + **Hu:** Hungary: Budapest, 3.10.2010, *P. Uotila 49008* (H).
- + **Ir:** Israel: N Negev, Arad, 15.10.1981, *A. Danin* (HUI).
- + **Lu:** Portugal: 16.6.1998, *J. Walter* (herb. Walter); Beira Alta, 12.9.1984, *M. Luceño* (UC).
- + **Po:** Poland: Kornik, 8.2007, *Boratyński*, (HUI); N Poland, Torun-Bielany, 20.9.1979, *M. Ceynowa-Gieldon 337* (H).
- + **Si(M):** Malta: NW of Mellieħa, 24.10.1974, *L. Y. T. Westra & J. V. Rooden* (E).
- + **Tu(E):** Turkey-in-Europe: A2(E) Istanbul, Kumburgaz, 2.7.2007, *A. Danin* (HUI); A1(E) Edirne, İpsala, 2.7.2007, *A. Danin* (HUI); A1(E) Çanakkale, Ge-libolu, 3.7.2007, *A. Danin* (HUI).

*Portulaca papillatostellulata* (Danin & H. G. Baker) Danin

- + **Ag:** Algeria: Algiers, Zéralda, 19.11.2005, *A. Zeddami* (B).
- + **Au(A):** Austria: Niederösterreich, Breitenau, 8.9.1992, *J. Walter* (WU); Wien, 9.9.1992, *J. Walter* (WU).
- + **Co:** Corsica (France): 5.7.2000, *C. Dobes* (herb. C. Dobes).
- A **Fe:** Finland: Al. Finström, Godby, 8.1980, *P. Johans-son* (H); N. Helsinki, Kaisaniemi, 5.9.1984, *P. Al-anko 49584* (H).
- + **Gr:** Greece: Macedonia, Halkidiki, Sithonia Peninsula, Sarti, 4.7.1997, *M. Koistinen 1997/275* (H).
- + **Hu:** Hungary: Zala, Karmacs, 27.9.2010, *P. Uotila 48906* (H).

- + **Ir:** Israel: N Negev, Arad, 15.10.1981, *A. Danin* (HUI).
- + **It:** Italy: Toscana, Valle Bandetta, c. 15 km E of Li-vorno, 13.9.2007, *A. Danin* (HUI); Toscana, Gab-bro, c. 15 km SE of Livorno, 13.9.2007, *A. Danin* (HUI); Lazio, Rome, “Ateneo Pontificio Regina Apostolorum”, 15.9.2007, *A. Danin* (HUI); To-scana, Elba, 9.2007, *A. Reyes-Betancort* (HUI).
- + **Rf(C):** Russia: Lipetsk oblast, Usman, 11.9.1917, *P. Smirnov* (MW).
- + **Sk:** Slovakia: Bratislava IV–Devín, Námestie práce square, 20.8.2010, *V. Feráková* (herb. V. Feráko-vá); Malé Karpaty Mts., castle Červený Kameň near Častá, 16.9.2010, *V. Feráková* (herb. V. Feráková).
- + **Tu(A):** Turkey: A2(A) Istanbul, Istanbul, Bostancı, 1.7.2007, *A. Danin* (HUI).
- + **Tu(E):** Turkey-in-Europe: A1(E) Kırklareli, Babaeski, 1.7.2007, *A. Danin* (HUI).

*Portulaca rausii* Danin

- + **Ag:** Algeria: Algiers, centre, 13.8.2006, *A. Zeddami* (B).
- + **Gr:** Greece: Attiki, Elefsis, 27.5.1939, *H. Lindberg* (H).
- + **Tu(A):** Turkey: B1 Aydın, Ephesos, 18.8.2004, *J. Wal-ter* (herb. J. Walter).
- + **Tu(E):** Turkey-in-Europe: A1(E) Kırklareli, Lülebur-gaz, 1.7.2007, *A. Danin* (HUI).

*Portulaca sativa* Haw.

- C **Fe:** Finland: N. Helsinki, Viikki, Institute of Horticul-ture, cultivated, 19.9.1969, *P. Alanko 14153* (H); N. Helsinki, Kaisaniemi, Bot. Garden, 11.9.1913, *V. Pesola* (H).
- C **Tu(A):** Turkey: Erzurum, seed bought from bazar in 1972, cult. in Finland, Helsinki, 4.9.1972, *P. Uo-tila 21090* (H).

*Portulaca trituberculata* Danin, Domina & Raimondo

- + **Ag:** Algeria: Algiers, Hussein Dey, 6.9.2006, *A. Zed-dami* (B); Algiers, Brossette, 13.8.2006, *A. Zed-dami* (B).
- + **Au(A):** Austria: Wien, 7.8.1995, *J. Walter* (WU); Nieder-österreich, Gramatneusiedl, 15.8.1994, *W. Re-hak* (WU).
- + **Co:** Corsica: Entre Ajaccio et la Cave de Povata[?], 19.7.1900, *E. Burnat, J. Briquet & F. Cavillier* (G).
- + **Eg:** Egypt: New Cairo, 30.6.2009, *M. Hassan EG-01* (B).
- N **Fe:** Finland: Ta. Hattula, Tenhola, 30.7.2005, *P. Uotila 45687* (H).
- + **Ga(F):** France: Isère, Heyrieux, 9.2000, *J.-M. Tison* (HUI); Bas-Rhin, Roppenheim, 13.8.1994, *A. Schneider* (H).
- + **Gr:** Greece: Macedonia, Grevena–Kozani, 18.8.1976, *W. Greuter 14477* (G); Mani Peninsula, 22.11.1991, *Kit Tan & G. Vold 10666* (G); 4 km from Paralia Tirou to Leonidion, 10.9.1997, *Kit Tan & A. Strid 18712* (HUI); Athens airport, 7.10.2004, *A. Da-*

- nin* (HUJ); Mesogia, 10 km N of Athens airport, 7.10.2004, A. Danin (HUJ).
- + **He**: Switzerland: Valais, 27.9.1897, P. Cheneverd (G); Vaud, 24.8.1929, O. Meylan (G).
  - + **Ho**: Netherlands: Gelderland, Wilp, 18.7.1955, W. C. van Heurn (H).
  - + **Hu**: Hungary: Zala, Karmacs, 27.9.2010, P. Uotila 48909 (H); Veszprem. Egyházaskesö, 29.9.2010, P. Uotila 48961 & al. (H).
  - + **Ir**: Israel: Sharon Plain, Netanya, 14.10.2005, A. Danin (HUJ); Judean Mts, Jerusalem, 10.10.2005, A. Danin (HUJ).
  - + **It**: Italy: Toscana, Pisa, 13.9.2007, A. Danin (HUJ); Lazio, Rome, "Ateneo Pontificio Regina Apostolorum", 15.9.2007, A. Danin (HUJ).
  - + **Ma**: Morocco: Sidi Hamed el Sunni, 15.7.1929, F. Quer 122 (G).
  - + **Po**: Poland: Kornik, 8.2007, Boratyński (HUJ).
  - + **Rf(C)**: Russia: Kursk city, 1906, V. Alyekhin (MW).
  - + **Rm**: Romania: Dobrogea, 1971, V. Feráková 27 (Herb. V. Feráková).
  - + **Si(M)**: Malta: Gnien Ingraw, just NW of Mellieħa, 24.10.1974, L. Y. T. Westra & J. V. Rooden 204 (G).
  - + **Sn**: Egypt: Sinai, 1930, A. Kaiser 853 (G).
  - + **Sr**: Serbia: Beograd, 10.9.2004, A. Danin (HUJ); Beograd, Kalemegra, 5.9.2004, A. Danin (HUJ); Niš–Dobrič Valley, 14.9.2009, P. Uotila 48451 (H).
  - + **Sy**: Syria: Damascus, 28.5.1951, H. Pabot (G); Baniyas, 22.10.1953, H. Pabot (G).
  - + **Tu(A)**: Turkey: B3 Eskişehir, Eskişehir, Anadolu University campus, 29.6.2007, A. Danin (HUJ); A2(A) Istanbul, Istanbul, Bostancı, 1.7.2007, A. Danin (HUJ); A1(A) Çanakkale, Denizkent, shore of the Sea of Marmara, 22.7.1981, P. Uotila 30390 (H); A2(A) Bursa, İznik Gölü, 10 km E of İznik, 21.7.1981, K. Hormia 453 (H).
  - + **Tu(E)**: Turkey-in-Europe: A1(E) Kırklareli, Babae-ski, 1.7.2007, A. Danin (HUJ); A1(E) Çanakkale, Gelibolu, 3.7.2007, A. Danin (HUJ).

### *Portulaca zaffranii* Danin

- + **Ag**: Algeria: Algiers, city centre, 19.11.2005, A. Zeddam (B).
- + **Gr**: Greece: Mesogia, 10 km N of Athens airport, 7.10.2004, A. Danin (HUJ).
- + **It**: Italy: Emilia-Romagna, Lugo (RA), Parco del Loto, 5.8.2010, G. Bosi 34 (MOD).
- + **Sy**: Syria: Damascus, 24.6.1927, G. von Frenckell (H).
- + **Tu(A)**: Turkey: B3 Eskişehir, Eskişehir, Anadolu University campus, 29.6.2007, A. Danin (HUJ).
- + **Tu(E)**: Turkey-in-Europe: A1(E) Edirne, Keşan, 3.7.2007, A. Danin (HUJ).

### Ranunculaceae

#### *Anemonastrum narcissiflorum* subsp. *crinitum* (Juz.)

Raus, **comb. nov.** ≡ *Anemone crinita* Juz. in Komarov,

Fl. URSS 7: 739. 1937 ≡ *Anemone narcissiflora* subsp. *crinita* (Juz.) Kitag., in Rep. Inst. Sci. Res. Manchoukuo 3, App. 1.: 213. 1939.

= *Anemone biarmiensis* Juz. in Komarov, Fl. URSS 7: 738. 1937 ≡ *Anemonastrum biarmiense* (Juz.) Holub in Folia Geobot. Phytotax. 12: 428. 1977 ≡ *Anemone narcissiflora* ["*narcissifolia*"] subsp. *biarmiensis* (Juz.) Jalas in Ann. Bot. Fenn. 25: 297. 1988.

#### *Anemonastrum narcissiflorum* subsp. *fasciculatum* (L.)

Raus, **comb. nov.** ≡ *Anemone fasciculata* L., Sp. Pl.: 542. 1753 ≡ *Anemonastrum fasciculatum* (L.) Holub in Folia Geobot. Phytotax. 8: 165. 1973 ≡ *Anemone narcissiflora* subsp. *fasciculata* (L.) Ziman & Fedor. in Ziman & al., Taxon. & Evol. *Anemone narcissiflora* complex: 34. 1997.

= *Anemone umbellata* Willd., Sp. Pl. 2: 1284. 1799.

= *Anemone speciosa* Adams ex Pritz. in Linnaea 15: 685. 1841 ≡ *Anemonastrum speciosum* (Pritz.) Galushko, Fl. Severn. Kavkaza Vopr. Ist. 3: 56. 1979.

= *Anemone narcissiflora* var. *willdenowii* Boiss., Fl. Orient. 1: 14. 1867 ≡ *Anemone narcissiflora* subsp. *willdenowii* (Boiss.) P. H. Davis in Notes Roy. Bot. Gard. Edinburgh 26: 175. 1965.

= *Anemone narcissiflora* subsp. *chrysanth* Ulbr. in Bot. Jahrb. Syst. 37: 266. 1906 ≡ *Anemone chrysanth* (Ulbr.) Grossh., Fl. Kavkaza 2: 105. 1930.

= *Anemone impexa* Juz. in Komarov, Fl. URSS 7: 737. 1937 ≡ *Anemonastrum impexum* (Juz.) Holub in Folia Geobot. Phytotax. 8: 165. 1973.

Fischer & al. (2008) and Buttler & Hand (2008), based on Ehrendorfer & Samuel (2001), accept that the generic segregation of *Hepatica* Mill. and *Pulsatilla* Mill. from *Anemone* L. s.l., as commonly accepted (Komarov 1937; Greuter & al. 1989; Tutin & al. 1993), necessitates the same procedure for *Anemonastrum* Holub in order to allow for taxonomic consistency. According to the recent in-depth revision of infraspecific taxa in *Anemone* sect. *Omalocarpus* DC. by Ziman & al. (2005), *Anemonastrum narcissiflorum* (L.) Holub is represented in the Euro+Med area by three subspecies, viz. subsp. *narcissiflorum* in Central and S Europe from N Spain to SW Ukraine (Jalas & Suominen 1989: 80, map 1638) including the Crimea (Ziman & al. 2005: 294), subsp. *fasciculatum* in N and E Asia Minor (Davis 1965: 135) and the Caucasus (Russia, Georgia and Armenia; see maps 39 & 40 in Grossheim 1950 under *Anemone fasciculata*, *A. impexa* and *A. speciosa*), and subsp. *crinitum* at its westernmost distribution limit in the Urals (mapped by Jalas & Suominen 1989; the total range of the latter subspecies extending through the whole of Siberia eastwards to Kamchatka and N Japan, Ziman & al. 2005: 295-296). Subspecific (plus ancillary varietal) instead of specific rank for taxa within the *Anemonastrum narcissiflorum* complex proves appropriate since they exhibit sufficient geographical separation (Hultén 1986: 1049–1050, map

829) being at the same time connected by numerous transitional forms, probably due to parallel radiations into similar habitats (Ehrendorfer 1995). Th. Raus

### Salicaceae

*Salix pentandrifolia* Sennikov, **nom. nov.** = *Salix skvortsovii* Sennikov in Komarovia 4: 140. 2006, non Y. L. Chang & Y. L. Chou (in Chou & al. 1955: 86 [n.v.]) [– *Salix pentandroides* A. Skvortsov in Feddes Repert. Spec. Nov. Regni Veg. 64: 73. 1961, nom. inval. (Art. 37.1 of the Code; McNeill & al. 2006), non *S. pentandroides* Rouy in Rev. Bot. Syst. Géogr. Bot. 2: 168. 1904]. – Holotype: Russia, “Caucasus septentr., Balkaria, in ripa rivuli Baschyl-sugusu, alt. 1950 m”, 1.9.1939, *R. Elenevsky* (MW).

*Salix skvortsovii* Sennikov was published as a new species name because the previous name used for this species, *S. pentandroides* A. Skvortsov, was found not to be validly published, with two types designated. It was dedicated to Alexei K. Skvortsov (1920–2008), the renowned Russian taxonomist of willows. Unfortunately, at that time I overlooked the existence of the earlier name *S. skvortzovii* Y. L. Chang & Y. L. Chou, which was dedicated to another Russian botanist with the same surname, Boris V. Skvortsov (1896–1980), who worked e.g. in Manchuria in NE China and used to transliterate his name as “Skvortzov”. The combinations of letters “tz” and “ts” stay for the same Russian letter “т” and have identical pronunciation, the first following the traditions of German (the dominant language of scientific writing and transliteration in B. Skvortsov’s times), the second, contemporary English style.

Following a request by Irina Belyaeva, the Nomenclature Committee for Vascular Plants recommended that *Salix skvortsovii* and *S. skvortzovii* be considered as confusingly similar and treated as homonyms (Brummitt 2009: 291). For this reason, a new name is needed for the Caucasian representative of the *Salix pentandra* L. group, distinguished by A. Skvortsov on the basis of its narrow and acute (not broad and obtuse) leaf-buds and concolorous leaves. A. Sennikov

### Gramineae

*Avena fatua* subsp. *cultiformis* Malzev

+ **Po**: Poland: N edge of Warthebruch, NW of Jenin, c. 0.5 km N of the main road, field, 52°42'06.2"N, 15°06'02.8"E, 40 m, 26.6.2010, *M. Ristow* 460/10 (B); edge of Warthetal, N above Santok, c. 1.2 km W of Warthe bridge, field border with *Tordylium maximum*, *Cyanus segetum*, *Cota tinctoria* and *Consolida regalis*, 52°44'05.6"N, 15°23'30.9"E, 40 m, 27.6.2010, *M. Ristow* 482/10 (B). – “This subspecies is closely related to *A. sativa* subsp. *orientalis* possibly being either a fatuoid mutation of the latter subspecies or its hybrid with *A. fatua* s.l.” (Tzvelev 1983: 350). M. Ristow & H. Scholz

*Dactylis glomerata* subsp. *merinoana* (Horjales & al.) H. Scholz, **comb. nov.** = *Dactylis izcoi* subsp. *merinoana* (“*merinoiana*”) Horjales & al. in Nova Acta Ci. Compostelana, Biol. 18: 166. 2010.

*Dactylis glomerata* subsp. *stebbinsii* (Horjales & al.) H. Scholz, **comb. nov.** = *Dactylis juncinella* subsp. *stebbinsii* (“*stebbinis*”) Horjales & al. in Nova Acta Ci. Compostelana, Biol. 17: 81. 2009.

*Dichanthium annulatum* (Forssk.) Stapf

**P Ca(F)**: Canary Islands (Spain): Fuerteventura, Costa Calma, centre of the village, 11.2008, *H. Langbehn* (B). – Second record from the Canary Islands. In this region first collected in 1998 in Tenerife (Cruz & al. 2008). Naturalised in, e.g. Madeira, Crete and countries of N Africa and the Near East. H. Scholz

*Eragrostis tenuifolia* (A. Rich.) Steud.

**A Ca(F)**: Canary Islands (Spain): Fuerteventura, Costa Calma, centre of the village, 11.2008, *H. Langbehn* (B). – A species of probably African origin, now widespread throughout the Tropics. Also known from Madeira. H. Scholz

*Festuca brachyphylla* Schult. & Schult. f.

+ **Su**: Sweden: Torne Lappmark; Fjäll, gravel- and moorplain between Alesjaurestugorne and Tjåktjastugan, c. 35 km SW of Abisko, 750–1000 m, 21.8.1990, *R. W. Bussmann* (STU). – The area of this fescue extends from Svalbard and N Norway to the arctic parts of European and Asian Russia. H. Scholz

*Holcus rigidus* Hochst.

**A Ge**: Germany, N. Bavaria: Construction site of a new highway bridge W of Waldau (5935), 12.7.2010, *M. Breitfeld* (UBT). – An endemic of the Azores, this species had never before been recorded as a casual alien. M. Breitfeld & H. Scholz

*Homalotrichon pubescens* subsp. *longifolium* (Boiss.)

H. Scholz & Valdés, **comb. nov.** = *Avenula pubescens* var. *longifolia* Boiss., Fl. Orient. 5: 545. 1884 = *Avenula pubescens* subsp. *longifolia* (Boiss.) H. Scholz & Valdés in Willdenowia 36: 662. 2006 = *Helictotrichon pubescens* subsp. *longifolium* (Boiss.) Doğan in Notes Roy. Bot. Gard. Edinburgh 40: 86. 1982.

*Poa poiformis* (Labill.) Druce (*Arundo poiformis* Labill.; “*P. caespitosa*” auct. non G. Forst.; “*P. labillardierei*” auct. non Steud.).

**P Ge**: Germany, Hesse: Frankfurt-Niedereschbach, fallow land on Homburger Landstraße (5717/44), 140 m, 3.7.1991, *H. Kramer* 91/45 (FR, 2 specimens). – For this densely tufted perennial from Australia,

compare Walsh & al. (2009: 335). Perhaps the records of *Poa labillardierei* Steud. from the British Isles, as a wool casual (Ryves & al. 1996), also belongs here. K. Baumann & H. Scholz

## References

- Akhani H., Edwards G. & Roalson E. H. 2007: Diversification of the Old World *Salsolae* s.l. (*Chenopodiaceae*): molecular phylogenetic analysis of nuclear and chloroplast data sets and a revised classification. – *Int. J. Pl. Sci.* **168**: 931–956.
- Artelari R. & Georgiou O. 2003: Biosystematic study of the genus *Limonium* (*Plumbaginaceae*) in the Aegean area, Greece III. *Limonium* on the islands Kithira and Antikithira and the surrounding islets. – *Nordic J. Bot.* **22**: 483–501.
- Beck G. 1887: Flora von Südbosnien und der angrenzenden Hercegovina. – *Ann. K. K. Naturhist. Hofmus.* **2**: 142–183.
- Bosi G., Guarrera P. M., Rinaldi R. & Bandini Mazzanti M. 2009: Ethnobotany of purslane (*Portulaca oleracea* L.) in Italy and morfo-biometric analyses of seeds from archaeological sites of Emilia Romagna (Northern Italy). – Pp. 129–139 in: Morel J. P. & Mercuri A. M. (ed.), Plants and culture: seeds of the cultural heritage of Europe. – Bari: Edipuglia.
- Botschantzev V. P. 1968: Obzor vidov sekcii *Belanthera* Iljin roda *Salsola* L. – *Bot. Zhurn.* **53**: 1440–1450.
- Brummitt R. K. 2009: Report of the Nomenclature Committee for Vascular Plants: 60. – *Taxon* **58**: 280–292.
- Buttler K. P. & Hand R. 2008: Liste der Gefäßpflanzen Deutschlands. – *Kochia Beih.* **1**: 1–107.
- Chaix D. 1785: Plantae vapincenses sive enumeratio in agro vapincensi observatarum stirpium. – Baux: [the author].
- Chaix D. 1786: Plantae vapincenses, sive Enumeratio Plantarum in agro vapincensi a valle le Valgaudemar, ad amniculum le Buech, prope Segesteronem, sponte nascentium, aut œconomice cicurum. – Pp. 309–382 in: Villars D., Histoire des plantes de Dauphiné **1**. – Grenoble: L'auteur [Reprint of Chaix 1785]
- Chou Y. L., Huang T. C., Zhang Y. L., Li C. T. & Zhao T. C. 1955: Xiao Xinganling muben zhiwu [Woody plants of the Xiao Hingan Mountains]. – Beijing: Zhongguo linye chubanshe.
- Cowan R., Ingrouille M. J. & Lledó M. D. 1998: The taxonomic treatment of agamosperms in the genus *Limonium* Mill. (*Plumbaginaceae*). – *Folia Geobot.* **33**: 353–366.
- Cristofolini G. 1991: Taxonomic revision of *Cytisus* Desf. sect. *Tubocytisus* DC. (*Fabaceae*) – *Webbia* **45**: 187–219.
- Cruz Trujillo G. M., Rodríguez Delago O. & Wildpret de la Torre W. 2008: *Dichanthium annulatum* (Forssk.) Stapf, una gramínea nueva para las Islas Canarias. – *Bot. Macaronés.* **27**: 137–140.
- Danin A., Baker I. & Baker H. G. 1978: Cytogeography and taxonomy of the *Portulaca oleracea* L. polyploid complex. – *Israel J. Bot.* **27**: 177–211.
- Danin A., Domina G. & Raimondo F. M. 2008: Microspecies of the *Portulaca oleracea* aggregate found on major Mediterranean islands (Sicily, Cyprus, Crete, Rhodes). – *Fl. Medit.* **18**: 89–108.
- Danin A., Marrosu M., Caria M. C. & Bagella S. 2011: The status of *Portulaca oleracea* aggregate in Sardinia (Italy) and a key to its 19 microspecies. – *Pl. Biosystems* (in press).
- Danin A. & Reyes-Betancort J. A. 2006: The status of *Portulaca oleracea* L. in Tenerife, the Canary Islands. – *Lagascalia* **26**: 71–81.
- Davis P. H. (ed.) 1965: Flora of Turkey and the East Aegean Islands **1**. – Edinburgh: Edinburgh University.
- Demiri M. 1983: Flora ekskursioniste e Shqipërisë. – Tiranë: Shtëpia Botuese e Librit Shkollor.
- Diklić N. 1972: *Chamaecytisus* Link. – Pp. 497–515 in: Josifović M. (ed.), Flora SR Srbije **4**. – Beograd: Srpska Akad. Nauka i Umetnosti.
- Domina G., Schäfer P. A. & Danin A. 2010: Typification and taxonomic status of *Portulaca macrantha* (*Portulacaceae*). – *Fl. Medit.* **20**: 187–191.
- Ehrendorfer F. 1995: Evolutionary trends and patterns in the *Anemoninae*. – *Pl. Syst. Evol., Suppl.* **9**: 283–293.
- Ehrendorfer F. & Samuel R. 2001: Contributions to a molecular phylogeny and systematics of *Anemone* and related genera (*Ranunculaceae-Anemoninae*). – *Acta Phytotax. Sin.* **39**: 293–307.
- Ehrhart J. F. 1783: Bestimmung einiger Bäume und Sträucher aus unsern Lustgebüschchen. – *Gartenkalender* **1783**: 189–192.
- Erben M. 1979: Karyotype differentiation and its consequences in Mediterranean *Limonium*. – *Webbia* **34**: 409–417.
- Euro+Med 2006+ [continuously updated]: Euro+Med Plantbase – the information resource for Euro-Mediterranean plant diversity. – Published at <http://ww2.bgbm.org/EuroPlusMed> [accessed 23.3.2011].
- Fischer M., Oswald K. & Adler W. (ed.) 2008: Exkursionsflora für Österreich, Liechtenstein und Südtirol, ed. 3. – Linz: Oberösterreichische Landesmuseen.
- Fodor S. S. 1974: Flora Zakarpatija. – L'viv: Vidavn. Ob'edinenie Vishcha Shkola.
- Freitag H. & Özhatay E. 1997: A new subspecies of *Salsola canescens* (*Chenopodiaceae*) from SW Anatolia, Turkey. – *Willdenowia* **27**: 185–190.
- Freitag H. & Rilke S. 1997: *Salsola*. – Pp. 154–255 in: Rechinger K. H. (ed.), Flora iranica **173**. – Graz: Akademische Druck- und Verlagsanstalt.
- Gibbs P. E. 1969: *Chamaecytisus* Link – Pp. 16–21 in: Davis P. H. (ed.), Flora of Turkey and the East Aegean islands **3**. – Edinburgh: Edinburgh University.
- Greuter W., Burdet H. M. & Long G. 1984, 1989: Med-Checklist **1**, **4**. – Genève: Conservatoire et Jardin

- botaniques, Ville de Genève; Berlin: Botanischer Garten & Botanisches Museum Berlin-Dahlem.
- Greuter W. & Raab-Straube E. von 2005: Euro+Med Notulae, 1. – *Willdenowia* **35**: 223–239.
- Greuter W. & Raab-Straube E. von 2009: Euro+Med Notulae, 4. – *Willdenowia* **39**: 327–333.
- Grossheim A. A. 1950: Flora Kavkasa. **4**. – Moskva & Leningrad: Izdatel'stvo Akademii Nauk SSSR.
- Hand R. & Buttler K. P. 2006: Taxonomische und nomenklatorische Neuigkeiten zur Flora Deutschlands **1**. – *Kochia* **1**: 147–155.
- Hartig T. 1842–51: Vollständige Naturgeschichte der forstlichen Culturpflanzen Deutschlands. Heft 1–3 (1842), 4–10 (1843), 11–12 (1851). – Berlin: Förstner.
- Heywood V. H. & Frodin D. G. 1968: *Chamaecytisus* Link. – Pp. 90–93 in: Tutin T. G., Heywood V. H., Burges N. A., Moore D. M., Valentine D. H., Walters S. M. & Webb D. A. (ed.), *Flora europaea* **2**. – Cambridge: Cambridge University.
- Holub J. 1986: Comments on the “Med-Checklist 1”. – *Preslia* **58**: 289–306.
- Horjales M., Laso G. & Redondo N. 2009: *Dactylis glomerata* L. (*Poaceae*) en la Península Ibérica. – *Nova Acta Ci. Compostelana, Biol.* **17**: 65–85.
- Hultén E. & Fries M. 1986: Atlas of North European vascular plants north of the tropic of cancer **1–3**. – Königstein: Koeltz.
- Jalas J. & Suominen J. (ed.) 1989: Atlas Florae europaeae. Distribution of vascular plants in Europe **8**. – Helsinki: Committee for Mapping and the Flora of Europe.
- Jonsell B. (ed.) 2000: *Flora nordica* **1**. – Stockholm: Bergius Foundation.
- Kadereit G. & Freitag H. 2011: Molecular phylogeny of *Camphorosmeae* (*Camphorosmoideae*, *Chenopodiaceae*): implications for biogeography, evolution of C<sub>4</sub> photosynthesis and taxonomy. – *Taxon* **60**: 51–78.
- Kadereit G., Borsch T., Weising K. & Freitag H. 2003: Phylogeny of *Amaranthaceae* and *Chenopodiaceae* and the evolution of C<sub>4</sub> photosynthesis. – *Int. J. Pl. Sci.* **164**: 959–986.
- Kagalo A. A. 1999: Checklist of *Fabaceae* of the western Ukraine. – *Acta Bot. Fenn.* **162**: 137–140.
- Koch K. 1872: Dendrologie. Bäume, Sträucher und Halbsträucher, welche in Mittel- und Nord-Europa im Freien kultiviert werden, kritisch beleuchtet. **2/1**, die Mono- und Apetalen, mit Ausnahme der Cupuliferen, enthaltend. – Erlangen: Enke.
- Komarov V. L. (ed.) 1936, 1937: *Flora SSSR* **5, 7**. – Moskva & Leningrad: Izdatel'stvo Akademii Nauk SSSR.
- Kuzmanov B. 1976: *Chamaecytisus* Link. – Pp. 74–119 in: Jordanov D. (ed.): *Flora na Narodna Republika Bălgarija* **6**. – Sofija: Izdatelstvo na Bălgarskata Akademija na Naukite.
- Lambinon J. & Kerguélen L. 1988: Trois combinaisons nomenclaturales nouvelles relatives à la flore corse. – *Candollea* **43**: 405–406.
- Lledó M. D., Crespo M. B., Fay M. F. & Chase M. W. 2005: Molecular phylogenetics of *Limonium* and related genera (*Plumbaginaceae*): biogeographical and systematic implications. – *Amer. J. Bot.* **92**: 1189–1198.
- Manitz H. 1975: Friedrich Ehrhart's botanische Publikationen. – *Taxon* **24**: 469–474.
- McNeill J., Barrie F., Burdet H. M., Demoulin V., Hawksworth D. L., Marhold K., Nicolson D. H., Prado J., Silva P. C., Skog J. E., Wiersema J. H. & Turland N. J. 2006: International Code of Botanical Nomenclature (Vienna Code) adopted by the Seventeenth International Botanical Congress Vienna, Austria, July 2005. – *Regnum Veg.* **146**.
- Micevski K. 2001: *Flora na Republika Makedonija* **1(5)**. – Skopje: Makedonska Akademija na Naukite i Umetnostite.
- Morin N. R. & Flora of North America Editorial Committee (ed.) 1997: *Flora of North America north of Mexico* **3**. – New York: Oxford University.
- Oprea A. 2005: Lista critică a plantelor vasculare din România. – Iași: Univ. “Alexandru Ioan Cuza”.
- Papatsou S. & Phitos D. 1975: Two new taxa from the eastern Aegean. – *Notes Roy. Bot. Gard. Edinburgh* **34**: 203–204.
- Perret P. & Burdet H. M. 1981: 2. Les „Plantae Vapincenses“ de Dominique Chaix et les travaux floristiques de Dominique Villars en Dauphiné. – Pp. 400–408 in Burdet, H. M. (ed.), *Med-Checklist notulae bibliographicae*, 1 et 2. – *Candollea* **36**: 397–408.
- Pifkó D. 2009: Schur's *Cytisus* taxa. – *Stud. Bot. Hung.* **40**: 143–163.
- Pignatti S. 1962: Studi sui *Limonium*, V. Note sulla sistematica delle specie iberiche di *Limonium*. – *Collect. Bot. (Barcelona)* **6**: 293–330.
- Pouzar Z. 1982: Otázka správného jména pro olši zelednou. – *Čas. Nár. Muz. Praze, Rada Přír.* **151**: 20.
- Qosja X. (ed.) 1992: *Flora e Shqipërisë* **2**. – Tiranë: Akademia e Shkencave e Republikës së Shqipërisë.
- Ricceri C. & Arrigoni P. V. 2000: L'aggregato di *Portulaca oleracea* L. (*Portulacaceae*) in Italia. – *Parlatorea* **4**: 91–97.
- Ryves T. B., Clement E. J. & Foster M. 1996: *Alien Grasses of the British Isles*. – London: Botanical Society of the British Isles.
- Sennikov A. 2006: Miscellaneous nomenclatural notes (II). – *Komarovia* **4**: 138–154.
- Täckholm V. & Boulos L. 1974: Supplementary notes to the Students' Flora of Egypt, Second Edition. – *Publ. Cairo Univ. Herb.* **5**.
- Tolmachev A. I., Packer J. G. & Griffiths G. C. D. (ed.) 2000: *Flora of the Russian Arctic. A critical review of the vascular plants occurring in the Arctic region of the former Soviet Union*. **3**. – Berlin: Cramer.
- Tutin T. G., Burges N. A., Chater A. O., Edmondson J. R., Heywood V. H., Moore D. M., Valentine D. H., Walters S. M. & Webb D. A. (ed.) 1993: *Flora europaea*, ed. 2, **1**. – Cambridge: Cambridge University.

- Tzvelev N. N. 1983: Grasses of the Soviet Union **1**. – New Delhi: Amerind [Engl. transl. of: Cvelev, N. N. 1976: Zlaki SSSR. – Leningrad].
- Walsh N. G., Weiller C. M. & Thompson I. R. 2009: *Poa*. – Pp. 301–338 in: Wilson A. (ed.), Flora of Australia **44A**. – Melbourne: Australian Government Publ. Service.
- Yakovlev G. P., Sytin A. K. & Roskov Yu. 1996: Legumes of northern Eurasia. – Kew: Royal Botanic Gardens.
- Ziman S. N., Ehrendorfer F., Kadota Y., Keener C. S., Tsarenko O. N., Bulakh E. & Dutton B. E. 2005: A taxonomic revision of *Anemone* L. section *Omalocarpus* DC. sensu lato (*Ranunculaceae*): Part I. – J. Jap. Bot. **80**: 282–302.