

Current knowledge of trematodes (Platyhelminthes: Digenea, Aspidogastrea) in Chile

Authors: Oyarzún-Ruiz, Pablo, and González-Acuña, Daniel

Source: Revue suisse de Zoologie, 129(1) : 1-17

Published By: Muséum d'histoire naturelle, Genève

URL: <https://doi.org/10.35929/RSZ.0059>

BioOne Complete (complete.bioone.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Current knowledge of trematodes (Platyhelminthes: Digenea, Aspidogastrea) in Chile

Pablo Oyarzún-Ruiz^{1,2*} & Daniel González-Acuña¹

¹ Laboratorio de Parásitos y Enfermedades en Fauna Silvestre, Facultad de Ciencias Veterinarias, Universidad de Concepción, Chillán, Chile

² Becario ANID Programa de Doctorado en Ciencias Veterinarias, Facultad de Ciencias Veterinarias, Universidad de Concepción, Chillán, Chile

* Corresponding author: pablooyarzunruiz@gmail.com

Abstract: The aim of the present review was to evaluate the state of knowledge of trematodes (Platyhelminthes: Trematoda) in Chile, covering taxonomy, biology, geographic distribution, and a checklist of all taxa reported in the country. A total of 277 articles published between 1849 and December 2020, have been analyzed. A total of 215 taxa belonging to the subclasses Digenea and Aspidogastrea have been reported from Chile, 119 of them identified to species level and only three species with completely described life cycles. Trematodes from Chile were found parasitizing both native and exotic species, however, no records from insects and most crustacean classes exist. Overall, 159 out of 3989 potential host native species were reported as hosts for trematodes. Molecular tools were used in more recent taxonomic studies. Although research on trematodes in Chile is on the rise, there is a clear need for more detailed taxonomic studies to include integrative taxonomy approach, deposition of helminths in collections and the training of new generation of parasitologists to better understand the diversity, ecology and evolutionary relationships of trematodes in the region.

Keywords: Trematoda, Digenea, Aspidogastrea, biodiversity, checklist.

INTRODUCTION

The class Trematoda is subdivided in two subclasses, Aspidogastrea and Digenea. The former group is small and comprises only four families, 12 genera and about 80 species (Roberts *et al.*, 2013; Collins, 2017). On the other hand, digeneans are a very diverse group of parasites with more than 2500 genera and 18 000 nominal species (Kostadinova & Pérez-del-Olmo, 2014; Collins, 2017). However, there is a significant number of yet undescribed species, with some estimates suggesting that the total number of digenean species may exceed 44 000 (Carlson *et al.*, 2020a).

Gay (1849) was the first to mention a trematode in Chile. It was a report of the liver fluke *Fasciola hepatica* (Fasciolidae) in cattle. Since then, trematodes have been reported from a variety of hosts across the country. Previously published checklists were restricted to particular groups of hosts or types of ecosystems (see Garín & González-Acuña, 2008; Muñoz & Olmos, 2008; Oyarzún-Ruiz & González-Acuña, 2021), while an overview covering the whole trematode fauna in the country is currently lacking. The aim of the present review was to analyze the data on trematodes in Chile

and summarize of knowledge on their diversity, taxonomy, biology, and geographic distribution. We also outline the areas most urgently needing, attention and/or improvement in future research.

MATERIAL AND METHODS

Literature search

We performed an exhaustive bibliographic review of trematodes reported from domestic and wild animals, including humans, in Chile. Data from freshwater, marine and terrestrial environments, and all invertebrate and vertebrate hosts were included in this review. To achieve this objective, we analyzed 275 research articles, one book and one book chapter from physical libraries and electronic databases such as NCBI, ScholarGoogle, Scopus and Web of Science (WOS). The time cut-off for the published data was until December 2020. The present review did not consider “grey” literature (i.e. technical reports, conference proceedings, congress posters, theses) because they are not subjected to a peer review process.

Manuscript accepted 23.08.2021

DOI: 10.35929/RSZ.0059

Appendices 1 and 2 are available as supplementary material S1

DOI: 10.35929/RSZ.0059.S1

Taxonomy of hosts and parasites

The taxonomic classification for host species was stated as follows: Avibase (<https://avibase.bsc-eoc.org/>) for birds; FishBase (<https://www.fishbase.in/search.php>) for fishes; GBIF (<https://www.gbif.org>) and ITIS (Integrated Taxonomic Information System; <https://www.itis.gov/>), in this order of priority, for freshwater mollusks, reptiles, amphibians and mammals; and WoRMS (World Register of Marine Species; <http://www.marinespecies.org/>) and SeaLifeBase (<https://www.sealifebase.ca/>), in this order of priority, for marine mollusks and crustaceans. We followed Gibson *et al.* (2002), Jones *et al.* (2005), Bray *et al.* (2008), Roberts *et al.* (2013), GBIF and WoRMS for the taxonomy of trematodes.

Hosts

The categorization of hosts as native, endemic, or exotic species followed Ministerio de Medio Ambiente (MMA, 2018a, b) and PNUD (2017). In addition, the diversity of native, endemic, and exotic animal species for every host group present in Chile was stated according to Valdovinos (1999), Martínez & González (2017), PNUD (2017) and Ministerio de Medio Ambiente (2018a, b; <http://especies.mma.gob.cl>).

Locations

The collection sites were attributed to Chilean regions represented by roman numbers and ordered according its geographical distribution from northern Chile to Chilean Patagonia following the Biblioteca del Congreso Nacional de Chile (BCN) (<https://www.bcn.cl/siit/mapoteca/regiones>) and Ministerio de Medio Ambiente (MMA) (2018a): (i) *Northern Chile*: XV= Arica y Parinacota region ($18^{\circ}22'4.8''$ S $70^{\circ}13'1.199''$ W), I= Tarapacá region ($20^{\circ}9'49.202''$ S $69^{\circ}32'46.841''$ W), II= Antofagasta region ($23^{\circ}36'14.873''$ S $69^{\circ}5'3.401''$ W), III= Atacama region ($27^{\circ}33'25.842''$ S $70^{\circ}0'56.477''$ W), IV= Coquimbo region ($30^{\circ}45'16.795''$ S $70^{\circ}54'1.992''$ W); (ii) *Central Chile*: V= Valparaíso region ($32^{\circ}35'51.392''$ S $70^{\circ}51'10.711''$ W), RM= Metropolitan region ($33^{\circ}30'13.172''$ S $70^{\circ}45'44.692$ W), VI= Libertador General Bernardo O'Higgins region ($34^{\circ}32'4.337''$ S $71^{\circ}2'7.735''$ W), VII= Maule region ($35^{\circ}35'50.022''$ S $71^{\circ}29'19.248''$ W); (iii) *Southern Chile*: XVI= Ñuble region ($36^{\circ}37'13.518''$ S $72^{\circ}6'5.983''$ W), VIII= Biobío region ($37^{\circ}20'20.907''$ S $72^{\circ}24'38.456''$ W), IX= Araucanía region ($38^{\circ}40'4.424''$ S $72^{\circ}15'39.623''$ W), XIV= Los Ríos region ($39^{\circ}58'27.448''$ S $72^{\circ}40'2.92''$ W), X= Los Lagos region ($42^{\circ}18'3.038''$ S $73^{\circ}6'19.393''$ W); (iv) *Chilean Patagonia*: XI= Aysén del General Carlos Ibáñez del Campo region ($46^{\circ}28'16.16''$ S

$73^{\circ}31'14.417''$ W), XII= Magallanes y de la Antártica Chilena region ($53^{\circ}21'9.906''$ S $71^{\circ}33'17.201''$ W).

RESULTS

Over the last 120 years, there is a clear tendency for the increase of number of publications reporting trematodes in Chile, with a first peak in the 1980s and a second peak in the 2010s (Fig. 1).

Trematodes

A total of 215 taxa of trematodes belonging to two subclasses, three orders, 47 families and 127 genera have been recorded in Chile. Of these, 119 (55.35 %; 119/215) taxa were identified to species level, with three species belonging to Aspidogastrea subclass and 116 to Digenea subclass (Table 1).

The most common digenetic families mentioned for particular host groups were Fellodistomidae for mollusks, Microphallidae for crustaceans, Opecoelidae for fishes, Gorgoderidae for amphibians, Echinochasmidae and Echinostomatidae for birds, and Fasciolidae for mammals. Only family Opisthognomidae was reported from reptiles.

Complete life cycles are known for only 3 out 119 species of trematodes reported in the country, namely *Fasciola hepatica*, *Proctoeces lintoni* (Fellodistomidae) and *Prosorhynchoides carvajali* (Bucephalidae).

Hosts

Trematodes were reported from 159 out of 3989 potential native animal species (including 14 out of 1081 endemic species) and 12 out of 92 exotic animal species known in Chile (Table 2). So far, there are no records of trematodes from insects or classes of crustaceans other than Malacostraca.

The highest diversity of trematodes was found in representatives of the following host orders: Mytilida (9 taxa) in mollusks, order Decapoda (3 taxa) in crustaceans, order Perciformes (75 taxa) in fishes, order Anseriformes (21 taxa) in birds, and orders Carnivora and Cetacea (7 taxa each) in mammals. One order of reptiles and one order of amphibians were reported as hosts.

Geographical distribution

Taxonomic diversity of trematodes at family level across different regions of Chile is presented in Figs 2-4. A total of 428 records were plotted for all trematode families. These records represent the sum of all taxa reported for every trematode family in every region of the country. Our analysis showed certain patterns in the distribution of several digenetic families. For instance, families Allocreadidae, Brauninidae, Dicrocoeliidae,




Fig. 1. Temporal tendency in the number of publications related to trematodes in Chile.

Table 1. Parasitic taxa isolated from each host group.

	Moll‡	Crus	Fishes*	Am	Rep	Birds	Mam
Species	8	1	74	3	0	27	17
New species	3†	1†	22	3	0	0	3
Incompletely described	21	4	59	0	1	14	5
Genus sp.	7	0	49	0	1	11	4
Family gen. sp.	10	2	9	0	0	3	0
Subfamily gen. sp.	0	2	0	0	0	0	0
<i>incertae sedis</i>	4	0	1	0	0	0	1

Symbols and abbreviations: Moll= Mollusks; Crus= Crustaceans; Am= Amphibians; Rep= Reptiles; Mam= Mammals; ‡Including classes Gastropoda and Bivalvia; *including classes Actinopterygii, Holocephali and Elasmobranchii; †These species were also described from fishes.

Himasthlidae, Multicalycidae, Phaneropsidae and Pleurogenidae have been recorded only in Southern Chile. Meanwhile, the Fasciolidae has been reported throughout the entire country.

The highest diversity of trematodes was recorded on Biobío region (VIII), Southern Chile, with 98 taxa belonging to 36 families, followed by Valparaíso region (V), Central Chile, with 62 taxa belonging to 20 families. On the other hand, the region with the lower diversity of trematodes was Libertador General Bernardo O'Higgins region (VI), Central Chile, with only 1 taxon (see Fig. 3).

Checklist

The checklist of the trematodes and their associated hosts is provided in the Appendix 1 [supplementary material S1]. The stage of development, parasitized organs, localities and references are also stated. An additional table provides a host-parasite list the Appendix 2 [supplementary material S1].

Table 2. Number of host species per host group in Chile (top 3 rows), and number and percentage, between parenthesis, of recorded species hosting trematodes in Chile (bottom 3 rows).

		Moll	Crus	Fishes	Am	Rep	Birds	Mam†	Total
Hosts biodiversity	Native species	1187	606	1401	57	135	443	160	3989
	Endemic species	886	21	35	34	82	10	13	1081
	Exotic species	35	0	26	1	2	9	19	92
Present Review	Native species	24 (2.02)	12 (1.98)	84 (6.00)	3 (5.26)	1 (0.74)	19 (4.29)	16 (10.00)	159 (3.99)
	Endemic species	0	0	9 (25.71)	2 (5.88)	1 (1.22)	0	2 (15.38)	14 (1.30)
	Exotic species	0	0	2 (7.69)	0	0	0	10 (52.63)	12 (13.04)

Abbreviations: Moll= Mollusks; Crus= Crustaceans; Am= Amphibians; Rep= Reptiles; Mam= Mammals; †= excluding human being.




Fig. 2. Number of taxa per trematode family in Northern Chile (NC). Every color in the pie chart represents a particular family. The highest number of taxa for trematode families in every region is given between parentheses. The name of every region represented as a roman number is detailed in Material and Methods.




Fig. 3. Number of taxa per trematode family in Central Chile (CC) and Southern Chile (SC), regions XVI, VIII and IX. Every color in the pie chart represents a particular family. The highest number of taxa for trematode families in every region was stated between parentheses. The name of every region represented as a roman number is detailed in Material and Methods.

DISCUSSION

A significant trematode diversity has been recorded from Chile since the first publication by Gay (1849). However, the identification of a number of reported taxa is questionable, therefore, the collection of new quality specimens as well as the thorough re-examination of specimens deposited in museums are necessary. A number of species reported from Chile do not have descriptions or illustrations. For instance, *Proctoeces lintoni* (Felodistomidae) (originally described from Europe) was reported in native fishes from Chile without adequate descriptions (Valdivia *et al.*, 2010; Oliva *et al.*, 2018). In the absence of DNA sequence data, comparisons with the two native species from Chile, *P. humboldti* and *P. sicyases*, is not possible (George-Nascimento & Quiroga, 1983; Valdivia *et al.*, 2010; Oliva *et al.*, 2018). Taking into account the uncertainty of the identification, the information on life cycle and presence of *P. lintoni*

in Chile should be considered with caution (Valdivia *et al.*, 2010; Oliva *et al.*, 2018). The same is true for *Helicometrina nimia* (Opecoelidae), another European fluke reported in Chile, which needs additional analysis to establish its identity (Oliva *et al.*, 2015).

Although complete life cycles are known for only 3 species of digenetic trematodes reported from Chile, larval stages of some other species have been reported. For instance, metacercariae of *Stephanoprorra uruguayense* (Echinochasmidae), parasitic as an adult in seabirds, have been reported from two native fish species (Torres *et al.*, 1982, 1983, 1991, 1992a, 1993a, 2017; Viozzi *et al.*, 2008). Although in Chile the first intermediate host of this digenetic trematode remains unknown, aquatic snails of family Hydrobiidae have been recorded as its first intermediate host in Argentina (Ostrowski de Núñez, 2007). Members of family Hydrobiidae are distributed in Chile (Valdovinos, 1999), therefore, we anticipate that




Fig. 4. Number of taxa per trematode family in Southern Chile (SC), regions XIV and X, and Chilean Patagonia (CP). Every color in the pie chart represents a particular family. The highest number of taxa for trematode families in every region was stated between parentheses. The name of every region represented as a roman number is detailed in Material and Methods.

future surveys may reveal the local transmission pathway of this trematode.

The insufficient coverage of host species diversity by parasitological studies in Chile (see Garín & González-Acuña, 2008; Muñoz & Olmos, 2008; Oyarzún-Ruiz & González-Acuña, 2021) might underestimate the complete richness of trematode species in the country. This echoes the situation of the coverage of South American fishes by parasitological studies with only 5 % of their total diversity examined for parasites so far (Choudhury *et al.*, 2016).

Among introduced animals, cattle, and lagomorphs have been reported as hosts of *F. hepatica* (e.g. Alcaíno & Gorman, 1999; Courtin *et al.*, 1975, 1979; Oyarzún-Ruiz *et al.*, 2019b), and some salmonid fishes were reported as hosts of digenleans (Arata *et al.*, 1977; Torres, 1995; Torres *et al.*, 1992a). While publications on the helminth fauna of other exotic species such synanthropic rodents (*Rattus rattus*, *Rattus norvegicus*, and *Mus musculus*)

do exist, there are no records of trematodes on these vertebrates (see Rojas *et al.*, 1971; Landaeta-Aqueveque *et al.*, 2014, 2018; Seguel *et al.*, 2017).

Only a small proportion of studies on trematodes in Chile were performed using experimental infections (e.g. Tagle, 1944; Fernández, 1987b; Rebolledo *et al.*, 2014; Muñoz *et al.*, 2015). Thus, authors want to emphasize the need for performing this kind of studies, because these would help to a better understanding of the relationships between parasites and their hosts, mechanisms of transmission, pathological changes, behavioral manipulation, among others (Fried, 2000; Ostrowski de Núñez *et al.*, 2004; Morley & Lewis, 2014; Blasco-Costa & Poulin, 2017; Karvonen *et al.*, 2017).

The present review highlights the clear need for the broader use of molecular tools in the taxonomic studies of trematodes in Chile. Although the first study using DNA sequencing was published in 2010 (Valdivia *et al.*, 2010), the number of works applying molecular techniques is

still limited. The combined use of morphological and molecular approaches (aka integrative taxonomy) is the most promising approach to advance our knowledge of biodiversity, classification, diagnostics, and phylogenetic interrelationships of trematodes and other parasites (Kostadinova & Gibson, 2000; Kostadinova & Pérez-del-Olmo, 2014; Hoberg *et al.*, 2015).

Another interesting and currently discussed topic is the conservation of parasites (Gómez & Nichols, 2013; Carlson *et al.*, 2020b). Some trematodes reported in Chile have been found in endemic species, e.g. *Gorgoderina valdiviensis* (Gorgoderidae) in the helmeted water toad (*Calyptocephalella gayi*) (Puga, 1979, 1982) and *Rudolphitrema chilensis* (Plagiorchiidae) in the rosy ground frog (*Eupsophus roseus*) (Puga, 1986; Puga & Torres, 1999), with no additional reports in other hosts (see Fernandes & Kohn, 2014). In addition, hosts of these particular flukes are considered threatened species with declining populations (IUCN, 2020). Considering their host specificity, these parasites could be classified both as endemic and threatened species, and the concept of co-threatened species should be applied to these parasites (Gómez & Nichols, 2013). Likely, there are additional trematode species in not yet studied endemic hosts that would fall into the same category.

The deposition of parasites into publicly accessible museum collections is critically important. This is especially true for the descriptions of new species (Krell, 2016). As discussed above, an illustration of such need is the situation with *P. lintoni* and *H. nimia*, (Valdivia *et al.*, 2010; Oliva *et al.*, 2015, 2018). Thus, the authors strongly encourage researchers to deposit their specimens, both types and vouchers, in public collections. In Chile, the museums Museo Nacional de Historia Natural and Museo de Zoología at Universidad de Concepción, have helminthological collections receiving and curating specimens.

Although there has been an increase in the number of publications on trematodes in Chile in the past decades, there is a clear need for training parasitologists who would apply modern integrative approaches, and more broadly use experimental models. There is an obvious need for geographic and taxonomic expansion of trematological research in the country in order to investigate not yet studied regions and host species.

ACKNOWLEDGEMENTS

We are thankful of Ms Silvia Oliveros and Angie Figueroa, Drs Fabiana Drago, Melissa Querido, Patricio Torres, Paulina Sepúlveda and Javier Paineán who provided us with publications. Also, to Drs C. Landaeta-Aqueveque, F. Drago and P. Torres for their revision and comments on the draft of this manuscript, to Sebastián Muñoz-Leal who reviewed the English text and to Richard Thomas for his support in the construction of Figs 2-4. Also, to the

reviewers and subject editor for their useful suggestions. This publication is dedicated to professor D. González-Acuña, who passed away recently.

REFERENCES

- Aburto N., Torres P., Lamilla J. 2008. Infection of the Yellownose Skate, *Dipturus chilensis*, by *Otodistomum cestoides* (Trematoda, Digenea) in Three Coastal Localities of Southern Chile. *Comparative Parasitology* 75(1): 150-152.
- Achatz T.J., Pulis E.E., González-Acuña D., Tkach V. V 2020. Phylogenetic Relationships of *Cardiocephalooides* spp. (Digenea, Diplostomoidea) and the Genetic Characterization of *Cardiocephalooides physalis* from Magellanic Penguin, *Spheniscus magellanicus*, in Chile. *Acta Parasitologica* 65(2): 525-534.
- Alcaíno H. 1985. Epizootiología de la distomatosis bovina en Chile. *Parasitología al Día* 9: 22-26.
- Alcaíno H.A., Aguilar A.M. 1985. Actividad antihelmíntica del closantel y de la combinación febantel + triclorfón en caballos fina sangre de carrera. *Archivos de Medicina Veterinaria* 17(2): 103-109.
- Alcaíno H., Apt W. 1989. Algunos antecedentes sobre la fascioliasis animal y humana. *Monografías de Medicina Veterinaria* 11: 14-29.
- Alcaíno H., Gorman T. 1999. Parásitos de los animales domésticos en Chile. *Parasitología al Día* 23(1-2): 33-41.
- Alcaíno H., Mozo P. 1983. Aspectos del parasitismo en bovinos de la provincia de Linares (VII Región de Chile). *Parasitología al Día* 7: 108-112.
- Alcaíno H., Silva J. 1988. Algunos aspectos del parasitismo en bovinos de la provincia de Colchagua (VI Región de Chile). *Parasitología al Día* 12: 62-67.
- Alcaíno H., Gorman T., Phillips J. 1983a. Distomatosis en equinos fina sangre de carrera en haras e hipódromos de las regiones V y Metropolitana de Chile. *Parasitología al Día* 7: 37-40.
- Alcaíno H.A., Gorman T.R., Guevara G., Fernández J.D. 1983b. Distomatosis y parasitosis del intestino grueso de equinos de la zona centro-sur de Chile. *Archivos de Medicina Veterinaria* 15(1): 27-36.
- Alcaíno H., Vega F., Klein P., Gorman T., Apt W. 1988. Fascioliasis en caballos y conejos silvestres (*Oryctolagus cuniculus*) en la Provincia de Curicó, Chile. *Parasitología al Día* 12(3): 136-140.
- Alcaíno H., Gorman T., González V., Vega F., Apt W. 1990. Fascioliasis en caballos, cerdos y conejos silvestres en la provincia de Talca, VII región, Chile. *Parasitología al Día* 14: 9-13.
- Alcaíno H., Apt W., Vega F., Gorman T., Apt P. 1992. Fascioliasis animal en la VII Región de Chile: áreas de distribución e infecciones en caballos y conejos silvestres. *Parasitología al Día* 16(1-2): 11-16.
- Alcaíno H., Vega F., Gorman T. 1993. Epidemiología de la Fascioliasis hepática en la VII Región, Chile. *Parasitología al Día* 17: 99-106.
- Alcaíno H., Parra L., Gorman T.R. 2005. Fasciolosis en equinos fina sangre de carrera de los hipódromos de la zona central de Chile: 2002-2003. *Parasitología Latinoamericana* 60(1-2): 61-64.
- Aldana M., González K., Loot G., Pulgar J., Marquet P.A. 2009.

- First intermediate host of the digenetic trematode *Proctoeces lintoni* (Fellodistomidae) in Chile. *Journal of Parasitology* 95(6): 1408-14.
- Aldana M., Pulgar J.M., Orellana N., Patricio Ojeda F., García-Huidobro M.R. 2014. Increased parasitism of limpets by a trematode metacercaria in fisheries management areas of central Chile: Effects on host growth and reproduction: Management areas and parasitism. *EcoHealth* 11(2): 215-226.
- Aldana M., Pulgar J., Hernández B., George-Nascimento M., Lagos N.A., García-Huidobro M.R. 2020. Context-Dependence in parasite effects on keyhole limpets. *Marine Environmental Research* 157: 104923.
- Álvarez V., Pefaur J. 1970. Parasitismo hepático por tremátodos en "Rorqual de Rudolphi" (Cetacea) capturados en Chile. *Boletín Chileno de Parasitología* 25: 2-5.
- Andres M.J., Curran S.S., Fayton T.J., Pulis E.E., Overstreet R.M. 2015. An additional genus and two additional species of forticulcitinae (Digenea: Haploporidae). *Folia Parasitologica* 62(1): 25.
- Aparicio-Rizzo P., Muñoz G. 2017. Variaciones espacio temporales de la estructura comunitaria de endoparasitos en peces intermareales de Chile central: descriptores parasitológicos, composición de especies y anidamiento. *Latin American Journal of Aquatic Research* 45(4): 737-747.
- Apt W., Margulis P., Andreis M. 1984. Coexistencia de Abceso hepático amebiano y fascioliasis en un adulto con hipogamaglobinemia E. *Revista médica de Chile* 112: 152-156.
- Apt W., Klein P., Vega F., Alcaíno H., Retamal C. 1988. Fascioliasis humana en población rural de la provincia de Curicó (VII Región), Chile. *Parasitología al Día* 12: 155-164.
- Apt W., Aguilera X., Vega F., Alcaíno H., Zulantay I., Apt P., González V., Retamal C., Rodríguez J., Sandoval J. 1993. Prevalencia de fascioliasis en humanos, caballos, cerdos y conejos silvestres, en tres provincias de Chile. *Boletín de la Oficina Sanitaria Panamericana* 115(5): 405-414.
- Apt W., Aguilera X., Vega F., Miranda C., Zulantay I., Pérez C., Gabor M., Apt P. 1995. Treatment of Human Chronic Fascioliasis with Triclabendazole: Drug Efficacy and Serologic Response. *The American Journal of Tropical Medicine and Hygiene* 52(6): 532-535.
- Arata N., Luengo M., Segovia P., Luengo J., Garrido V. 1977. Parasitismo por *Diplostomulum* en truchas (*Salmo gairdneri*). *Boletín Chileno de Parasitología* 32: 89-92.
- Atías A., Fernández C. 1965. Un caso de fascioliasis (distomatosis) hepática con posible localización errática pulmonar. *Boletín Chileno de Parasitología* 20: 47-49.
- Babero B.B., Cabello C., Kinoed J.E. 1979. Helmintoфаuna de Chile. V. parásitos del coipo, *Myocastor coypus* (Molina, 1782). *Boletín Chileno de Parasitología* 34: 26-31.
- Balboa L., George-Nascimento M. 1998. Variaciones ontogenéticas y entre años en las infracomunidades de parásitos metazoos de dos especies de peces marinos de Chile. *Revista Chilena de Historia Natural* 71(1): 27-37.
- Balboa L., George-Nascimento M., Ojeda F.P. 2001. Differential precocious sexual development of *Proctoeces lintoni* (Digenea: Fellodistomidae) in three sympatric species of keyhole limpets *Fissurella* spp. may affect transmission to the final host. *Journal of Parasitology* 87(5): 1164-1167.
- Barbosa F.S., Barbosa I. 1958. *Tropicorbis chilensis* from Santiago, Chile, a potential intermediate host of *Schistosoma mansoni*. *Boletín Chileno de Parasitología* 13(1): 7-9.
- Bay-Schmidt E. 1972. Un nuevo género y especie de Lecithodendriidae (Trematoda) parásito de *Lasiurus borealis bonaerensis* (Lesson y Garnot, 1826) (Mammalia, Chiroptera). *Boletín Chileno de Parasitología* 27(1): 36-39.
- Bell J.A., González-Acuña D., Tkach V. V. 2018. First Record of *Gyrabascus* (Digenea, Pleurogenidae) from *Dromiciops bozinovici* D'Elia et al., 2016 (Marsupialia: Microbiotheriidae) in Chile and its Phylogenetic Relationships. *Comparative Parasitology* 85(1): 58-65.
- Blasco-Costa I., Poulin R. 2017. Parasite life-cycle studies: a plea to resurrect an old parasitological tradition. *Journal of Helminthology* 91(6): 647-656.
- Bray R.A. 2002a. Three species of plagioporine opecoelids (Digenea), including a new genus and two new species, from marine fishes from off the coast of Chile. *Systematic Parasitology* 51(3): 227-236.
- Bray R.A. 2002b. Family Fellodistomidae Nicoll, 1909 [pp. 261-293]. In: Bray R., Gibson D., Jones A. Keys to the Trematoda, Volume 1. *CABI Publishing and The Natural History Museum, London*, 550 pp.
- Bray R.A., Gibson D.I., Jones A. 2008. Keys to the Trematoda, Volume 3. *CABI Publishing and The Natural History Museum, London*, 848 pp.
- Bretos M., Chihuailaf R.H. 1993. Studies on the reproduction and gonadal parasites of *Fissurella pulchra* (Gastropoda: Prosobranchia). *The Veliger* 36(3): 245-251.
- Bretos M., Jirón C. 1980. Trematodes in Chilean Fissurellid Molluscs. *The Veliger* 22(3): 293.
- Bretos M., Tesorieri I., Álvarez L. 1983. The Biology of *Fissurella maxima* Sowerby (Mollusca: Archaeogastropoda) in Northern Chile. 2. Notes on Its Reproduction. *Biological Bulletin* 165(3): 559-568.
- Brieva L.M., Oporto J.A. 1991. Prevalencia e intensidad de la infección por tremátodos del género *Nasitrema* en el delfín chileno *Cephalorhynchus eutropis* (Cetacea: Delphinidae). *Archivos de Medicina Veterinaria* 23(1): 97-100.
- Cañas P., Lozada E. 1987. Nota sobre un trematode digenético endoparásitico del loco *Concholepas concholepas* Bruguière 1789 (Gastropoda: Muricidae). *Parasitología al Día* 11: 173-176.
- Carlson C.J., Dallas T.A., Alexander L.W., Phelan A.L., Phillips A.J. 2020a. What would it take to describe the global diversity of parasites? *Proceedings of the Royal Society B: Biological Sciences* 287(1939): 20201841.
- Carlson C.J., Hopkins S., Bell K.C., Doña J., Godfrey S.S., Kwak M.L., Lafferty K.D., Moir M.L., Speer K.A., Strona G., Torchin M., Wood C.L. 2020b. A global parasite conservation plan. *Biological Conservation* 250: 108596.
- Carvajal J., Durán L.E., George-Nascimento M. 1983. *Ogmogaster heptalineatus* n. sp. (Trematoda: Notocotylidae) from the Chilean sea lion *Otaria flavescens*. *Systematic Parasitology* 5(3): 169-173.
- Castro-Rojas M., Cid-Moya K., George-Nascimento M. 2015. Consumo de oxígeno en *Perumytilus purpuratus* según el parasitismo por Digenea, y efectos de la temperatura y del ciclo de emersión-inmersión sobre la liberación de cercarias. *Revista de Biología Marina y Oceanografía* 50(2): 315-321.
- Cerdeira C., Veloso-Frías J., Lobos-Chávez F., Oyarzún-Ruiz P., Henríquez A., Loyola M., Silva-de la Fuente M.C., Ortega R., Letelier R., Landaeta-Aqueveque C. 2019. Morphological and molecular identification with frequency analysis of *Calicophoron microbothrioides* in central Chile.

- Brazilian Journal of Veterinary Parasitology* 28(4): 582-591.
- Chávez R.A., González M.T., Oliva M.E., Valdivia I.M. 2012. Endoparasite fauna of five Gadiformes fish species from the coast of Chile: Host ecology versus phylogeny. *Journal of Helminthology* 86(1): 10-15.
- Choudhury A., Aguirre-Macedo M.L., Curran S.S., Ostrowski-De Núñez M., Overstreet R.M., Pérez-Ponce de León G., Santos C.P. 2016. Trematode diversity in freshwater fishes of the Globe II: "New World." *Systematic Parasitology* 93(3): 257-269.
- Collins J.J. 2017. Platyhelminthes. *Current Biology* 27(7): R252-R256.
- Corrêa F., Hidalgo C., Stoore C., Jiménez M., Hernández M., Paredes R. 2020. Cattle co-infection of *Echinococcus granulosus* and *Fasciola hepatica* results in a different systemic cytokine profile than single parasite infection. *Plos One* 15(9): e0238909.
- Cortés Y., Muñoz G. 2008. Infracomunidades de parásitos eumetazoos del bagre de mar *Aphos porosus* (Valenciennes, 1837) (Actinopterygii: Batrachoidiformes) en Chile central. *Revista de Biología Marina y Oceanografía* 43(2): 255-263.
- Cortés Y., Muñoz G. 2009. Metazoan Parasite Infracommunities of the Toadfish *Aphos porosus* (Pisces: Batrachoidiformes) in Central Chile: How Variable Are They over Time? *Journal of Parasitology* 95(3): 753-756.
- Courtin S., Ferriere G., Cerda J. 1975. Primer estudio de *Fasciola hepatica* en el conejo silvestre (*Oryctolagus cuniculus*) de la precordillera de Nahuelbuta. *Boletín Chileno de Parasitología* 30(3-4): 65-67.
- Courtin S., Alcaíno H., Plaza J., Ferriere G. 1979. Platelmintos del conejo silvestre (*Oryctolagus cuniculus*) en la cordillera de Nahuelbuta, Chile. *Archivos de Medicina Veterinaria* 11(1): 23-26.
- Curran S.S., Pulis E.E., Andres M.J., Overstreet R.M. 2018. Two New Species of *Saccocoelioidea* (Digenea: Haploporidae) with Phylogenetic Analysis of the Family, Including Species of *Saccocoelioidea* from North, Middle, and South America. *Journal of Parasitology* 104(3): 221-239.
- Díaz F., George-Nascimento M. 2002. Estabilidad temporal de las infracomunidades de parásitos en la borrachilla *Scartichthys viridis* (Valenciennes, 1836) (Pisces: Blenniidae) en la costa central de Chile. *Revista Chilena de Historia Natural* 75(4): 641-649.
- Díaz P.E., Muñoz G., George-Nascimento M. 2016. A new species of *Hemipera* Nicoll, 1913 (Digenea: Derogenidae) from fishes of the intertidal rocky zone of Chile. *Acta Parasitologica* 61(3): 516-522.
- Dioni W. 1947. *Gorgoderina chilensis* n. sp., Tremátodo de la vejiga urinaria del sapito vaquero, *Rhinoderma darwini* D. B. *Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo* 2: 1-11.
- Drago F.B., Lunaschi L.I., Hinojosa-Saez A.C., González-Acuña D. 2007. First record of *Australapatemon burti* and *Paramonostomum pseudalveatum* (Digenea) from *Anas georgica* (Aves, Anseriformes) in Chile. *Acta Parasitologica* 52(3): 201-205.
- Dubois G. 1981. Notes Helminthologiques V: Strigeidae Railliet, Diplostomidae Poirier et Proterodiplostomidae Dubois. *Revue suisse de Zoologie* 88(1): 227-232.
- Dubois G., Rausch R. 1960. Quatrième contribution à l'étude des Strigeides (Trematoda) nord-américains. *Bulletin de la Société Neuchâteloise des Sciences Naturelles* 83: 79-92.
- Ebmer D., Navarrete M.J., Muñoz P., Flores L.M., Gärtner U., Brabec J., Poppert S., Taubert A., Hermosilla C. 2020. Anthropozoonotic parasites circulating in synanthropic and pacific colonies of south american sea lions (*Otaria flavescens*): non-invasive techniques data and a review of the literature. *Frontiers in Marine Science* 7: 847.
- Espínola-Novelo J.F., Escribano R., Oliva M.E. 2018. Metazoan parasite communities of two deep-sea elasmobranchs: the southern lanternshark, *Etmopterus granulosus*, and the largenose catshark, *Apristurus nasutus*, in the Southeastern Pacific Ocean. *Parasite* 25: 53.
- Faigenbaum J., Agosin M., Tamargo A. 1950. Distomatosis humana. *Revista Médica de Chile* 78(6): 383-387.
- Faigenbaum J., Feres A., Donckaster R., Atías A., Jarpa A., Niedmann G., Donoso F., Rubio M., Meruane J. 1962. Fascioliasis hepática humana. *Boletín Chileno de Parasitología* 17: 7-12.
- Fernandes B.M.M., Kohn A. 2014. South American trematodes parasites of amphibians and reptiles. *Oficina de Livros, Rio de Janeiro*, 225 pp.
- Fernández I., Oyarzún C., Valenzuela A., Burgos C., Guaquín V., Campos V. 2016. Parásitos del pez luna *Mola mola* (Pisces: Molidae). Primer registro en aguas de la costa centro sur de Chile. *Gayana* 80(2): 192-197.
- Fernández I., Moraga R., Yáñez F., Mansilla M., Smith C., Campos V. 2019. Gastrointestinal helminths of Humboldt penguin (*Spheniscus humboldti* Meyen, 1834) from south central Chile. *Latin American Journal of Aquatic Research* 47(1): 206-211.
- Fernández J. 1985. Estudio parasitológico de *Merluccius australis* (Hutton, 1872) (Pisces: Merlucciidae). Aspectos sistemáticos, estadísticos y zoogeográficos. *Boletín de la Sociedad de Biología de Concepción* 56: 31-41.
- Fernández J. 1987a. Nuevos registros de parásitos en mamíferos marinos chilenos. *Parasitología al Día* 11: 120-126.
- Fernández J. 1987b. Los parásitos de la lisa *Mugil cephalus* L. en Chile: sistemática y aspectos poblacionales (Perciformes: Mugilidae). *Gayana Zoología* 51: 3-58.
- Fernández J., Durán L. 1985. *Aporocotyle australis* n. sp. (Digenea: Sanguinicolidae), parásito de *Merluccius australis* en Chile y su relación con la filogenia de *Aporocotyle* Odhner, 1900 en *Merluccius* spp. *Revista Chilena de Historia Natural* 58(2): 121-126.
- Fernández J.C., Villalba C., Alviña A. 1986. Parásitos del pejegallo, *Callorhynchus callorhynchus* (L.), en Chile: aspectos biológicos y sistemáticos. *Biología Pesquera* 15: 63-73.
- Fernández V., Semenov L., Viozzi G.P. 2012. Parasites of the "Peladilla," *Aplochiton zebra* (Osmeriformes: Galaxiidae), from Patagonia (Argentina and Chile). *Comparative Parasitology* 79(2): 231-237.
- Fernández-Cisternas I., George-Nascimento M., Ojeda F.P. 2017. Comparison of parasite diversity of intertidal fish assemblages from central California and central Chile. *Revista de Biología Marina y Oceanografía* 52(3): 505-521.
- Figueroa L. 1988. *Synthesium tursionis* (Marchi, 1873) Price, 1932 (Digenea: Compulidae) en *Cephalorhynchus eutropis* Gray, 1846 (Cetacea: Odontoceti) en la costa chilena. *Parasitología al Día* 12: 185-186.
- Figueroa L., Franjola R. 1988. *Braunina cordiformis* Wolf,

- 1903 (Digenea: Brauninidae) en delfín chileno *Cephalorhynchus eutropis* Gray, 1846 (Cetacea: Odontoceti). *Boletín Chileno de Parasitología* 43(3-4): 71-72.
- Figueroa L., Puga S. 1991. *Lecithaster pacificum* n. sp. (Digenea: Hemiriorchidae), parásito del “lenguado de ojos chicos” *Paralichthys microps* (Pleuronectiformes: Bothidae) en la costa de Valdivia, Chile. *Archivos de Medicina Veterinaria* 23(1): 101-104.
- Flores K., George-Nascimento M. 2009. Las infracomunidades de parásitos de dos especies de *Scartichthys* (Pisces: Blenniidae) en localidades cercanas del norte de Chile. *Revista Chilena de Historia Natural* 82(1): 63-71.
- Flores K., López Z., Levicoy D., Muñoz-Ramírez C.P., González-Wevar C., Oliva M.E., Cárdenas L. 2019. Identification assisted by molecular markers of larval parasites in two limpet species (Patellogastropoda: *Nacella*) inhabiting Antarctic and Magellan coastal systems. *Polar Biology* 4: 1175-1182.
- Franjola R., Gallardo C. 1991. Estados larvales de Digenea en *Kingiella chilensis* (Mollusca, Bivalvia, Cyamidae). *Boletín Chileno de Parasitología* 46(1-2): 34-35.
- Fried B. 2000. Maintenance, Cultivation, and Excystation of Echinostomes [pp. 100-118]. In: Fried B., Graczyk T. K. Echinostomes as Experimental Models for Biological Research. Springer-Science+Business Media, B. V., USA, 273 pp.
- García-Huidobro M.R., Varas O., George-Nascimento M., Pulgar J., Aldana M., Lardies M.A., Lagos N.A. 2019. Role of temperature and carbonate system variability on a host-parasite system: Implications for the gigantism hypothesis. *International Journal for Parasitology: Parasites and Wildlife* 9: 7-15.
- García-Huidobro M.R., Aldana M., Varas O., Pulgar J., García-Herrera C., Abarca-Ortega A., Grenier C., Rodríguez-Navarro A.B., Lagos N.A. 2020. Geographical variability and parasitism on body size, reproduction and shell characteristics of the keyhole limpet *Fissurella crassa* (Mollusca: Vetigastropoda). *Marine Environmental Research* 161: 105060.
- Garcías F., Mendoza R., George-Nascimento M. 2001. Variación entre años de las infracomunidades de parásitos metazoos de la corvina *Cilus gibberita* (Pisces: Sciaenidae) en Chile. *Revista Chilena de Historia Natural* 74(4): 833-840.
- Gardner A.L., Handley Jr. C.O. 2007. Genus *Lasiurus* [pp. 457-467]. In: Gardner A.L. Mammals of South America, Volume 1: Marsupials, Xenarthrans, Shrews, and Bats. The University of Chicago Press, London, 669 pp.
- Garin C.F., González-Acuña D. 2008. Parásitos de anfibios y reptiles [pp. 303-332]. In: Vidal M.A., Labra A. Herpetología de Chile. Gráficas Andes, Santiago de Chile, 593 pp.
- Gay C. 1849. Historia física y política de Chile - Zoología, Tomo Tercero. Imprenta de Maulde y Benou, Paris, 547 pp.
- Gehrung E., Veit O., Velasco M., Faiguenbaum J., Apablaza A. 1967. Un caso de distomatosis (fascioliasis) hepática sin eosinofilia sanguínea. *Boletín Chileno de Parasitología* 23: 37-39.
- George-Nascimento M. 1996. Populations and assemblages of parasites in hake, *Merluccius gayi*, from the southeastern Pacific Ocean: stock implications. *Journal of Fish Biology* 48(4): 557-568.
- George-Nascimento M. 2000. Geographical variations in the jack mackerel *Trachurus symmetricus murphyi* populations in the southeastern Pacific Ocean as evidenced from the associated parasite communities. *The Journal of Parasitology* 86(5): 929-932.
- George-Nascimento M., Arancibia H. 1992. Stocks ecológicos del jurel (*Trachurus symmetricus murphyi* Nichols) en tres zonas de pesca frente a Chile, detectados mediante comparación de su fauna parasitaria y morfometría. *Revista Chilena de Historia Natural* 65(4): 453-470.
- George-Nascimento M., Arancibia H. 1994. La fauna parasitaria y la morfometría de la merluza austral *Merluccius australis* (Hutton) como indicadoras de unidades de stock. *Biología Pesquera* 23: 31-47.
- George-Nascimento M., Carvajal J. 1981. Helmintos parásitos del lobo marino común *Otaria flavescens* en el Golfo de Arauco, Chile. *Boletín Chileno de Parasitología* 36: 72-73.
- George-Nascimento M., Huet B. 1984. Una aproximación ecológica al estudio del parasitismo en el “congrio negro” *Genypterus maculatus* (Tschudi) (Pisces: Ophidiidae). *Biología Pesquera* 13: 23-30.
- George-Nascimento M., Iriarte J.L. 1989. Las infracomunidades de parásitos metazoos del chancharro *Helicolenus lengerichi* Norman, 1937 (Pisces, Scorpaenidae): un ensamble no interactivo de especies. *Revista Chilena de Historia Natural* 62(2): 217-227.
- George-Nascimento M., Moscoso D. 2013. Variación local y geográfica de las infracomunidades de parásitos de la anchoveta *Engraulis ringens* en Chile. *Revista de Biología Marina y Oceanografía* 48(1): 207-212.
- George-Nascimento M., Quiroga G. 1983. Descripción de una nueva especie de trematodo *Proctoeces humboldti* n. sp. (Fellodistomidae) parásito de las lapas *Fissurella* spp. Brugière, 1789 (Mollusca: Archaeogastropoda). *Parasitología al Día* 7: 100-103.
- George-Nascimento M., Balboa L., Aldana M., Olmos V. 1998. Las lapas *Fissurella* spp. (Mollusca: Archaeogastropoda) y el pejesapo *Sicyases sanguineus* (Pisces: Gobiesocidae) son hospedadores secundarios de *Proctoeces lintoni* (Digenea: Fellodistomidae) en Chile. *Revista Chilena de Historia Natural* 71(2): 169-176.
- George-Nascimento M., Garcías F., Muñoz G. 2002. Parasite body volume and infracommunity patterns in the southern pomfret *Brama australis* (Pisces: Bramidae). *Revista Chilena de Historia Natural* 75(4): 835-839.
- George-Nascimento M., Mellado A., Saavedra S., Carvajal J. 2009. Variabilidad de las comunidades de parásitos metazoos del róbalo *Eleginops maclovinus* (Cuvier & Valenciennes, 1830) (Pisces: Eleginopidae) en Chile. *Revista Chilena de Historia Natural* 82(2): 199-207.
- George-Nascimento M., Moscoso D., Niklitschek E., González K. 2011. Variación geográfica de las comunidades de parásitos de la merluza de tres aletas *Micromesistius australis* al sur de Sudamérica. *Revista de Biología Marina y Oceanografía* 46(1): 53-58.
- George-Nascimento M., López-Rodríguez R., Górska K. 2020. Geographic variation in composition of metazoan parasite infracommunities in *Galaxias maculatus* Jenyns 1842 (Osmeriformes: Galaxiidae) in southern Chile (38-47° S). *Revista Chilena de Historia Natural* 93(1): 2.
- Gibson D.I., Jones A., Bray R.A. 2002. Keys to the Trematoda, Volume 1. CABI Publishing and The Natural History Museum, London, 550 pp.
- Gil L.C., Díaz A., Rueda C., Martínez C., Castillo D., Apt W. 2014. Fascioliasis hepática humana: Resistencia al tratamiento con triclabendazol. *Revista Médica de Chile* 142(10): 1330-1333.

- Gómez A., Nichols E. 2013. Neglected wild life: Parasitic biodiversity as a conservation target. *International Journal for Parasitology: Parasites and Wildlife* 2: 222-227.
- González D., Daugschies A., Rubilar L., Pohlmeyer K., Skewes O., Mey E. 2004. Fauna parasitaria de la tórtola común (*Zenaida auriculata*, de Murs 1847) (Columbiformes: Columbidae) en Ñuble, Chile. *Parasitología Latinoamericana* 59(1-2): 37-41.
- González H. 1982. Distomatosis hepática. *Parasitología al Día* 6: 47-48.
- González H., Plaza J. 1966. Hallazgo de trematodes de la familia Paramphistomidae en bovinos importados desde Australia. *Boletín Chileno de Parasitología* 21: 19-21.
- González L., Carvajal J. 1994. Estudio parasitológico de *Merluccius australis* (Hutton, 1972) del mar interior de Aysén. *Investigación Pesquera* 38: 75-85.
- González M.T., Acuña E. 1998. Metazoan Parasites of the Red Rockfish *Sebastodes capensis* off Northern Chile. *Journal of Parasitology* 84(4): 783-788.
- González M.T., Oliva M.E. 2009. Is the nestedness of metazoan parasite assemblages of marine fishes from the southeastern Pacific coast a pattern associated with the geographical distributional range of the host? *Parasitology* 136(4): 401-409.
- González M.T., Poulin R. 2005a. Spatial and temporal predictability of the parasite community structure of a benthic marine fish along its distributional range. *International Journal for Parasitology* 35(13): 1369-1377.
- González M.T., Poulin R. 2005b. Nested patterns in parasite component communities of a marine fish along its latitudinal range on the Pacific coast of South America. *Parasitology* 131(4): 569-577.
- González M.T., Acuña E., Oliva M.E. 2001. Metazoan Parasite Fauna of the Bigeye Flounder, *Hippoglossina macrops*, from Northern Chile. Influence of Host Age and Sex. *Memórias do Instituto Oswaldo Cruz* 96(8): 1049-1054.
- González M.T., Barrientos C., Moreno C.A. 2006. Biogeographical patterns in endoparasite communities of a marine fish (*Sebastodes capensis* Gmelin) with extended range in the Southern Hemisphere. *Journal of Biogeography* 33(6): 1086-1095.
- González M.T., Vásquez R., Acuña E. 2008. Biogeographic Patterns of Metazoan Parasites of the Bigeye Flounder, *Hippoglossina macrops*, in the Southeastern Pacific Coast. *Journal of Parasitology* 94(2): 429-435.
- González M.T., Henríquez V., López Z. 2013. Variations in the fecundity and body size of digenean (Opecoelidae) species parasitizing fishes from Northern Chile. *Revista de Biología Marina y Oceanografía* 48(3): 421-429.
- González T., Acuña E. 2000. Influence of Host Size and Sex on the Endohelminth Infracommunities of the Red Rockfish *Sebastodes capensis* off Northern Chile. *Journal of Parasitology* 86(4): 854-857.
- González-Acuña D., Kinsella J.M., Lara J., Valenzuela-Dellarossa G. 2008. Parásitos gastrointestinales en pingüino de Humboldt (*Spheniscus humboldti*) y pingüino de Magallanes (*Spheniscus magellanicus*) en las costas del centro y centro sur de Chile. *Parasitología Latinoamericana* 63(1-4): 58-63.
- González-Acuña D., Cerdá F., López J., Ortega R., Mathieu C., Kinsella M. 2009. Checklist of the helminths of the kelp gull, *Larus dominicanus* (Aves: Laridae), with new records from Chile. *Zootaxa* 2297(1): 27-43.
- González-Acuña D., Moreno L., Cicchino A., Mironov S., Kinsella M. 2010. Checklist of the parasites of the black-necked swan, *Cygnus melanocoryphus* (Aves: Anatidae), with new records from Chile. *Zootaxa* 2637(1): 55-68.
- González-Acuña D., Lohse E., Cicchino A., Mironov S., Figueroa R.A., Ardiles K., Kinsella M. 2011a. Parasites of the American Kestrel (*Falco sparverius*) in South-Central Chile. *The Journal of Raptor Research* 45(2): 188-193.
- González-Acuña D., Silva C., Soto M., Mironov S., Moreno L., González-Gómez P.L., Badrul H., Kinsella M. 2011b. Parasites of the Green-backed Firecrown (*Sephanoides sephanioides*) in Chile. *Revista Mexicana de Biodiversidad* 82(4): 1333-1336.
- González-Acuña D., Llanos-Soto S., Oyarzún-Ruiz P., Kinsella J.M., Barrientos C., Thomas R., Cicchino A., Moreno L. 2020a. Parasites of the Neotropic cormorant *Nannopterum (Phalacrocorax) brasiliensis* (Aves, Phalacrocoracidae) in Chile. *Brazilian Journal of Veterinary Parasitology* 29(3): e003920.
- González-Acuña D., Veloso-Frías J., Missene C., Oyarzún-Ruiz P., Cicchino A., Fuentes-Castillo D., Kinsella J.M., Mironov S., Barrientos C., Moreno L. 2020b. External and gastrointestinal parasites of the Franklin's Gull, *Leucophaeus pipixcan* (Charadriiformes: Laridae), in Talcahuano, central Chile. *Brazilian Journal of Veterinary Parasitology* 29(4): e016420.
- Gorman T.R., Alcaíno H.A., Vidal E.H., Berardi L. 1979. Prevalencias y tendencias de la distomatosis, cisticercosis y triquinosis en la VII Región. *Archivos de Medicina Veterinaria Suppl.* 1: 123-128.
- Gorman T.R., Plaza J., Cortés J. 1980. Tendencia de algunas zoonosis parasitarias en porcinos y bovinos de la Zona Centro Sur de Chile. *Archivos de Medicina Veterinaria* 12(1): 30-43.
- Gorman T., Bravo J., Lorca M., Alcaíno H., Ibarra L. 1991. Diagnóstico de la Fasciolasis de equinos y porcinos mediante las pruebas de doble difusión, contrainmunoelectroforesis y hemoaglutinación indirecta. *Archivos de Medicina Veterinaria* 23(2): 123-130.
- Gorman T., Valdés A., Fredes F., Ferreira A., Alcaíno H. 1995. Reconocimiento antigenético de ovinos naturalmente infectados con *Fasciola hepatica*, monitoreado a través de inmunoelectrotransferencia enzimática (Western Blotting). *Archivos de Medicina Veterinaria* 27(2): 33-40.
- Grandón-Ojeda A., Valdebenito J.O., Moreno L., Kinsella J.M., Mironov S., Cicchino A., Barrientos C., González-Acuña D. 2018. Gastrointestinal and external parasitism in the Magellanic Horned Owl *Bubo magellanicus* (Strigiformes: Strigidae) in Chile. *Brazilian Journal of Veterinary Parasitology* 27(2): 161-168.
- Henríquez V., González M.T. 2014. Patterns of variation in parasitic component communities and infracommunities of a littoral fish species from the northern coast of Chile. *Journal of Helminthology* 88(1): 89-96.
- Henriquez V.P., Gonzalez M.T., Licandeo R., Carvajal J. 2011. Metazoan parasite communities of rock cod *Eleginops maclovinus* along southern Chilean coast and their use as biological tags at a local spatial scale. *Journal of Fish Biology* 79(7): 1851-1865.
- Hermosilla C., Silva L.M.R., Navarro M., Taubert A. 2016. Anthropozoonotic endoparasites in free-ranging “urban” South American sea lions (*Otaria flavescens*). *Journal of Veterinary Medicine* 2016: 7507145.

- Hidalgo C., Stoore C., Hernández M., Paredes R. 2020. *Fasciola hepatica* coinfection modifies the morphological and immunological features of *Echinococcus granulosus* cysts in cattle. *Veterinary Research* 51(1): 76.
- Hinojosa-Sáez A., González-Acuña D., George-Nascimento M. 2009. Parásitos metazoos de *Anas georgica* Gmelin, 1789 (Aves: Anseriformes) en Chile central: especificidad, prevalencia y variaciones entre localidades. *Revista Chilena de Historia Natural* 82(3): 337-345.
- Hoberg E.P., Agosta S.J., Boeger W.A., Brooks D.R. 2015. An integrated parasitology: Revealing the elephant through tradition and invention. *Trends in Parasitology* 31(4): 128-133.
- Hurtado-Ormeño C., George-Nascimento M. 2016. Factors accounting for variations in the parasitism by metacercariae (Digenea, Microphallidae) in the amphipod *Apohyale hirtipalma* (Gammaridea) in the Southeastern Pacific coast. *Revista de Biología Marina y Oceanografía* 51(1): 81-88.
- Inzunza Á., Torres P., Franjola R. 1989. Nuevos hospedadores para *Helicometrina nimia* Linton, 1910 (Digenea: Opcoelidae) en la costa centro sur de Chile. *Boletín Chileno de Parasitología* 44: 41-43.
- IUCN. 2020. The IUCN Red List of Threatened Species. Version 2020-3. <https://www.iucnredlist.org>. Downloaded on 20 February 2020.
- Jiménez M., Stoore C., Hidalgo C., Corrêa F., Hernández M., Benavides J., Ferreras M.C., Sáenz L., Paredes R. 2020. Lymphocyte populations in the adventitial layer of hydatid cysts in cattle: relationship with cyst fertility status and *Fasciola hepatica* co-infection. *Veterinary Pathology* 57(1): 108-114.
- Jones A., Bray R.A., Gibson D.I. 2005. Keys to the Trematoda, Volume 2. CABI Publishing and The Natural History Museum, London, 768 pp.
- Kanev I., Vassilev I., Dimitrov V., Radev V. 1994. Life-cycle, delimitation and redescription of *Catatropis verrucosa* (Frölich, 1789) Odhner, 1905 (Trematoda: Notocotylidae). *Systematic Parasitology* 29(2): 133-148.
- Karvonen A., Faltýnková A., Choo J.M., Valtonen E.T. 2017. Infection, specificity and host manipulation of *Australapatemon* sp. (Trematoda, Strigeidae) in two sympatric species of leeches (Hirudinea). *Parasitology* 144(10): 1346-1355.
- Kostadinova A., Gibson D. 2000. The systematics of the echinostomes [pp. 31-58]. In: Fried B., Gracyk T.K. Echinostomes as experimental models for biological research. Springer Science+Business Media, B. V., USA, 273 pp.
- Kostadinova A., Pérez-del-Olmo A. 2014. The systematics of the Trematoda [pp. 21-44]. In: Toledo R., Fried B. Digenetic Trematodes. Springer Science+Business Media, New York, 474 pp.
- Krell F.T. 2016. Taxonomy: Preserve specimens for reproducibility. *Nature* 539(7628): 168-168.
- Lam S. 1982. Distomatosis hepática masiva. *Revista Chilena de Pediatría* 53: 39-41.
- Landaeta-Aqueveque C., Notarnicola J., Correa J.P., Yáñez-Meza A., Henríquez A., Cattan P.E., Botto-Mahan C., Torres-Pérez F. 2014. First record of *Litomosoides pardinasi* (Nematoda: Onchocercidae) in native and exotic rodents from Chile. *Revista Mexicana de Biodiversidad* 85(4): 1032-1037.
- Landaeta-Aqueveque C., Robles M.D.R., Henríquez A., Yáñez-Meza A., Correa J.P., González-Acuña D., Cattan P.E. 2018. Phylogenetic and ecological factors affecting the sharing of helminths between native and introduced rodents in Central Chile. *Parasitology* 145(12): 1570-1576.
- Lasiak T. 1991. Bucephalid trematode infections in mytilid bivalves from the rocky intertidal of Southern Chile. *Journal of Molluscan Studies* 58(1): 29-36.
- Leiva N., George-Nascimento M., Muñoz G. 2015. Carga parasitaria en crustáceos decápodos de la costa central de Chile: ¿existe alguna asociación con la abundancia de los hospedadores definitivos? *Latin American Journal of Aquatic Research* 43(4): 726-738.
- Leiva N.V., López Z., González M.T., Muñoz G. 2017. Determining Intermediate Hosts for Opcoelidae and Microphallidae Species (Platyhelminthes: Trematoda) in the Southeastern Pacific Coast, Using Molecular Markers. *Journal of Parasitology* 103(1): 132-137.
- Leiva N. V., Muñoz G., González M.T. 2020. Geographic and ontogenetic variations in parasite communities of intertidal fish species from the south-eastern Pacific coast. *Journal of Helminthology* 94: e124.
- Leyan V., Franjola R., Oberg C. 1979. Hallazgo de trematodos del género *Echinochasmus* en un perro doméstico (*Canis familiaris*) en Valdivia, Chile. *Boletín Chileno de Parasitología* 34: 44-46.
- Llanos-Soto S., Córdoba M., Moreno L., Kinsella J.M., Mironov S., Cicchino A., Barrientos C., Martín-Ordenes J.S., González-Acuña D. 2019. External and intestinal parasites of the Austral thrush *Turdus falcklandii* (Aves, Turdidae) in central Chile. *Brazilian Journal of Veterinary Parasitology* 28(3): 432-442.
- Loot G., Aldana M., Navarrete S.A. 2005. Effects of human exclusion on parasitism in intertidal food webs of central Chile. *Conservation Biology* 19(1): 203-212.
- Loot G., Blanchet S., Aldana M., Navarrete S.A. 2008. Evidence of Plasticity in the Reproduction of a Trematode Parasite: The Effect of Host Removal. *Journal of Parasitology* 94(1): 23-27.
- López Z., Cárdenas L., González M.T. 2014. Metazoan Symbionts of the Yellow Clam, *Mesodesma donacium* (Bivalvia), in Southern Chile: Geographical Variations. *Journal of Parasitology* 100(6): 797-804.
- Lotz J.M., Font W.F. 2008. Family Phaneropsidae Mehra, 1935 [pp. 545-562]. In: Bray R., Gibson D., Jones A. Keys to the Trematoda, Volume 3. CABI Publishing and The Natural History Museum, London, 848 pp.
- Luengo J., Arata N., Luengo M., Araya N. 1984. Estudio anatomopatológico de hígado de equinos con fascioliasis. *Boletín Chileno de Parasitología* 39: 43-46.
- Luque J., Oliva M. 1993. Trematodes of marine fishes from the Peruvian Faunistic Province (Peru and Chile), with description of *Lecithochirium callaoensis* n. sp. and new records. *Revista de Biología Marina* 28(2): 271-286.
- Mackenzie K., Longshaw M. 1995. Parasites of the hakes *Merluccius australis* and *M. hubbsi* in the waters around the Falkland Islands, southern Chile, and Argentina, with an assessment of their potential value as biological tags. *Canadian Journal of Fisheries and Aquatic Sciences* 52(Suppl. 1): 213-224.
- MacKenzie K., Brickle P., Hemmingsen W., George-Nascimento M. 2013. Parasites of hoki, *Macruronus magellanicus*, in the Southwest Atlantic and Southeast Pacific Oceans, with an assessment of their potential value as biological tags. *Fisheries Research* 145: 1-5.

- Manfredi M.T., Oneto M. 1997. *Phagicola longa* (Heterophyidae) in dogs from Chile: morphological findings and taxonomical problems. *Parassitologia* 39(1): 9-11.
- Martínez D., González G. 2017. Aves De Chile: Guía de Campo y breve Historia Natural. *Ediciones del Naturalista, Santiago*, 540 pp.
- Millán M., Wagenknecht R., Cárdenas A., Carrasco C. 2008. Parásitos de *Fasciola hepatica* intracoledociano. *Revista Chilena de Cirugía* 60(4): 332-335.
- MMA 2018a. Biodiversidad de Chile: Patrimonio y desafíos, Tomo I. Tercera edición. In: Figueroa A., Rovira J., Flores S., Tala C., Avilés R., Orellana J.C., Ferreyra J., Díaz P., Cohen R. *Ministerio de Medio Ambiente, Santiago de Chile*, 393 pp.
- MMA 2018b. Biodiversidad de Chile: Patrimonio y desafíos, Tomo II. Tercera edición. In: Figueroa A., Rovira J., Flores S., Tala C., Avilés R., Orellana J.C., Ferreyra J., Díaz P., Cohen R. *Ministerio de Medio Ambiente, Santiago de Chile*, 264 pp.
- Montenegro D., Jones B., González M.T. 2012. Report of pathogens and parasites in *Perumytilus purpuratus* from San Jorge Bay, Antofagasta, Chile. *Revista de Biología Marina Y Oceanografía* 47(2): 345-350.
- Montenegro D., Romero M.S., González M.T. 2020. Morphological and molecular characterization of larval digenean trematodes (*Parvatrema*: Gymnophallidae) and their pathological effects on the clam *Leukoma thaca* (= *Protothaca thaca*) (Bivalvia: Veneridae) (Molina, 1782) from northern Chile. *Parasitology International* 80: 102238.
- Morales M.A., Luengo J., Vasquez J. 2000. Distribución y tendencia de la fasciolosis en ganado de abasto en Chile, 1989-1995. *Parasitología al Día* 24(3-4): 115-118.
- Morley N.J., Lewis J.W. 2014. Temperature stress and parasitism of endothermic hosts under climate change. *Trends in Parasitology* 30: 221-227.
- Muñoz G. 2005. Metacercarias de la familia Microphallidae (Trematoda: Digenea) en el anfípodo *Hyale grandicornis* en la costa de Maule, Chile Central. *Parasitología Latinoamericana* 60(3-4): 165-169.
- Muñoz G. 2014. Parasites communities in the clingfish *Gobiesox marmoratus* from central Chile. *Acta Parasitologica* 59(1): 108-114.
- Muñoz G. 2019. Parasite community of the tadpole codling *Salilota australis* (Günther, 1878) (Gadiformes: Moridae) from the southern Chile and its comparison with its closest relatives. *Neotropical Helminthology* 13(2): 181-191.
- Muñoz G. 2020. Comunidades de parásitos de *Patagonotothen cornucola* (Pisces: Nototheniidae) de pozas rocosas intermareales del estrecho de Magallanes. *Anales del Instituto de la Patagonia* 48(1): 37-45.
- Muñoz G., Bott N.J. 2011. A new species of *Prosorhynchoides* (Trematoda, Bucephalidae) from the intertidal rocky zone of central Chile. *Acta Parasitologica* 56(2): 140-146.
- Muñoz G., Castro R. 2012. Comunidades de parásitos eumetazoos de peces labrisómidos de Chile central. *Revista de Biología Marina y Oceanografía* 47(3): 565-571.
- Muñoz G., Delorme N. 2011. Variaciones temporales de las comunidades de parásitos en peces intermareales de Chile central: hospedadores residentes vs temporales. *Revista de Biología Marina y Oceanografía* 46(3): 313-327.
- Muñoz G., Olmos V. 2008. Revisión bibliográfica de especies endoparásitas y hospedadoras de sistemas acuáticos de Chile. *Revista de Biología Marina y Oceanografía* 43(2): 173-245.
- Muñoz G., Randhawa H.S. 2011. Monthly variation in the parasite communities of the intertidal fish *Scartichthys viridis* (Blenniidae) from central Chile: Are there seasonal patterns? *Parasitology Research* 109(1): 53-62.
- Muñoz G., Zamora L. 2011. Ontogenetic Variation in Parasite infracommunities of the Clingfish *Sicyases sanguineus* (Pisces: Gobiesocidae). *Journal of Parasitology* 97(1): 14-19.
- Muñoz G., Garcías F., Valdebenito V., George-Nascimento M. 2001. Parasitofauna y alimentación de *Notothenia c.f. angustata* Hutton, 1875 (Pisces: Nototheniidae) en el intermareal de dos localidades del Golfo de Arauco, Chile. *Boletín Chileno de Parasitología* 56(1-2): 29-33.
- Muñoz G., Valdebenito V., George-Nascimento M. 2002. La dieta y la fauna de parásitos metazoos del torito *Bovichthys chilensis* Regan 1914 (Pisces: Bovichthyidae) en la costa de Chile centro-sur: Variaciones geográficas y ontogenéticas. *Revista Chilena de Historia Natural* 75(4): 661-671.
- Muñoz G., López Z., Cárdenas L. 2013a. Morphological and molecular analyses of larval trematodes in the intertidal bivalve *Perumytilus purpuratus* from central Chile. *Journal of Helminthology* 87(3): 356-363.
- Muñoz G., Torres P., Valdés J., Rodríguez A. 2013b. Spatio-temporal variation in the prevalence of trematodes in the bivalve *Perumytilus purpuratus*. *Acta Parasitologica* 58(2): 155-166.
- Muñoz G., Valdivia I., López Z. 2015. The life cycle of *Prosorhynchoides carvajali* (Trematoda: Bucephalidae) involving species of bivalve and fish hosts in the intertidal zone of central Chile. *Journal of Helminthology* 89(5): 584-592.
- Muñoz G., George-Nascimento M., Bray R.A. 2017. Two new species of digeneans (Lecithasteridae and Haploporidae) of the intertidal blenny *Scartichthys viridis* (Valenciennes) from the central coast of Chile. *Acta Parasitologica* 62(1): 50-62.
- Muñoz G., Vildoso F., George-Nascimento M. 2018. Parasite community of a temporal notothen fish from intertidal rocky pools in south-central Chile: Is it similar to other fish from the same habitat? *Revista de Biología Marina y Oceanografía* 53(2): 209-221.
- Muñoz L., Rubilar L., Zamora D., Sepúlveda O., Rehhof C., Ortiz R. 2008. Fasciolosis en equinos fina sangre de carrera del Club Hípico Concepción, Chile. *Parasitología Latinoamericana* 63(1-2-3-4): 88-91.
- Muñoz L., Sepúlveda-Calderón P., Villaguala-Pacheco C., Aqueveque C.L. 2020. Copro-prevalence of *Fasciola hepatica* in Chilean breed horses in the province of Concepción, Chile. *Brazilian Journal of Veterinary Parasitology* 29(2): e019019.
- Muñoz P., Fredes F., Raffo E., González-Acuña D., Muñoz L., Cid C. 2011. New report of parasite-fauna of the free-tailed bat (*Tadarida brasiliensis*, Geoffroy, 1824) in Chile. *Veterinary Research Communications* 35: 61-66.
- Muñoz-Muga P., Muñoz G. 2010. Parasite communities of *Scartichthys viridis* (Pisces: Blenniidae) from Central Chile: locality vs. host length. *Revista de Biología Marina y Oceanografía* 45(1): 165-169.
- Niklitschek E.J., Secor D.H., Toledo P., Lafon A., George-Nascimento M. 2010. Segregation of SE Pacific and SW Atlantic southern blue whiting stocks: Integrating evidence

- from complementary otolith microchemistry and parasite assemblage approaches. *Environmental Biology of Fishes* 89(3): 399-413.
- Ñacari L.A., Oliva M.E. 2016. Metazoan parasites of deep-sea fishes from the South Eastern Pacific: Exploring the role of ecology and host phylogeny. *Deep-Sea Research Part I: Oceanographic Research Papers* 115: 123-130.
- Ñacari L.A., Sepulveda F.A., Escribano R., Bray R.A., Oliva M.E. 2018. Morphological and molecular characterisation of digenetic parasites of the Galápagos sheepshead *Semicossyphus darwini* (Jenyns) with the re-description of *Labriter secundus* Manter, 1940 (Lepidapedidae) from the Humboldt Current Large Marine Eco. *Systematic Parasitology* 95(4): 391-401.
- Oberg C., Diaz L., Valenzuela G. 1974. Parásitos identificados en bovinos, ovinos, suinos y equinos en el laboratorio de enfermedades parasitarias de la escuela de Medicina Veterinaria de la Universidad Austral de Chile, 1963-1973. *Boletín Chileno de Parasitología* 29(3-4): 99-102.
- Oberg C., Franjola R., Leyán V. 1979. Helmintos del perro doméstico (*Canis familiaris*) en la ciudad de Valdivia, Chile. *Boletín Chileno de Parasitología* 34: 21-26.
- Oliva M.E. 1982. Parásitos en peces marinos de la zona de Antofagasta. *Ciencia y Tecnología del Mar* 6: 45-51.
- Oliva M. 1984a. Nuevos registros de trematodos digenéticos en peces marinos de Antofagasta. *Ciencia y Tecnología del Mar* 8: 9-15.
- Oliva M. 1984b. *Proctoeces chilensis* nueva especie (Trematoda, Strigeatoidea, Fellodistomidae), parásito en *Sicyas sanguineus* Muller & Troschel, 1843 (Pisces: Teleostei). *Boletín de la Sociedad de Biología de Concepción* 55: 87-92.
- Oliva M., Carvajal J. 1984. *Lobatostoma anisotremum* new species, (Trematoda: Aspidogastrea), parasitic in the teleost fish *Anisotremus scapularis* from Chile. *Bulletin of Marine Science* 35(2): 195-199.
- Oliva M., Diaz M. 1988. Aspectos cuantitativos de la infección por *Proctoeces humboldti* (Trematoda: Fellodistomidae) en la lapa *Fissurella crassa* (Mollusca: Archaeogastropoda). *Revista Chilena de Historia Natural* 61(1): 27-33.
- Oliva M., Guerra C. 1987. Infección por *Lecithochirium* sp. (Trematoda: Hemiuridae) en *Gymnothorax porphyreus* (Pisces: Teleostei) del Archipiélago de Juan Fernández. *Estudios Oceanológicos* 6: 103-107.
- Oliva M., Muñoz M.A. 1985. *Helicometra fasciata* (Rudolphi, 1819) y *Helicometrina nimia* Linton, 1910 (Trematoda: Opecoelidae) en peces marinos de la II Región, Chile. *Parasitología al Día* 9: 107-111.
- Oliva M., Zegers J. 1988. Variaciones intraespecíficas del adulto de *Proctoeces lintoni* Siddiqi & Cable, 1960 (Trematoda: Fellodistomidae) en hospedadores vertebrados e invertebrados. *Studies on Neotropical Fauna and Environment* 23(3): 189-195.
- Oliva M.E. 1993. Effect of an adult trematode, *Proctoeces lintoni* (Fellodistomidae), on the gonadosomatic index of *Fissurella limbata* (Archaeogastropoda). *Acta Parasitologica* 38(4): 155-156.
- Oliva M.E. 1999. Metazoan parasites of the jack mackerel *Trachurus murphyi* (Teleostei, Carangidae) in a latitudinal gradient from South America (Chile and Peru). *Parasite* 6(3): 223-230.
- Oliva M.E. 2001. Metazoan parasites of *Macruronus magellanicus* from southern Chile as biological tags. *Journal of Fish Biology* 58(6): 1617-1622.
- Oliva M.E. 1994. Parasites of the chilean jack mackerel *Trachurus symmetricus murphyi* (Pisces: Carangidae). *Memórias do Instituto Oswaldo Cruz* 89(3): 363-364.
- Oliva M.E., Alvarez C. 2011. Is a vertebrate a better host for a parasite than an invertebrate host? Fecundity of *Proctoeces cf lintoni* (Digenea: Fellodistomidae), a parasite of fish and gastropods in northern Chile. *Parasitology Research* 109(6): 1731-1734.
- Oliva M.E., Ballón I. 2002. Metazoan parasites of the Chilean hake *Merluccius gayi gayi* as a tool for stock discrimination. *Fisheries Research* 56(3): 313-320.
- Oliva M.E., Diaz M.A. 1992. An ecological approach to the study of infection of *Proctoeces lintoni* (Digenea, Fellodistomidae) in the keyhole limpet, *Fissurella limbata* Sowerby, 1835 (Archaeogastropoda) from northern Chile. *Acta Parasitologica* 37(3): 115-118.
- Oliva M.E., González M.T. 2004. Metazoan parasites of *Sebastodes capensis* from two localities in northern Chile as tools for stock identification. *Journal of Fish Biology* 64(1): 170-175.
- Oliva M.E., Huaquin L.G. 2000. Progenesis in *Proctoeces lintoni* (Fellodistomidae), a parasite of *Fissurella crassa* (Archaeogastropoda) in a latitudinal gradient in the Pacific coast of South America. *The Journal of Parasitology* 86(4): 768-772.
- Oliva M.E., Luque J.L. 1989. The genus *Lobatostoma* (Trematoda: Aspidocotylea) in the pacific coast of South America, with description of *Lobatostoma veranoi* new species, parasite of *Menticirrhus ophycephalus* (Teleostei: Sciaenidae). *Memórias do Instituto Oswaldo Cruz* 84(2): 167-170.
- Oliva M.E., Sánchez M.F. 2005. Metazoan parasites and commensals of the northern Chilean scallop *Argopecten purpuratus* (Lamarck, 1819) as tools for stock identification. *Fisheries Research* 71(1): 71-77.
- Oliva M.E., Vásquez A.M. 1999. Effects of the Digenea *Proctoeces lintoni* (Fellodistomidae) in the proportion of hemolymphatic cells in *Fissurella crassa* (Mollusca: Archaeogastropoda). *Memórias do Instituto Oswaldo Cruz* 94(6): 827-828.
- Oliva M.E., Vega A.A. 1994. Effect of *Proctoeces lintoni* (Digenea) on the fecundity of *Fissurella crassa* (Archaeogastropoda). *Memórias do Instituto Oswaldo Cruz* 89(2): 225.
- Oliva M.E., Castro R.E., Burgos R. 1996. Parasites of the Flatfish *Paralichthys adspersus* (Steindachner, 1867) (Pleuronectiformes) from Northern Chile. *Memórias do Instituto Oswaldo Cruz* 91(3): 301-306.
- Oliva M.E., Olivares A.N., Diaz C.D., Pasten M. V. 1999. Parasitic castration in *Concholepas concholepas* (Gastropoda: Muricidae) due to a larval digenetic in northern Chile. *Diseases of Aquatic Organisms* 36(1): 61-65.
- Oliva M.E., González M.T., Acuña E. 2006. Metazoan parasite fauna as a biological tag for the habitat of the Flounder *Hippoglossina macrops* from Northern Chile, in a depth gradient. *Journal of Parasitology* 90(6): 1374-1377.
- Oliva M.E., Fernández I., Oyarzún C., Murillo C. 2008. Metazoan parasites of the stomach of *Dissostichus eleginoides* Smitt 1898 (Pisces: Notothenidae) from southern Chile: A tool for stock discrimination? *Fisheries Research* 91(2-3): 119-122.
- Oliva M.E., Valdivia I.M., Cárdenes L., George-Nascimento M., Gonzalez K., Guiñez R.E., Cuello D. 2010. Molecular

- and experimental evidence refuse the life cycle of *Proctoeces lintoni* (Fellodistomidae) in Chile. *Parasitology Research* 106(3): 737-740.
- Oliva M.E., Valdivia I.M., Chavez R.A., Molina H., Cárdenas L. 2015. Molecular and Morphological Evidence Demonstrating Two Species of *Helicometrina* Linton 1910 (Digenea: Opecoelidae) in Northern Chile. *Journal of Parasitology* 101(6): 694-700.
- Oliva M.E., Espinola J.F., Ñacari L.A. 2016. Metazoan parasites of *Brama australis* from southern Chile: A tool for stock discrimination? *Journal of Fish Biology* 88(3): 1143-1148.
- Oliva M.E., Valdivia I.M., Cárdenas L., Muñoz G., Escribano R., George-Nascimento M. 2018. A new species of *Proctoeces* and reinstatement of *Proctoeces humboldti* George-Nascimento and Quiroga 1983 (Digenea: Fellodistomidae) based on molecular and morphological evidence. *Parasitology International* 67(2): 159-169.
- Oliva-Moreno M., Guerra-Correa C. 1988. *Lecithochirium selkiriensis* n. sp. (Trematoda: Hemimuridae), parásito de *Gymnothorax porphyreus* (Pisces: Teleostei). *Revista Ibérica de Parasitología* 48: 373-378.
- Olmos V., George-Nascimento M. 1997. El gremio de larvas de Digenea en el caracol del sur de Chile *Chilina dombeyana*: ¿qué indica la tasa metabólica de los hospedadores parasitados? *Revista Chilena De Historia Natural* 70(2): 109-118.
- Olmos V.L., Victoriano P., Habit E., Valdovinos C. 2003. Parásitos de peces nativos de la cuenca del Río Laja (Chile Central) y alcances sobre sus ciclos de vida. *Archivos de Medicina Veterinaria* 35(2): 195-203.
- Olmos V., Muñoz G. 2006. Estado de conocimiento de los parásitos Eumetazoos de organismos Dulceacuícolas de Chile. *Gayana* 70(1): 122-139.
- Osorio C., Castillo M. 1984. Almejas comestibles (*Venus antiqua*) infectadas por tremátodos. *Parasitología al Día* 8: 117-118.
- Osorio C., Díaz H., Ríos M. 1986. Prevalencia de *Proctoeces humboldti* George-Nascimento & Quiroga 1983 (Digenea, Fellodistomidae) en la población de *Fissurella maxima* Sowerby 1835 (Mollusca, Gastropoda) de Los Vilos, Chile. *Revista de Biología Marina* 22(2): 157-168.
- Ostrowski de Núñez M., Flores V., Viozzi G., Kreiter A. 2004. *Stephanopora uruguayense* Holcman-Spector et Olagüe, 1989 (Digenea, Echinostomatidae) from Argentina, and comments on species of *Stephanopora* from birds of the Neotropical Region. *Acta Parasitologica* 49(4): 292-299.
- Ostrowski de Núñez M. 2007. Life cycle of *Stephanopora uruguayense* (Digenea: Echinostomatidae) in Argentina. *Journal of Parasitology* 93(5): 1090-1096.
- Oyarzún-Ruiz P., González-Acuña D. 2021. Checklist and state of knowledge of helminths in wild birds from Chile: an update. *Austral Journal of Veterinary Sciences* 53(1): 63-72.
- Oyarzún-Ruiz P., Muñoz-Alvarado P., Raffo E. 2016. Helminths of *Milvago chimango temucoensis* (Aves: Falconiformes) from Los Ríos Region, Chile: New Records for Neotropical Raptors. *Helminthologia* 53(4): 336-353.
- Oyarzún-Ruiz P., Barrientos V., Rodríguez R., Almonacid A., Barrientos O., Paineau J., Ortiz C., Ratto M. 2017. Identificación y cuantificación de parásitos en Llamas (*Lama glama*) de Valdivia, Chile. *Parasitología Latinoamericana* 66(3): 364.
- Oyarzún-Ruiz P., Muñoz P., Paredes E., Valenzuela G., Ruiz J. 2019a. Gastrointestinal helminths and related histopathological lesions in black-necked swans *Cygnus melanocoryphus* from the Carlos Anwandter Nature Sanctuary, Southern Chile. *Brazilian Journal of Veterinary Parasitology* 28(4): 613-624.
- Oyarzún-Ruiz P., Alvelo C., Vera F., Moroni M. 2019b. Histopathological findings of *Fasciola hepatica* infection in non-native European hare (*Lepus europaeus*) in Southern Chile. *Brazilian Journal of Veterinary Parasitology* 28(1): 145-150.
- Oyarzún-Ruiz P., Di Cataldo S., Cevidanes A., Millán J., González-Acuña D. 2020. Endoparasitic fauna of two South American foxes in Chile: *Lycalopex culpaeus* and *Lycalopex griseus*. *Brazilian Journal of Veterinary Parasitology* 29(3): e006220.
- Pardo-Gandarillas M.C., Garcías F., George-Nascimento M. 2004. La dieta y fauna de endoparásitos del pejeshapo *Gobiesox marmoratus* Jenyns, 1842 (Pisces: Gobiesocidae) en el litoral central de Chile están conectadas pero no correlacionadas. *Revista Chilena de Historia Natural* 77(4): 627-637.
- Pardo-Gandarillas M.C., González K., Ibáñez C.M., George-Nascimento M. 2008. Parasites of two deep-sea fish *Coelorynchus chilensis* (Pisces: Macrouridae) and *Notacanthus sexspinis* (Pisces: Notacanthidae) from Juan Fernández Archipelago, Chile. *Marine Biodiversity Records* 1: e76.
- Paré J.A., Black S.R. 1999. Schistosomiasis in a Collection of Captive Chilean Flamingos (*Phoenicopterus chilensis*). *Journal of Avian Medicine and Surgery* 13(3): 187-191.
- Pequeño G., Valdebenito V., Muñoz G. 2011. *Pseudopentacerus richardsoni* (Osteichthyes: Pentacerotidae) en el Océano Pacífico suroriental: Morfometría, parásitos y comentarios sobre la familia en el área. *Revista de Biología Marina y Oceanografía* 46(2): 275-280.
- Pinto H.A., Brant S. V., de Melo A.L. 2014. *Physa marmorata* (Mollusca: Physidae) as a natural intermediate host of *Trichobilharzia* (Trematoda: Schistosomatidae), a potential causative agent of avian cercarial dermatitis in Brazil. *Acta Tropica* 138: 38-43.
- PNUD. 2017. Catálogo de las especies exóticas asilvestradas/naturalizadas en Chile. *Laboratorio de Invasiones Biológicas (LIB)* Universidad de Concepción, Proyecto GEF/MMA/PNUD Fortalecimiento de los Marcos Nacionales para la Gobernabilidad de las Especies Exóticas Invasoras: Proyecto Piloto en el Archipiélago de Juan Fernández, Santiago de Chile, 61 pp.
- Povilitis A. 1998. Characteristics and conservation of a fragmented population of huemul *Hippocamelus bisulcus* in central Chile. *Biological Conservation* 86(1): 97-104.
- Puente S. 1963. Distomatosis coledociana. Primer caso observado en Arica. *Boletín Chileno de Parasitología* 18: 107-108.
- Puga S. 1979. *Gorgoderina valdiviensis* sp. nov., un nuevo trematóodo digénico (Gorgoderidae) parásito de la rana chilena (*Caudiverbera caudiverbera*). *Studies on Neotropical Fauna and Environment* 14(4): 227-232.
- Puga S. 1982. Two platyhelminths from *Caudiverbera caudiverbera* (Anura: Leptodactylidae) in southern Chile. *Herpetological Review* 13(1): 13-14.
- Puga S. 1986. *Rudolphitrema chilensis* sp. nov. un nuevo trematóodo digénico parásito del anuro chileno *Eusophus roseus* (Leptodactylidae). *Boletín Chileno de Parasitología* 41(1-2): 13-16.

- Puga S., Figueroa L. 1989. *Pirupalkia queulensis* n. gen., n. sp. (Digenea: Calyptostomidae) en el pez *Cilus montti* (Pomadasytidae). *Boletín Chileno de Parasitología* 44: 43-45.
- Puga S., Figueroa L. 1993. *Lepocreadium valdiviensis* nov. sp. (Digenea: Lepocreadiidae) parásito de *Paralichthys microps* "lenguado de ojos chicos" en el sur de Chile. *Boletín Chileno de Parasitología* 48: 55-57.
- Puga S., Torres P. 1999. Helminths parasites of *Eupsophus roseus* (Anura: Leptodactylidae) from Southern Chile. *Memórias do Instituto Oswaldo Cruz* 94(6): 725-726.
- Rebolledo M., Landaeta M.F., Muñoz G. 2014. Efecto del endoparásito *Prostorhynchoides* sp. (Trematoda: Bucephalidae) en la capacidad de nado sostenido del baunco *Girella laevifrons* (Osteichthyes: Kynphosidae). *Revista de Biología Marina y Oceanografía* 49(3): 625-630.
- Reyes P., Hüne M. 2012. Peces del Sur de Chile. *Ocho Libros Editores, Santiago de Chile*, 500 pp.
- Riffo R. 1991. La fauna de parásitos metazoos del lenguado de ojo grande *Hippoglossina macrops* Steindachner, 1876 (Pisces: Bothidae): una aproximación ecológica. *Medio Ambiente* 11(2): 54-60.
- Riffo R. 1994. Composición taxonómica y características cuantitativas de la fauna de parásitos metazoos del congro dorado *Genypterus blacodes* Schneider 1801. *Medio Ambiente* 12(1): 27-31.
- Riffo R. 1995. Análisis comparativo de la fauna de parásitos metazoos de dos especies de lenguados congénéricos y sintópicos: *Paralichthys microps* Gunther 1881 y *Paralichthys adspersus* Steindachner 1867 (Pleuronectiformes: Bothidae) en la Bahía Concepción. *Medio Ambiente* 12(2): 51-59.
- Riffo R., Nuñez M. 2000. La fauna de parásitos metazoos del rollizo *Mugilooides chilensis* (Pisces: Mugiloididae): la importancia de las relaciones ecológicas del hospedador. *Medio ambiente (Valdivia)* 13(2): 89-96.
- Roberts L.S., Janovy Jr. J., Nadler S. 2013. Foundations of Parasitology. 9th ed. *McGraw-Hill Company, New York*, 697 pp.
- Rodríguez L., George-Nascimento M. 1996. La fauna de parásitos metazoos del bacalao de profundidad *Dissostichus eleginoides* Smitt, 1898 (Pisces: Nototheniidae) en Chile central: aspectos taxonómicos, ecológicos y zoogeográficos. *Revista Chilena de Historia Natural* 69(1): 21-33.
- Rojas A., Villarroel F., Díaz F., Rubio P., Schenone H. 1971. Investigación de triquinosis y capilariasis hepática en *Rattus norvegicus* del Matadero Municipal de Santiago (Chile). *Boletín Chileno de Parasitología* 26(1-2): 65-66.
- Rojas J.R., Torres P., Pequeño G. 1999. Acanthocolpid metacercariae in the sea bass from Alejandro Selkirk Island, Chile. *Journal of Wildlife Diseases* 35(1): 90-93.
- Romero J., Villaguala C., Quiroz F., Landaeta-Aqueveque C., Alfaro G., Pérez R. 2019. Flukicide efficacy against *Fasciola hepatica* of Triclabendazole and Nitroxynil in cattle of the central valley of Chile. *Brazilian Journal of Veterinary Parasitology* 28(1): 164-167.
- Salinas X., González M.T., Acuña E. 2008. Metazoan parasites of the thumb grenadier *Nezumia pulchella*, from the south-eastern Pacific, off Chile, and their use for discrimination of host populations. *Journal of Fish Biology* 73(3): 683-691.
- Sanhueza C., Álvarez M. 1981. Zoonosis comprobadas en animales beneficiados en los mataderos del país durante 1979. *Boletín de Vigilancia Epidemiológica* 7: 2-4.
- Sapunar J., Donckaster R. 1966. Peritoneoscopías en fascioliasis hepática humana. *Boletín Chileno de Parasitología* 21: 21.
- Sapunar J., Donckaster R. 1967. Fascioliasis hepática humana. Laparoscopía en períodos de invasión y estado. *Boletín Chileno de Parasitología* 22: 90-96.
- Sapunar J., Gallo G., Csendes A., Sapunar J. 1983. Fascioliasis hepática diagnosticada por colangiografía endoscópica. *Boletín Chileno de Parasitología* 38: 17-20.
- Sapunar J., Braghetto I., Díaz J.C., Brahm J., Apt W., Carreño L., Poniachick J. 2001. Fascioliasis hepática que simularon tumores hepáticos. *Boletín Chileno de Parasitología* 56(3-4): 59-65.
- Schell S.C. 1985. Handbook of Trematodes of North America, North of Mexico. *University Press of Idaho, Idaho*, 263 pp.
- Schenone H., Rojas A. 1988. Epidemiología de la fascioliasis animal en Chile. Tendencia de las tasas de infección, por regiones, en cinco especies de mamíferos de consumo beneficiados en mataderos del país 1977-1986. *Boletín Chileno de Parasitología* 43: 68-70.
- Seguel M., Muñoz F., Paredes E., Navarrete M.J., Gottdenker N.L. 2017. Pathological Findings in Wild Rats (*Rattus rattus*) Captured at Guafo Island, Northern Chilean Patagonia. *Journal of Comparative Pathology* 157(2-3): 163-173.
- Sepúlveda F., Marín S.L., Carvajal J. 2004. Metazoan parasites in wild fish and farmed salmon from aquaculture sites in southern Chile. *Aquaculture* 235(1-4): 89-100.
- Sepúlveda M.S., Alcaíno H. 1993. Fauna helmintológica en el lobo fino de Juan Fernández, *Arctocephalus philippii* (Peters, 1866). *Parasitología al Día* 17: 19-24.
- Shimazu T., Urawa S., Coria C.O. 2000. Four species of digeneans, including *Allocreadium patagonicum* sp. n. (Allocreadiidae), from freshwater fishes of Patagonia, Argentina. *Folia Parasitologica* 47(2): 111-117.
- Siegmund I., Franjola R., Torres P. 1997. Diplostomatid metacercariae in the brain of silversides from Lake Riñihue, Chile. *Journal of Wildlife Diseases* 33(2): 362-364.
- Stoore C., Andrade C., Hidalgo C., Corrêa F., Jiménez M., Hernandez M., Paredes R. 2018. *Echinococcus granulosus* hydatid cyst location is modified by *Fasciola hepatica* infection in cattle. *Parasites & vectors* 11(1): 542.
- Suárez-Villota E.Y., Quercia C.A., Nuñez J.J., Gallardo M.H., Himes C.M., Kenagy G.J. 2018. Monotypic status of the South American relictual marsupial *Dromiciops gliroides* (Microbiotheria). *Journal of Mammalogy* 99(4): 803-812.
- Subercaseaux B., Tapia S., Guglielmetti A., Stanley W., Muñoz N. 1985. Brote epidémico de fascioliasis hepática humana en Valparaíso. *Parasitología al Día* 9(1): 10-14.
- Szidat L. 1962. *Steganoderma oviformis* n. sp. (Trematoda) del intestino de *Aplochiton zebra* Jenyns. *Neotropica* 8(26): 67-72.
- Szidat L. 1969. Structure, Development, and Behaviour of New Strigeatoid Metacercariae from Subtropical Fishes of South America. *Journal of the Fisheries Research Board of Canada* 26(4): 753-786.
- Tagle I. 1942. Parásitos de los animales domésticos en Chile determinados en el Instituto de Investigaciones Veterinarias. *Agricultura Técnica* 13(2): 93-108.
- Tagle I. 1944. Observaciones sobre la evolución de la *Fasciola hepatica* Linneo 1758. Comprobación del huésped intermedio en Chile. *Revista Chilena de Historia Natural* 46-47(1): 232-241.
- Tagle I. 1966. Parásitos de los animales domésticos en Chile. *Boletín Chileno de Parasitología* 21(4): 118-123.
- Threlfall W., Carvajal J. 1986. *Otodistomum cestoides* (van Beneden, 1871) from two species of skates taken in Chilean

- waters. *Memórias do Instituto Oswaldo Cruz* 81(3): 341-342.
- Torres P. 1995. Some Trematode, Nematode, and Acanthocephalan parasites of rainbow trout, *Oncorhynchus mykiss*, introduced into Chile. *Journal of the Helminthological Society of Washington* 62(2): 257-259.
- Torres P., Neira A. 1991. A new species of *Limnoderecrema* (Trematoda, Digenea) from the freshwater Atherinid fish *Basilichthys australis* Eigenmann, 1927 from the south of Chile. *Memórias do Instituto Oswaldo Cruz* 86(4): 433-436.
- Torres P., Hott A., Boehnwald H. 1972. Protozoos, helmintos y artrópodos en gatos de la ciudad de Valdivia y su importancia en el hombre. *Archivos de Medicina Veterinaria* 4: 20-29.
- Torres P., Ramos M., Carrasco L., Neumann M., Franjola R., Navarrete N., Figueroa L. 1974. Protozoos, helmintos y artrópodos del perro doméstico en la ciudad de Valdivia, Chile. *Boletín Chileno de Parasitología* 29: 18-23.
- Torres P., Hauser M., Santibáñez J., Marín F., Gesche W., Montefusco A. 1980. Búsqueda de *Diphyllobothrium* y otros parásitos intestinales en la población humana y carnívoros domésticos del sector del lago Calafquén, Chile. *Boletín Chileno de Parasitología* 35(3-4): 55-61.
- Torres P., Figueroa L., Saldívia A., Barrientos J. 1982. Gastrointestinal Helminths of Fish-eating birds from the Valdivia River, Chile. *Journal of Parasitology* 68(6): 1157.
- Torres P., Figueroa L., Saldívia A. 1983. *Stephanopraora denticulata* (Trematoda, Echinostomatidae) en gaviotas del sur de Chile. *Boletín Chileno de Parasitología* 38(1-2): 33-34.
- Torres P., Franjola R., Cubillos V., Miranda J.C., Vera R. 1988. Parasitismo en ecosistemas de agua dulce en Chile 1. Presencia de metacercarias del género *Stephanostomum* (Digenea: Acanthocolpidae) en peces. *Journal of Veterinary Medicine, Series B* 35(1-10): 169-177.
- Torres P., Ruiz E., Gesche W., Montefusco A. 1991. Gastrointestinal helminths of fish-eating birds from Chiloe Island, Chile. *Journal of Wildlife Diseases* 27(1): 178-179.
- Torres P., Contreras A., Cubillos V., Gesche W., Montefusco A., Rebolledo C., Mira A., Arenas J., Miranda J.C., Asenjo S., Schlatter R. 1992a. Parasitismo en peces, aves piscívoras y comunidades humanas ribereñas de los lagos Yelcho y Tagua-Tagua, X Región de Chile. *Archivos de Medicina Veterinaria* 24(1): 77-92.
- Torres P., Oporto J.A., Brieva L.M., Escare L. 1992b. Gastrointestinal helminths of the cetaceans *Phocoena spinipinnis* (Burmeister, 1865) and *Cephalorhynchus europia* (Gray, 1846) from the southern coast of Chile. *Journal of Wildlife Diseases* 28(2): 313-315.
- Torres P., Schlatter R., Montefusco A., Gesche W., Ruiz E., Contreras A. 1993a. Helminth parasites of piscivorous birds from lakes in the south of Chile. *Memórias do Instituto Oswaldo Cruz* 88(2): 341-343.
- Torres P., Contreras A., Revenga J., Fritz N. 1993b. Helminth parasites in fishes from Valdivia and Tornagaleones river estuaries in the south of Chile. *Memórias do Instituto Oswaldo Cruz* 88(3): 491-492.
- Torres P., Franjola R., Montefusco A. 1996. Infección estacional por metacercarias de *Diplostomum (Austrodiplostomum) mordax* (Szidat y Nani, 1951) y *Tyloodelphys destructor* Szidat y Nani, 1951 en el pejerrey chileno, *Basilichthys australis* Eigenmann, 1927 (Pisces: Atherinidae) en el lago Riiñihue, Chile. *Boletín Chileno de Parasitología* 51(1): 15-19.
- Torres P., Leyan V., Lamilla J. 2017. Cyst Stages of Gordiids (Nematomorpha) and Other Eukaryotic Parasites from the Inanga, *Galaxias maculatus* (Osmeriformes: Galaxiidae), in the Lingue River, Southern Chile. *Comparative Parasitology* 84(1): 72-79. <https://doi.org/10.1654/1525-2647-84.1.72>
- Valderrama K., Oliva M., Campos B., Brown D. 2004. Parasitic castration of *Eurhomalea lenticularis* (Bivalvia: Veneridae) by a digenetic trematode: Quantitative histological analysis. *Diseases of Aquatic Organisms* 59(2): 151-158.
- Valdivia I.M., Cardenas L., Gonzalez K., Jofré D., George-Nascimento M., Guiñez R., Oliva M.E. 2010. Molecular evidence confirms that *Proctoeces humboldti* and *Proctoeces chilensis* (Digenea: Fellodistomidae) are the same species. *Journal of Helminthology* 84(4): 341-347.
- Valdivia I.M., Criscione C.D., Cárdenas L., Durán C.P., Oliva M.E. 2014. Does a facultative precocious life cycle predispose the marine trematode *Proctoeces cf. lintoni* to inbreeding and genetic differentiation among host species? *International Journal for Parasitology* 44(3-4): 183-188.
- Valdovinos C. 1999. Biodiversidad de moluscos chilenos: base de datos taxonómica y distribucional. *Gayana* 63(2): 111-164.
- Valdovinos C., Balboa C. 2008. Cercarial dermatitis and lake eutrophication in south-central Chile. *Epidemiology & Infection* 136(3): 391-394.
- Valenzuela G. 1979. Estudio epidemiológico sobre el desarrollo de huevos de *Fasciola hepatica* en el medio ambiente de Valdivia, Chile. *Boletín Chileno de Parasitología* 34(1-2): 31-35.
- Valenzuela G., Quintana I. 1998. Evolución de huevos de *Fasciola hepatica* en el medio ambiente en Temuco, IX Región de Chile. *Archivos de Medicina Veterinaria* 30(1): 109-114.
- Valenzuela G., Fuentealba L., Henríquez O. 1979. Acción de la temperatura ambiente sobre el desarrollo de huevos de *Fasciola hepatica* en Valdivia. Período Invierno-Primavera. *Archivos de Medicina Veterinaria Suplem.* 1: 134-139.
- Valenzuela G., Araya A., Oyarzún-Ruiz P., Muñoz P. 2018. Helminto fauna del cisne de cuello negro *Cygnus melancoryphus* (Aves: Anatidae) del Santuario de la Naturaleza Carlos Anwandter, Valdivia, Chile. *Revista Mexicana de Biodiversidad* 89(2): 568-571.
- Venturelli A., Monje M., Assef V., Venturelli F. 2003. Fasciolasis hepática. *Cuadernos de Cirugía* 17(1): 43-46.
- Vergara L.A., George-Nascimento M. 1982. Contribución al estudio del parasitismo en el congrio colorado *Genypterus chilensis* (Güichenot, 1848). *Boletín Chileno de Parasitología* 37(1-2): 9-14.
- Villalba C., Fernández J. 1985. Parásitos de *Mola ramsayi* (Giglioli, 1883) (Pisces: Molidae) en Chile. *Boletín de la Sociedad de Biología de Concepción* 56: 71-78.
- Villalba C., Fernández J. 1986a. Tres nuevas especies de *Aporocotyle* Odhner, 1900 (Digenea: Sanguinicolidae) parásitas de *Genypterus* spp. en Chile (Pisces: Ophidiidae). *Revista de Biología Marina* 22(2): 125-139.
- Villalba C., Fernández J. 1986b. Dos nuevas especies de tremátodos parásitos de peces marinos en Chile. *Parasitología al Día* 10: 45-51.
- Viozzi G., Flores V., Marín S.L., Mancilla M., Carvajal J. 2008. Parasites of the Red Jollytail, *Brachygalaxias bullocki* (Osmeriformes: Galaxiidae), from the Maullín River, Patagonia, Chile. *Comparative Parasitology* 75(2): 326-328.