

## **Bats (Mammalia, Chiroptera) from a cave area in NW Italy**

Authors: Godio, Ilaria, Toffoli, Roberto, Trombin, Denise, and Cucco, Marco

Source: Revue suisse de Zoologie, 129(1) : 137-145

Published By: Muséum d'histoire naturelle, Genève

URL: <https://doi.org/10.35929/RSZ.0068>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Bats (Mammalia, Chiroptera) from a cave area in NW Italy

Ilaria Godio<sup>1</sup>, Roberto Toffoli<sup>2</sup>, Denise Trombin<sup>2</sup>, Marco Cucco<sup>1\*</sup>

<sup>1</sup> University of Piemonte Orientale, Vercelli, Italy

<sup>2</sup> Chiroptera, Association for the Study and Protection of Bats and the Environment, Santena, Italy

\* Corresponding author: marco.cucco@uniupo.it, ORCID: 0000-0002-4749-3653

### Abstract

Caves play an important role in bat ecology, offering natural roosting and wintering sites. In the reproductive period, bats need to find an effective foraging environment away from the cave entrance, but only few studies have considered the use of foraging habitats in the immediate proximity of the caves. We provide a list of 18 bat species and 5 bat acoustic groups observed in a system of 81 caves in NW Italy. Winter censuses carried out during 17 years showed an increase of the Greater Horseshoe bat, a species of conservation concern. In the summer period this rare species was also observed foraging in nearby areas. Recording of bat calls in the summer period was performed both at cave entrances and in three foraging habitats present in the area surrounding the cave (woods, agricultural mosaic with artificial light, dark mosaic). Light mosaics and woods showed the highest foraging activity (160.3 and 72.0 pass/hour), while woods and cave entrances were accessed by the largest number of bat species. Our survey improves the inventory of bat species in a poorly sampled area, and we recommend a reduced tourism access, thereby promoting preservation of bats.

**Keywords:** Monte Fenera karst caves - *Rhinolophus ferrumequinum* - Vespertilionidae - extra-cave habitats.

### INTRODUCTION

Although caves may be used by bats throughout all year (e.g. as thermally favourable breeding habitat), it is well known that their most important role in ecology of temperate bats is as hibernacula (Zukal *et al.*, 2017). However, even if there are a multitude of studies examining the presence of bats wintering in caves as part of bat monitoring programs (Loeb *et al.*, 2015; Collins, 2016), to date there are surprisingly few systematic investigations of bat activity in habitats nearby hypogeal sites compared to other habitats (Wieser *et al.*, 2020).

Hibernacula have a key role in the life cycle of bats (Altringham, 2011; Toffoli & Calvini, 2021) and the survival of many bat species worldwide depends on the availability of natural caves and other underground sites such as mines (Mickleburgh *et al.*, 2002) that provide relatively stable above-freezing temperature and high humidity (Zukal *et al.*, 2017). Subterranean winter temperatures are warmer and less variable further from the surface (McClure *et al.*, 2020) and are close to the mean annual surface temperature for the nearby area (Zukal *et al.*, 2017). Furthermore, hibernaculum humidity impacts overwinter survival since dry conditions may

lead to reduced length of torpor bouts (Thomas & Geiser, 1997) and may stimulate arousals (Ben-Hamo *et al.*, 2013; Klüg-Baerwald & Brigham, 2017), increasing death probability. Highly stable climate conditions in caves thus promote energy-saving torpor and enable hibernating bats to overcome cold winters (McClure *et al.*, 2020; Speakman & Thomas, 2003).

Identifying and protecting hibernacula is crucial for bats conservation (Glover & Altringham, 2008) and their numbers and diversity in hypogeal environments are influenced by many factors including human disturbance, cave dimensions, structural complexity and microclimate, availability of food in the surrounding landscape, parasite and predation pressure. Historical use by bats and maneuverability of flight and interactions between species have a relevant impact on roost selection as well (Furey & Racey, 2015). Intraspecific needs and tolerance may vary geographically (Klüg-Baerwald & Brigham, 2017) and they also determine the suitability of a site. Selection of microclimate conditions by bats during hibernation is complex (McClure *et al.*, 2020) and not easily predictable.

External environments surrounding caves are equally important for bats all year round, even though foraging

habitat preferences may vary among species. Forests provide cover for emerging bats reducing flight costs and predation risk (Verboom & Spoelstra, 1999; Coleman & Barclay, 2013) and represent a source of invertebrate prey (Kofoky *et al.*, 2006), whilst forest edges contribute to landscape connectivity enhancing the movement of species (Heim *et al.*, 2015). Scattered trees and hedgerow in farmland and prairie riparian areas harbour large numbers of insects and, consequently, they are used as foraging habitats (Lumsden & Bennett, 2005; Holloway & Barclay, 2000; Toffoli, 2016). In Mediterranean regions, wetlands constitute an essential ecosystem for bats and a higher bat foraging activity is reported near more accessible water bodies (Rainho, 2007). Anthropogenic environments are used by bats as well but, although they might be particularly attractive to *Pipistrellus*, *Eptesicus* and *Hypsugo* species, they are much less so to *Myotis* and *Plecotus* and they can even cause habitat fragmentation (Barataud, 2016).

While the study of the relationships between habitats and bat presence, or bat activity, has attracted much attention in a multitude of areas (Kunz & Fenton, 2003), the study of bat activity near the area surrounding the cave entrance has drawn little attention. We carried out systematic recordings of bat calls at cave entrance and three other habitat types (wood, dark open mosaic, open mosaic with light) to better understand the importance of these habitats for bats.

We examined the Monte Fenera Natural Park area, in NW Italy, which includes 81 caves (Lana & Sella, 1986). Previous biospeleological surveys conducted by Gruppo Speleologico Biellese C.A.I. (GSBi-CAI) from 1993 to 1999 reported the presence of Mouse-eared bats *Myotis myotis/blythii*, Lesser Horseshoe bats *Rhinolophus hipposideros* and Greater Horseshoe bats *Rhinolophus ferrumequinum* not only inside the major caves of the massif but also in several small cavities (Pascutto & Balestrieri, 2001). Further observations in subsequent




Fig. 1. Map of the study area in NW Italy. Star: sampled sites of the Fenera's caves area.

years confirmed this first list of species and revealed the attendance of a cave by Long-eared bats (*Lana et al.*, 2021). The availability of different habitat types nearby caves makes the area suitable as breeding, hunting or swarming sites not only for cave-dwelling bats, extremely favoured by the geological features of Monte Fenera, but for tree-dwelling bats as well.

Our study aims to fill the lack of data about bat fauna in this area by utilizing winter censuses carried out along a 17-year period, some captures with mist nets at cave entrance, several occasional reports, and an intense monitoring through bat call recordings collected in 12 sites pertaining to four habitat categories inside the Fenera area.

## MATERIAL AND METHODS

### Study Area

The Monte Fenera, in NW Italy, is a protected area of 3378 ha (Fig. 1), included in the Sites of Community Importance list (code IT1120003). Evidence of the use of the caves since the Riss-Würm interglacial was documented by the description of fossils of at least 9 species of bats (*Berto et al.*, 2016). Our sample sites are comprised in the municipalities of Borgosesia, Valduggia and Grignasco. Most of the protected area is a carbonate-built hill that is home to 81 karstic caves at an altitude of 600-800 m a.s.l. The environment surrounding the caves' entrance is mainly composed of woods, with some cultivated areas and small villages from 2-3 km henceforth.

### Winter areas

We inspected the four largest caves, namely:

- Bondaccia (Italian cave registry code PI2505); WGS84 45.7114438°N, 8.3118285°E, 690 m a.s.l., 500 m-long.
- Ciota Ciara (code PI2507); WGS84 45.7108912°N, 8.3112443°E, 675 m a.s.l., 202 m-long.
- Ciutarun (code PI2506); WGS84 45.710138°N, 8.3102257°E, 655 m a.s.l., 66 m-long.
- Arenarie (code PI2509); WGS84 45.711919°N, 8.3145207°E, 770 m a.s.l., 3 km-long.

### Summer-autumn areas

We inspected the environment surrounding the Monte Fenera cave area, grouping the observations among four habitat categories:

- cave entrance: the sampling covers the space (about 20-30 m) in front of the entrance of three large caves, Bondaccia, Ciota Ciara and Ciutarun.
- woods: five sampling points. The woods are dominated by chestnut which is accompanied by other species such as ash, English oak, sessile oak, Turkey oak, birch, maple, aspen, wild cherry, willow, and mountain ash; black locust has invaded the southern hills, while in the humid valleys and along the streams there is alder and

black poplar. To the north, on the cooler slopes, beech is abundant, while to the south, on rocky walls and calcareous soils, there is manna ash, a typical essence of Mediterranean environment to which butcher's broom and juniper also belong, which are found in abundance. Natural conifers (Scots pine) are present sporadically while the most common are those of planting, preferred by man for their rapid growth (such as strobe pine). Among typical shrubs in the woods are hazelnut, dogwood, elder, hawthorn, barberry and privet.

- agricultural mosaic with light: two sampling points. Open spaces with grass vegetation and orchards, with small buildings and the presence of light coming from the houses.
- dark agricultural mosaic: two sampling points. Open spaces with grass vegetation and orchards, with no buildings nor light.

### Areas with incidental observations

We utilized some occasional data collected by speleologists in smaller caves:

- La Caudrola PI2735: WGS84 45.7045478°N, 8.3101146°E, 515 m a.s.l.
- Buco delle Marmite della cava Antoniotti PI2550: WGS84 45.7006103°N, 8.312411°E, 402 m a.s.l.
- Buco delle Radici PI2540: WGS84 45.7116969°N, 8.3225534°E, 804 m a.s.l.
- Bell'ingresso PI2539: WGS84 45.7104921°N, 8.3242896°E, 754 m a.s.l.

Other data were collected from citizens reporting dead specimens, and in a single event placing mist nets at the Bondaccia cave entrance in October 2012.

## Field surveys

### Winter census

Hibernating individuals were monitored through standardized counts carried out in their underground roosts between the last five days of December and the first ten days of February (Battersby, 2010). Hibernation site surveys were repeated once every year at each site in order to minimize disturbance. Bats were counted directly or with binoculars along all human accessible tracts in caves, or with the aid of images in case of aggregated groups that required the use of photography or video cameras (Toffoli & Calvini, 2021).

### Summer-autumn census

In our study area, the year-round reproductive life cycle of bats typically starts with the autumnal swarming-mating period, followed in spring and early-summer by a female-lactating period. In late summer, both adults and recently born bats can be observed foraging. In this study we inspected the environment utilized by foraging bats from early July to early October, thus encompassing both the lactating period and the periods when young bats start to forage independently, with some overlapping with the mating period.

Bat activity was investigated using AudioMoth bat detectors (Hill *et al.*, 2019), with their microphones placed 1.5 m above the ground. Real time recordings were made all-night-long at a sampling frequency of 256 kHz. We started bat acoustic sampling half an hour after sunset until half an hour before sunrise. This time interval includes the main peaks of activity for most insectivorous bats of the temperate zone (Erkert, 1982). Recordings were made on nights without precipitation, with wind strength below 6 m/s and temperature above 10 °C at sunset, to avoid measuring low bat activity due to weather factors. All recordings were then divided in segments of 5 sec duration using the Kaleidoscope program 3.1.1 (Wildlife Acoustics, Maynard, Massachusetts, USA).

To describe activity levels, the number of passes per hour of each different species and acoustic groups was used (Barataud, 2016). A pass is defined as a recording consisting of at least one echolocation signal (call) up to a total length of 5 sec (Thomas & West, 1989; Barataud, 2016). In the results, we reported the average number of bat passes / hour in order to take into account the difference in the number of sampled sites and the recording time in the different habitat categories.

## Data analysis

### Winter census

Population trend was analyzed with the TRIM software (Pannekoek & Van Strien, 2001) through the package “Rtrim” 2.0.6 within the R statistical environment (Bogaart *et al.*, 2016). TRIM implements log-linear Poisson regression models to analyse time series of count data (Gregory *et al.*, 2005), and was developed for the analysis of time series of counts with missing values (Pannekoek & van Strien, 2001). It has been utilized to analyze trends in hibernating bats (Uhrin *et al.*, 2010; Van der Meij *et al.*, 2015; Toffoli & Calvini, 2019).

### Summer-autumn census

Species identification was obtained by analyzing each bat call sequence following the methodological approach provided by Barataud (2016) and comparing the data with those published by Russo & Jones (2002). The identification method is based on the signal shape (FM: frequency modulation; QCF: quasi constant frequency), the energy peak distribution and on the measurement of the following parameters: start frequency (SF), end frequency (EF), bandwidth (BW), frequency of maximum energy (FME) in kHz, duration (D) and inter-pulse interval in ms (IPI). To reduce the time spent examining sonograms, in a first step we utilized the SonoChiro® 3.3.2 software (Biotopie Society, France) for automatic identification of echolocation signals of European bats (Azam *et al.*, 2015; Charbonnier *et al.*, 2016). This software identifies signals in a recorded sequence and classifies them at a homogeneous species group level according to characteristics of signals. The software attributes a reliability index from 0 to 10 for each

identification. In a second step, we controlled manually all sequences with a score lesser than 5, as indicated by Toffoli & Rughetti (2017), to attribute sequences to one species or one acoustic group according to the characters reported in Barataud (2016), Russo & Jones (1999) and Middleton *et al.* (2014).

Signals belonging to the genus *Myotis*, excluding those classified at a specific level, were incorporated into the two groups *Myotis* LF (*Myotis myotis* and *Myotis blythii*) and *Myotis* HF (all other species), due to the difficulties encountered in identifying them at species level (Russo & Jones, 2002). Due to a lack of firm discrimination criteria for bioacoustic distinction between *Plecotus auritus*, *Plecotus austriacus* and *Plecotus macrobullaris*, we attributed all signals of the genus to the *Plecotus* sp. group.

With regard to *Pipistrellus kuhlii* and *Pipistrellus nathusii*, whose identification is difficult (Barataud, 2016), only sequences with a score equal to or greater than 9 were attributed to one of the two species. All others were controlled manually and only those with the characteristic social calls were identified at species level (Russo & Jones, 1999; Middleton *et al.*, 2014), while the remnants were referred to the acoustic group *Pipistrellus kuhlii/nathusii*.

Sequences with FME signals from 20 to 30 kHz and the presence of characteristic alternate FM/QCF and QCF signals of different FME were associated to the *Nyctalus* genus (Waters & Jones, 1995). Sequences of primarily FM/QCF signals either longer than 17 ms and with end frequency between 21 and 24 kHz, or shorter than 5 ms and with EF between 29 and 32 kHz were associated to *Eptesicus serotinus* (Barataud, 2016). Concerning the identification of European Free-tailed Bat *Tadarida teniotis*, in order to exclude *Nyctalus lasiopterus* and the social calls of *Nyctalus leisleri*, reference was made to the guidelines of Barataud (2016). In particular, in order to exclude *Nyctalus lasiopterus*, sequences without alternating structure between QCF and FM signals and with a maximum energy frequency between 9 and 13 kHz have been attributed to *Tadarida teniotis*, while as regard the exclusion with the social calls of *Nyctalus leisleri*, reference was made to the initial form of the characteristic signal of the social calls of *Nyctalus leisleri*.

## RESULTS

### Winter census

The winter censuses in the four largest caves of Fenera along the 17-years monitoring period confirmed the cumulated presence of 534 *Rhinolophus ferrumequinum* (501 at Bondaccia, 18 at Ciota Ciara, 11 at Arenarie, 4 at Ciutarun), while other species were much less abundant: 3 large *Myotis* (*M. myotis/blythii* at Bondaccia), 1 small *Myotis* (*M. alcatoe/brandtii/mystacinus* at Bondaccia), 1 *Myotis crypticus* and 1 *Pipistrellus* sp. at Ciota Ciara.



The trend of observed and estimated number of wintering *Rhinolophus ferrumequinum* is reported in Table 1, with a clear and steady increase of census individuals.

The TRIM population analysis model shows a significant overall gradient of 1.22 (Wald test;  $p < 0.001$ ), which corresponds to an average annual increase of 22 % over the period considered (Fig. 2).

Table 1. Number of hibernating *Rhinolophus ferrumequinum* bats in Fenera caves, and trends (TRIM index, TRIM estimates and standard errors) along 17 winters. The dashes refer to winters when no inspections were done.

Winter	Observed individuals	TRIM index	Estimated individuals	SE
2004/2005	5	1	5	5
2005/2006	-	1.2192	6	3
2006/2007	-	1.4987	8	3
2007/2008	-	1.8424	10	4
2008/2009	-	2.2648	12	4
2009/2010	-	2.7841	15	4
2010/2011	-	3.4225	18	5
2011/2012	24	4.3923	24	10
2012/2013	-	5.2987	27	6
2013/2014	36	6.9713	36	12
2014/2015	-	7.6223	41	6
2015/2016	-	9.6077	51	6
2016/2017	63	11.9867	63	16
2017/2018	65	13.0798	69	18
2018/2019	92	17.5044	92	20
2019/2020	101	20.2935	107	22
2020/2021	148	29.4829	155	25




Fig. 2. Variation of wintering *Rhinolophus ferrumequinum* bats in the period 2004/2005 - 2020/2021.

### Summer-autumn census

Data on the activity of bat species, or bat acoustic groups, in the four different habitats surrounding the cave area are reported in Table 2. The most utilized habitat was the agricultural mosaic with light, which showed a 2-3-fold activity level respect to the other habitats. These high activity levels were mainly due to the most ubiquitous species of the genus *Pipistrellus* and of the acoustic group *Eptesicus/Nyctalus/Vespertilio*.

The habitats utilized by the higher number of species were the woods and the cave entrances. The most active species in all four habitat categories was *Pipistrellus pipistrellus*, while the acoustic group *kuhlii/nathusii* was particularly active in the agricultural mosaic with light, as well as *Nyctalus leisleri*.

Several species were rarely observed (Table 2), except for the high activity of *Rhinolophus ferrumequinum* recorded in the area surrounding the cave entrances.

### Incidental observations

A checklist of all species observed in the Fenera cave area is reported in Table 3. The list includes the incidental observations reported below:

- *Tadarida teniotis*: recorded with a bat-detector at the entrance of Ciota Ciara cave on 6 October 2012;
- *Rhinolophus ferrumequinum*: a single individual roosting in a small building at Colma (summer 2018);
- *Pipistrellus pipistrellus*: 1 ind. recorded with a bat-detector at the entrance of Ciota Ciara cave (summer 2018); 1 ind. found dead at Colma (summer 2018); 2 ind. roosting in the roof of a house at Maretti Valduggia (summer 2020);
- *Plecotus austriacus*: 1 ind. found dead at Zuccaro Valduggia on 17 June 2021;
- *Plecotus auritus*: 2 ind. captured by mist net at the entrance of Ciota Ciara cave on 6 October 2012.

### DISCUSSION

Our study reports the presence of 18 species of bats found wintering or flying in the area surrounding cave entrances in a karstic area with several caves in NW Italy. In the winter period, according to visual inspections of caves, hibernating bats were almost exclusively *Rhinolophus ferrumequinum*. This species is scored as Near Threatened in the European Mammal Assessment (Temple & Terry, 2007). Situations of local extinction of the species have been reported in parts of England, Holland, Israel. Negative demographic trends were recorded in Austria, Belgium, Germany, Bulgaria, France, and Switzerland, followed only recently by a positive trend (Van der Meij *et al.*, 2015). In Great Britain, numerical decreases have been documented in the early 60s and 80s, while current populations seem stable. In Italy the species was probably abundant in the last century (Gulino & Dal Piaz, 1939), but surveys

Table 2. Activity of bat species or bat acoustic groups in four environments surrounding the cave entrances in Fenera's area. Activity is reported as mean number of bat passes per hour.

Species/Acoustic groups	Woods	Cave entrances	Dark agricultural mosaic	Agricultural mosaic with light
<i>Barbastella barbastellus</i>	0.03	0	0	0.03
<i>Eptesicus serotinus</i>	0.38	0.12	0.47	5.32
<i>Eptesicus/Nyctalus/Vespertilio</i>	0.45	0.10	0.21	17.97
<i>Hypsugo savii</i>	1.13	2.90	0.44	0.35
<i>Myotis crypticus</i>	0.03	0.38	0.02	0
<i>Myotis daubentonii</i>	0.10	0	0	0
<i>Myotis emarginatus</i>	0	0.18	0	0
<i>Myotis</i> HF	0.86	1.26	0.54	0.85
<i>Myotis</i> LF <i>myotis/blythii</i>	0.04	0.03	0.36	0.65
<i>Nyctalus leisleri</i>	0.05	0.03	3.44	23.82
<i>Nyctalus noctula</i>	0	0.05	0	0
<i>Pipistrellus kuhlii</i>	1.85	0.86	1.38	2.88
<i>Pipistrellus kuhlii/nathusii</i>	6.38	2.90	5.56	50.94
<i>Pipistrellus pipistrellus</i>	60.55	35.43	38.12	51.09
<i>Pipistrellus pygmaeus</i>	0.15	0.49	0	0
<i>Plecotus</i> sp.	0.01	0.02	0.48	6.38
<i>Rhinolophus ferrumequinum</i>	0.01	17.48	0.03	0
<i>Tadarida teniotis</i>	0	0	0.03	0
<b>TOTAL BAT ACTIVITY (passes/hour)</b>	<b>72.0</b>	<b>62.2</b>	<b>51.1</b>	<b>160.3</b>
<b>Sampling effort (hours)</b>	<b>79.5</b>	<b>115.5</b>	<b>66.5</b>	<b>34.0</b>

carried out recently show a notable rarefaction compared to the past (Agnelli *et al.*, 2004). The population is in decline due to the loss of foraging habitats, due to the intensification of agriculture and the use of pesticides, as well as to the reduction of adequate refuge sites. Large colonies are very rare (usually few individuals per colony, rarely more than 100 individuals). Our data highlight the importance of the Fenera caves as one of the most important wintering areas in NW Italy for this species. The large numerical increase that occurred during the last decade, as well as the high number of individuals counted in the last five years, are probably related both to the recent trend observed in NW Italy (Toffoli & Calvini, 2021) and to the conservation actions which have been put in place since the 1998, with the assembly of closed metal gates that prevent access to unauthorized tourists but allow the entry and exit of bats.

Our study highlights the interest of the Fenera as a foraging area utilized by *Rhinolophus ferrumequinum* during the summer and autumn seasons too. In NW Italy, during the 90s, observations of reproductive sites utilized in spring or summer were completely lacking (Sindaco *et al.*, 1992). To avoid possible disturbance of maternity roosts, we did not explore the caves in the

summer period, but the presence of the species with high levels of activity is a good indication of reproduction in the area, considering that females usually have a much reduced home range during the lactating period (Jeon *et al.*, 2018). Alongside, there may have been a substantial contribution to the high activity rate of the area due to the presence of commuting males (Downs *et al.*, 2016) who possibly had daytime rest areas nearby. The presence of woods is also important for *Rhinolophus ferrumequinum*, the preferred foraging habitats of which are broad-leaved woodland interspersed with open spaces (Duvergé & Jones, 1994; Ransome & Hutson, 2000; Flanders & Jones, 2009).

Our study provides the first data of foraging activity in a NW Italy karstic area. The most utilized habitat was the agricultural mosaic with light, which showed a high activity level mainly due to the most ubiquitous species of the genus *Pipistrellus* and of the acoustic group *Eptesicus/Nyctalus/Vespertilio*. Besides, the habitats utilized by the higher number of species were the woods and the cave entrances. However, we underline that a fine comparison of the number of species in the different habitats would have required a much more extensive sampling, with the calculation of accumulation curves (Moreno & Halfpeter,

Table 3. Checklist of bats species observed in the caves of Fenera.

\*: data reported in Pascutto & Balestrieri (2001) for the period 1993-1999, with supplementation from Lana *et al.* (2021) and the authors up to 2021.

Sites	<i>Tadarida teniotis</i>	<i>Rhinolophus ferrumequinum</i>	<i>Rhinolophus hipposideros</i>	<i>Barbastella barbastellus</i>	<i>Eptesicus serotinus</i>	<i>Hypsugo savii</i>	<i>M. alcatraz/brandtii/mystacinus</i>	<i>Myotis daubentonii</i>	<i>Myotis emarginatus</i>	<i>Myotis myotis/blythii</i>	<i>Myotis crypticus</i>	<i>Nyctalus leisleri</i>	<i>Nyctalus noctula</i>	<i>Pipistrellus kuhlii</i>	<i>Pipistrellus pipistrellus</i>	<i>Pipistrellus pygmaeus</i>	<i>Plecotus auritus</i>	<i>Plecotus austriacus</i>
<b>Years 1993-1999 (2021)*</b>																		
Bondaccia Cave		x	*							*							*	
Ciutarun Cave		x																
Ciota Ciara Cave		x	*															
Arenarie Cave		x																
Small caves		x	x							x								
Small building																		*
<b>Years 2004-2020 (winter)</b>																		
Bondaccia Cave		x					x			x								
Ciutarun Cave		x																
Ciota Ciara Cave		x									x							
Arenarie Cave		x																
<b>Year 2020 (acoustic)</b>																		
Cave entrances		x			x	x			x	x	x	x	x	x	x	x	x	
Woods		x		x	x	x		x		x	x	x		x	x	x		
Agricultural mosaic with light				x	x	x				x		x		x	x			
Dark agricultural mosaic	x	x			x	x				x	x	x		x	x			

2001; Skalak *et al.*, 2012). These habitats were utilized by some rare species, i.e. *Barbastella barbastellus* (Toffoli & Cucco, 2020) and all species of the genus *Myotis*. Our results are in line with the observations of Zúkal & Řehák (2006) in the Moravian karst, with the exception of the high bat activity noted over Moravian ponds, a habitat that was not present in the Fenera area. The importance of karstic areas outside of the hibernating period has been underlined in several tropical zones (Struebig *et al.*, 2009; Racey & Furey, 2014), but has been little studied in temperate areas. Our results underline the importance of protecting the cave underground system as well as the habitats surrounding cave entrance for an effective conservation of bat biodiversity.

## ACKNOWLEDGEMENTS

We thank Paola Culasso, Enrico Lana, Tiziano Pascutto, Tito Princisvalle, Luciano Rossi, the Gruppo Speleologico Biellese and Gruppo Speleologico C.A.I. Varallo for their help and support in the fieldwork.

## REFERENCES

- Agnelli P., Martinoli A., Patriarca E., Russo D., Scaravelli D., Genovesi P. 2004. Linee guida per il monitoraggio dei Chiroteri: indicazioni metodologiche per lo studio e la conservazione dei pipistrelli in Italia. *Quaderni di Conservazione della Natura, Ministero dell'Ambiente e della Tutela del Territorio*, Roma.



- Altringham J.D. 2011. Bats: from evolution to conservation. *Oxford University Press, Oxford*.
- Azam C., Kerbiriou C., Vernet A., Julien J.-F., Bas Y., Plichard L., Maratrat J., Le Viol I. 2015. Is part-night lighting an effective measure to limit the impacts of artificial lighting on bats? *Global Change Biology* 21(12): 4333-4341. DOI: 10.1111/gcb.13036
- Barataud M. 2016. Acoustic ecology of European bats: species identification, study of their habitats and foraging behaviour. *Biotope Editions, Muséum Histoire Naturelle, Paris*.
- Battersby J. 2010. Guidelines for surveillance and monitoring of European bats. Eurobats publication series (no.5). *UNEP/EUROBATS secretariat, Bonn, Germany*.
- Ben-Hamo M., Muñoz-García A., Williams J.B., Korine C., Pinshow B. 2013. Waking to drink: rates of evaporative water loss determine arousal frequency in hibernating bats. *Journal of Experimental Biology* 216: 573-577. DOI: 10.1242/jeb.078790
- Berto C., Bertè D., Luzi E., López-García J.M., Pereswiet-Soltan A., Arzarello M. 2016. Small and large mammals from the Ciota Ciara cave (Borgosesia, Vercelli, Italy): an Isotope Stage 5 assemblage. *Comptes Rendus Palevol* 15: 669-680. DOI: 10.1016/j.crpv.2015.05.014
- Bogaart P., Loo M., Pannekoek J. 2016. Rtrim: trends and indices for monitoring data. *R package, version 2.0.6*. <https://CRAN.R-project.org/package=rtrim>
- Charbonnier Y.M., Barbaro L., Barnagaud J.-Y., Ampoorter E., Nezan J., Verheyen K., Jactel H. 2016. Bat and bird diversity along independent gradients of latitude and tree composition in European forests. *Oecologia* 182: 529-537. DOI: 10.1007/s00442-016-3671-9
- Coleman J.L., Barclay R.M.R. 2013. Prey availability and foraging activity of grassland bats in relation to urbanization. *Journal of Mammalogy* 94: 1111-1122. DOI: 10.1644/12-MAMM-A-217.1
- Collins J. 2016. Bat surveys for professional ecologists: good practice guidelines (3rd edn). *The Bat Conservation Trust, London*.
- Duvergé P.L., Jones G. 1994. Greater horseshoe bats activity, foraging behaviour and habitat use. *British Wildlife* 6: 69-77.
- Downs N.C., Cresswell W.J., Reason P., Sutton G., Wells D., Wray S. 2016. Sex-specific habitat preferences of foraging and commuting lesser horseshoe bats *Rhinolophus hipposideros* (Borkhausen, 1797) in lowland England. *Acta Chiropterologica* 18: 451-465. DOI: 10.3161/15081109ACC2016.18.2.012
- Erkert H.G. 1982. Ecological aspects of bat activity rhythms (pp. 201). In: Kunz T.H. (ed.), *Ecology of bats*. Plenum Press, New York.
- Flanders J., Jones G. 2009. Roost use, ranging behavior, and diet of Greater Horseshoe bats (*Rhinolophus ferrumequinum*) using a transitional roost. *Journal of Mammalogy* 90: 888-896. DOI: 10.1644/08-MAMM-A-352.1
- Furey N. M., Racey P. A. 2015. Conservation Ecology of Cave Bats (pp. 463-500). In: Voigt C. C., Kingston T. (eds), *Bats in the Anthropocene: Conservation of Bats in a Changing World*. Springer.
- Glover A. M., Altringham J.D. 2008. Cave selection and use by swarming bat species. *Biological Conservation* 141: 1493-1504. DOI: 10.1016/j.biocon.2008.03.012
- Gregory R.D., Van Strien A.J., Vorisek P., Gmelig Meyling A.W., Noble D.G., Foppen R.P.B., Gibbons D.W. 2005. Developing indicators for European birds. *Philosophical Transactions of the Royal Society of London, B* 360: 269-288. DOI: 10.1098/rstb.2004.1602
- Gulino G., Dal Piaz 1939. I Chiroterri italiani. *Bollettino Musei di Zoologia ed Anatomia Comparata Regia Università di Torino* 47: 63-103.
- Heim O., Treitler J.T., Tschapka M., Knörnschild M., Jung K. 2015. The importance of landscape elements for bat activity and species richness in agricultural areas. *Plos One* 10(7): 1-13. DOI: 10.1371/journal.pone.0134443
- Hill A.P., Prince P., Snaddon J.L., Doncaster C.P., Rogers A. 2019. AudioMoth: a low-cost acoustic device for monitoring biodiversity and the environment. *HardwareX* 6: e00073. DOI: 10.1016/j.ohx.2019.e00073
- Holloway G.L., Barclay R.M.R. 2000. Importance of prairie riparian zones to bats in southeastern Alberta. *Ecoscience* 7: 115-122. DOI: 10.1080/11956860.2000.11682579
- Klüg-Baerwald B.J., Brigham R.M. 2017. Hung out to dry? Intraspecific variation in water loss in a hibernating bat. *Oecologia* 183: 977-985. DOI: 10.1007/s00442-017-3837-0
- Kofoky A., Andriafidison D., Ratrimomanarivo F., Razafimanahaka H.J., Rakotondravony D., Racey P.A., Jenkins R.K.B. 2006. Habitat use, roost selection and conservation of bats in Tsingy de Bemaraha National Park, Madagascar (pp. 213-227). In: Hawksworth D.L. & Bull A.T. (eds), *Vertebrate conservation and biodiversity*. Springer, Dordrecht, 494 pp.
- Kunz T.H., Fenton M.B. 2003. *Bat Ecology*. University of Chicago Press, Chicago, 779 pp.
- Jeon Y.S., Kim S.C., Han S.H., Chung C.U. 2018. Characteristics of the home range and habitat use of the Greater Horseshoe bat (*Rhinolophus ferrumequinum*) in an urban landscape. *Journal of Environmental Science International* 27: 665-675. DOI: 10.5322/jesi.2018.27.8.665
- Lana E., Sella R. 1986. Le grotte del Monte Fenara e la loro fauna. *Rivista Piemontese di Storia Naturale* 37: 225-297.
- Lana E., Giachino P.M., Casale A. 2021. Fauna Hypogaea Pedemontana. Grotte e ambienti sotterranei del Piemonte e della Valle d'Aosta. *WBA Monographs 6, WBA Project Ed., Verona*, 1044 pp.
- Loeb S.C., Rodhouse T.J., Ellison L.E., Lausen C.L., Reichard J.D., Irvine K.M., Ingersoll T.E., Coleman J.T.H., Thogmartin W.E., Sauer J.R., Francis C.M., Bayless M.L., Stanley T.R., Johnson D.H. 2015. A plan for the North American Bat Monitoring Program (NABat). Gen. Tech. Rep. SRS-208. Asheville, NC. *U.S. Department of Agriculture Forest Service, Southern Research Station*, 100 pp.
- Lumsden L.F., Bennett A.F. 2005. Scattered trees in rural landscapes: foraging habitat for insectivorous bats in south-eastern Australia. *Biological Conservation* 122: 205-222. DOI: 10.1016/j.biocon.2004.07.006
- McClure M.L., Crowley D., Haase C.G., McGuire L.P., Fuller N.W., Hayman D.T.S., Lausen C.L., Plowright R.K., Dickson B.G., Olson S.H. 2020. Linking surface and subterranean climate: implications for the study of hibernating bats and other cave dwellers. *Ecosphere* 11 (10): e03274. DOI: doi.org/10.1002/ecs2.3274
- Mickleburgh S. P., Hutson A. M., Racey P. A. 2002. A review of the global conservation status of bats. *Oryx* 36: 18-34. DOI: S0030605301000011
- Middleton N., Froud A., French K. 2014. Social calls of the bats of Britain and Ireland. *Pelagic Publishing UK, Exeter*.

- Moreno C. E., Halffter G. 2001. Assessing the completeness of bat biodiversity inventories using species accumulation curves. *Journal of Applied Ecology* 37: 149-158. DOI: 10.1046/j.1365-2664.2000.00483.x
- Pannekoek J., Van Strien A.J. 2001. TRIM3 manual. Trends and indices for monitoring data. *Statistics Netherlands Voorburg, the Netherlands*.
- Pascutto T., Balestrieri A. 2001. Note sui Chiroterri troglodili osservati in alcune cavità del Piemonte. *Orso Speleo Biellese* XXVII 21: 22-28.
- Racey P.A., Furey N.M. 2014. Are bats karst-dependent? *Vespertilio* 17: 157-159.
- Rainho A. 2007. Summer foraging habitats of bats in a Mediterranean region of the Iberian Peninsula. *Acta Chiropterologica* 9: 171-181. DOI: 10.3161/150811007781694408
- Ransome R.D., Hutson A.M. 2000. Action plan for the conservation of the Greater Horseshoe Bat in Europe (*Rhinolophus ferrumequinum*). Convention on the Conservation of European Wildlife and Natural Habitats, Nature and Environment 109. *Council of Europe Publishing*.
- Russo D., Jones G. 1999. The social calls of Kuhl's pipistrelles *Pipistrellus kuhlii* (Kuhl, 1819): structure and variation (Chiroptera: Vespertilionidae). *Journal of Zoology* 249: 476-481. DOI: 10.1111/j.1469-7998.1999.tb01219.x
- Russo D., Jones G. 2002. Identification of twenty-two bat species (Mammalia: Chiroptera) from Italy by analysis of time-expanded recordings of echolocation calls. *Journal of Zoology* 258: 91-103. DOI: 10.1017/S0952836902001231
- Sindaco R., Baratti N., Boano G. 1992. Bats of Piedmont and the Aosta Valley (NW Italy). *Hystrix Italian Journal of Mammalogy* 4: 1-40. DOI: 10.4404/hystrix-4.1-3977
- Skalak S.L., Sherwin R.E., Brigham R.M. 2012. Sampling period, size and duration influence measures of bat species richness from acoustic surveys. *Journal of Applied Ecology* 3: 490-502. DOI: 10.1111/j.2041-210X.2011.00177.x
- Speakman J.R., Thomas D.W. 2003. Physiological ecology and energetics of bats (pp. 430-490). In: Kunz T.H. & Fenton M.B. (eds), *Bat biology*. University of Chicago Press.
- Struebig M.G., Kingston T., Zubaid A., Le Comber S.C., Mohd-Adnan A., Turner A., Kelly K., Božek M., Rossiter S.J. 2009. Conservation importance of limestone karst outcrops for Palaeotropical bats in a fragmented landscape. *Biological Conservation* 142: 2089-2096. DOI: 10.1016/j.biocon.2009.04.005
- Temple H.J., Terry A. 2007. The status and distribution of European mammals. *Office for Official Publications of the European Communities*. pp. VIII + 48.
- Thomas D.W., Geiser F. 1997. Periodic arousals in hibernating mammals: is evaporative water loss involved? *Functional Ecology* 11: 585-591. DOI: 10.1046/j.1365-2435.1997.00129.x
- Thomas D.W., West S.D. 1989. Sampling methods for bats. General Technical Report PNW-GTR-243. Portland, OR: *U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station*. 20 pp.
- Toffoli R. 2016. The importance of linear landscape elements for bats in a farmland area: the influence of height on activity. *Journal of Landscape Ecology* 9(1): 49-62. DOI: 10.1515/jlecol-2016-0004
- Toffoli R., Calvini M. 2019. Long term trend of an endangered bat species in north-western Italy. *Folia Zoologica* 68(2): 95-99. DOI: 10.25225/fozo.006.2019
- Toffoli R., Calvini M. 2021. Long term trends of hibernating bats in North-Western Italy. *Biologia* 76(2): 633-643. DOI: 10.2478/s11756-020-00584-x
- Toffoli R., Cucco M. 2020. Habitat suitability, connection analysis, and effectiveness of protected areas for conservation of the Barbastelle bat *Barbastella barbastellus* in NW Italy. *Acta Chiropterologica* 22: 281-291. DOI: 10.3161/15081109ACC2020.22.2.004
- Toffoli R., Rugghetti M. 2017. Bat activity in rice paddies: organic and conventional farms compared to unmanaged habitat. *Agriculture, Ecosystems & Environment* 249: 123-129. DOI: 10.1016/j.agee.2017.08.022
- Uhrin M., Benda P., Obuch J., Urban P. 2010. Changes in abundance of hibernating bats in central Slovakia (1992-2009). *Biologia* 65: 349-361. DOI: 10.2478/s11756-010-0020-z
- Van der Meij T., Van Strien A.J., Haysom K.A., Dekker J. et al. 2015. Return of the bats? A prototype indicator of trends in European bat populations in underground hibernacula. *Mammalian Biology* 80: 170-177. DOI: 10.1016/j.mambio.2014.09.004
- Verboom B., Spoelstra K. 1999. Effects of food abundance and wind on the use of tree lines by an insectivorous bat, *Pipistrellus pipistrellus*. *Canadian Journal of Zoology* 77: 1393-1401. DOI: 10.1139/z99-116
- Waters D.A., Jones G. 1995. Echolocation call structure and intensity in five species of insectivorous bats. *Journal of Experimental Biology* 198: 475-489. DOI: 10.1242/jeb.198.2.475
- Wieser D., Mixanig H., Krainer K., Bruckner A., Reiter G. 2020. Importance of inland cliffs and quarries for bats. *Acta Chiropterologica* 22 (2): 391-402. DOI: 10.3161/15081109ACC2020.22.2.013
- Zukal J., Řehák Z. 2006. Flight activity and habitat preference of bats in a karstic area, as revealed by bat detectors. *Folia Zoologica* 55(3): 273-281.
- Zukal J., Berková H., Bandouchová H., Kováčová V., Pikula J. 2017. Bats and caves: activity and ecology of bats wintering in caves. In: Karabulut S., Cinku M.C. (eds). *Cave investigation*. InTech, Rijeka, pp. 51-75.